

REVIEW

Sex, parturition and motherhood without oxytocin?

J A Russell and G Leng

Department of Physiology, The University of Edinburgh Medical School, Teviot Place, Edinburgh EH8 9AG, UK
(Requests for offprints should be addressed to J A Russell)

Our understanding of the functions of oxytocin in mammals has recently been challenged by findings in transgenic mice in which the oxytocin gene has been knocked out. Mammals generally have only two posterior pituitary nonapeptide genes, for oxytocin and vasopressin, while some marsupials do not express oxytocin but do express a closely related peptide, mesotocin, which acts at the oxytocin receptor (Acher *et al.* 1995). Birds have arginine vasotocin and mesotocin (Acher *et al.* 1995), while among other vertebrates a subclass of cartilaginous fishes evidently produces oxytocin (Michel *et al.* 1993).

Mice homozygous for deletion of exon 1 of the oxytocin gene, containing the oxytocin nonapeptide sequence, were produced by homologous recombination in mouse embryonic stem cells. Offspring from matings of the heterozygotes were in the correct Mendelian frequency, indicating no lethal developmental defects in the homozygotes, which later showed normal sexual maturation, with both males and females showing sexual behaviour and normal fertility (Nishimori *et al.* 1996). Young and colleagues (1996) used gene targeting to generate a mouse with most of the first intron and the last two of the three exons of the oxytocin gene eliminated. The amount of oxytocin gene transcripts in the supraoptic and paraventricular nuclei of homozygotes was 1% of the wild-type level, with less than 0.4% of the wild-type content of oxytocin in the pituitary gland, and no oxytocin detectable in blood plasma by RIA. In both types of oxytocin gene disablement, the only evident defect was a complete failure of postpartum homozygotes to transfer milk to the suckling young, which consequently did not survive unless the mothers were treated with exogenous oxytocin. Thus parturition and maternal behaviour proceeded without oxytocin, although these processes were not studied in detail.

These findings are surprising in view of the vast literature on a wide range of species suggesting important roles for oxytocin in regulating gonadal function, in expression of sexual behaviour, in parturition and initiation of maternal behaviour as well as in lactation. It is appropriate to consider whether the results from the oxytocin knockout mice arise from peculiarities in the roles of oxytocin in the mouse in comparison with other species, or

indicate redundancy in the mechanisms in which oxytocin normally has an important role; apart that is, from the milk-ejection reflex in which there is no evident redundancy, at least in the mouse.

Evidence for a role for oxytocin in parturition

Initiation of delivery

One of the first actions of posterior pituitary extracts to be described was the ability to stimulate contractions of the uterus (Dale 1906). Because of this uterotonic action, posterior pituitary extracts were soon used in clinical practice (see Robinson & Amico (1985) for a review). In 1948 a midwife reported her observations of a woman in labour with her second child while still in lactation following the birth of her first child; during each uterine contraction, beads of milk stood out from the nipples, as from a lactating breast during milk let-down (Gunther 1948). The observant midwife speculated that 'labour proceeds from a series of discharges of pituitrin-like substance which acts on the prepared uterus and which is rapidly rendered inactive'.

Synthetic oxytocin has now been used in clinical and veterinary practice to induce or assist parturition for many years (see Owen & Hanth 1992, Vivrette 1994), and a role for oxytocin is fitted into all models that seek to describe the mechanisms involved in the onset and maintenance of parturition in a range of placental mammals. The classic account of parturition has it that, in late pregnancy, the high circulating concentrations of progesterone induce uterine quiescence while the uterus acquires contractile ability. Shortly before term, plasma progesterone concentrations fall steeply, oxytocin receptor expression in the uterus increases markedly, and uterine contractile activity increases. At term, the contractions rise to a crescendo, resulting in delivery as a result of two interacting positive feedback loops. The first is a local uterine loop: within the uterus itself, prostaglandins and other uterotonic factors are produced and released in response to uterine contractions. The second loop involves the hypothalamus: in response to uterine contractions and vaginal and cervical distension, magnocellular oxytocin neurones in the hypothalamus




Figure 1 Positive feedback stimulation of maternal oxytocin secretion during parturition in the rat. Schema based on a variety of data. The initial events at term are in the uterus: withdrawal of progesterone permits myometrial oxytocin receptor expression, gap junctions are formed between myometrial cells, oxytocin may be released from the uterine epithelium and act locally, and prostaglandin production increases. As a result, the myometrium is more excitable and the contractions of early labour begin. The contractions increase afferent nerve activity, especially when the cervix is distended by the presenting part of the fetus. Via afferents relaying in the nucleus tractus solitarii, the cell bodies of the magnocellular neurones in the supraoptic and paraventricular nuclei of the hypothalamus receive a barrage of impulses, originating in the uterus, cervix and, when a fetus is expelled, vagina. The increased continuous background electrical activity (action potentials) of the oxytocin neurones drives secretion of oxytocin from the expanded store in the nerve terminals in the posterior pituitary gland, but intermittently each of these neurones fires in a high-frequency burst (*), synchronously with other oxytocin neurones, to cause the secretion of a pulse (†), or bolus, of oxytocin into the circulation; this is immediately followed by the birth of a pup (Summerlee 1981). I.U.P., intrauterine pressure; OXT, oxytocin; NTS, nucleus tractus solitarii; PGF_{2α}, prostaglandin F_{2α}.

increase their activity resulting in the release of oxytocin from their axon terminals in the posterior pituitary; the released oxytocin acts upon the uterus both to stimulate the further production of prostaglandins and to contribute further to the contractions of the uterus (Fig. 1).

During parturition, elevated oxytocin secretion has been measured in all placental mammals thus far studied, including the rabbit (Fuchs & Dawood 1980), sheep (Glatz *et al.* 1981), cow (Landgraf *et al.* 1983), rat (Higuchi *et al.* 1986), goat (Currie *et al.* 1988), pony (Haluska & Currie 1988), human (Fuchs *et al.* 1991), rhesus monkey (Hirst *et al.* 1993) and pig (Gilbert *et al.* 1994); representing Australian marsupials, the wallaby shows increased secretion of mesotocin, an oxytocin-like peptide, during parturition (Parry *et al.* 1996); for the birds, in the chicken the secretion of arginine vasotocin (which is a more potent

oxytocic on chicken uterus than the other avian neurohypophysial nonapeptide, mesotocin) increases abruptly at oviposition (Koike *et al.* 1988).

Again, in all mammals thus far studied, there is a marked increase in uterine responsiveness to oxytocin at term, and oxytocin is the most powerful uterotonic agent identified to date. In the rat, sensitivity to the contractile actions of oxytocin of at least the circular layer of the myometrium greatly increases at the end of pregnancy (Fuchs *et al.* 1983, Crankshaw 1987), and the sensitivity of mouse and guinea-pig myometrium to oxytocin is increased at the end of gestation (Bell 1941, Suzuki & Kuriyama 1975*b*), with a similar change and increase in maximal response to oxytocin in the rabbit (Riemer *et al.* 1986). The sensitivity of the human uterus to oxytocin increases in late pregnancy (Takahashi *et al.* 1980). Sensitivity to the uterotonic

action of oxytocin increases in late pregnancy in the quokka, a macropodial marsupial (Heller 1973), and in the wallaby (see Renfree 1994), while the endogenous mesotocin can be expected to have similar effects to oxytocin (Bathgate *et al.* 1995). In the chicken, the sensitivity of the uterus to contractile stimulation by arginine vasotocin, acting via vasopressin-like receptors, peaks at oviposition (Koike *et al.* 1988, Saito & Koike 1992).

The increased responsiveness of the myometrium to oxytocin in late pregnancy is related to increased oxytocin receptor density, measured as specific radioligand binding. Oxytocin receptor density in the myometrium sharply increases at the end of pregnancy in the rat, and this is localised to myometrial cell membranes, including the longitudinal muscle layer (Soloff *et al.* 1979, Kaneko *et al.* 1995, Kawarabayashi *et al.* 1996). Myometrial oxytocin receptor density increases in the guinea-pig towards the end of pregnancy (Alexandrova & Soloff 1980), and similarly oxytocin receptor density increases greatly at term in the rabbit and sheep myometrium and endometrium (Maggi *et al.* 1988, 1991, Wathes *et al.* 1996) and in the human myometrium and decidua (Fuchs *et al.* 1984). Endometrial oxytocin receptor concentration increases in pregnancy in the cow (Fuchs *et al.* 1996). The receptor for arginine vasotocin in the chicken uterus is up-regulated a few hours before oviposition (Takahashi *et al.* 1994).

The increased oxytocin receptor density in the uterus at the end of pregnancy is likely to be a result of stimulated gene expression. Thus there is a large increase in uterine oxytocin receptor mRNA content at the end of pregnancy in the rat (Liu *et al.* 1996), and in the sheep there is a clear correlation between myometrial and endometrial oxytocin receptor mRNA and protein expression, and with myometrial contractile activity (Wu *et al.* 1996). Similarly, oxytocin receptor mRNA and protein content in the human myometrium are greatly increased at the end of pregnancy (Kimura *et al.* 1996). In the cow, oxytocin receptor mRNA expression in the myometrium and endometrium is increased in the third trimester, although there is no evident further increase at term (Ivell *et al.* 1995). Considering the bulk of the uterus, this enhanced expression must be the most massive increase in peptide receptor production to be seen in the mature mammal.

The temporal pattern of oxytocin receptor expression in the uterus and cervix clearly indicates that endometrial and myometrial oxytocin receptor expression may be a key factor in the timing of parturition in ruminants (Wathes *et al.* 1996, Wu *et al.* 1996). In sheep there is a progressive increase in uterine contractile activity toward the end of pregnancy, and, in late pregnancy, infusion of oxytocin antagonists decreases the frequency of uterine contractures (Owiny *et al.* 1992), involving a blockade of stimulation of uterine prostaglandin production (Jenkin *et al.* 1994). Similarly in the rhesus monkey and baboon, a nocturnal increase in uterine contractions late in gestation appears to

be caused by maternal oxytocin (Hirst *et al.* 1993), since a selective oxytocin antagonist attenuates the contractions (Honnebier *et al.* 1989, Wilson *et al.* 1990). In the rhesus monkey, an increase in nocturnal fetal Δ^4 -androstenedione production, entrained by maternal mechanisms, is proposed to increase placental oestrogen secretion, and thence to increase oxytocin secretion, and thereby myometrial contractile activity (Mecenas *et al.* 1996, Nathanielsz *et al.* 1995, Giussani *et al.* 1996), with the further increase in Δ^4 -androstenedione at term triggering parturition (Mecenas *et al.* 1996).

For the rat, as for many species, there is little or no evidence of a rise in oxytocin release before the first delivery, although, since parturition in the rat is preceded by an abrupt and dramatic increase in uterine oxytocin receptor expression, no increase in oxytocin release may be necessary for oxytocin to initiate delivery. Nevertheless, it is generally recognised that, at least for many species, parturition is not necessarily initiated by an increase in oxytocin secretion. Thus, for instance in the guinea-pig, oxytocin antagonist treatment does not delay the start of parturition but does decrease uterine motility at term and prolongs delivery (Schellenberg 1995). In the rat, even very high doses of oxytocin antagonists do not consistently delay the onset of birth in the rat (Chan & Chen 1992), and the delays are relatively short (Antonijevic *et al.* 1995a). Indeed, in the rat, administration of low doses of oxytocin produces a much more effective delay to subsequent birth than administration of high doses of antagonist (Antonijevic *et al.* 1995b), possibly reflecting desensitisation of uterine oxytocin receptors, with decreased receptor density on myometrial cells and reduced oxytocin receptor gene expression (Phaneuf *et al.* 1997), but perhaps also reflecting fatigue of other uterotonic control mechanisms within the uterus. Nonetheless, the onset of parturition in the rat can be advanced by several hours by intravenous infusion of oxytocin, especially if given in pulses (Randolph & Fuchs 1989), and labour is initiated, with variable progression to parturition, at the end of gestation by intermittent electrical stimulation of the neurohypophysis in the rat or rabbit (Lincoln 1971, Boer *et al.* 1975).

So what other mechanisms are known to be involved in the initiation of parturition? In the pregnant rat, about 24 h before term, circulating progesterone secretion falls dramatically, with the collapse of the corpus luteum (Sanyal 1978). This fall leads to increased intrauterine prostaglandin production (Chan 1983), perhaps partly through the action of local oxytocin on decidual oxytocin receptors (Chan *et al.* 1993, Larcher *et al.* 1995), both of which are up-regulated by oestrogen (Lefebvre *et al.* 1994), and probably involving cytokines (Rozen *et al.* 1995). Endotoxin given *in vivo* sensitises the pregnant uterus to oxytocin tested *in vitro* (Suzuki *et al.* 1995). The oxytocin gene in human, mouse and rat has an oestrogen response element (also responding to thyroid hormone and retinoic

acid; Richard & Zingg 1990), while the decidua contain oestrogen receptors (Chibbar *et al.* 1995), but other factors are doubtless involved (Burbach *et al.* 1995), especially since the equivalent sequence in the sheep and bovine oxytocin genes is mutated such that the oestrogen receptor cannot bind (Ivell *et al.* 1990, Adan *et al.* 1991). The human oxytocin receptor gene has half-palindromic oestrogen receptor elements and interleukin-6-binding sequences like the rat and mouse (Inoue *et al.* 1994, Kubota *et al.* 1996); however, these half-sites are unlikely to interact with oestrogen receptor *in vivo*, although the oestrogen antagonist tamoxifen inhibits uterine oxytocin receptor expression at the end of pregnancy in the rat and delays parturition (Fang *et al.* 1996).

The fall in progesterone will also remove progesterone's influence on the inhibitory paths within the myometrial cells (Elwardy-Merezak *et al.* 1994, Cohen-Tannoudji *et al.* 1995, Tezuka *et al.* 1995, Orsino *et al.* 1996), and reduced NO production may also help to activate the myometrium (Izumi & Garfield 1995). In the human, changes in a paracrine network within the fetal membranes and especially the decidua may initiate parturition (Mitchell & Chibbar 1995). Within these tissues a shift to progesterone inactivation and to synthesis of more potent oestrogens will both increase local synthesis of prostaglandins, stimulating contractions, and sensitise the myometrial cells. This sensitisation is achieved by induction of gap junctions between the myometrial cells by oestrogen or relaxin (Tabb *et al.* 1992, Burghardt *et al.* 1996), or, in the guinea-pig, blockade of progesterone action thereby increasing gap junction expression (Chwalisz *et al.* 1991). The marked increase in the expression of the oxytocin receptor gene in the myometrium to greatly increase receptor B_{max} will also increase sensitivity to oxytocin (Fuchs *et al.* 1984). In addition, both oxytocin and the oxytocin receptor gene are expressed in the human decidua (Mitchell & Chibbar 1995), and the release of oxytocin from the luminal surface may act in an autocrine or paracrine fashion to stimulate production of prostaglandins (Wilson *et al.* 1988), which may then act on the underlying myometrium. A further consequence of the decline in local progesterone production is decreased expression of $G_{\alpha s}$ protein in the myometrial cells (Europe-Finner *et al.* 1994), thus weakening inhibitory pathways acted on by β -adrenergic agonists, for example. It is also likely that cytokines, produced by cells in the decidua, act on the decidua or the myometrial cells to stimulate prostaglandin production (Mitchell *et al.* 1990, 1991, Todd *et al.* 1996). Other mechanisms involving, *inter alia*, endothelin, NO and relaxin (Goldsmith *et al.* 1995, Dong & Yallampalli 1996) may also regulate myometrial contractility. In the rat, a decline in luteal relaxin production will contribute to increased excitability, removing relaxin's inhibitory actions, mediated by protein kinase A, on myometrial cells (Meera *et al.* 1995); this may involve interference with $G_{\alpha q/11}$ which mediates oxytocin actions

(Sanborn *et al.* 1995). Thus, while oxytocin may play some role in the initiation of labour, there are certainly other uterine mechanisms that can substitute for it in its absence, and which indeed may play a more dominant role. However, prostaglandins are less effective than oxytocin at inducing parturition in rats at term (Fuchs 1972).

Progress of delivery

Thus it is broadly recognised that oxytocin is not generally essential for the initiation of labour, but there is much stronger evidence that it plays an important role in the progress of labour in many species. In the rabbit, oxytocin secretion is maximal with delivery of the first fetus (Fuchs & Dawood 1980). In generally monotocous species, peak oxytocin secretion is seen at the birth of the singleton (sheep (Glatz *et al.* 1981), cow (Landgraf *et al.* 1983), rhesus monkey (Hirst *et al.* 1993)). In ruminants, enhanced pulsatile oxytocin secretion is readily demonstrable during parturition (Fuchs 1985, Fuchs *et al.* 1995). In the pig, a pulse of oxytocin is secreted at the birth of each piglet (Gilbert *et al.* 1994), while environmental disturbance during parturition both decreases oxytocin secretion and slows down the birth of the piglets (Lawrence *et al.* 1992). Again though, it is clear that parturition is not necessarily disrupted by the absence of oxytocin: in the rat, oxytocin antagonists have a consistent and marked effect on the progress of parturition when administered after the birth of the first pup (Fig. 2), but, when administered after the birth of the fourth pup, they have no effect in about half of the animals studied (Antonijevic *et al.* 1995a).

In the human, cystine aminopeptidase from the decidua breaks down oxytocin, and circulates as oxytocinase (Burd *et al.* 1987), supposedly protecting the myometrium from all but large pulses of oxytocin from the maternal posterior pituitary, and ensuring intermittent stimulation of the myometrium. As in the rat, pulsatile infusion is the most efficient way to stimulate the myometrium with exogenous oxytocin (Cummiskey & Dawood 1990). The power of oxytocin is demonstrated by the association of uterine rupture in multiparae with its inappropriate use to promote labour (Golan *et al.* 1980). Because of the pulsatile secretion of oxytocin (Fuchs *et al.* 1991), coupled with the action of the circulating oxytocinase, it has been difficult to consistently show increased oxytocin secretion in women in labour; however, this has been shown in some studies, including early in pre-term labour (Vavra *et al.* 1993), with a pattern of increasing frequency of pulses (Fuchs *et al.* 1991), with the most consistent increases measured during the expulsive phase, when the sustained distension of the uterine cervix and vagina leads to strong reflex stimulation of oxytocin secretion (Vasicka *et al.* 1978, Leake *et al.* 1981, Thornton *et al.* 1992). Nonetheless, the importance of oxytocin even in early labour is suggested by the effectiveness in clinical trials of an oxytocin antagonist (Atosiban) in threatened pre-term labour (Goodwin *et al.*




Figure 2 Progress of parturition in the rat and the slowing actions of oxytocin antagonist or morphine. All rats were fitted with a jugular venous cannula 1 or 2 days before term for injections after birth of the second pup. The control group was given i.v. vehicle, and the differently shaded bars indicate the duration of each subsequent interbirth interval. The i.v. injection of a peptide oxytocin antagonist (Ferring 382; 30 μ g/kg) prolonged the intervals between the births of the subsequent four pups, thus doubling the time to deliver five pups after injection. The i.v. injection of morphine sulphate (1 mg/kg) markedly prolonged the interval to the birth of pup 3, and lengthened the next interval, thus also extending the time to deliver five pups after injection. Pulsatile i.v. injection of oxytocin (OXT 1 mU (2 ng; 2 pmoles) every 10 min) after a single injection of morphine after the birth of pup 2 fully restored the progress of normal parturition; continuous infusion (not shown) of the same total dose of oxytocin was ineffective (data from Luckman *et al.* 1993, Antonijevic *et al.* 1995a). The prolonged inhibitory action of the oxytocin antagonist clearly indicates an important role for oxytocin in promoting parturition. The inhibitory effects of morphine are a consequence of inhibition of the secretion of oxytocin from the posterior pituitary gland: morphine inhibits the firing of magnocellular oxytocin neurones via μ -opioid receptors on their cell bodies or inputs, and like other opiates it decreases oxytocin secretion in parturition (Russell *et al.* 1989, Douglas *et al.* 1993). The restoration of the normal progress of parturition by i.v. pulses of oxytocin supports both a singular action of morphine to inhibit oxytocin secretion and the conclusion that it is secretion of oxytocin from the posterior pituitary in a pulsatile pattern that is important in driving parturition.

1996). In the baboon in labour, intravenous oxytocin antagonist greatly decreases uterine contractile force (Wilson *et al.* 1990).

Since the pituitary oxytocin content is dramatically depleted during parturition in rat, mouse and guinea-pig, the major source of oxytocin release is generally likely to be the maternal pituitary, but it may also be released from the fetal posterior pituitary gland in some species, including the sheep (Dawood *et al.* 1983), pig (Sander-Richter *et al.* 1988) and human (Chard 1989), although not the rat (Higuchi *et al.* 1985). In some species, oxytocin may also be released from the subjacent endometrium or decidua where the gene is strongly expressed at the end of pregnancy, although the amount of peptide produced is very small relative to the posterior pituitary content (human, rat, but not bovine, mouse or sheep: Chibbar *et al.* 1993, Lefebvre *et al.* 1993, Ivell *et al.* 1995, Murphy & Ho 1995, Wathes *et al.* 1996).

So is the massive release of oxytocin from the maternal pituitary at term an anachronism? A remnant perhaps of an evolutionary history in which the ancestral hormone from

which oxytocin evolved played an important role in egg laying, a role played by arginine vasotocin in modern birds (Takahashi *et al.* 1994). Australian marsupials produce mesotocin rather than oxytocin, and mesotocin in Australian marsupials may not be an ancestor of mammalian oxytocin, but a descendant, since South American marsupials have oxytocin (Acher *et al.* 1995).

Marsupials such as the tammar wallaby secrete mesotocin in circumstances, including parturition and lactation (see Bathgate *et al.* 1995, Parry *et al.* 1996), when eutherian mammals secrete oxytocin, and with similar consequences (Lincoln & Renfree 1981a). Selection pressure has thus been for a neutral mutation in the peptide ligand for the oxytocin receptor. Mesotocin differs from oxytocin by the substitution of isoleucine for leucine at position 8, and involving a single nucleotide substitution, unlike vasopressin which differs from oxytocin by two amino acids (Chauvet *et al.* 1984, Acher *et al.* 1995), and furthermore acts via a receptor that has a similar pharmacological profile to the mammalian oxytocin receptor. In the tammar wallaby, mesotocin secretion is elevated




Figure 3 Diurnal rhythm in time of onset of parturition in the rat. Meta-analysis of observations on parturition in 293 rats, caged singly and housed under controlled conditions, with lights on for 14 h and off for 10 h each day. The time of birth of the first pup shows a clear bimodal distribution across days 21 and 22 of pregnancy, with fewest first births in the lights-off phase. Litter size does not seem to be an important factor in determining time or day of birth. Since the time of birth can be advanced by shifting the lighting cues in mid-pregnancy (Lincoln & Porter 1976), events early in pregnancy do not determine the time of birth. The likely explanation for the diurnal variation in the timing of the first birth is that a maternal signal late in pregnancy triggers the process. In primates, there is a diurnal variation in uterine contractile activity that follows a diurnal rhythm in oxytocin secretion from the maternal posterior pituitary (Hirst *et al.* 1993, Giussani *et al.* 1996), and this is proposed as the explanation for the pattern of timing of first births seen in the rat. Oxytocin infusion can advance the onset of parturition by a few hours, and oxytocin antagonist infusion can delay the onset by a few hours (Antonijevic *et al.* 1995a,b). Infusion of an ineffective dose of oxytocin also delays the onset of parturition (Antonijevic *et al.* 1995b), probably by desensitising the myometrium. In the mouse, in contrast with the rat, most births occur during the dark phase (around day 19), but a similar explanation may hold (i.e. diurnal variation in maternal oxytocin secretion), although in the oxytocin knockout mouse the precise timing of parturition has not yet been reported (Nishimori *et al.* 1996).

immediately after delivery of the neonate, consistent with release during or immediately after delivery (Parry *et al.* 1996), and again, as in eutherian mammals, uterine sensitivity to oxytocin is maximal at term (see Renfree 1994). In the tamar wallaby, oxytocin antagonists delay the onset of parturition, suggesting that this mechanism is crucially important (Renfree *et al.* 1996). Parturition in the tamar wallaby (Hearn 1973) and possum (Hinds 1990) is prevented by hypophysectomy, and this is a consequence of loss of posterior rather than anterior pituitary hormones (see Bathgate *et al.* 1995).

Parturition in the rat

Probably the most complete account of the role of oxytocin during parturition comes from studies in the rat.

During pregnancy in the rat, the posterior pituitary content of oxytocin increases by about 50%; during the 90 min or so of normal parturition, this accumulated excess of some 2 µg (1000 mU, sufficient for 1000 to 2000 milk ejections) is secreted into the circulation. Parturition is thus the most powerful physiological stimulus known for the release of oxytocin (Fuchs & Saito 1971). Parturition normally begins, and is completed, during the second half of the light cycle on day 21 of gestation (mated on day 0), otherwise the onset will tend to be delayed until the late morning of the following day; for example, meta-analysis of the start times of 293 parturitions in our laboratory (lights off 1900 h, on 0500 h) shows a mode between 1500 and 1700 h on day 21 ($n=59$) and a second mode between 0900 and 1100 h ($n=15$) on day 22, with a nadir between 0300 and 0500 h ($n=3$) on day 22 (Fig. 3).

The mechanisms underlying this timing are not known, but the supraoptic and paraventricular nuclei both receive a direct afferent projection from the suprachiasmatic nucleus (Cui *et al.* 1997), which is the principal circadian oscillator in mammals (Hastings 1995). Oxytocin secretion is continuously elevated throughout parturition, and superimposed on this elevated baseline secretion are large pulses of oxytocin release (Higuchi *et al.* 1986). There is clear evidence that the pulsatile pattern of oxytocin secretion is important in driving parturition. The pulsatile secretion derives from the synchronised intense bursting activity of oxytocin neurones in the hypothalamus in the rat and rabbit (Summerlee 1981, O'Byrne *et al.* 1986), activity similar to that in the rat during reflex milk ejection in response to suckling. The increased electrical activity is accompanied by induction in these neurones of the immediate-early gene *c-fos*, an indicator of trans-synaptic activation of the neurones, and consequent prominent expression of the protein product Fos in the nuclei of oxytocin cells (Luckman *et al.* 1993). Furthermore, expression of the oxytocin gene in supraoptic neurones is stimulated at parturition (Douglas *et al.* 1998), while it is controversial whether expression is increased in pregnancy (Horwitz *et al.* 1994, Douglas *et al.* 1998). Fos is also expressed during parturition in brainstem neurones, which project to the oxytocin cells, and these brainstem neurones are activated by uterine contractions even in the absence of parturition itself (Antonijevic *et al.* 1995b). The secreted oxytocin enters the circulation at a time when uterine responsiveness to oxytocin is maximal as a result of a rapid increase in oxytocin receptor expression in the myometrium at the end of pregnancy (Rozen *et al.* 1995). Inhibition of oxytocin secretion by exogenous opiates (which act to inhibit the oxytocin neurones directly) slows down the early progress of parturition, and this can be reversed by oxytocin infusion (Russell *et al.* 1989), with pulsatile delivery of oxytocin being particularly effective (Luckman *et al.* 1993, Antonijevic *et al.* 1995b) (Fig. 2). Similarly, mild environmental stress slows down the early progress of parturition and reduces oxytocin secretion, and these effects are fully reversed by the opioid antagonist naloxone, which increases oxytocin secretion and restores the normal progress of parturition (Leng *et al.* 1988). Blocking the actions of oxytocin by specific antagonists similarly leads to impaired early progress of parturition (Chan & Chen 1992, Antonijevic *et al.* 1995a).

However, notably, magnocellular vasopressin neurones are also activated during parturition (Lin *et al.* 1995), and vasopressin release from the posterior pituitary is enhanced during parturition in the rat (Hartman *et al.* 1986) and in other species (e.g. cow (Landgraf *et al.* 1983), sheep (Kendrick *et al.* 1991), pig (Lawrence *et al.* 1995)). Vasopressin may act weakly at myometrial oxytocin receptors in the rat (Chan *et al.* 1996), while in the human and rabbit an action on myometrial vasopressin (V_{1a}) receptors is also possible (Bossmar *et al.* 1994, Maggi *et al.* 1988, 1990). Since vasopressin can activate oxytocin receptors,

its actions would be expected to supplement those of oxytocin during parturition. Is it possible then that, in the absence of oxytocin, vasopressin would be an effective substitute? One might think this unlikely given the much lower potency of vasopressin at the oxytocin receptor (Berde & Boissonas 1968), but vasopressin is quite effective at stimulating prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) production by the decidua (Chen *et al.* 1994). Furthermore, if in the absence of endogenous oxytocin the uterine oxytocin receptors are up-regulated, then the actions of vasopressin alone might indeed provide an effective substitute for the absent oxytocin.

In the oxytocin knockout mouse, vasopressin mRNA content in the supraoptic and paraventricular nuclei is normal (Nishimori *et al.* 1996) or reduced (Young *et al.* 1996), although this has not been measured in pregnancy.

Parturition in the mouse

Strictly, the outcome of studies on the oxytocin knockout mice ought to be considered in the context of previous experiments on the role of oxytocin in parturition in the mouse. As in the rat and other species, the content of oxytocin in the posterior pituitary decreases during parturition in the mouse (Fuchs 1985), but other than this little is known. Parturition can be interrupted by environmental disturbance, indicating a central mechanism influencing parturition (Newton *et al.* 1968), but it is not known if this is due to inhibition of oxytocin secretion as it is in the rat. Yet transgenic mice lacking oxytocin have no obvious defects in parturition (Nishimori *et al.* 1996, Young *et al.* 1996). In both lines, the expected numbers of live young were delivered, but few details are given on the timing of parturition, although one of the studies (Nishimori *et al.* 1996) reports that homozygous females mated with homozygous males delivered 18.5–19.5 days post coitum, which is approximately as expected of normal mice.

The sequences of the human, rat, mouse, bovine and sheep oxytocin receptor genes are known (Kimura *et al.* 1992, Adan *et al.* 1995, Kubota *et al.* 1996, Ivell *et al.* 1995, Riley *et al.* 1995, Rozen *et al.* 1995), and there is a striking conservation of the extracellular sequences of the receptor between species so far studied. The mouse oxytocin receptor gene, detected with a specific probe or probe for the rat receptor mRNA, is expressed in the uterus and in particular in the myometrium in pregnancy, peaking just before parturition (Kubota *et al.* 1996, Mahendroo *et al.* 1996). Functional receptor is expressed since contractile activity is stimulated by oxytocin, to which the mouse uterus becomes more sensitive by the end of pregnancy (Suzuki & Kuriyama 1975b), and comparison of actions of oxytocin and vasopressin *in vitro* are consistent with expression of, and action through, the oxytocin receptor (Stepke *et al.* 1994); sensitivity to vasopressin is also increased at the end of pregnancy (Stepke *et al.* 1994).

Unlike the rat or human (but like the cow and sheep), the uterus of the pregnant mouse does not express the oxytocin gene (Murphy & Ho 1995).

In other respects, parturition in the mouse involves mechanisms identified or proposed for other mammals. Progesterone secretion falls sharply on the day before parturition, presumably triggering events leading to parturition (Soares & Talamantes 1984). A synthetic gestagen (Org 5933) prolongs pregnancy (Gao & Short 1993). For reasons that are not clear, transgenic mice overexpressing the oestrogen receptor have prolonged gestation and dystocia (Davis *et al.* 1994). In late gestation, enzymes synthesising the 5α -reduced androgen, 3α -Adiol, (5α -androstan- $3\alpha,17\beta$ -diol) are induced in the uterus, and transgenic mice with disruption of the gene of the key enzyme 5α -reductase type 1 also have a prolonged pregnancy, then a long labour with dystocia: the patterns of uterine oxytocin receptor gene expression and plasma progesterone and oestrogen profiles are normal and these animals can be rescued by injection of either progesterone antagonist (RU486) or oxytocin (Mahendroo *et al.* 1996).

Other local uterine mechanisms involved in the onset of parturition in the mouse bear comparison with other species. Uterine cytokines, especially interleukin 1 or 2, probably acting via stimulation of decidual PGE₂ production (Dudley *et al.* 1993), are implicated in the initiation of parturition in the mouse (Romero *et al.* 1991, Romero & Tartakovsky 1992, Fidel *et al.* 1994, Hirsch *et al.* 1995, Kaga *et al.* 1996), particularly in pre-term parturition provoked by bacterial endotoxin (Kaga *et al.* 1996); although not via direct acute actions on the myometrium (Oshiro *et al.* 1993), there is circumstantial evidence for a role for neutrophils in the uterus in parturition onset (Kasik & Rice 1995). The pregnant mouse uterus contracts in response to PGE₂, and depolarisation sensitivity is much greater in late pregnancy than in early pregnancy (Suzuki & Kuriyama 1975a). PGF_{2 α} , histamine and 5-hydroxytryptamine, all potentially from uterine mast cells, together stimulate uterine contractions (Rudolph *et al.* 1992) and so does endothelin 1, although sensitivity does not change in pregnancy, unlike the large increase in sensitivity to oxytocin (Gong *et al.* 1994). Uterine histamine content increases prepartum, and is abruptly reduced postpartum (Padilla *et al.* 1990). Adrenergic agonists inhibit uterine contractions, via the β_2 -adrenoreceptor (Chen *et al.* 1994), becoming less effective near term (Cruz *et al.* 1990). Relaxin from the corpora lutea inhibits contractions of the mouse uterus via a selective receptor (Fields *et al.* 1980, Yang *et al.* 1992, Evans *et al.* 1993).

An important role for PGF_{2 α} in the initiation of parturition in the mouse is indicated by a study of parturition in mice with knockout of the receptor for PGF_{2 α} (Sugimoto *et al.* 1997). These homozygous mice are fertile, but do not deliver their young spontaneously, do not show luteolysis at term (and thus have high circulating levels of proges-

terone) and the uterus is insensitive to oxytocin, not showing the normal increase in oxytocin receptor mRNA expression at term. However, ovariectomy on day 19, precipitating a fall in circulating progesterone, is followed by induction of uterine oxytocin receptor mRNA expression 12 h later, and by parturition after a further 12 h. Clearly any action of PGF_{2 α} on the myometrium is not essential for parturition, provided that oxytocin is secreted to act on up-regulated oxytocin receptors in the myometrium. Instead, PGF_{2 α} , or rather its receptor, is essential in mice to induce luteolysis at term, which then allows, through enhanced oxytocin receptor expression, uterine sensitisation to oxytocin. If a myometrial action of PGF_{2 α} compensates for oxytocin in the oxytocin knockout mice, and oxytocin compensates for lack of any uterine actions of PGF_{2 α} in the PGF receptor knockout mice, then study of crosses between these two knockout strains should provide information about whether there are other redundant mechanisms regulating parturition in mice.

Despite some differences between the mouse and other species studied, it seems unlikely that species differences really explain the lack of apparent effect of the oxytocin knockout on parturition. What is clear though, from work across many species, is that several mechanisms concurrently regulate myometrial contractility in parturition, providing a high degree of redundancy among these mechanisms.

Given this, is there likely to be any single aspect of parturition in which oxytocin plays an essential role which cannot be compensated for by other known systems? Parturition as described above is regulated by both a uterine feedback loop and a hypothalamic feedback loop, of which oxytocin is normally the effector molecule. Clearly the uterine loop alone can effect successful parturition; however, without oxytocin, it seems likely that the progress of parturition may be largely outside the control of the maternal central nervous system. This control seems likely to be important in the circadian timing of parturition, and in regulating the initiation and progress of parturition with regard to environmental circumstances. In many species the onset and progress of parturition are critically influenced by environmental stress (dog (Bleicher 1962), mouse (Newton *et al.* 1968), rat (Leng *et al.* 1988), pig (Lawrence *et al.* 1992, 1995)). It would certainly be instructive to establish whether the knockout mouse shows any alterations in the pattern or progress of parturition in these respects. However, as suggested above, it is possible that vasopressin may, in the chronic absence of oxytocin, be an effective substitute. It is notable that mice hypophysectomised in mid-pregnancy were observed to start parturition at the appropriate day, but almost half (of sixteen) had a 'difficult and prolonged parturition' (Gardner & Allen 1942). Studies of the effects of hypophysectomy in pregnancy on parturition in several other eutherian species have produced similarly variable results (Amoroso & Porter 1966).

Lactation

Clearly, the failure of the oxytocin knockout mice to transfer milk to the suckling young, and the repair of this deficit by oxytocin injection (Nishimori *et al.* 1996, Young *et al.* 1996), is powerful evidence for an essential role for oxytocin in effecting the milk-ejection reflex. This finding is entirely consistent with many previous studies in several species. In marsupials, mesotocin has this role (Bathgate *et al.* 1995), substituting for oxytocin (Lincoln & Renfree 1981b).

In particular, oxytocin is effective in the rat when secreted during suckling in pulses a few minutes apart, and this is a consequence of synchronised burst-firing of the magnocellular oxytocin neurones projecting into the posterior pituitary gland (Belin & Moos 1986). Although bolus injection of vasopressin can elicit milk ejections, it is much less potent than oxytocin (Bisset *et al.* 1967), and vasopressin neurones do not show synchronised burst-firing, and few otherwise respond, during suckling (Lincoln & Wakerley 1974). For this reason, vasopressin, or rather vasopressin neurones, cannot substitute for oxytocin in lactation. To do so in the oxytocin knockout mouse would require altered 'wiring' of vasopressin neurones, and expression of the distinctive and specialised electrophysiological properties normally only seen in oxytocin neurones. Evidently, and not surprisingly, these adaptations do not occur in the oxytocin knockout mice. Furthermore, it is obvious that, at least in the knockout mouse, there are no other mechanisms that can effect milk transfer in the absence of oxytocin. This may reflect the fact that lactation, or specifically milk ejection, is recent in evolution, in contrast with egg laying, the precursor of parturition.

There is evidence from studies in the rat that oxytocin, possibly from the posterior pituitary although it is also secreted from magnocellular axons in the median eminence into the hypothalamohypophysial portal blood (Horn *et al.* 1985), stimulates prolactin secretion in lactation. Thus the consequences of removal or denervation of the posterior pituitary include reduced stimulation of prolactin secretion by suckling (Murai & Ben-Jonathan 1987, Vecsernyés *et al.* 1997). Furthermore, there is a dramatic increase in oxytocin receptor mRNA expression in lactotropes at the end of pregnancy (Breton *et al.* 1995). Suppression by suckling of the release of dopamine into portal blood and the stimulation of vasoactive intestinal peptide (VIP) or thyrotrophin-releasing hormone secretion are also involved in regulation of prolactin secretion (see Mogg & Samson 1990), so the evident capacity to produce milk in the oxytocin knockout mice (demonstrated by milk transfer when injected with oxytocin) implies adequate prolactin secretion through action of these mechanisms, a clear case of redundancy, and involving differential expression of receptors by subsets of lactotropes, so that the subset of lactotropes expressing oxytocin

receptors also expresses VIP receptors (Samson & Schell 1995). However, this has not been studied in detail: reduced prolactin secretion in response to suckling could be a feature of the knockout mouse. This leads to consideration of the usefulness of the oxytocin knockout mice to analyse the importance of other factors in mechanisms in which oxytocin has a role, but in which there is redundancy. Thus, in lactation, appropriate experiments could address the issue of the relative importance of suppression of dopamine secretion or stimulation of VIP secretion in suckling stimulation of prolactin secretion. Similarly, the importance of other factors in parturition, freed from contamination by any effects of oxytocin, can be examined, for example, assessing a role for vasopressin.

Oxytocin and maternal behaviour

Whereas all mammalian species express oxytocin receptors at high levels in the term pregnant uterus (see above), the distribution of oxytocin receptors in the brain shows very wide interspecies variation (Tribollet *et al.* 1992, Insel *et al.* 1993), and considerable variation is apparent even between rat and mouse. In the rat, oxytocin, produced and acting in the brain, is important in stimulating the expression of female receptive behaviour (Caldwell *et al.* 1990) and in penile erection in the male (Melis *et al.* 1994); in particular, centrally administered oxytocin facilitates lordosis by acting in the ventromedial nucleus of the hypothalamus, through specific oxytocin receptors positively regulated by oestrogen or testosterone (Johnson 1992, McCarthy *et al.* 1994, Bale & Dorsa 1995). By contrast, in the normal (and homozygous oxytocin knockout) mouse, there is a low density of oxytocin receptors in the ventromedial nucleus (Nishimori *et al.* 1996), and in the male mouse testosterone negatively regulates oxytocin receptor binding in the ventromedial nucleus, the reverse of the effects of gonadal steroid in the rat (Insel *et al.* 1993). Also, adult male mouse brain has fewer immunoreactive oxytocin neurones than the female, consistent with a suppressor action of testosterone (Haussler *et al.* 1990), and again unlike the rat, which shows no sex difference in neuronal oxytocin expression.

Notwithstanding considerable species variability in the suggested central actions of oxytocin, there is evidence across distantly related species of a role for central oxytocin in maternal behaviour. Lactating rats display a range of behaviours, together described as maternal behaviour, which are not normally shown by virgin rats, unless they are exposed to young over many days (Numan 1994). These component behaviours include nest building, gathering young into the nest, and licking and crouching over the young, and lactating rats will readily extend these attentions towards alien foster pups as well as to their own young. Maternal behaviour in the rat appears abruptly

immediately after parturition, and is maintained subsequently throughout lactation. Oxytocin, released within the brain at parturition, is thought to be involved in stimulating the rapid onset of maternal behaviour acting on up-regulated oxytocin receptors (Insel 1986), and i.c.v. injections of oxytocin antagonists are effective in preventing the induction of maternal behaviour (van Leengoed *et al.* 1987), in particular acting through the ventral tegmental and medial preoptic areas and olfactory bulbs (Pedersen *et al.* 1994, Yu *et al.* 1996). In contrast, once maternal behaviour has been induced, i.c.v. injection of oxytocin antagonists has generally been found to be ineffective, although olfactory signals are of key importance (Kolunie & Stern 1995), and antagonist administration into the olfactory bulbs does reduce maternal behaviour (Yu *et al.* 1995). Similar actions of oxytocin have been demonstrated in sheep, in which the central release of oxytocin is triggered by stimulation of the uterine cervix and vagina (Kendrick *et al.* 1988). Rapid expression of maternal behaviour can be induced in virgin rats by oestrogen treatment ('priming') followed by i.c.v. oxytocin (Pedersen & Prange 1979), provided that the testing is in a novel environment (Fahrbach *et al.* 1985). In this model, i.c.v. oxytocin antiserum disrupts maternal behaviour (Pedersen *et al.* 1985).

Against this background, what is already known about the expression of maternal behaviour in the mouse? There is an important difference between the rat and the laboratory mouse. 'Only a few studies have been directed at examining the hormonal basis of pup-directed maternal behaviour in mice, and this is because the virgin laboratory mouse usually shows *spontaneous maternal responsiveness* to test pups' (Numan 1994; our italics), and full maternal behaviour is shown by virtually all late-pregnant laboratory mice (Gandelman 1973). In the light of this, it is less surprising that postpartum maternal behaviour is normal in the oxytocin knockout mouse, since its induction in the laboratory mouse clearly cannot depend on oxytocin release during parturition. Nonetheless, there are detectable quantitative differences in maternal behaviour between virgin and primiparous mice (Laviola *et al.* 1994). Such subtleties have evidently not been sought in the oxytocin knockout mice. Because lactating mice share the nursing of their litters under natural conditions, unlike rats (Gandelman *et al.* 1970, see Numan 1994), it is likely that oxytocin deficiency in an individual lactating mouse could be compensated for by this social behaviour. Thus, in the mouse's natural social context, oxytocin could be considered to be redundant for an individual, even with respect to its role in milk ejection. Disabling natural mutations of the oxytocin gene may be prevalent in such communal nursing species.

In marked contrast with the behaviour of virgin laboratory mice, virgin wild mice usually cannibalise strange pups. Infanticide by pregnant wild mice ceases at the birth of the young, and returns only after the young are weaned.

Interestingly, infanticide in virgin wild mice is suppressed after systemic or i.c.v. oxytocin injection (McCarthy *et al.* 1986, McCarthy 1990). Olfactory processing is again implicated, since prepartum destruction of the noradrenergic input to the olfactory bulbs induces postpartum cannibalism in laboratory mice (Dickinson & Keverne 1988); the mouse olfactory bulb is rich in oxytocin receptors (Insel *et al.* 1993).

Postpartum mice (including laboratory strains) display aggression toward conspecific intruders. The aggressive behaviour is suckling-dependent (Garland & Svare 1988), but not dependent on prolactin secretion (Mann *et al.* 1980). Lactational aggressiveness towards intruders is seen also in the rat; in neither rat nor mouse has a role for central oxytocin in this behaviour been tested, but micro-injection of oxytocin into the amygdala in the hamster increases aggressiveness of lactating females toward intruder males (Ferris *et al.* 1992). In the rat, maternal aggression is independent of suckling (Mayer *et al.* 1987), prolactin secretion (Erskine *et al.* 1980) and probably oxytocin secretion (Factor *et al.* 1992), but is dependent on somatosensory inputs to the snout and ventral body surface, and is influenced by volatile odours acting via the olfactory epithelium and olfactory bulb (Ferreira *et al.* 1987, Stern & Kolunie 1993, Kolunie & Stern 1995). This behaviour in the rat also involves lateral connections of the ventromedial nucleus, a region rich in oxytocin receptors (Bale & Dorsa 1995), which are not involved in pup-directed behaviour (Hansen 1989).

In male oxytocin knockout mice, there is reduced aggressiveness toward intruder males, with no evident sensorimotor deficits in the homozygous knockouts, although reversal by centrally administered oxytocin has not been tested (De Vries *et al.* 1997).

With regard to this group of actions of oxytocin, in facilitating receptive or copulatory or maternal and related behaviours, for the knockout mice, the fact that the oxytocin knockout is lifelong leaves the possibility for compensation by recruitment of other neuropeptides acting on oxytocin receptors; an obvious candidate is vasopressin. With respect to actions in the brain, matching of oxytocin receptor distribution to sites of vasopressin release, but under appropriate conditions, would be required, while oxytocin receptors in the brain bind vasopressin with a similar affinity for oxytocin (see Barberis & Tribollet 1996). In the rat, centrally administered vasopressin can, like oxytocin, induce maternal behaviour in appropriately primed virgin rats (Pedersen *et al.* 1982).

In short, the studies to date on normal or oxytocin knockout mice do not allow a critical evaluation of the role of central oxytocin in the expression of maternal behaviour in the mouse. This is made difficult by the facility with which the neural circuitry for this behaviour is activated even in virgin female laboratory mice: in wild mice, and other species, facilitation by release of oxytocin within the brain is likely to be important, with a broad spectrum of

necessity, while the oxytocin gene product will interact with other neurotransmitters in shaping and evoking maternal behaviour.

Oxytocin and gonadal function

In ruminants, oxytocin from the corpus luteum provides an essential signal to the non-pregnant endometrium to trigger PGF_{2α} release, which in turn initiates luteolysis and drives further luteal oxytocin secretion in a positive feedback circuit (Flint *et al.* 1990); in early pregnancy in the sheep, suppression of endometrial oxytocin receptor by interferon τ from the embryo is essential to prevent luteolysis (Lamming *et al.* 1995). Many species show some evidence of gonadal expression of either vasopressin or oxytocin; in the human, oxytocin and oxytocin receptors are expressed in cumulus cells around the oocytes (Furuya *et al.* 1995), and oxytocin is expressed in Sertoli cells of the bovine testis (Ang *et al.* 1991). However, there are considerable interspecies differences in the regulation of luteolysis, and there is no evidence that gonadal oxytocin plays a role in rodents similar to that documented for ruminants. Previous studies in the mouse on the role of oxytocin in gonadal function indicate a capacity to stimulate ovulation (Robinson *et al.* 1985) and weak expression in granulosa cells; at the anterior pituitary, oxytocin can advance the luteinizing hormone preovulatory surge in the rat (Robinson & Evans 1990). Evidence has also been put forward that oxytocin can stimulate blastocyst development *in vitro* (Furuya *et al.* 1995), act on seminiferous tubule motility (Nicholson *et al.* 1986) and possibly stimulate, via vasopressin receptors, testosterone production (Tahri-Joutei & Pointis 1989), although the oxytocin gene is not naturally expressed in the mouse testis (Ang *et al.* 1991).

Reconciliation

If the above accounts of studies on the roles of oxytocin in mice and other mammals appear to be inconsistent with observations on oxytocin knockout mice, the inconsistency may be more apparent than real. Specifically, with regard to parturition, it is perfectly reasonable to affirm that oxytocin is released from the pituitary in large pulses, which act upon a uterus that is expressing an abundance of oxytocin receptors, and thereby influences uterine contractility and the progress of parturition, while at the same time affirming that, in the absence of oxytocin, other mechanisms may be substituted to ensure ultimately successful parturition. Indeed, it would have been brave to deny the likely truth of any part of this, for there have been many previous well-described examples of normal labour in humans and experimental animals with apparently complete posterior pituitary dysfunction (Amoroso &

Porter 1966). Even in birds, egg laying has been observed to occur after acute neurohypophysectomy (Nakada *et al.* 1993). With regard to other reproductive roles of oxytocin, it seems clear that species differences are considerable and obscure the interpretation of results from the knockout mouse.

Conclusion

Despite the reservations expressed in this review, there is much that can be learned about the roles of oxytocin in the oxytocin knockout mouse. In particular, even if the results from the knockout mouse can tell us little about the role played by oxytocin in reproductive functions, they can certainly be invaluable in telling us how these are fulfilled in the absence of oxytocin.

If nothing else, it is now evident that oxytocin has physiological actions that are strictly redundant. Natural selection is harsh in culling features that are counter-adaptive, but clearly the actions of oxytocin on the uterus at term, which involve co-ordinated up-regulation of uterine oxytocin receptor expression, if not essential are not counteradaptive. Indeed, we may expect that any local uterine, neural or endocrine mechanism that will tend to favour successful delivery at term will be preserved by selection, provided that the costs of maintaining a redundant mechanism do not exceed the costs of eliminating it. The neurohypophysial system regulates the delivery of progeny in all vertebrates, and, during its long evolutionary history, other mechanisms may have evolved convergent roles simply by a process of exclusion. When everything that opposes the actions of oxytocin in parturition is excluded, the things that remain are neutral, assist oxytocin or, in dogging the footsteps of oxytocin, can substitute for it. What these key substitutes are in the oxytocin knockout mouse are at present not known.

Acknowledgements

Research in our laboratory is supported by the BBSRC, MRC, Wellcome Trust, British Council/ARC, Ferring. Data in Fig. 3 are from collaborative studies with Dr R J Bicknell, The Babraham Institute, Cambridge, Cambs, UK.

References

- Acher R, Chauvet J & Chauvet MT 1995 Man and the chimaera: selective versus neutral oxytocin evolution. *Advances in Experimental Medicine and Biology* **395** 615–627.
- Adan RA, Walther N, Cox JJ, Ivell R & Burbach JP 1991 Comparison of the estrogen responsiveness of the rat and bovine oxytocin gene promoters. *Biomedical and Biophysical Research Communications* **175** 117–122.

- Adan RA, Van Leeuwen FW, Sonnemans MA, Brouns M, Hoffman G, Verbalis JG & Burbach JP 1995 Rat oxytocin receptor in brain, pituitary, mammary gland, and uterus: partial sequence and immunocytochemical localization. *Endocrinology* **136** 4022–4028.
- Alexandrova M & Soloff MV 1980 Oxytocin receptors and parturition in the guinea pig. *Biology of Reproduction* **22** 1106–1111.
- Amoroso EC & Porter DG 1966 Anterior pituitary function in pregnancy. In *The Pituitary Gland*, vol 2, pp 364–411. Eds GW Harris & BT Donovan. London: Butterworths.
- Ang HL, Ungefroren H, De Bree F, Foo NC, Carter D, Burbach JP, Ivell R & Murphy D 1991 Testicular oxytocin gene expression in seminiferous tubules of cattle and transgenic mice. *Endocrinology* **128** 2110–2117.
- Antonijevic IA, Douglas AJ, Dye S, Bicknell RJ, Leng G & Russell JA 1995a Oxytocin antagonists delay the initiation of parturition and prolong its active phase in rats. *Journal of Endocrinology* **145** 97–103.
- Antonijevic I, Leng G, Luckman SM, Douglas AJ, Bicknell RJ & Russell JA 1995b Induction of uterine activity with oxytocin in late pregnant rats replicates the expression of *c-fos* in neuroendocrine and brainstem neurones as seen during parturition. *Endocrinology* **136** 154–163.
- Bale TL & Dorsa DM 1995 Regulation of oxytocin receptor messenger ribonucleic acid in the ventromedial hypothalamus by testosterone and its metabolites. *Endocrinology* **136** 5135–5138.
- Barberis C & Tribollet E 1996 Vasopressin and oxytocin receptors in the central nervous system. *Critical Reviews in Neurobiology* **10** 119–154.
- Bathgate RA, Parry LJ, Fletcher TP, Shaw G, Renfree MB, Gemmill RT & Sernia C 1995 Comparative aspects of oxytocin-like hormones in marsupials. *Advances in Experimental Medicine and Biology* **395** 639–655.
- Belin V & Moos F 1986 Paired recording from supraoptic and paraventricular oxytocin cells in suckled rats: recruitment and synchronisation. *Journal of Physiology* **377** 369–390.
- Bell GH 1941 The behaviour of the pregnant uterus of the guinea-pig. *Journal of Physiology* **100** 263–274.
- Berde B & Boissonnas RA 1968 Basic pharmacological properties of synthetic analogues and homologues of the neurohypophysial hormones. In *Handbook of Experimental Pharmacology XXIII*, pp 802–870. Ed B Berde. Berlin: Springer-Verlag.
- Bisset GW, Clark BJ, Haldar J, Harris MC, Lewis GP & Roche e Silva MJ 1967 The assay of milk-ejecting activity in the lactating rat. *British Journal of Pharmacology and Chemotherapy* **31** 537–549.
- Bleicher N 1962 Behavior of the bitch during parturition. *Journal of the American Veterinary Medical Association* **140** 1076–1082.
- Boer K, Lincoln DW & Swaab DF 1975 Effects of electrical stimulation of the neurohypophysis on labour in the rat. *Journal of Endocrinology* **65** 163–176.
- Bossmar T, Akerlund M, Fantoni G, Szamatowicz J, Melin P & Maggi M 1994 Receptors for and myometrial responses to oxytocin and vasopressin in preterm and term human pregnancy: effects of the oxytocin antagonist atosiban. *American Journal of Obstetrics and Gynecology* **171** 1634–1642.
- Breton C, Pechoux C, Morel G & Zingg HH 1995 Oxytocin receptor messenger ribonucleic acid: characterization, regulation, and cellular localization in the rat pituitary gland. *Endocrinology* **136** 2928–2936.
- Burbach JPH, van Schaick HSA, de Bree FM, Lopes da Silva S & Adan RAH 1995 Functional domains in the oxytocin gene for regulation of expression and biosynthesis of gene products. *Advances in Experimental Medicine and Biology* **395** 9–21.
- Burd JM, Davison J, Weightman DR & Baylis PH 1987 Evaluation of enzyme inhibitors of pregnancy associated oxytocinase: application to the measurement of plasma immunoreactive oxytocin during human labour. *Acta Endocrinologica* **114** 458–464.
- Burghardt RC, Barhouni R, Stickney M, Monga M, Ku CY & Sanborn BM 1996 Correlation between connexin 43 expression, cell-cell communication, and oxytocin-induced Ca^{2+} responses in an immortalized human myometrial cell line. *Biology of Reproduction* **55** 433–438.
- Caldwell JD, Barakat AS, Smith DD, Hrubby VJ & Pedersen CA 1990 A uterotonic antagonist blocks the oxytocin-induced facilitation of female sexual receptivity. *Brain Research* **512** 291–296.
- Chan WY 1983 Uterine and placental prostaglandins and their modulation of oxytocin sensitivity and contractility in the parturient uterus. *Biology of Reproduction* **29** 680–688.
- Chan WY & Chen DL 1992 Myometrial oxytocin receptors and prostaglandin in the parturition process in the rat. *Biology of Reproduction* **46** 58–64.
- Chan WY, Chen DL & Manning M 1993 Oxytocin receptor subtypes in the pregnant rat myometrium and decidua: pharmacological differentiations. *Endocrinology* **132** 1381–1386.
- Chan WY, Wo NC & Manning M 1996 The role of oxytocin receptors and vasopressin V1a receptors in uterine contractions in rats: implications for tocolytic therapy with oxytocin antagonists. *American Journal of Obstetrics and Gynecology* **175** 1331–1335.
- Chard T 1989 Fetal and maternal oxytocin in human parturition. *American Journal of Perinatology* **6** 145–152.
- Chauvet J, Hurpet D, Chauvet MT & Acher R 1984 Divergent neuropeptide evolutionary drifts between American and Australian marsupials. *Bioscience Reports* **4** 245–252.
- Chen DL, Chan WY & Manning M 1994 Agonist and antagonist specificities of decidual prostaglandin-releasing oxytocin receptors and myometrial uterotonic oxytocin receptors in pregnant rats. *Journal of Reproduction and Fertility* **102** 337–343.
- Chibbar R, Miller FD & Mitchell BF 1993 Synthesis of oxytocin in amnion, chorion and decidua may influence the timing of human parturition. *Journal of Clinical Investigation* **91** 185–192.
- Chibbar R, Wong S, Miller FD & Mitchell BF 1995 Estrogen stimulates oxytocin gene expression in human chorio-decidua. *Journal of Clinical Endocrinology and Metabolism* **80** 567–572.
- Chwalisz K, Fahrenholz F, Hackenberg M, Garfield R & Elger W 1991 The progesterone antagonist onapristone increases the effectiveness of oxytocin to produce delivery without changing the myometrial oxytocin receptor concentrations. *American Journal of Obstetrics and Gynecology* **165** 1760–1770.
- Cohen-Tannoudji J, Mhaouty S, Elwardy-Merezak J, Lecrivain JL, Robin MT, Legrand C & Maltier JP 1995 Regulation of myometrial Gi2, Gi3, and Gq expression during pregnancy. Effects of progesterone and estradiol. *Biology of Reproduction* **53** 55–64.
- Crankshaw DJ 1987 The sensitivity of the longitudinal and circular muscle layers of the rat's myometrium to oxytocin *in vitro* during pregnancy. *Canadian Journal of Physiology and Pharmacology* **65** 773–777.
- Cruz MA, Sepulveda WH & Rudolph MI 1990 Changes in the response to adrenergic drugs on mouse uterine contractions during pregnancy. *Life Sciences* **46** 99–104.
- Cui L-N, Saerb-Parsy K & Dyball REJ 1997 Neurones in the supraoptic nucleus of the rat are regulated by a projection from the suprachiasmatic nucleus. *Journal of Physiology* **502** 149–159.
- Cummiskey KC & Dawood MY 1990 Induction of labor with pulsatile oxytocin. *American Journal of Obstetrics and Gynecology* **163** 1868–1874.
- Currie WB, Gorewit RC & Michel FJ 1988 Endocrine changes, with special emphasis on oestradiol-17beta, prolactin and oxytocin, before and during labour and delivery in goats. *Journal of Reproduction and Fertility* **82** 299–308.
- Dale HH 1906 On some physiological actions of ergot. *Journal of Physiology* **34** 163–206.
- Davis VL, Couse JF, Goulding EH, Power SG, Eddy EM & Korach KS 1994 Aberrant reproductive phenotypes evident in transgenic mice expressing the wild-type mouse estrogen receptor. *Endocrinology* **135** 379–386.

- Dawood MY, Khan-Dawood FS, Ayromlooi J & Tobias M 1983 Maternal and fetal plasma oxytocin levels during pregnancy and parturition in the sheep. *American Journal of Obstetrics and Gynecology* **147** 584–588.
- De Vries AC, Young WS & Nelson RJ 1997 Reduced aggressive behaviour in mice with targeted disruption of the oxytocin gene. *Journal of Neuroendocrinology* **9** 363–368.
- Dickinson C & Keverne EB 1988 Importance of noradrenergic mechanisms in the olfactory bulbs for the maternal behaviour of mice. *Physiology and Behavior* **43** 313–316.
- Dong YL & Yallampalli C 1996 Interaction between nitric oxide and prostaglandin E2 pathways in pregnant rat uteri. *American Journal of Physiology* **270** E471–476.
- Douglas AJ, Clarke G, MacMillan SJA, Bull PM, Neumann I, Way SA, Wright DM, McGrory BG & Russell JA 1993 Effects of the kappa-opioid agonist U50,488 on parturition in rats. *British Journal of Pharmacology* **109** 251–258.
- Douglas AJ, Meeren HKM, Johnstone LE, Pfaff DW, Russell JA & Brooks PJ 1998 Stimulation of expression of the oxytocin gene in rat supraoptic neurons at parturition. *Brain Research* **782** 167–174.
- Dudley DJ, Chen CL, Branch DW, Hammond E & Mitchell MD 1993 A murine model of preterm labor: inflammatory mediators regulate the production of prostaglandin E2 and interleukin-6 by murine decidua. *Biology of Reproduction* **48** 33–39.
- Elwardy-Merezak J, Maltier JP, Cohen-Tannoudji J, Lecrivain JL, Vivat V & Legrand C 1994 Pregnancy-related modifications of rat myometrial Gs proteins: ADP ribosylation, immunoreactivity and gene expression studies. *Journal of Molecular Endocrinology* **13** 23–37.
- Erskine MS, Barfield RJ & Goldman BD 1980 Postpartum aggression in rats. I. Effects of hypophysectomy. *Journal of Comparative and Physiological Psychology* **94** 484–494.
- Europe-Finner GN, Phaneuf S, Tolkovsky AM, Watson SP & López Bernal A 1994 Down-regulation of G_{as} in human myometrium in term and preterm labor: a mechanism for parturition. *Journal of Clinical Endocrinology and Metabolism* **79** 1835–1839.
- Evans BA, John M, Fowler KJ, Summers RJ, Cronk M, Shine J & Tregear GW 1993 The mouse relaxin gene: nucleotide sequence and expression. *Journal of Molecular Endocrinology* **10** 15–23.
- Factor EM, Mayer AD & Rosenblatt JS 1992 Preventing suckling-induced release of oxytocin does not inhibit maternal aggression in lactating rats. *Annals of the New York Academy of Sciences* **652** 423–424.
- Fahrbach SE, Morrell JI & Pfaff DW 1985 Role of oxytocin in the onset of estrogen-facilitated maternal behavior. In *Oxytocin: Clinical and Laboratory Studies*, pp 372–388. Eds JA Amico & AG Robinson. Amsterdam: Excerpta Medica.
- Fang X, Wong S & Mitchell BF 1996 Relationships among sex steroids, oxytocin, and their receptors in the rat uterus during late gestation and at parturition. *Endocrinology* **137** 3213–3219.
- Ferreira A, Dahlof LG & Hansen S 1987 Olfactory mechanisms in the control of maternal aggression, appetite, and fearfulness: effects of lesions to olfactory receptors, mediodorsal thalamic nucleus, and insular prefrontal cortex. *Behavioral Neuroscience* **101** 709–717.
- Ferris CF, Foote KB, Meltzer HM, Plenby MG, Smith KL & Insel TR 1992 Oxytocin in the amygdala facilitates maternal aggression. *Annals of the New York Academy of Sciences* **652** 456–457.
- Fidel PL, Romero R, Wolf N, Cutright J, Ramirez M, Araneda H & Cotton DB 1994 Systemic and local cytokine profiles in endotoxin-induced preterm parturition in mice. *American Journal of Obstetrics and Gynecology* **170** 1467–1475.
- Fields MJ, Fields PA, Castro-Hernandez A & Larkin LH 1980 Evidence for relaxin in corpora lutea of late pregnant cows. *Endocrinology* **107** 869–876.
- Flint APF, Sheldrick EL, McCann TJ & Jones DSC 1990 Luteal oxytocin: characteristics and control of synchronous episodes of oxytocin and PGF_{2α} secretion at luteolysis in ruminants. *Domestic Animal Endocrinology* **7** 111–124.
- Fuchs AR 1972 Prostaglandin effects on rat pregnancy. I. Failure of induction of labour. *Fertility and Sterility* **23** 410–416.
- Fuchs AR 1985 Oxytocin in animal parturition. In *Oxytocin: Clinical and Laboratory Studies*, pp 207–235. Eds JA Amico & AG Robinson. Amsterdam: Elsevier, Excerpta Medica.
- Fuchs AR & Dawood MY 1980 Oxytocin release and uterine activation during parturition in rabbits. *Endocrinology* **107** 1117–1126.
- Fuchs AR & Saito S 1971 Pituitary oxytocin and vasopressin content of pregnant rats before, during and after parturition. *Endocrinology* **88** 574–578.
- Fuchs AR, Periyasamy S, Alexandrova M & Soloff MS 1983 Correlation between oxytocin receptor concentration and responsiveness to oxytocin in pregnant rat myometrium: effects of ovarian steroids. *Endocrinology* **113** 742–749.
- Fuchs AR, Fuchs F, Husslein P & Soloff MS 1984 Oxytocin receptors in the human uterus during pregnancy and parturition. *American Journal of Obstetrics and Gynecology* **150** 734–741.
- Fuchs AR, Romero R, Keefe D, Parra M, Oyarzun E & Behnke E 1991 Oxytocin secretion and human parturition: pulse frequency and duration increase during spontaneous labor in women. *American Journal of Obstetrics and Gynecology* **165** 1515–1523.
- Fuchs AR, Fields MJ, Freidman S, Shemesh M & Ivell R 1995 Oxytocin and the timing of parturition. *Advances in Experimental Biology and Medicine* **395** 405–433.
- Fuchs AR, Rollyson MK, Meyer M, Fields MJ, Mixin JM & Randel RD 1996 Oxytocin induces prostaglandin F_{2α} release in pregnant cows: influence of gestational age and oxytocin receptor concentrations. *Biology of Reproduction* **54** 647–653.
- Furuya K, Mizumoto Y, Makimura N, Mitsui C, Murakami M, Tokuoka S, Ishikawa N, Nagata I, Kimura T & Ivell R 1995 A novel biological aspect of ovarian oxytocin: gene expression of oxytocin and oxytocin receptor in cumulus/luteal cells and the effect of oxytocin on embryogenesis in fertilized oocytes. *Advances in Experimental Medicine and Biology* **395** 523–528.
- Gandelman R 1973 The ontogeny of maternal responsiveness in female Rockland-Swiss albino mice. *Hormones and Behavior* **4** 257–268.
- Gandelman R, Paschke RE, Zarrow MX & Denenberg VH 1970 Care of young under communal conditions in the mouse (*Mus musculus*). *Developmental Psychobiology* **3** 245–250.
- Gao Y & Short RV 1993 Use of an oestrogen, androgen or gestagen as a potential chemosterilant for control of rat and mouse populations. *Journal of Reproduction and Fertility* **97** 39–49.
- Gardner WU & Allen E 1942 Effect of hypophysectomy at mid-pregnancy in the mouse. *Anatomical Record* **83** 75–98.
- Garland M & Svare B 1988 Suckling stimulation modulates the maintenance of postpartum aggression in mice. *Physiology and Behavior* **44** 301–305.
- Gilbert CL, Goode JA & McGrath TJ 1994 Pulsatile secretion of oxytocin during parturition in the pig: temporal relationship with fetal expulsion. *Journal of Physiology* **475** 129–137.
- Giussani DA, Jenkins SL, Mecenas CA, Winter JA, Honnebler BO, Wu W & Nathanielsz PW 1996 Daily and hourly temporal association between delta4-androstenedione-induced preterm myometrial contractions and maternal plasma estradiol and oxytocin concentrations in the 0.8 gestation rhesus monkey. *American Journal of Obstetrics and Gynecology* **174** 1050–1055.
- Glatz TH, Weitzman RE, Eliot RJ, Klein AH, Nathanielsz PW & Fisher DA 1981 Ovine maternal and fetal plasma oxytocin concentrations before and during parturition. *Endocrinology* **108** 1328–1332.
- Golan A, Sandbank O & Rubin A 1980 Rupture of the pregnant uterus. *Obstetrics and Gynecology* **56** 549–554.
- Goldsmith LT, Weiss G & Steinetz BG 1995 Relaxin and its role in pregnancy. *Endocrinology and Metabolism Clinics of North America* **24** 171–186.

- Gong QY, Chen LA, Li JZ, Xu WH & Yang ZC 1994 Effects of endothelin-1 on isolated uterine horns in estrogen-primed and pregnant mice. *Chung-Kuo Yao Li Hsueh Pao-Acta Pharmacologica Sinica* **15** 129–132.
- Goodwin TM, Valenzuela GJ, Silver H & Creasy G 1996 Dose ranging study of the oxytocin antagonist atosiban in the treatment of preterm labor. Atosiban Study Group. *Obstetrics and Gynecology* **88** 331–336.
- Gunther M 1948 The posterior pituitary and labour. *British Medical Journal* **1** 567.
- Haluska GJ & Currie WB 1988 Variation in plasma concentrations of oestradiol-17 beta and their relationship to those of progesterone, 13,14-dihydro-15-keto-prostaglandin F-2 alpha and oxytocin across pregnancy and at parturition in pony mares. *Journal of Reproduction and Fertility* **84** 635–646.
- Hansen S 1989 Medial hypothalamic involvement in maternal aggression of rats. *Behavioral Neuroscience* **103** 1035–1046.
- Hartman RD, Rosella-Dampman LM, Emmert SE & Summy-Long JY 1986 Inhibition of release of neurohypophysial hormones by endogenous opioid peptides in pregnant and parturient rats. *Brain Research* **382** 352–359.
- Hastings M 1995 Circadian rhythms: peering into the molecular clockwork. *Journal of Neuroendocrinology* **7** 331–340.
- Hausler HU, Jirikowski GF & Caldwell JD 1990 Sex differences among oxytocin-immunoreactive neuronal systems in the mouse hypothalamus. *Journal of Chemical Neuroanatomy* **3** 271–276.
- Hearn JP 1973 Pituitary inhibition of pregnancy. *Nature* **241** 207–208.
- Heller H 1973 The effects of oxytocin and vasopressin during the oestrous cycle and pregnancy on the uterus of a marsupial species, the quokka (*Setonix brachyurus*). *Journal of Endocrinology* **58** 657–671.
- Higuchi T, Uchide K, Honda K & Negoro H 1985 Functional development of the oxytocin release mechanism and its role in the initiation of parturition in the rat. *Journal of Endocrinology* **106** 311–316.
- Higuchi T, Tadokoro Y, Honda K & Negoro H 1986 Detailed analysis of blood oxytocin levels during suckling and parturition in the rat. *Journal of Endocrinology* **110** 251–256.
- Hinds LA 1990 Control of pregnancy, parturition and luteolysis in marsupials. *Reproduction, Fertility and Development* **2** 535–552.
- Hirsch E, Saotome I & Hirsh D 1995 A model of intrauterine infection and preterm delivery in mice. *American Journal of Obstetrics and Gynecology* **172** 1598–1603.
- Hirst JJ, Haluska GJ, Cook MJ & Novy MJ 1993 Plasma oxytocin and nocturnal uterine activity: maternal but not fetal concentrations increase progressively during late pregnancy and delivery in rhesus monkeys. *American Journal of Obstetrics and Gynecology* **169** 415–422.
- Honnabier MB, Figueroa JP, Rivier J, Vale W & Nathanielsz PW 1989 Studies on the role of oxytocin in late pregnancy in the pregnant rhesus monkey: plasma concentrations of oxytocin in the maternal circulation throughout the 24-h day and the effect of the synthetic oxytocin antagonist [1-beta-Mpa(beta-(CH₂)₅)¹, (OMe)Tyr², Orn³] oxytocin on spontaneous nocturnal myometrial contractions. *Journal of Developmental Physiology* **12** 225–232.
- Horn AM, Robinson IC & Fink G 1985 Oxytocin and vasopressin in rat hypophysial portal blood: experimental studies in normal and Brattleboro rats. *Journal of Endocrinology* **104** 211–224.
- Horwitz MJ, Bloch KD, Kim NB & Amico JA 1994 Expression of the endothelin-1 and oxytocin genes in the hypothalamus of the pregnant rat. *Brain Research* **648** 59–64.
- Inoue T, Kimura T, Azuma C, Inazawa J, Takemura M, Kikuchi T, Kubota Y, Ogita K & Saji F 1994 Structural organization of the human oxytocin receptor gene. *Journal of Biological Chemistry* **269** 32451–32456.
- Insel TR 1986 Post partum increases in brain oxytocin binding. *Neuroendocrinology* **44** 515–518.
- Insel TR, Young L, Witt DM & Crews D 1993 Gonadal steroids have paradoxical effects on brain oxytocin receptors. *Journal of Neuroendocrinology* **5** 619–628.
- Ivell R, Hunt N, Abend N, Brackman B, Nollmeyer D, Lamsa JC & McCracken JA 1990 Structure and expression of the oxytocin gene in the sheep. *Reproduction, Fertility and Development* **2** 703–711.
- Ivell R, Rust W, Einspanier A, Hartung S, Fields MJ & Fuchs AR 1995 Oxytocin and oxytocin receptor gene expression in the reproductive tract of the pregnant cow: rescue of luteal oxytocin production at term. *Biology of Reproduction* **51** 551–558.
- Izumi H & Garfield RE 1995 Relaxant effects of nitric oxide and cyclic GMP on pregnant rat uterine longitudinal smooth muscle. *European Journal of Obstetrics, Gynecology and Reproductive Biology* **60** 171–180.
- Jenkin G, Buttress D & Harding R 1994 Oxytocin receptor blockade and prostaglandin release in late pregnant sheep. *European Journal of Obstetrics, Gynecology and Reproductive Biology* **53** 1–6.
- Johnson AE 1992 The regulation of oxytocin receptor binding in the ventromedial hypothalamic nucleus by gonadal steroids. *Annals of the New York Academy of Sciences* **682** 357–373.
- Kaga N, Katsuki Y, Obata M & Shibutani Y 1996 Repeated administration of low-dose lipopolysaccharide induces preterm delivery in mice: a model for human preterm parturition and for assessment of the therapeutic ability of drugs against preterm delivery. *American Journal of Obstetrics and Gynecology* **174** 754–759.
- Kaneko Y, Kawarabayashi T, Sugimori H & Tsukamoto T 1995 Changes in kinetic properties of oxytocin receptors in longitudinal muscle membranes of rat uterus during gestation. *Journal of Molecular Recognition* **8** 179–183.
- Kasik JW & Rice EJ 1995 An increase in expression of the lipocalin 24p3 is found in mouse uterus coincident with birth. *American Journal of Obstetrics & Gynecology* **173** 613–617.
- Kawarabayashi T, Kobayashi M, Akahane M & Ajisawa Y 1996 Comparison of *in vitro* and *in vivo* inhibitory effects of peptide and nonpeptide oxytocin antagonists on radioligand binding and uterine contractility of rats during pregnancy. *American Journal of Obstetrics and Gynecology* **175** 1348–1355.
- Kendrick KM, Keverne EB, Chapman C & Baldwin BA 1988 Microdialysis measurement of oxytocin, aspartate, γ -aminobutyric acid and glutamate release from the olfactory bulb of the sheep during vaginocervical stimulation. *Brain Research* **442** 171–174.
- Kendrick KM, Keverne EB, Hinton MR & Goode JA 1991 Cerebrospinal fluid and plasma concentrations of oxytocin and vasopressin during parturition and vaginocervical stimulation in the sheep. *Brain Research Bulletin* **26** 803–807.
- Kimura T, Tanizawa O, Mori K, Brownstein MJ & Okayama H 1992 Structure and expression of a human oxytocin receptor. *Nature* **356** 526–529.
- Kimura T, Takemura M, Nomura S, Nobunaga T, Kubota Y, Inoue T, Hashimoto K, Kumazawa I, Ito Y, Ohashi K *et al.* 1996 Expression of oxytocin receptor in human pregnant myometrium. *Endocrinology* **137** 780–785.
- Koike TI, Shimada K & Cornett LE 1988 Plasma levels of immunoreactive mesotocin and vasotocin during oviposition in chickens: relationship to oxytocic action of the peptides *in vitro* and peptide interaction with myometrial membrane binding sites. *General and Comparative Endocrinology* **70** 119–126.
- Kolonie JM & Stern JM 1995 Maternal aggression in rats: effects of olfactory bulbectomy, ZnSO₄-induced anosmia, and vomeronasal organ removal. *Hormones and Behavior* **29** 492–518.
- Kubota Y, Kimura T, Hashimoto K, Tokugawa Y, Nobunaga K, Azuma C, Saji F & Murata Y 1996 Structure and expression of the mouse oxytocin receptor gene. *Molecular and Cellular Endocrinology* **124** 25–32.
- Lamming GE, Wathes DC, Flint APF, Payne JM, Stevenson KR & Vallet JL 1995 Trophoblast interferons act locally to suppress oxytocin and oestradiol receptor development in ovine endometrium. *Journal of Reproduction and Fertility* **105** 165–175.

- Landgraf R, Schulz J, Eulenberger K & Wilhelm J 1983 Plasma levels of oxytocin and vasopressin before, during and after parturition in cows. *Experimental and Clinical Endocrinology* **81** 321–328.
- Larcher A, Neculcea J, Breton C, Arslan A, Rozen F, Russo C & Zingg HH 1995 Oxytocin receptor gene expression in the rat uterus during pregnancy and the estrous cycle and in response to gonadal steroid treatment. *Endocrinology* **136** 5350–5356.
- Laviola G, Petrucci S, Rankin J & Alleva E 1994 Induction of maternal behavior by mouse neonates: influence of dam parity and prenatal oxazepam exposure. *Pharmacology, Biochemistry and Behavior* **49** 871–876.
- Lawrence AB, Petherick JC, McLean K, Gilbert C, Chapman C & Russell JA 1992 Naloxone prevents interruption of parturition and increases plasma oxytocin following environmental disturbance in parturient sows. *Physiology and Behavior* **52** 917–923.
- Lawrence AB, Petherick JC, McLean KA, Deans L, Chirnside J, Vaughan A, Gilbert CL, Forsling ML & Russell JA 1995 The effects of chronic environmental stress on parturition and on oxytocin and vasopressin secretion in the pig. *Animal Reproduction Science* **38** 251–264.
- Leake RD, Weitzman RE, Glatz TH & Fisher DA 1981 Plasma oxytocin concentrations in men, nonpregnant women, and pregnant women before and during spontaneous labor. *Journal of Clinical Endocrinology and Metabolism* **53** 730–733.
- van Leengoed E, Kerker E & Swanson HH 1987 Inhibition of post partum maternal behaviour in the rat by injecting an oxytocin antagonist into the cerebral ventricles. *Journal of Endocrinology* **112** 275–282.
- Lefebvre DL, Lariviere R & Zingg HH 1993 Rat amnion: a novel site of oxytocin production. *Biology of Reproduction* **48** 632–639.
- Lefebvre DL, Farookhi R, Giaid A, Neculcea J & Zingg HH 1994 Uterine oxytocin gene expression. II. Induction by exogenous steroid administration. *Endocrinology* **134** 2562–2566.
- Leng G, Mansfield S, Bicknell RJ, Blackburn RE, Brown D, Chapman C, Dyer RG, Hollingsworth S, Shibuki K & Way S 1988 Endogenous opioid actions and effects of environmental disturbance on parturition and oxytocin secretion in rats. *Journal of Reproduction and Fertility* **84** 345–356.
- Lin SH, Miyata S, Itoh T, Kawarabayashi T, Nakashima T & Kiyohara T 1995 Fos expression in the hypothalamic magnocellular neurons of rats during pregnancy, parturition and lactation. *Neuroscience Research* **23** 29–34.
- Lincoln DW 1971 Labour in the rabbit: effect of electrical stimulation applied to the infundibulum and median eminence. *Journal of Endocrinology* **50** 607–618.
- Lincoln DW & Porter DG 1976 Timing of the photoperiod and the hour of birth in rats. *Nature* **260** 780–781.
- Lincoln DW & Renfree MB 1981a Milk ejection in a marsupial, *Macropus agilis*. *Nature* **289** 504–506.
- Lincoln DW & Renfree MB 1981b Mammary growth and milk ejection in the agile wallaby, *Macropus agilis*, displaying concurrent asynchronous lactation. *Journal of Reproduction and Fertility* **63** 193–203.
- Lincoln DW & Wakerley JB 1974 Electrophysiological evidence for the activation of supraoptic neurones during the release of oxytocin. *Journal of Physiology* **242** 533–554.
- Liu CX, Takahashi S, Murata T, Hashimoto K, Agatsuma T, Matsukawa S & Higuchi T 1996 Changes in oxytocin receptor mRNA in the rat uterus measured by competitive reverse transcription-polymerase chain reaction. *Journal of Endocrinology* **150** 479–486.
- Luckman SM, Antonijevic I, Leng G, Dye S, Douglas AJ, Russell JA & Bicknell RJ 1993 The maintenance of normal parturition in the rat requires neurohypophysial oxytocin. *Journal of Neuroendocrinology* **5** 7–12.
- McCarthy MM 1990 Oxytocin inhibits infanticide in female house mice (*Mus domesticus*). *Hormones and Behavior* **24** 365–375.
- McCarthy MM, Bare JE & vom Saal FS 1986 Infanticide and parental behavior in wild female house mice: effects of ovariectomy, adrenalectomy and administration of oxytocin and prostaglandin F2 alpha. *Physiology and Behavior* **36** 17–23.
- McCarthy MM, Kleopoulos SP, Mobbs CV & Pfaff DW 1994 Infusion of antisense oligodeoxynucleotides to the oxytocin receptor in the ventromedial hypothalamus reduces estrogen-induced sexual receptivity and oxytocin receptor binding in the female rat. *Neuroendocrinology* **59** 432–440.
- Maggi M, Genazzani AD, Giannini S, Torrisci C, Baldi E, di Tomasso M, Munson PJ, Rodbard D & Serio M 1988 Vasopressin and oxytocin receptors in vagina, myometrium, and oviduct of rabbits. *Endocrinology* **122** 2970–2980.
- Maggi M, Del Carlo P, Fantoni G, Giannini S, Torrisci C, Casparis D, Massi G & Serio M 1990 Human myometrium during pregnancy contains and responds to V1 vasopressin receptors as well as oxytocin receptors. *Journal of Clinical Endocrinology and Metabolism* **70** 1142–1154.
- Maggi M, Peri A, Giannini S, Fantoni G, Guardabasso V & Serio M 1991 Oxytocin and V1 vasopressin receptors in rabbit endometrium during pregnancy. *Journal of Reproduction and Fertility* **91** 575–581.
- Mahendroo MS, Cala KM & Russell DW 1996 5 Alpha-reduced androgens play a key role in murine parturition. *Molecular Endocrinology* **10** 380–392.
- Mann M, Michael SD & Svare B 1980 Ergot drugs suppress plasma prolactin and lactation but not aggression in parturient mice. *Hormones and Behavior* **14** 319–328.
- Mayer AD, Carter L, Jorge WA, Mota MJ, Tannu S & Rosenblatt JS 1987 Mammary stimulation and maternal aggression in rodents: thelectomy fails to reduce pre- or postpartum aggression in rats. *Hormones and Behavior* **21** 501–510.
- Mecenas CA, Giussani DA, Owiny JR, Jenkins SL, Wu WX, Honnebler BO, Lockwood CJ, Kong L, Guller S & Nathanielsz PW 1996 Production of premature delivery in pregnant rhesus monkeys by androstenedione infusion. *Nature Medicine* **2** 443–448.
- Meera P, Anwer K, Monga M, Oberti C, Stefani E, Toro L & Sanborn BM 1995 Relaxin stimulates myometrium calcium-activated potassium channel activity via protein kinase A. *American Journal of Physiology* **269** C312–C317.
- Melis MR, Stancampiano R & Argiolas A 1994 Penile erection and yawning induced by paraventricular NMDA injection in male rats are mediated by oxytocin. *Pharmacology Biochemistry and Behavior* **48** 203–207.
- Michel G, Chauvet J, Chauvet MT, Clarke C, Bern H & Acher R 1993 Chemical identification of the mammalian oxytocin in a holocephalian fish, the ratfish (*Hydrolagus collie*). *General and Comparative Endocrinology* **92** 260–268.
- Mitchell BF & Chibbar R 1995 Synthesis and metabolism of oxytocin in late gestation in human decidua. *Advances in Experimental Medicine and Biology* **395** 365–380.
- Mitchell MD, Edwin S & Romero RJ 1990 Prostaglandin biosynthesis by human decidual cells: effects of inflammatory mediators. *Prostaglandins Leukotrienes and Essential Fatty Acids* **41** 35–38.
- Mitchell MD, Dudley DJ, Edwin SS & Schiller SL 1991 Interleukin-6 stimulates prostaglandin production by human amnion and decidual cells. *European Journal of Pharmacology* **192** 189–191.
- Mogg RJ & Samson WK 1990 Interactions of dopaminergic and peptidergic factors in the control of prolactin release. *Endocrinology* **126** 728–735.
- Murai I & Ben-Jonathan N 1987 Posterior pituitary lobectomy abolishes the suckling-induced rise in prolactin (PRL): evidence for a PRL-releasing factor in the posterior pituitary. *Endocrinology* **121** 205–211.
- Murphy D & Ho M-Y 1995 Oxytocin transgenic mice. *Advances in Experimental Medicine and Biology* **395** 67–78.

- Nakada T, Koja Z & Tanaka K 1993 The influence of hypophysectomy on the time of oviposition in hens. *Japanese Poultry Science* **30** 306–309.
- Nathanielsz PW, Giussani DA, Mecenas CA, Wu W, Winter JA, Garcia-Villar R, Baguma-Nibasheka M, Honnebiel MB & McDonald TJ 1995 Regulation of the switch from myometrial contractures to contractions in late pregnancy: studies in the pregnant sheep and monkey. *Reproduction, Fertility and Development* **7** 595–602.
- Newton N, Peeler D & Newton M 1968 Effect of disturbance on labor. An experiment with 100 mice with dated pregnancies. *American Journal of Obstetrics and Gynecology* **101** 1096–1102.
- Nicholson HD, Worley RT, Charlton HM & Pickering BT 1986 LH and testosterone cause the development of seminiferous tubule contractile activity and the appearance of testicular oxytocin in hypogonadal mice. *Journal of Endocrinology* **110** 159–167.
- Nishimori K, Young LJ, Guo Q, Wang Z, Insel TR & Matzuk MM 1996 Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. *Proceedings of the National Academy of Sciences of the USA* **93** 11699–11704.
- Numan M 1994 Maternal behavior. In *The Physiology of Reproduction*, vol 2, pp 221–302. Eds E Knobil & JD Neill. New York: Raven Press.
- O'Byrne KT, Ring JP & Summerlee AJ 1986 Plasma oxytocin and oxytocin neurone activity during delivery in rabbits. *Journal of Physiology* **370** 501–513.
- Orsino A, Taylor CV & Lye SJ 1996 Connexin-26 and connexin-43 are differentially expressed and regulated in the rat myometrium throughout late pregnancy and with the onset of labor. *Endocrinology* **137** 1545–1553.
- Oshiro BT, Monga M, Eriksen NL, Graham JM, Weisbrodt NW & Blanco JD 1993 Endotoxin, interleukin-1 beta, interleukin-6, or tumor necrosis factor-alpha do not acutely stimulate isolated murine myometrial contractile activity. *American Journal of Obstetrics and Gynecology* **169** 1424–1427.
- Owen J & Hanth JC 1992 Oxytocin for the induction or augmentation of labor. *Clinical Obstetrics and Gynecology* **35** 464–475.
- Owiny JR, Mitchell M & Nathanielsz PW 1992 Effect of 48-hour infusion of the synthetic oxytocin antagonist, [1-beta-mercaptopropyl-(beta-(CH₂)₅)¹(OMe)Tyr², Orn⁸]-oxytocin, on myometrial activity of pregnant sheep at 139–140 days of gestation. *Biology of Reproduction* **47** 436–440.
- Padilla L, Reinicke K, Montesino H, Villena F, Asencio H, Cruz M & Rudolph MI 1990 Histamine content and mast cells distribution in mouse uterus: the effect of sexual hormones, gestation and labor. *Cellular and Molecular Biology* **36** 93–100.
- Parry LJ, Guymer FJ, Fletcher TP & Renfree MB 1996 Release of an oxytocic peptide at parturition in the marsupial, *Macropus eugenii*. *Journal of Reproduction and Fertility* **107** 191–198.
- Pedersen CA & Prange AJ 1979 Induction of maternal behavior in virgin rats after intracerebroventricular administration of oxytocin. *Proceedings of the National Academy of Sciences of the USA* **76** 6661–6665.
- Pedersen CA, Ascher JA, Monroe YL & Prange AJ 1982 Oxytocin induces maternal behavior in virgin female rats. *Science* **216** 648–650.
- Pedersen CA, Caldwell JD, Johnson MF, Fort SA & Prange AJ 1985 Oxytocin antiserum delays onset of ovarian steroid-induced maternal behavior. *Neuropeptides* **6** 175–182.
- Pedersen CA, Caldwell JD, Walker C, Ayers G & Mason GA 1994 Oxytocin activates the postpartum onset of rat maternal behavior in the ventral tegmental and medial preoptic areas. *Behavioral Neuroscience* **108** 1163–1171.
- Phaneuf S, Asbóth G, Carrasco MP, Europe-Finner GN, Saji F, Kimura T, Harris A & López Bernal A 1997 The desensitization of oxytocin receptors in human myometrial cells is accompanied by down-regulation of oxytocin receptor messenger RNA. *Journal of Endocrinology* **154** 7–18.
- Randolph GW & Fuchs AR 1989 Pulsatile administration enhances the effect and reduces the dose of oxytocin required for induction of labor. *American Journal of Perinatology* **6** 159–166.
- Renfree MB 1994 Endocrinology of pregnancy, parturition and lactation in marsupials. In *Marshall's Physiology of Reproduction*, edn 4, pp 677–766. Ed GE Lamming. London: Chapman & Hall.
- Renfree MB, Parry LJ & Shaw G 1996 Infusion with an oxytocin receptor antagonist delays parturition in a marsupial. *Journal of Reproduction and Fertility* **108** 131–137.
- Richard S & Zingg HH 1990 The human oxytocin gene promoter is regulated by estrogen. *Journal of Biological Chemistry* **265** 6098–6103.
- Riemer RK, Goldfien AC, Goldfien A & Roberts JM 1986 Rabbit uterine oxytocin receptors and *in vitro* contractile response: abrupt changes at term and the role of eicosanoids. *Endocrinology* **119** 699–709.
- Riley PR, Flint AP, Abayasekara DR & Stewart HJ 1995 Structure and expression of an ovine endometrial oxytocin receptor cDNA. *Journal of Molecular Endocrinology* **15** 195–202.
- Robinson AG & Amico JA 1985 Remarks on the history of oxytocin. In *Oxytocin: Clinical and Laboratory Studies*, pp xvii–xxxiii. Eds JA Amico & AG Robinson. Amsterdam: Excerpta Medica International Congress Series.
- Robinson G & Evans JJ 1990 Oxytocin has a role in gonadotrophin regulation in rats. *Journal of Endocrinology* **125** 425–432.
- Robinson G, Evans JJ & Forster ME 1985 Oxytocin can affect follicular development in the adult mouse. *Acta Endocrinologica* **108** 273–276.
- Romero R & Tartakovsky B 1992 The natural interleukin-1 receptor antagonist prevents interleukin-1-induced preterm delivery in mice. *American Journal of Obstetrics and Gynecology* **167** 1041–1045.
- Romero R, Mazor M & Tartakovsky B 1991 Systemic administration of interleukin-1 induces preterm parturition in mice. *American Journal of Obstetrics and Gynecology* **165** 969–971.
- Rozen F, Russo C, Banville D & Zingg H 1995 Structure, characterization, and expression of the rat oxytocin receptor gene. *Proceedings of the National Academy of Sciences of the USA* **92** 200–204.
- Rudolph MI, Bardisa L, Cruz M & Reinicke K 1992 Mast cells mediators evoke contractility and potentiate each other in mouse uterine horns. *General Pharmacology* **23** 833–836.
- Russell JA, Gosden RG, Humphreys EM, Cutting R, Fitzsimons N, Johnston V, Liddle S, Scott S & Stirland JA 1989 Interruption of parturition in rats by morphine: a result of inhibition of oxytocin secretion. *Journal of Endocrinology* **121** 521–536.
- Saito N & Koike TI 1992 Alterations in uterine contractility during the oviposition cycle in domestic hens. *British Poultry Science* **33** 671–676.
- Samson WK & Schell DA 1995 Oxytocin and the anterior pituitary gland. *Advances in Experimental Medicine and Biology* **395** 355–364.
- Sanborn BM, Qian A, Ku CY, Wen Y, Anwer K, Monga M & Singh SP 1995 Mechanisms regulating oxytocin receptor coupling to phospholipase C in rat and human myometrium. *Advances in Experimental Medicine and Biology* **395** 469–479.
- Sander-Richter H, Schams D, Ellendorff F 1988 Fetal and postnatal oxytocin secretion in the pig before and after a prostaglandin challenge. *Journal of Developmental Physiology* **10** 63–76.
- Sanyal MK 1978 Secretion of progesterone during gestation in the rat. *Journal of Endocrinology* **79** 179–190.
- Schellenberg JC 1995 The effect of oxytocin receptor blockade on parturition in guinea pigs. *Journal of Clinical Investigation* **95** 13–19.
- Soares MJ & Talamantes F 1984 Pre-parturitional changes in serum prolactin, placental lactogen, growth hormone, progesterone, and corticosterone in the C3H/HeN mouse. *Journal of Developmental Physiology* **6** 423–429.
- Soloff MS, Alexandrova M & Fernstrom MJ 1979 Oxytocin receptors: triggers for parturition and lactation? *Science* **204** 1313–1315.

- Stepke MT, Schwenger N & Eichhorn W 1994 Vasoconstrictors during pregnancy: *in vitro* trial on pregnant and nonpregnant mouse uterus. *International Journal of Oral and Maxillofacial Surgery* **23** 440–442.
- Stern JM & Kolunje JM 1993 Maternal aggression of rats is impaired by cutaneous anesthesia of the ventral trunk, but not by nipple removal. *Physiology and Behavior* **54** 861–868.
- Sugimoto Y, Yamasaki A, Sergi E, Tsuboi K, Aze Y, Nishimura T, Oida H, Yoshida N, Tanaka T, Katsuyama M, Hasumoto K-y, Murata T, Hirata M, Ushikubi F, Negishi M, Ichikawa A & Narumiya S 1997 Failure of parturition in mice lacking the prostaglandin receptor. *Science* **277** 681–683.
- Summerlee AJ 1981 Extracellular recordings from oxytocin neurones during the expulsive phase of birth in unanaesthetized rats. *Journal of Physiology* **321** 1–9.
- Suzuki H & Kuriyama H 1975a Comparison between prostaglandin E2 and oxytocin actions on pregnant mouse myometrium. *Japanese Journal of Physiology* **25** 345–356.
- Suzuki H & Kuriyama H 1975b Effects of prostaglandin E2 on the electrical property of the pregnant mouse myometrium. *Japanese Journal of Physiology* **25** 201–215.
- Suzuki H, Okawa T, Endo C, Hoshi K & Sato A 1995 Influence of endotoxin on sensitivity to oxytocin in pregnant rat myometrium. *Acta Obstetrica et Gynecologica Japonica* **47** 377–384.
- Tabb T, Thilander G, Grover A, Hertzberg E & Garfield R 1992 An immunohistochemical and immunocytochemical study of the increase in myometrial gap junctions (and connexin-43) in rats and humans during pregnancy. *American Journal of Obstetrics and Gynecology* **167** 559–567.
- Tahri-Joutei A & Pointis G 1989 Developmental changes in arginine vasopressin receptors and testosterone stimulation in Leydig cells. *Endocrinology* **125** 605–611.
- Takahashi K, Diamond F, Bieniarz J, Yen H & Burd L 1980 Uterine contractility and oxytocin sensitivity in preterm, term, and postterm pregnancy. *American Journal of Obstetrics and Gynecology* **136** 774–779.
- Takahashi T, Kawashima M, Kamiyoshi M & Tanaka K 1994 Arginine vasotocin receptor binding in the hen uterus (shell gland) before and after oviposition. *European Journal of Endocrinology* **130** 366–372.
- Tezuka N, Ali M, Chwalisz K & Garfield RE 1995 Changes in transcripts encoding calcium channel subunits of rat myometrium during pregnancy. *American Journal of Physiology* **269** C1008–C1017.
- Thornton S, Davison JM & Baylis PH 1992 Plasma oxytocin during the first and second stages of spontaneous human labour. *Acta Endocrinologica* **126** 425–429.
- Todd HM, Dundoo VL, Gerber WR, Cwiak CA, Baldassare JJ & Hertelendy F 1996 Effect of cytokines on prostaglandin E2 and prostacyclin production in primary cultures of human myometrial cells. *Journal of Mass Spectrometry* **5** 161–167.
- Tribollet E, Barberis C, Dubois-Dauphin M & Dreifuss JJ 1992 Localization and characterization of binding sites for vasopressin and oxytocin in the brain of the guinea pig. *Brain Research* **589** 15–23.
- Vasicka A, Kumaresan P, Han GS & Kumaresan M 1978 Plasma oxytocin in initiation of labor. *American Journal of Obstetrics and Gynecology* **130** 263–273.
- Vavra N, Eppel W, Sevela P, Fitz R, Fuchs AR, Fuchs F & Husslein P 1993 Serum prostaglandin F2 alpha (PGFM) and oxytocin levels correlate with sonographic changes in the cervix in patients with preterm labor. *Archives of Gynecology and Obstetrics* **253** 33–36.
- Vecsernyés M, Krempels K, Tóth BE, Julesz J, Makara GB & Nagy GM 1997 Effect of posterior pituitary denervation (PPD) on prolactin (PRL) and α -melanocyte stimulating hormone (α MSH) secretion of lactating rats. *Brain Research Bulletin* **43** 313–319.
- Vivrette S 1994 The endocrinology of parturition in the mare. *Veterinary Clinics of North America-Equine Practice* **10** 1–17.
- Wathes DC, Smith HF, Leung ST, Stevenson KR, Meier S & Jenkin G 1996 Oxytocin receptor development in ovine uterus and cervix throughout pregnancy and at parturition as determined by *in situ* hybridization analysis. *Journal of Reproduction and Fertility* **106** 23–31.
- Wilson LJ, Parsons MT & Flouret G 1990 Inhibition of spontaneous uterine contractions during the last trimester in pregnant baboons by an oxytocin antagonist. *American Journal of Obstetrics and Gynecology* **163** 1875–1882.
- Wilson T, Liggins GC & Whittaker DJ 1988 Oxytocin stimulates the release of arachidonic acid and prostaglandin F_{2a} from human decidual cells. *Prostaglandins* **35** 771–780.
- Wu WX, Verbalis JG, Hoffman GE, Derks JB & Nathanielsz PW 1996 Characterization of oxytocin receptor expression and distribution in the pregnant sheep uterus. *Endocrinology* **137** 722–728.
- Yang S, Rembisa B, Bullesbach EE & Schwabe C 1992 Relaxin receptors in mice: demonstration of ligand binding in symphyseal tissues and uterine membrane fragments. *Endocrinology* **130** 179–185.
- Young WS, Shepard E, Amico J, Hennighausen L, Wagner K-U, La Marca ME, McKinney C & Ginns EI 1996 Deficiency in mouse oxytocin prevents milk ejection, but not fertility or parturition. *Journal of Neuroendocrinology* **8** 847–853.
- Yu G-Z, Kaba H & Higuchi T 1995 The action of oxytocin on the olfactory bulb is required for both initiation and maintenance of maternal behavior. [Abstract] *First Joint World Congress of Neurohypophysiology and Vasopressin, Nasu, Japan*, P-54.
- Yu G-Z, Kaba H, Okutani F, Takahashi S & Higuchi T 1996 The olfactory bulb: a critical site of action for oxytocin in the induction of maternal behaviour in the rat. *Neuroscience* **72** 1083–1088.

Received 28 July 1997

Revised manuscript received 22 December 1997

Accepted 16 January 1998