The antiprogesterin RU486 dissociates LH and FSH secretion in male rats: evidence for direct action at the pituitary level

J E Sánchez-Criado, C Bellido, M Tébar, A Ruiz and D Gonzalez

Department of Physiology, Faculty of Medicine, University of Córdoba, Spain

(Requests for offprints should be addressed to J E Sánchez-Criado, Department of Physiology, Faculty of Medicine, Avda. Menendez Pidal s/n, University of Córdoba, 14004 Córdoba, Spain)

Abstract

Administration of 4 mg of the antisteroid RU486 over 8 consecutive days to adult male rats dissociated *in vivo* and *in vitro* gonadotrophin secretion, increasing FSH and decreasing LH secretion. In subsequent experiments we evaluated the involvement of testicular or adrenal secretory products, as well as hypothalamic LHRH, in the effects of 4 consecutive days of RU486 treatment on the secretion of gonadotrophins. The first day of RU486 injection was designated day 1, subsequent days being numbered consecutively. Groups of rats injected with oil (0.2 ml) or RU486 (4 mg) were: (i) injected s.c. from day 1 to day 4 with the antiandrogen flutamide (10 mg/kg); (ii) bilateral orchidectomized (ORCH) on day 1; and (iii) bilateral adrenalectomized (ADX) on day 1. Controls were given flutamide vehicle or were sham operated. To ascertain whether the secretion of LHRH is involved in the effects of RU486 on gonadotrophin secretion, we measured the LHRH secretion into the pituitary stalk blood vessels at 1100 h on day 5 in oil- or RU486-treated rats. Additional oil- and RU486-treated rats were injected i.p. with 100 ng LHRH at 1000 h on day 5, or s.c. with 1 mg

LHRH antagonist (LHRH-ANT) at 1000 h on days 2 and 4. Controls were given saline. All animals were decapitated at 1100 h on day 5, trunk blood collected and serum stored frozen until FSH, LH and testosterone assays.

While ADX had no effect on FSH and LH secretion in either oil- or RU486-treated rats, the removal of androgen negative feedback with flutamide treatment or by ORCH substantially increased serum levels of FSH and LH in both oil- and RU486-treated rats, and thus annulled the effects of RU486. No differences in pituitary stalk plasma LHRH concentrations were found between oil- and RU486-treated rats. Injection of LHRH increased serum FSH and LH concentrations in oil-treated rats but only, and to a lesser extent, LH concentrations in RU486-treated rats. Treatment with LHRH-ANT decreased serum concentrations of FSH and LH in both oil- and RU486-treated rats. These results suggest that RU486 inhibited LHRH-stimulated LH secretion at the pituitary level, and that FSH secretion increased in response to a reduction in the negative feedback of androgen.

Journal of Endocrinology (1999) **160**, 197–203

Introduction

The antiprogesterin RU486 is a synthetic steroid with strong antiprogesterone and antiglucocorticoid activity (Philibert 1984, Baulieu 1989). It has been shown that administration of RU486 to female cyclic rats dissociates dioestrous secretion of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) (Sánchez-Criado *et al.* 1990, 1992, 1993), reduces the magnitude of the preovulatory LH surge (Rao & Mahesh 1986, Sánchez-Criado *et al.* 1990) and abolishes the primary and secondary surge of FSH (Knox & Schwartz 1992, Knox *et al.* 1993, Sánchez-Criado *et al.* 1994). These effects of RU486 on FSH and LH secretion could be the result of the blockade of progesterone and/or glucocorticoid action (Lusting *et al.* 1988, Tébar *et al.* 1995). Also, results from recent experiments indicate that RU486 may affect FSH (Szabo *et al.* 1996, Ringstrom *et al.* 1997) and LH (Waring

& Turgeon 1992, Turgeon & Waring 1994, Levine 1997) secretion through the blockade of ligand-independent activation of progesterone receptor.

In a pilot experiment we found that in male rats which have pituitary progesterone receptors (Kato *et al.* 1985, Negro-Vilar *et al.* 1984) and negligible progesterone levels (Döhler & Wuttke 1975, Schwartz & Justo 1977), administration of RU486 increases FSH, while decreasing LH secretion. This effect is opposite to that found in females, in which RU486 decreases FSH and increases LH baseline secretion (Sánchez-Criado *et al.* 1990, 1992, 1993, 1994). Aiming at elucidating this intriguing effect of RU486 in male rats, in the present study we characterized the involvement of the hypothalamus and the steroid-secreting glands in the effects of RU486 on FSH and LH secretion in male rats. The results indicated that RU486, acting at the pituitary, decreased basal luteinizing hormone-releasing hormone (LHRH)-stimulated LH secretion.

Furthermore, the results suggest that FSH secretion increased in response to a partial reduction of negative androgen feedback.

Materials and Methods

Animals

Adult male Wistar rats weighing 250–300 g were used. The rats were housed (four per cage) under standard lighting (lights on from 0500 to 1900 h) and temperature (20–23 °C) conditions and had free access to rat chow and tap water. Animals were treated in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Drugs, treatment and surgery

Antiprogesteragen RU486 (mifepristone, 11 β -(4-dimethylaminophenyl)-17 β -hydroxy-17 α -(1-propynyl)-estra-4,9-dien-3-one) was donated by Dr Sitruk-Ware (Exelgyn, Paris, France). This compound has a high affinity for progesterone and glucocorticoid receptors (Philibert 1984, Baulieu 1989). Injections consisted of 4 mg/0.2 ml oil (s.c.). Control injections consisted of oil.

Antiandrogen flutamide (4'-nitro-3'-trifluoro-methylisobutyranilide; Schering-Plough, Madrid, Spain) was dissolved in oil at a concentration of 20 mg/ml, and administered s.c. at 10 mg/kg per injection. This dose effectively blocks androgen action (Chandolia *et al.* 1991). Controls received oil.

The synthetic LHRH used (Peninsula 7201) was obtained from Peninsula Laboratories, Inc., Belmont, CA, USA. Immediately before use, the peptide was dissolved in saline to a concentration of 200 ng/ml. The animals were injected i.p. with 0.5 ml of this solution. Controls were given 0.5 ml saline.

The LHRH antagonist (LHRH-ANT) used was ORG.30276 (Ac-D-p-Cl-Phe-D-p-Cl-Phe-D-Trp-Ser-Tyr-D-Arg-Leu-Arg-Pro-D-Ala-NH₂-CH₃-COOH) (Organon International BV, Oss, The Netherlands). Immediately before use, the peptide was dissolved in saline to a concentration of 5 mg/ml. Injections consisted of 0.2 ml of this solution. This dosage causes maximal suppression of LH serum levels (van den Dungen *et al.* 1989, Sánchez-Criado *et al.* 1993). Controls were injected with saline.

Bilateral orchidectomy (ORCH) was performed under light ether anaesthesia through the scrotal route between 0900 and 1100 h on day 1. Sham-ORCH consisted of exposing but not removing the testes.

Adrenalectomy (ADX) was performed under ether anaesthesia between 0900 and 1100 h on day 1. Adrenal glands were removed through a subcostal 1 cm incision on both sides. Sham-ADX consisted of exposing but not removing the adrenals.

Pituitary stalk blood collection

The pituitary stalk was exposed following the methods of Worthington (1966) and Sarkar *et al.* (1976) under midatrene (Glaxo Laboratories, Enghien, France). This anaesthetic consists of 9 mg alphaxalone (3 α -hydroxy-5 α -pregnane-11,20-dione) and 3 mg alphadolone acetate (21-acetoxy-3 α -hydroxy-5 α -pregnane-11,20-dione) per ml. It was injected i.p. at a dosage of 0.5 ml/100 g body weight.

Pituitary incubation

Incubations were carried out as described previously (Rodríguez-Padilla *et al.* 1987). Briefly, the anterior pituitary was cut in half and the halves were incubated at 37 °C with constant shaking in an atmosphere of 95% CO₂-5% O₂. After 60 min of preincubation, FSH and LH concentrations were measured at 120 and 240 min of incubation.

Experiments

In the first experiment the effects of RU486 on FSH and LH secretion *in vivo* and *in vitro* were studied. Just after the collection of blood, rats were injected daily with RU486 or oil over 8 consecutive days. The first day of treatment was designated as day 1. Less than 0.5 ml blood was withdrawn at 1100 h by direct jugular venipuncture under light ether anaesthesia on days 1, 3, 5 and 7. At 1100 h on day 9 all animals were killed by decapitation, trunk blood was collected, and their pituitaries were rapidly collected for pituitary incubation. Serum was stored frozen until assayed for LH and FSH.

In the second experiment the possible involvement of testes or adrenal gland secretory products in the effects of RU486 on LH and FSH secretion were evaluated. Groups of rats injected with RU486 or oil were: (i) injected with flutamide from day 1 to day 4; (ii) ORCH; or (iii) ADX. Controls were sham-operated or injected with oil. All rats were decapitated at 1100 h on day 5. Trunk blood was collected and the serum was stored frozen until assayed for LH and FSH.

In the third experiment we explored the participation of the hypothalamic LHRH in the effects of RU486 on LH and FSH secretion. First, male rats received injections of oil or RU486 over 4 consecutive days and, on day 5, 144 $\pm$ 5.4 μ l portal blood were collected for 30 min between 1000 and 1200 h. Secondly, oil- or RU486-injected rats were given LHRH at 1000 h on day 5 or LHRH-ANT at 1000 h on days 2 and 4. All animals were decapitated at 1100 h on day 5. Trunk blood was collected and serum stored frozen until assayed for gonadotrophins.

RIA of LHRH

Pituitary stalk plasma concentrations of LHRH were measured in duplicate in a direct RIA using the LHRH

antisera HU60 supplied by Dr H F Urbanski (Neuroscience Division, Oregon Regional Primate Research Center, Beaverton, OR, USA). A complete characterization of this antiserum has been reported by Urbanski *et al.* (1990). Synthetic LHRH (Peninsula Laboratories) was used as the standard and labelled with ^{125}I by the chloramine-T method (Greenwood *et al.* 1963). The assay was routinely performed under disequilibrium conditions as reported by Sanchez-Criado *et al.* (1994). To avoid interference with other plasma proteins, bound and free LHRH were separated using a second antibody system (López *et al.* 1990). All samples were run in the same assay. Under these conditions, the sensitivity of the assay was 0.25 pg per tube with an intra-assay variability of 8%.

RIA of FSH and LH

The concentrations of FSH and LH were measured in duplicate in 25 μl samples by double-antibody RIA methods, using the RIA kits supplied by the National Institute of Diabetes, Digestive and Kidney Diseases (Baltimore, MD, USA) and following a microassay method described previously (Sánchez-Criado *et al.* 1993). Rat FSH-I-8 and LH-I-9 were labelled with ^{125}I using the chloramine-T method (Greenwood *et al.* 1963). Serum and medium FSH and LH concentrations were expressed as ng/ml and ng/hemihypophysis respectively of the reference preparations LH-rat-RP-3 and FSH-rat-RP-2. All samples were run in the same assay. The intra-assay coefficients of variation were 8 and 7% for LH and FSH respectively. The sensitivities of the assay were 20 and 7.5 pg/tube for FSH and LH respectively.

RIA of testosterone

Serum concentrations of testosterone were measured by the RIA method described previously (Rodríguez-Padilla *et al.* 1987), using antiserum supplied by Dr G D Niswender (Colorado State University, Fort Collins, CO, USA). All samples were run in the same assay. The intra-assay coefficient of variation was 6% and the sensitivity 5 pg/tube.

Data evaluation and statistical analysis

Results were expressed as means $\pm$ S.E.M. Data were analysed by one-way ANOVA using Duncan's multiple range test or Student's *t*-test when two means had to be compared. Results were considered significant at the $P < 0.05$ level.

Results

Effect of RU486 on FSH, LH and testosterone secretion in intact rats

Administration of RU486 over 8 consecutive days increased FSH serum concentrations throughout the ex-

Figure 1 Serum concentrations of (A) FSH, (B) LH and (C) testosterone at 1100 h on days 1, 3, 5, 7 and 9 in adult male rats injected daily with 4 mg RU486 (s.c.) over 8 consecutive days starting on day 1. Control injections consisted of 0.2 ml oil. Values are means $\pm$ S.E.M. (12 rats per group). a= $P < 0.05$ and b= $P < 0.01$ vs oil-injected rats (Student's *t*-test).

periment (Fig. 1A). In contrast, LH (Fig. 1B) and testosterone (Fig. 1C) serum concentrations decreased in rats injected with RU486. Also, *in vitro* FSH secretion after *in vivo* administration of RU486 significantly increased for at least 4 h (Fig. 2A). Secretion of LH was significantly lower for the pituitaries of rats treated *in vivo* with RU486 at 120 and 240 min of incubation than for controls (Fig. 2B).

Effect of flutamide, ORCH or ADX on serum concentration of FSH and LH in rats injected with RU486

Treatment with RU486 over 4 consecutive days increased and decreased, respectively, the serum concentrations of FSH and LH in rats decapitated at 1100 h on day 5

Figure 2 *In vitro* basal FSH (A) and LH (B) secretion after 120 and 240 min of incubation on day 9 from pituitaries of male rats injected *in vivo* with 4 mg RU486 (s.c.) over 8 consecutive days. Values are means $\pm$ S.E.M. $a=P<0.01$ vs oil-injected rats (Student's *t*-test).

(Fig. 3). While this effect of RU486 was also evident in ADX rats, it was absent in flutamide-treated rats and in ORCH rats, in which serum concentrations of both gonadotrophins substantially increased (Fig. 3).

LHRH secretion in RU486-treated rats: effect of LHRH and LHRH-ANT on FSH, LH and testosterone concentrations in rats injected with RU486

Plasma concentration of LHRH in the pituitary stalk on day 5 after 4 consecutive days of RU486 treatment (11.8 ± 0.9 ng/l, $n=10$) did not differ by the Student's *t*-test from that of oil-injected rats (12.1 ± 1.7 ng/l, $n=10$).

Injection of LHRH increased the serum concentrations of LH at 1100 h on day 5 in both oil- and RU486-treated rats. This effect of LHRH was greater in controls (Fig. 4B). The injection of LHRH increased FSH serum concentrations in oil-injected rats. No statistically significant effects of LHRH were noted in FSH serum concentrations in RU486-treated rats (Fig. 4A). LHRH-ANT treatment decreased serum concentrations of FSH and LH in both oil- and RU486-treated rats (Fig. 4A and B). Testosterone serum concentrations paralleled those of LH (Fig. 4C).

Figure 3 Effects of antiandrogen flutamide (FLU), orchidectomy (ORCH) or adrenalectomy (ADX) on serum concentrations of (A) FSH and (B) LH at 1100 h on day 5 in male rats injected with 4 mg RU486/0.2 ml oil over 4 consecutive days starting on day 1. ORCH, ADX or sham-operations were performed on day 1. FLU treatment consisted of daily injections of 10 mg/kg over 4 consecutive days. Values are means $\pm$ S.E.M. (8–12 rats). $b=P<0.01$ vs oil-injected rats; $*=P<0.01$ vs the corresponding control group (flutamide-vehicle+sham-operated rats) (one-way ANOVA and Duncan's multiple range test).

Discussion

Secretion of FSH and LH in male rats depends on the eliciting action of the hypothalamic decapeptide gonadotrophin-releasing hormone and on steroidal and non-steroidal inhibitory gonadal feedback (reviewed by Fink 1988). Administration of the antisteroid RU486 to adult male rats increased FSH secretion but decreased LH secretion. This dissociation of gonadotrophin secretion in male rats contrasts with the effect witnessed in females. In female rats blockade of progesterone actions by administration of RU486 decreases the magnitude of the pre-ovulatory LHRH surge, and increases LHRH secretion during the period of low secretion rate of gonadotrophin, without affecting LHRH pituitary responsiveness (Sánchez-Criado *et al.* 1994). Administration of RU486 to male rats decreased LH secretion without altering LHRH secretory rate (present results). Because of this, the effects of RU486 on FSH and LH secretion in males should be the result of either the blockade of the feedback effect of peripheral glands at the pituitary level and/or a change of pituitary responsiveness to LHRH.

Figure 4 Effects of LHRH or LHRH antagonist (LHRH-ANT) on serum concentrations of (A) FSH, (B) LH and (C) testosterone at 1100 h on day 5 in oil- or RU486-treated rats. RU486 (4 mg) or oil (0.2 ml) was injected (s.c.) over 4 consecutive days. LHRH (100 ng/0.5 ml saline) was injected (i.p.) at 1000 h on day 5. LHRH-ANT (1 mg/0.2 ml saline) was injected (s.c.) at 1000 h on days 2 and 4. Values are means $\pm$ S.E.M. (10 rats). b= P <0.01 vs oil-treated rats; *= P <0.01 vs saline-injected rats (one-way ANOVA and Duncan's multiple range test).

Besides the antiprogesteragenic activity, the synthetic steroid RU486 also has strong antiglucocorticoid and moderate antiandrogenic activity (Philibert 1984). In male rats adrenal secretory products play a modulatory role in the secretion of FSH and LH (Lorenzen *et al.* 1980). However, we do not consider that the effects of RU486 on FSH and LH secretion could be due to the blockade of glucocorticoid actions due to the following reasons: (i) glucocorticoids inhibit LHRH-stimulated gonadotrophin secretion in males (Ringstrom & Schwartz 1985); (ii) the antiglucocorticoid action of RU486 in the rat, as measured

by increased serum corticosterone concentrations, lasts only 24 h (Sánchez-Criado *et al.* 1997); and (iii) the present results have shown that the removal of circulating adrenal steroids by ADX did not modify serum levels of FSH and LH or interfere with the effects of RU486 on FSH and LH.

The finding that flutamide increased both FSH and LH secretion to the same extent as ORCH indicates, in agreement with the results of Decker *et al.* (1981) and Summerville & Schwartz (1981), that testosterone, where there is no marked elevation of serum FSH over LH, accounts for almost all of the FSH- and LH-suppressing activity of the testes. Thus, the increased serum FSH concentrations found in RU486-treated male rats was, probably, the consequence of a partial reduction in the negative feedback of androgen, although this interpretation is controversial (Rea *et al.* 1986, Paul *et al.* 1990). Also, the results show that LHRH injection of RU486-treated rats did not increase further FSH secretion and that treatment with LHRH-ANT reduced FSH secretion. This suggests that the reduction in the inhibitory feedback of testosterone and/or its metabolites at the pituitary allowed endogenous LHRH to stimulate the release of FSH but not of LH. Indeed, in agreement with previous data (Gay & Midgley 1969, Kotsuji *et al.* 1988), pituitary LH secretion appeared more sensitive to the inhibitory effects of androgen than of FSH since flutamide treatment or ORCH increased serum LH to values about 10-fold while FSH levels were only double those of control males. The exact mechanism by which RU486 stimulates FSH secretion in male rats is not known. However, because RU486 antagonizes the FSH-modulatory activity of activin in female rats (Szabo *et al.* 1998), and testosterone and LHRH affect FSH secretion through pituitary FSH-modulating proteins in males (Kirk *et al.* 1994, Besecke *et al.* 1996, Bilezikjian *et al.* 1996), direct or indirect effects of RU486 on FSH secretion are possible.

The above reasoning, however, does not explain the inhibitory effect of RU486 on LH secretion. Results showed that RU486 reduces basal and LHRH-stimulated LH secretion and that decreased testosterone levels cannot be involved since androgens produce an inhibitory effect on LH secretion (Drouin & Labrie 1976, Summerville & Schwartz 1981). Thus, the reduction in androgen negative feedback would be expected to enhance LH secretion. Additionally, the results showed that the effect of RU486 disappeared when the strong inhibitory influences of testosterone on LH were removed by ORCH or flutamide treatment (Luderer & Schwartz 1991, present results).

In female rats ovariectomized on the morning of pro-oestrus, treatment with RU486 was found to attenuate the LH surge (Tébar *et al.* 1996). This, and other findings (Levine 1997), have been explained by observing that, even in the absence of progesterone as a ligand, progesterone receptor in the gonadotroph can be trans-activated by LHRH through a progesterone-independent

activation of progesterone receptor (Waring & Turgeon 1992, Turgeon & Waring 1994). Similarly, in the male rat, the inhibitory effect of RU486 on LH secretion could be due, in the virtual absence of circulating progesterone (Döhler & Wuttke 1975, Schwartz & Justo 1977), to a progesterone receptor-mediated blockade of LHRH-stimulated LH secretion. Indeed, experiments in progress in our laboratory using oestrogen-primed anterior pituitary cell cultures from male rats indicate that progesterone receptor antagonist RU486 (2 nM) significantly reduces LHRH (10 nM)-augmented LH secretion. Also, although in most *in vivo* systems RU486 does not display any agonistic action (Baulieu 1991, Knox *et al.* 1996), the possibility that the effects of RU486 on FSH and LH secretion could be ascribed to a progesterone (Horwitz 1985, Meyer *et al.* 1990) or a glucocorticoid (Grul & Altschmied 1993) agonistic action should not be discarded. Nevertheless, the finding that RU486 inhibits both endogenous and exogenous LHRH-stimulated LH secretion while enhancing endogenous LHRH-stimulated FSH secretion only, cannot be explained unless a differential regulation of gonadotrophin synthesis and secretion by LHRH, RU486 or androgen is considered. Accurate interpretation of the present results is not possible until more direct *in vitro* examination of the effects of RU486 on FSH and LH secretion in males is performed.

In summary, the results are consistent with the hypothesis that RU486 acting directly at the pituitary level decreases basal and LHRH-stimulated LH secretion. Also, despite the fact that the RU486-increased FSH secretion in male rats could be the result of an effect of RU486 at the pituitary through FSH-modulating proteins, the results suggest that increased FSH secretion may be the result of a partial reduction of the negative androgen feedback at the pituitary.

Acknowledgements

The authors wish to thank Dr Sitruk-Ware (Exelgyn, Paris, France) and Dr H Kloosterboer (Organon, Oss, The Netherlands) for the supply of the antiprogestrone RU486 and the LHRH antagonist Org.30276 respectively. The help of Mr Leonardo I Padrón with pituitary stalk blood collection is also acknowledged with gratitude. They are grateful to the National Hormone and Pituitary Program (Baltimore, MD, USA) for the LH and FSH RIA kits. This work has been subsidized by a grant (PM98-0167) from DGESEIC (Spain).

References

- Baulieu EE 1989 Contraception and other clinical applications of RU486, an antiprogestagen at the receptor. *Science* **345** 1351–1357.
- Baulieu EE 1991 The antisteroid RU486. Its cellular and molecular mode of action. *Trends in Endocrinology and Metabolism* **2** 233–239.
- Besecke LM, Guendner MJ, Schneyer AL, Bauer-Dantoin AC, Jameson JL & Weiss J 1996 Gonadotropin-releasing hormone regulates follicle-stimulating hormone- β gene expression through an activin/follistatin autocrine or paracrine loop. *Endocrinology* **137** 3667–3673.
- Bilezikjian LM, Corrigan AZ, Blount AL & Vale W 1996 Pituitary follistatin and inhibin subunit messenger ribonucleic acid levels are differentially regulated by local and hormonal factors. *Endocrinology* **137** 4277–4284.
- Chandolia RK, Weinbauer GF, Fingscheidt U, Bartlett JMS & Neischlang E 1991 Effects of flutamide on testicular involution induced by an antagonist of gonadotrophin-releasing hormone and on stimulation of spermatogenesis by follicle-stimulating hormone in rats. *Journal of Reproduction and Fertility* **93** 313–323.
- Decker MH, Loriaux DL & Cutler GB 1981 A seminiferous tubular factor is not obligatory for regulation of plasma follicle-stimulating hormone in the rat. *Endocrinology* **108** 1035–1039.
- Döhler KD & Wuttke W 1975 Changes with age in levels of serum gonadotropins, prolactin, and gonadal steroids in prepubertal male and female rats. *Endocrinology* **97** 898–907.
- Drouin J & Labrie F 1976 Selective effect of androgens on LH and FSH release in anterior pituitary cells in culture *Endocrinology* **98** 1528–1534.
- van den Dungen HM, van Dieten JAM, Tilders FJH, van Rens GP & Schoemaker J 1989 Administration of GnRH-antagonist to immature rats affects subsequent female and male pubertal development differently. *Acta Endocrinologica* **120** 778–784.
- Fink G 1988 Gonadotropin secretion and its control. In *The Physiology of Reproduction*, pp 1349–1377. Eds E Knobil & J Neill. New York: Raven Press.
- Gay VL & Midgley AR 1969 Response of the adult rat to orchidectomy and ovariectomy as determined by LH radioimmunoassay. *Endocrinology* **84** 1359–1364.
- Greenwood FC, Hunter WH & Glover JS 1963 The preparation of 125 I-labelled human growth hormone of high specific radioactivity. *Biochemical Journal* **89** 114–123.
- Grul DJ & Altschmied J 1993 Synergistic induction of apoptosis with glucocorticoids and 3',5'-cyclic adenosine monophosphate reveals agonist activity by RU486. *Molecular Endocrinology* **7** 104–113.
- Horwitz KB 1985 The antiprogestin RU38486: receptor-mediated progestin versus antiprogestin actions screened in estrogen-insensitive T47D_{co} human breast cancer cells. *Endocrinology* **116** 233–239.
- Kato J 1985 Progesterone receptors in brain and hypophysis. In *Actions of Progesterone on the Brain*, pp 31–81. Eds D Ganten & D Pfaff. Berlin: Springer-Verlag.
- Kirk SE, Dalkin AC, Yasin M, Haisenleder DJ & Marshall JC 1994 Gonadotropin-releasing hormone pulse frequency regulates expression of pituitary follistatin messenger ribonucleic acid: a mechanism for differential gonadotrope function. *Endocrinology* **135** 876–880.
- Knox LK & Schwartz NB 1992 RU486 blocks the secondary surge of follicle-stimulating hormone in the rat without blocking the drop in serum inhibin. *Biology of Reproduction* **46** 220–225.
- Knox LK, Ringstrom SJ & Schwartz NB 1993 RU486 blocks the effects of inhibin antiserum or luteinizing hormone on the secondary follicle-stimulating hormone surge. *Endocrinology* **133** 277–283.
- Knox KL, Ringstrom SJ, Szabo M, Perlyn CA, Sutandi S & Schwartz NB 1996 RU486 on an estrogen background blocks the rise in serum follicle-stimulating hormone induced by antiserum to inhibin or ovariectomy. *Endocrinology* **137** 1226–1232.
- Kotsuji F, Winters SJ, Attardi B, Keeping H, Oshima H & Troen Ph 1988 Effects of gonadal steroids on gonadotropin secretion in males: studies with perfused rat pituitary cells. *Endocrinology* **123** 2683–2689.
- Levine JE 1997 New concepts of the neuroendocrine regulation of gonadotropin surges in rats. *Biology of Reproduction* **56** 293–302.

- López FJ, Meade EH & Negro-Vilar A 1990 Development and characterization of a specific and sensitive radioimmunoassay for rat galanine: measurement in brain tissue, hypophyseal portal and peripheral serum. *Brain Research Bulletin* **124** 395–399.
- Lorenzen JR, Schlepphorst C & Schwartz NB 1980 The interaction of castration and adrenalectomy on pituitary responses to loss of target gland negative feedback in the male rat. *Endocrinology* **106** 592–599.
- Luderer U & Schwartz NB 1991 Sex differences in acute luteinizing hormone responses to gonadectomy remain after progesterone antagonist and dopamine agonist treatment. *Biology of Reproduction* **45** 918–926.
- Lustig RH, Pfaff DW & Fishman J 1988 Induction of LH hypersecretion in cyclic rats during the afternoon of oestrus by oestrogen in conjunction with progesterone antagonism or opioidergic blockade. *Journal of Endocrinology* **117** 229–235.
- Meyer ME, Pornon A, Ji J, Bocquel MT, Chambon P & Gronemeyer H 1990 Agonistic and antagonistic activities of RU486 on the functions of the human progesterone receptor. *EMBO Journal* **9** 3923–3932.
- Negro-Vilar A, Tesone M, Johnston CA, DePaolo L & Justo SN 1984 Sex differences in regulation of gonadotropin secretion: involvement of central monoaminergic and peptidergic systems and brain steroid receptors. In *Sexual Differentiation: Basic and Clinical Aspects*, pp 107–118. Eds M Serio, M Zanisi, M Motta & L Martini. New York: Raven Press.
- Paul SJ, Ortolano GA, Halsenleder DJ, Stewart JM, Shupnick MA & Marshall JC 1990 Gonadotropin subunit messenger RNA concentrations after blockade of gonadotropin-releasing hormone action: testosterone selectively increases follicle-stimulating hormone β -unit messenger RNA by posttranscriptional mechanism. *Molecular Endocrinology* **4** 1943–1955.
- Philibert D 1984 RU38486: an original multifaceted antihormone *in vivo*. In *Adrenal Steroid Antagonism*, pp 77–101. Ed MK Agarwal. Berlin: Walter de Gruyter & Co.
- Rao IM & Mahesh VB 1986 Role of progesterone in the modulation of the preovulatory surge of gonadotropins and ovulation in the pregnant mare's serum gonadotropin-primed immature rat and the adult rat. *Biology of Reproduction* **35** 1154–1161.
- Rea MA, Weinbauer GF, Marshall GR & Nieschlag E 1986 Testosterone stimulates pituitary and serum FSH in GnRH antagonist-suppressed rats. *Acta Endocrinologica* **116** 487–492.
- Ringstrom SJ & Schwartz NB 1985 Effects of cortisol or adrenalectomy on orchidectomy-induced gonadotropin response: differential LH and FSH response. *Endocrinology* **116** 472–474.
- Ringstrom SJ, Szabo M, Kilen SM, Saberi S, Knox KL & Schwartz NB 1997 The antiprogesterins RU486 and ZK98299 affect follicle-stimulating hormone secretion differentially on estrus, but not on proestrus. *Endocrinology* **138** 2286–2290.
- Rodriguez-Padilla M, Bellido C, Pinilla L & Aguilar E 1987 Secretion of LH in spontaneously hypertensive rats. *Journal of Endocrinology* **113** 255–260.
- Sánchez-Criado JE, Bellido C, Galiot F, López FJ & Gaytan F 1990 A possible dual mechanism of the anovulatory action of anti-progesterone RU486 in the rat. *Biology of Reproduction* **42** 877–886.
- Sánchez-Criado JE, Bellido C, López FJ & Galiot F 1992 Antiprogesterone RU486 induces dissociation of LH and FSH secretion in the cyclic rat: effect of anti-inhibin serum. *Journal of Endocrinology* **134** 43–49.
- Sánchez-Criado JE, Galiot F, Bellido C, González, D & Tebar M 1993 Hypothalamus–pituitary–ovarian axis in cyclic rats lacking progesterone actions. *Biology of Reproduction* **48** 916–925.
- Sánchez-Criado JE, Hernandez G, Bellido C, Gonzalez D, Tébar M, Diaz-Cruz MA & Alonso R 1994 Perioviatory LHRH, LH and FSH secretion in cyclic rats treated with RU486: effects of exogenous LHRH and LHRH antagonist on LH and FSH secretion at early oestrus. *Journal of Endocrinology* **141** 7–14.
- Sánchez-Criado JE, Tébar M, Ruiz A & Padrón L 1997 The steroid antagonist RU486 given at pro-oestrus induces hypersecretion of follicle-stimulating hormone from oestrus afternoon to early metoestrus in the rat. *European Journal of Endocrinology* **137** 281–286.
- Sarkar DK, Chiappa SA, Fink G & Sherwood NM 1976 Gonadotropin-releasing hormone surge in pro-oestrus rats. *Nature* **264** 461–463.
- Schwartz NB & Justo SN 1977 Acute changes in serum gonadotrophins and steroids following orchidectomy in the rat: role of adrenal gland. *Endocrinology* **100** 1550–1556.
- Summerville JW & Schwartz NB 1981 Suppression of serum gonadotropin levels by testosterone and porcine follicular fluid in castrate male rats. *Endocrinology* **109** 1442–1447.
- Szabo M, Knox KL, Ringstrom SJ, Perlyn CA, Sutandi S & Schwartz NB 1996 Mechanism of the inhibitory action of RU486 on the secondary follicle-stimulating hormone surge. *Endocrinology* **137** 85–89.
- Szabo M, Kilen SM, Saberi S, Ringstrom SJ & Schwartz NB 1998 Antiprogesterin suppress basal and activin-stimulated follicle-stimulating hormone secretion in an estrogen-dependent manner. *Endocrinology* **139** 2223–2228.
- Tébar M, Bellido C & Sánchez-Criado JE 1995 Luteinizing hormone (LH) and corticosterone in proestrus restore the follicle-stimulating hormone (FSH) secretion at early estrus in adrenalectomized-LH releasing hormone (LHRH)-antagonist treated rats. *Biology of Reproduction* **52** 63–67.
- Tébar M, Ruiz A & Sánchez-Criado JE 1996 Hypersecretion of follicle-stimulating hormone (FSH) on estrous afternoon in rats treated with RU486 in proestrus. *Cellular and Molecular Neurobiology* **16** 421–426.
- Turgeon JL & Waring DW 1994 Activation of the progesterone receptor by the gonadotropin-releasing hormone self-priming signaling pathway. *Molecular Endocrinology* **8** 860–869.
- Urbanski HF, Kim SO & Connolly ML 1990 Influence of photoperiod and 6-methoxy-benzoxazolinone on the reproductive axis of inbred LHS/Ss lak male hamster. *Journal of Reproduction and Fertility* **90** 157–163.
- Waring DW & Turgeon JL 1992 A pathway for luteinizing hormone releasing-hormone self-potential: cross-talk with the progesterone receptor. *Endocrinology* **130** 3275–3282.
- Worthington WC 1966 Blood samples from the pituitary stalk of the rat: method of collection and factors determining volume. *Nature* **210** 710–712.

Received 19 November 1997

Revised manuscript received 16 February 1998

Accepted 22 September 1998