PDX-1 and Msx-2 expression in the regenerating and developing pancreas

M R Kritzik¹, E Jones¹, Z Chen², M Krakowski¹, T Krah¹,
A Good¹, C Wright³, H Fox² and N Sarvetnick¹

¹Department of Immunology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, USA

²Department of Neuropharmacology, The Scripps Research Institute, La Jolla, California 92037, USA

³Vanderbilt University Medical School, Nashville, Tennessee 37232, USA

(Requests for offprints should be addressed to N Sarvetnick; Email: noras@scripps.edu)

(E Jones is now at School of Medicine, Department of Physiology, University of Maryland, Baltimore, Maryland 21201, USA)

Abstract

We have observed pancreatic duct cell proliferation and islet regeneration in transgenic mice whose pancreata produce interferon γ (IFN γ mice). We have previously demonstrated that new islet cells derive from endocrine progenitor cells in the pancreatic ducts of this model. The current study was initiated to define these endocrine progenitor cells further and to identify novel markers associated with pancreatic regeneration. Importantly, we have found that PDX-1, a transcription factor required for insulin gene transcription as well as for pancreatic development during embryogenesis, is expressed in the duct cells of IFN γ mice. This striking observation suggests an important role for PDX-1 in the marked regeneration

observed in IFN γ mice, paralleling its critical function during ontogeny. Also demonstrated was elevated expression of the homeobox-containing protein Msx-2 in the pancreata of fetal mice as well as in adult IFN γ mice, identifying this molecule as a novel marker associated with pancreatic development and regeneration as well. The identification of PDX-1 and Msx in the ducts of the IFN γ transgenic pancreas but not in the ducts of the non-transgenic pancreas suggests that these molecules are associated with endocrine precursor cells in the ducts of the IFN γ transgenic mouse.

Journal of Endocrinology (1999) **163**, 523–530

Introduction

Insulin-dependent, or type I, diabetes mellitus is characterized by the selective destruction of the insulin-synthesizing and insulin-secreting beta cells of the pancreatic islets of Langerhans, rendering type I diabetic patients dependent on exogenous insulin for life. Therefore, experimental models of pancreatic regeneration are vital for developing any treatment strategy aimed at alleviating the consequences of this disease. This rationale propels our continuing studies of the interferon- γ -producing (IFN γ) transgenic mouse. These mice, which express IFN γ in the beta cells of their pancreatic islets, undergo ductal hyperplasia and destruction of islets by infiltrating lymphocytes (Sarvetnick *et al.* 1988, Gu & Sarvetnick 1993). Remarkably, though, such mice undergo considerable islet regeneration. The newly formed islets bud into the lumen of ducts. The aggressive growth of pancreatic ducts and the continuous differentiation of new endocrine cells during adult life indicate that islet stem cells responsible for this extraordinary regrowth occupy the transgenic animals' pancreata. Here, we

attempted to define the stem cells responsible for this pancreatic regrowth, as well as to identify markers associated with the regenerative process.

Elegant and important experiments recently demonstrated that transcription factors important for insulin gene expression are critical for the development of the pancreas during embryogenesis (see Sander & German 1997 for review). PDX-1 (also called IDX-1, IPF-1, or STF-1), a transcription factor that regulates insulin expression, is one important marker (reviewed in Peshavaria & Stein 1997). PDX-1 is expressed in the pancreatic islets of adult mice but is absent from duct cells and acinar tissue (Ohlsson *et al.* 1993). The fact that PDX-1-deficient mice lack a pancreas and die shortly after birth indicates the absolute requirement for PDX-1 during pancreatic development (Jonsson *et al.* 1994, Offield *et al.* 1996). Additionally, PDX-1 functions in determination and/or maintenance of the pancreatic identity of common precursor cells (Peshavaria & Stein 1997).

Like PDX-1, Msx-2 is a homeobox-containing transcription factor. Msx-2 is part of a conserved family of transcription factors that play critical roles in tissue

patterning and organogenesis during development (Davidson & Hill 1991, Davidson 1995). For example, the involvement of *Msx-2* in bone and tooth development has been well described (Ignelzi *et al.* 1995, Maas & Bei 1997). Notably, *Msx-2* is expressed at a wide variety of sites in the developing embryo, suggesting its involvement in the generation of a number of organ systems (Davidson & Hill 1991, Davidson 1995). Nevertheless, no specific role for *Msx-2* in pancreatic development is known.

PDX-1 and *Msx* expression patterns were studied to define and characterize further the progenitor cells that give rise to islet cells in the IFN γ transgenic model of pancreatic regeneration. Whereas *Msx-2* was selected for study after it was found to be dramatically elevated in the IFN γ regenerating pancreas in a differential cDNA analysis (described below), PDX-1 was selected for study based on its critical and proven role in pancreatic development. Indeed, Sharma *et al.* (1999) recently published a report demonstrating the induction of PDX-1 in rat ducts following partial pancreatectomy. These authors further suggested that PDX-1 might be involved in the derivation of beta cells from ductal cells. Importantly, we now report significant expression of PDX-1 and *Msx* proteins in the ducts of the IFN γ transgenic pancreas.

Materials and Methods

Animal husbandry

Animals were maintained in a specific pathogen-free facility at The Scripps Research Institute according to the rules and regulations governed and enforced by the Institutional Animal Care and Use Committee. Animals were housed under a controlled 12-h light/darkness cycle with food and water available *ad libitum*. The embryos used in these studies did not carry the IFN γ transgene.

Transgenic mouse generation

Transgenic mice expressing IFN γ have been described previously (Sarvetnick *et al.* 1988). The IFN γ transgenic mice used in these studies were on the non-obese diabetic (NOD) background. IFN γ mice that have been backcrossed onto the NOD/Shi strain for more than ten generations have a very low incidence of diabetes, <20%, compared with NOD mice which have an incidence of ~80% for females and ~25% for males.

Immunohistochemistry

Pancreata from test mice were fixed overnight in 10% neutral buffered formalin (3.6% formaldehyde) and embedded in paraffin. Paraffin sections (5 μ m) were either conventionally stained with hematoxylin and eosin for histological evaluation or stained for the presence of

insulin, PDX-1, or *Msx* using immunocytochemical techniques. Briefly, sections were deparaffinized and blocked with 2% normal goat serum before applying the primary antibodies for insulin (DAKO, Carpinteria, CA, USA), *Msx* (BAbCO, Richmond, CA, USA), or PDX-1 (a generous gift from Dr Chris Wright, Vanderbilt University Medical School, Nashville, TN, USA and Dr Helena Edlund, University of Umeå, Umeå, Sweden). Binding of the primary antibody was detected using the appropriate secondary antibody (Vector Laboratories, Burlingame, CA, USA), and the horseradish peroxidase (HRP)-labeled avidin-biotin complex (ABC kit, Vector Laboratories). HRP was visualized using 3,3'-diaminobenzidine as a substrate. Gill's hematoxylin was used as a counterstain.

Immunoelectron microscopy

Pancreatic tissue was fixed in 10% normal buffered formalin (3.6% formaldehyde) for 2 h at 25 °C. Fixed tissue was infused in 1.5 M sucrose-PBS for 0.5 h with gentle inversion periodically. Infused tissue was then quick-frozen in liquid nitrogen, embedded in OCT and 2-methylbutane and sectioned 30–40 μ m thick. These sections were incubated in glycine-PBS to quench aldehyde for 0.5 h, blocked in 10% normal goat serum for 0.5 h, and incubated for 1 h each in PDX-1 (primary antibody) and an HRP-conjugated goat anti-rabbit secondary antibody before refixing in 1% glutaraldehyde-PBS for 0.25 h and washing in PBS. The reaction product was visualized with diaminobenzidine (DAB) for 7 min and DAB+H₂O₂ for 4 min before treating with 1% OsO₄. Tissue was dehydrated in graded ethanol, cleared in propylene oxide and embedded in Spurr resin. Thin sections were viewed with a Hitachi HU 12A electron microscope.

Differential gene expression analysis

The Atlas Mouse cDNA Expression Array I (Clontech, Palo Alto, CA, USA) was used to screen the IFN γ NOD-severe combined immunodeficiency (SCID)-regenerating pancreas for upregulation of mRNAs relative to the non-transgenic NOD-SCID pancreas. The analysis was carried out according to the manufacturer's recommendations. *Msx-2* was one of nineteen transcripts found to be expressed in the regenerating pancreas but not in the non-transgenic pancreas.

Results

PDX-1 and *Msx* in the fetal pancreas

We first sought to characterize PDX-1, *Msx-2*, and insulin expression during fetal pancreatic development for

Table 1 Summary of PDX-1, *Msx*, and insulin staining patterns in fetal, IFN γ transgenic, and non-transgenic pancreata

	PDX-1	<i>Msx</i>	Insulin
Fetal Balb/c			
Duct (cord)	+++	+	+/-
Peri-epithelial	+++	++	++
Acinar	++	+/-	-
Adult IFN γ transgenic			
Duct	+++	++	+
Acinar	-	-	-
Islet	+++	+	+++
Non-transgenic			
Duct	-	-	-
Acinar	-	-	-
Islet	+++	+	+++

Plus signs refer to the extent of staining, from low (+/-) to extensive (+++). Peri-epithelial staining refers to staining abutting the ductal cord region.

comparison with that during regeneration, with the results summarized in Table 1. After staining of pancreata from Balb/c embryos at 14.5 days post conception, PDX-1 reactivity was most notable in the cord region of expanding epithelial tissue from which the ducts and endocrine tissue develop, and was also observed in the acinar tissue, consistent with previous reports (Fig. 1A; Ohlsson *et al.* 1993, Guz *et al.* 1995). Nuclear staining was clearly observed in the expanding epithelia, as well as in the acinar tissue. Additionally, diffuse non-nuclear staining was apparent within the region of PDX-1 staining. Based upon a number of other studies which clearly demonstrate the nuclear localization of PDX-1, we believe that this non-nuclear staining is probably an artifact due to PDX-1 antigen leakage during fixation of the embryos. Importantly, although less extensive than PDX-1 staining, *Msx* displayed considerable staining in the expanding epithelia of the developing pancreas where PDX-1 was also

detected (Fig. 1B). Both nuclear and diffuse cytoplasmic staining were observed with this antibody as well, and faint nuclear staining in the acinar tissue was also detected (Fig. 1B and Fig. 2A). Finally, insulin-producing cells were also found in the region of expanding epithelia in the embryonic pancreas (Fig. 2B). Compared with PDX-1 and *Msx* staining, its expression was not as widespread within the cords, but rather was restricted more to the peri-epithelial region (Fig. 2 illustrates this point with a comparison of *Msx* and insulin staining patterns).

PDX-1 expression in pancreata of IFN γ transgenic mice

To determine if PDX-1 is present in the pancreas during regeneration as well as embryonic development, we screened regenerating pancreata of adult IFN γ mice for PDX-1 expression. As shown in Fig. 3A, we found striking expression of PDX-1 in the ducts of the IFN γ transgenic mouse. Specifically, we found nuclear localization of PDX-1 in islet cells (as expected), as well as in a proportion of cells within the duct wall. Staining was observed in both small and large ducts, as well as in ducts with and without intraductal islets. Staining of adjacent sections with antibody to insulin revealed that these PDX-1-positive ducts also contained a subpopulation of cells that were positive for insulin (Fig. 3B). PDX-1 expression has not previously been observed in murine pancreatic duct cells (Ohlsson *et al.* 1993) and, as expected, the ducts from non-transgenic control mice did not express PDX-1 or insulin (Fig. 3C and D). The expression of PDX-1 but not insulin in a subpopulation of ductal cells in IFN γ transgenic mice suggests that such cells represent pre-endocrine progenitor cells.

Ultrastructural analysis of PDX-1-expressing duct cells

We used immunoelectron microscopy to determine the ultrastructure of PDX-1-expressing cells in ducts of the

Figure 1 PDX-1 and *Msx* immunostaining of the fetal pancreas. Serial sections of an E14.5 embryo were stained with antibody to either PDX-1 (A) or *Msx* (B). Note the expression of both proteins, albeit at different levels, in the region of epithelial expansion. Original magnification $\times 20$.

Figure 2 Msx and insulin immunostaining of the fetal pancreas. Serial sections of an E14.5 embryo were stained with antibody to either Msx (A) or insulin (B). Note that Msx is expressed in both the ductal and peri-epithelial regions, while insulin is found mainly in the peri-epithelial region. Original magnification $\times 50$.

IFN γ transgenic mouse. Consistent with our observations that not all PDX-1-expressing cells also expressed insulin, we identified cells which expressed PDX-1 but which were devoid of endocrine granules (Fig. 4). Moreover, we

found no significant structural differences among duct cells, regardless of PDX-1 expression. In fact, ductal cells in the regenerating pancreas exhibited many of the morphological features characteristic of ductal epithelial cells,

Figure 3 PDX-1 immunostaining in the IFN γ transgenic pancreas. Pancreatic sections from an adult IFN γ transgenic mouse (A, B) and an adult non-transgenic mouse (C, D) were stained with antibody to either PDX-1 (A, C) or insulin (B, D). Panels A and B are serial sections. Note the presence of both insulin-positive and insulin-negative PDX-1-expressing ductal cells. Original magnification $\times 50$.

Figure 4 Ultrastructural analysis of PDX-1-expressing ductal cells in the IFNg transgenic pancreas. Immunoelectron microscopy was used to characterize PDX-1-expressing ductal cells in the pancreas of an adult IFNg transgenic mouse. The figure shows a PDX-1-positive cell (left) adjacent to a PDX-1-negative cell (right). Note the similar morphology of these cells. Original magnification $\times 8500$.

including centrally located, irregularly shaped nuclei, numerous interdigitating processes on the lateral aspects of the cells, unspecialized cytoplasm containing mitochondria, sparse elements of rough endoplasmic reticulum, scattered stacks of Golgi lamellae, and a variety of vesicular profiles. The apical plasma membrane possessed typical apical ultrastructural specializations including microvilli and junctional complexes complete with tight junctions (Yamamoto & Kataoka 1988).

Msx expression in IFNg mice

We next screened sections of regenerating pancreata from adult IFNg mice for the presence of the *Msx* proteins by staining with a polyclonal antibody directed against the *Msx* homeodomain. The only antibody available for this purpose recognizes both the *Msx-1* and *Msx-2* proteins and probably *Msx-3* as well. Our analyses demonstrated diffuse *Msx* staining throughout the islets of all mice assessed. In contrast, acinar tissue did not stain with the *Msx* antibody. Importantly, a significant proportion of duct cells in the regenerating pancreas were positive for *Msx* expression (Fig. 5B and C). We observed staining in the nucleus, as expected for members of this transcription factor family; however, cytoplasmic staining was also observed in many instances. The extent of staining varied among ducts; some ducts had many positive cells and other ducts had few, if any, *Msx*-positive cells. No *Msx* expression was evident in the ducts of non-transgenic control mice (data not shown). Furthermore, *Msx* staining was less extensive than PDX-1 staining in the transgenic ducts, and comparison of serial section staining patterns suggested that *Msx*-expressing cells also expressed PDX-1 (Fig. 5A

and B). Subsequent analyses further suggested coincident staining between insulin and *Msx* in many instances. Although the similarity in *Msx* and insulin expression patterns could reflect a basal level of cross-reactivity between the *Msx* antibodies and endocrine cells (supported by the diffuse islet staining we observe with the *Msx* antibody), we identified *Msx*-positive cells that did not stain for insulin in a number of ducts; insulin-positive cells lacking *Msx* expression were also observed (Fig. 5C and D). Thus, based on PDX-1, *Msx*, and insulin staining patterns, we suggest that at least four populations of ductal cells, not present in the non-transgenic pancreas, can be identified in the IFNg transgenic pancreas: PDX-1⁺ *Msx*⁻ insulin⁻ ductal cells, PDX-1⁺ *Msx*⁺ insulin⁻ ductal cells, PDX-1⁺ *Msx*⁺ insulin⁺ ductal cells, and PDX-1⁺ *Msx*⁻ insulin⁺ ductal cells.

Discussion

We have observed the regeneration of islets that bud into the ductal lumen in the IFNg transgenic mouse, where they are protected from destruction by infiltrating lymphocytes. In this study, we used histological analyses to characterize the progenitor cells responsible for the remarkable ductal proliferation and islet regeneration. Indeed, we have now defined markers associated with ducts during regeneration in the IFNg transgenic mouse: PDX-1 and *Msx-2*. These results suggest that these proteins are associated with endocrine progenitor cells in the ducts of the IFNg transgenic mouse.

The ductal epithelium, which consists of intercalated (small), intralobular (medium) and interlobular (large)

Figure 5 *Msx* immunostaining in the IFN γ transgenic pancreas. Serial sections of the pancreas from an adult IFN γ transgenic mouse were stained with antibody to either PDX-1 (A), *Msx* (B, C), or insulin (D). Note the significant expression of *Msx* as well as PDX-1 in panels A and B (the arrows in panel B highlight *Msx*-positive cells in the duct wall). *Msx*-positive cells lacking insulin expression are indicated by arrows in panel C; insulin-positive cells lacking *Msx* expression are indicated by arrows in panel D. Panels C and D represent serial sections somewhat offset and at slightly different magnifications. Original magnification A, B, D $\times 40$; C $\times 50$.

ducts, has been designated as the site of exocrine and endocrine development in the pancreas (Pictet *et al.* 1972, Argent *et al.* 1992, Slack 1995). A specific pathway for islet development in these animals, defined by the expression of duct, exocrine, and endocrine specific antigens, is thought to involve the derivation of endocrine cells from duct cells, which progress through a series of intermediate cell types (Gu *et al.* 1994). In promoting regeneration and new islet formation, we believe that progenitor stem cells in the ducts of the IFN γ transgenic mouse recapitulate the early development of the pancreas. Indeed, parallels exist between normal ontogeny and IFN γ -mediated regeneration, and in both cases, endocrine gene expression is an early event. Individual endocrine cells are initially scattered in the duct wall; these cells subsequently migrate to form clusters, which develop into fully differentiated islets. Thus, we hypothesized that endocrine cell precursors would be abundant in the ducts of mice undergoing islet regeneration, as they are in the fetal pancreas.

PDX-1 is thought to be expressed in the earliest pancreatic progenitor cell, from which both exocrine and endocrine cells develop (reviewed in Peshavaria & Stein 1997). Indeed, whereas PDX-1 is present only in the endocrine cells of the adult, PDX-1-deficient mice have no pancreas at all, indicating the critical role PDX-1 plays in the development of all pancreatic tissue (Jonsson *et al.* 1994, Offield *et al.* 1996). Using light and immunoelectron microscopy, we have now found that ductal cells in the transgenic pancreas also express PDX-1. In addition, only a subset of the PDX-1-expressing cells also express insulin. These observations are striking, because they define distinct populations of endocrine precursor cells during pancreatic regeneration and new islet growth. Indeed, we believe that duct cells expressing PDX-1 but not insulin represent early pre-endocrine progenitor cells.

We have also observed substantial expression of PDX-1 in the developing pancreas during ontogeny, as expected given the critical role of this molecule in pancreatic

development and consistent with reports by others (Ohlsson *et al.* 1993, Guz *et al.* 1995). Although its precise functions during ontogeny and regeneration are not clear, the expression of PDX-1 in the IFN γ regenerating pancreas appears to recapitulate the requirement for this protein in pancreatic development during ontogeny. Indeed, the ductal derivation of new endocrine cells coupled with PDX-1 expression in the ducts of the IFN γ transgenic mouse suggests that the new endocrine cell formation we observe during regeneration in the IFN γ transgenic mouse proceeds through mechanisms similar to those active during fetal development.

Recent studies in rats have found weak expression of PDX-1 in the pancreatic ducts of normal rats (Sharma *et al.* 1999). While PDX-1 expression is not detected in the ducts of normal mice, the expression of PDX-1 in normal rats is weak, and the differences between ductal expression patterns might be attributed to subtle differences between the species. Interestingly, Sharma *et al.* (1999) have further demonstrated that PDX-1 expression is significantly induced in ductal cells during the regeneration that ensues following partial pancreatectomy. Based on the kinetics of PDX-1 expression, these authors suggested that PDX-1 might be involved in the derivation of new endocrine cells from ductal cells. These studies, therefore, parallel and support our own observations, which suggest a role for PDX-1 in the extensive new endocrine cell formation exhibited by the IFN γ transgenic mouse. The ductal expression of PDX-1 in two distinct models of pancreatic regeneration and new islet formation is striking and suggests that PDX-1 serves a central and critical function in the differentiation of endocrine cells, during both ontogeny and regeneration.

Msx-2 is part of a conserved family of homeobox-containing transcription factors that regulate tissue growth and patterning during embryogenesis (reviewed in Davidson 1995). Using the Atlas cDNA array to study differential gene expression patterns, we found that *Msx-2* is expressed in the regenerating pancreas of the IFN γ transgenic mouse. We used immunohistochemistry to confirm these results and to follow expression of the *Msx* protein directly. These experiments utilized an antibody which detects both *Msx-1* and *Msx-2*, preventing us from drawing conclusions regarding *Msx-2* expression specifically from this staining alone. Despite this limitation, we detected significant expression of *Msx* in the pancreatic ducts of IFN γ transgenic mice. This is in contrast to the work of others as well as our own observations in non-transgenic mice demonstrating that the *Msx* proteins are not expressed in the normal adult mouse pancreas (Maas *et al.* 1996). Strikingly, we also observed significant expression of *Msx* in the developing pancreas during embryogenesis. This expression was localized to the growing epithelia from which ducts and endocrine cells arise. Coupled with the fact that we did not observe enhanced *Msx-1* expression in the transgenic pancreas using the

Atlas cDNA array, these results support our identification of *Msx-2* as a marker associated with endocrine progenitor cells both in the developing and regenerating pancreas. Furthermore, our staining demonstrated considerable cytoplasmic as well as nuclear localization of *Msx*. As a transcription factor, *Msx* is expected to be localized in the nucleus, although others have reported its presence as a diffuse cytoplasmic stain as well (Stelnicki *et al.* 1997). Although this cytoplasmic staining might represent an artifact due to the fixation procedures, it is possible that *Msx* might be localized to the cytoplasm under certain conditions, as is the case for NF κ B prior to its activation (Baeuerle & Baltimore 1988a,b).

A number of studies indicate that *Msx-2* can be induced through a signaling network involving members of the transforming growth factor (TGF) beta superfamily, including BMP-4 (Vainio *et al.* 1993, Marazzi *et al.* 1997). Although we have not yet explored regulatory aspects of *Msx* induction in the regenerating pancreas, possibly the cytokine network induced by IFN γ in this model contributes directly to the induction of *Msx* protein. Indeed, previous work in our laboratory has demonstrated significant changes in the cytokine profile of the regenerating pancreas. For example, TGF α , interleukin 1 β , and tumor necrosis factor α are elevated in the regenerating pancreas (Arnush *et al.* 1996 and M Arnush & N Sarvetnick, unpublished data). At this time, the functional significance of *Msx-2* expression in pancreatic regeneration is not clear. *Msx-2* is expressed at many sites during development, including the cranial neural crest, neural tube, tooth germs, eyes, ears, nose, limb buds, pituitary, and heart, and, in particular, *Msx-2* expression appears to be associated with sites of epithelial-mesenchymal interactions (reviewed in Davidson & Hill 1991, Davidson 1995). *Msx* proteins are also thought to be crucial to pattern formation during the development of diverse organs (reviewed in Davidson & Hill 1991, Davidson 1995). A number of reports have also implicated *Msx-2* in the apoptotic program during development (Graham *et al.* 1993, 1994, Marazzi *et al.* 1997, Winograd *et al.* 1997, Ferrari *et al.* 1998). Although the expression of *Msx-2* in the pancreas during development has not been reported previously, the evidence presented here suggests that *Msx-2* might play a critical role in regulating the pancreatic developmental program as well.

The identification of markers associated with endocrine progenitor cells in the IFN γ transgenic pancreas is clearly of value, with regards to both defining these precursor cells and in an analysis of the regenerative process. In this study we have correlated the expression of two such molecules, PDX-1 and *Msx-2*, with the striking pancreatic regeneration exhibited by the IFN γ transgenic mouse. Each of these homeodomain proteins appears to play a critical role in organ formation during ontogeny, and each is expressed in the developing as well as the regenerating pancreas. While future studies will be aimed at defining the precise

contributions of these proteins during pancreatic development and regeneration, their association with pancreatic progenitor cells will be valuable in the isolation and characterization of this critical cell type.

Acknowledgements

We would like to thank Dr Cheng-ming Chang for his valuable contribution to the electron microscopy studies. We would also like to thank Dr Helena Edlund for her generous gift of PDX-1 antibody, and Phyllis Minick for editorial assistance.

References

- Argent BE, Githens S, Kalser S, Longnecker DS, Metzgar R & Williams JA 1992 The pancreatic duct cell. *Pancreas* **7** 403-419.
- Arnush M, Gu D, Baugh C, Sawyer SP, Mroczkowski B, Krahl T & Sarvetnick N 1996 Growth factors in the regenerating pancreas of g-interferon transgenic mice. *Laboratory Investigation* **74** 985-990.
- Baeuerle PA & Baltimore D 1988a I κ B: a specific inhibitor of the NF- κ B transcription factor. *Science* **242** 540-546.
- Baeuerle PA & Baltimore D 1988b Activation of DNA-binding activity in an apparently cytoplasmic precursor of the NF- κ B transcription factor. *Cell* **53** 211-217.
- Davidson D 1995 The function and evolution of Msx genes: pointers and paradoxes. *Trends in Genetics* **11** 405-411.
- Davidson DR & Hill RE 1991 Msh-like genes: a family of homeobox genes with wide-ranging expression during vertebrate development. *Seminars in Developmental Biology* **2** 405-412.
- Ferrari D, Lichter AC, Pan ZZ, Dealy CN, Upholt WB & Kosher RA 1998 Ectopic expression of Msx-2 in posterior limb bud mesoderm impairs limb morphogenesis while inducing BMP-4 expression, inhibiting cell proliferation, and promoting apoptosis. *Developmental Biology* **197** 12-24.
- Graham A, Heyman I & Lumsden A 1993 Even-numbered rhombomeres control the apoptotic elimination of neural crest cells from odd-numbered rhombomeres in the chick hindbrain. *Development* **119** 233-245.
- Graham A, Francis-West P, Brickell P & Lumsden A 1994 The signalling molecule BMP4 mediated apoptosis in the rhombencephalic neural crest. *Nature* **372** 684-686.
- Gu D & Sarvetnick N 1993 Epithelial cell proliferation and islet neogenesis in IFN- γ transgenic mice. *Development* **118** 33-46.
- Gu D, Lee MS, Krahl T & Sarvetnick N 1994 Transitional cells in the regenerating pancreas. *Development* **120** 1873-1881.
- Guz Y, Montminy MR, Stein R, Leonard J, Gamer LW, Wright CV & Teitelman G 1995 Expression of murine STF-1, a putative insulin gene transcription factor, in beta cells of pancreas, duodenal epithelium, and pancreatic exocrine and endocrine progenitors during ontogeny. *Development* **121** 11-18.
- Ignelzi Jr MA, Liu Y-H, Maxson Jr RE & Snead ML 1995 Genetically engineered mice: tools to understand craniofacial development. *Critical Reviews in Oral Biology and Medicine* **6** 181-201.
- Jonsson J, Carlsson L, Edlund T & Edlund H 1994 Insulin-promoter factor 1 is required for pancreas development in mice. *Nature* **371** 606-609.
- Maas R & Bei M 1997 The genetic control of early tooth development. *Critical Reviews in Oral Biology and Medicine* **8** 4-39.
- Maas R, Chen YP, Bei M, Woo I & Satokata I 1996 The role of Msx genes in mammalian development. *Annals of the New York Academy of Sciences* **785** 171-181.
- Marazzi G, Wang Y & Sassoon D 1997 Msx2 is a transcriptional regulator in the BMP4-mediated programmed cell death pathway. *Developmental Biology* **186** 127-138.
- Offield MF, Jetton TL, Labosky PA, Ray M, Stein RW, Magnuson MA, Hogan BL & Wright CV 1996 PDX-1 is required for pancreatic outgrowth and differentiation of the rostral duodenum. *Development* **122** 983-995.
- Ohlsson H, Karlsson K & Edlund T 1993 IPF-1, a homeodomain-containing transactivator of the insulin gene. *EMBO Journal* **12** 4251-4259.
- Peshavaria M & Stein R 1997 PDX-1: an activator of genes involved in pancreatic development and islet gene expression. In *Pancreatic Growth and Regeneration*, edn1, pp 96-107. Ed N Sarvetnick. Basel, Switzerland: Karger Landes Systems.
- Pictet RL, Clark WR, Williams RH & Rutter WJ 1972 An ultrastructural analysis of the developing embryonic pancreas. *Developmental Biology* **29** 436-467.
- Sander M & German MS 1997 The β cell transcription factors and development of the pancreas. *Journal of Molecular Medicine* **75** 327-340.
- Sarvetnick N, Liggitt D, Pitts SL, Hansen SE & Stewart TA 1988 Insulin-dependent diabetes mellitus induced in transgenic mice by ectopic expression of class II MHC and interferon- γ . *Cell* **52** 773-782.
- Sharma A, Zangen DH, Reitz P, Taneja M, Lissauer ME, Miller CP, Weir GC, Habener JF & Bonner-Weir S 1999 The homeodomain protein IDX-1 increases after an early burst of proliferation during pancreatic regeneration. *Diabetes* **48** 507-513.
- Slack J 1995 Developmental biology of the pancreas. *Development* **121** 1569-1580.
- Stelnicki EJ, Komuves LG, Holmes D, Clavin W, Harrison MR, Adzick NS & Largman C 1997 The human homeobox genes MSX-1, MSX-2, and MOX-1 are differentially expressed in the dermis and epidermis in fetal and adult skin. *Differentiation* **62** 33-41.
- Vainio S, Karavanova I, Jowett A & Thesleff I 1993 Identification of BMP-4 as a signal mediating secondary induction between epithelial and mesenchymal tissues during early tooth development. *Cell* **75** 45-58.
- Winograd J, Reilly MP, Roe R, Lutz J, Laughner E, Xu X, Hu L, Asakura T, vander Koli C, Strandburg JD & Semenza GL 1997 Perinatal lethality and multiple craniofacial malformations in MSX2 transgenic mice. *Human Molecular Genetics* **6** 369-379.
- Yamamoto M & Kataoka K 1988 An electron microscope study of the development of the exocrine and endocrine pancreas with special reference to intercellular junctions. *Archives of Histology and Cytology* **51** 315-325.

Received 14 April 1999

Accepted 13 July 1999