Differential regulation of 3 β -hydroxysteroid dehydrogenase type II and 17 α -hydroxylase/lyase P450 in human adrenocortical carcinoma cells by epidermal growth factor and basic fibroblast growth factor

J Doi, H Takemori, M Ohta¹, Y Nonaka² and M Okamoto

Department of Molecular Physiological Chemistry, Osaka University Medical School, Suita, Osaka, Japan

¹Laboratory of Nutrition, Koshien College, Nishinomiya, Hyogo, Japan

²College of Nutrition, Koshien University, Takarazuka, Hyogo, Japan

(Requests for offprints should be addressed to M Okamoto, Department of Molecular Physiological Chemistry, Osaka University Medical School, 2-2 Yamadaoka, Suita, Osaka, 565-0871, Japan; Email: mokamoto@mr-mbio.med.osaka-u.ac.jp)

Abstract

Epidermal growth factor (EGF) and basic fibroblast growth factor (bFGF) are pluripotent growth factors that stimulate both the proliferation and steroidogenesis of adrenocortical cells. Here we demonstrate that EGF and bFGF specifically induce mRNA of 3 β -hydroxysteroid dehydrogenase type II (3 β HSD II) and suppress that of 17 α -hydroxylase/lyase P450 (CYP17) in human adrenocortical H295R cells. The induction of 3 β HSD II mRNA did not occur until 6 h after the growth factor treatment and was completely abolished in the presence of a protein synthesis inhibitor, cycloheximide (CHX), suggesting that the induction required *de novo* protein synthesis. The CYP17 mRNA suppression began at almost the same time as the induction of the 3 β HSD II mRNA. Interestingly, the CYP17 mRNA level was increased by the CHX treat-

ment. Both the 3 β HSD II and CYP17 mRNAs were repressed by treatment with a calmodulin kinase II (CaMK II) inhibitor, KN-93, and were enhanced by a mitogen-activated protein kinase (MAPK) inhibitor, PD98059. The PD98059-mediated induction of the 3 β HSD II mRNA was completely blocked by the CHX treatment. Interestingly, treatment with EGF in the presence of both PD98059 and CHX produced a greater increase in the CYP17 mRNA than did treatment in the presence of PD98059 alone. These results suggest that CHX-sensitive factor(s) and CaMK II- and MAPK-signaling pathways may have important roles in both induction of 3 β HSD II and suppression of CYP17 by EGF or bFGF in H295R cells.

Journal of Endocrinology (2001) **168**, 87–94

Introduction

Numerous investigators have reported that epidermal growth factor (EGF) and basic fibroblast growth factor (bFGF) influence steroidogenesis in adrenocortical cells (Gospodaeowicz & Handley 1975, 1986, McAllister & Hornsby 1987a, Fisher & Lakshmanan 1990, Mesiano *et al.* 1993, Mesiano & Jaffe 1997). Singh and coworkers reported that EGF stimulated cortisol secretion from cultured bovine (Singh & Waters 1983) and sheep (Singh *et al.* 1985) adrenal cortical cells. The stimulation was abolished when inhibitors of cholesterol biosynthesis, compactin and AY9944, were added. Further studies showed that the activity of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase and the rate of cholesterol synthesis were increased in the EGF-treated cells. In contrast, the rates of conversion of radioactive pregnenolone to steroid intermediates were little influenced by the growth factor. On the basis of these results, they

concluded that EGF activated the biosynthesis of the steroid hormone precursor, rather than the steroidogenic enzymes themselves.

bFGF has been suggested as a physiological regulator of adrenocortical functions both *in vivo* (Mesiano *et al.* 1991, Basile & Holzwarth 1994) and *in vitro* (Crickard *et al.* 1981, Hornsby *et al.* 1983, Hotta & Baird 1986, Ho & Vinson 1995). It was purified from extracts of bovine adrenals, and a large amount was detected in the zona fasciculata and the medulla (Basile & Holzwarth 1993). It was also detected immunochemically and by northern blot analysis in bovine (Gospodarowicz *et al.* 1986), rat (Basile & Holzwarth 1993) and human (Mesiano *et al.* 1991) adrenal cortex. Basile & Holzwarth (1993) observed that, in a unilaterally adrenalectomized rat, the remaining adrenal gland underwent compensatory hypertrophy, and they suggested that bFGF might be involved in this phenomenon. Recently, Thomas *et al.* (1997) showed that transplantation of bovine adrenocortical cells with

bFGF-overproducing mouse fibroblasts into immunodeficient mice resulted in both cell proliferation and activation of steroidogenesis of the transplants. Receptors for bFGF were also shown to exist in the rat zona glomerulosa (Basile & Holzwarth 1994). These previous reports strongly suggest that EGF and bFGF act as paracrine effectors in the adrenal cortex and may modulate steroidogenesis in adrenocortical cells.

H295R cells were originally isolated from human adrenal carcinoma cells (Gazdar *et al.* 1990, Rainey *et al.* 1994). The cells were reported to possess five steroidogenic P450s – namely, side-chain cleavage P450 (CYP11A), 17 α -hydroxylase/lyase P450 (CYP17), 21-hydroxylase P450 (CYP21), 11 β -hydroxylase P450 (CYP11B1) and aldosterone synthase P450 (CYP11B2) – and 3 β -hydroxysteroid dehydrogenase type II (3 β HSD II) (Bird *et al.* 1996). Because of this, these cells have often been used for studying the molecular mechanism underlying human adrenocortical steroidogenesis. In the present study, we investigated the effects of EGF and bFGF on gene expression of steroidogenic enzymes in H295R cells.

Materials and Methods

Cell culture and treatment with EGF and bFGF

H295R cells were generous gifts from J Ian Mason at University of Edinburgh, UK. The cells were maintained in Dulbecco's Modified Eagle's/Ham's F-12 medium (DMEM/F-12; Gibco BRL, Grand Island, NY, USA) containing 1% ITS plus (insulin/transferrin/selenium/linoleic acid; Becton Dickinson Labware, Lincoln Park, NJ, USA), 2% Ultrosor G (BioSeptra Inc., France) and antibiotics, at 37 °C under an atmosphere of 5% CO₂–95% air, as described before (Bird *et al.* 1996). The cells (5 × 10⁵) were cultured in 100 mm dishes for 48 h and then incubated for further 24 h in serum-free medium (DMEM/F-12 containing antibiotics and 0.01% BSA). To initiate the experiments, the medium was exchanged for one containing EGF (Austral Biochemicals, San Ramon, CA, USA) or bFGF (Austral Biochemicals) at the final concentration of 10 ng/ml. After the 24 h incubation, the cells were washed with PBS, and the total RNA was extracted. Protein kinase inhibitors (bisindolylmaleimide I, H-8, H-89, PD98059 and KN-93) and a protein synthesis inhibitor (cycloheximide; CHX) were purchased from Calbiochem Ltd (La Jolla, CA, USA) and Sigma, respectively, dissolved in dimethylsulfoxide, and stored at –20 °C until required for use. Cells that had been maintained in the serum-free medium for 24 h, were first incubated with or without bisindolylmaleimide I, H-8, H-89, PD98059 or KN-93 for 1 h, and then stimulated with EGF or bFGF dissolved in the freshly prepared inhibitor-containing medium.

RNA extraction

Isolation of the total RNA was performed according to the method of Chomczynski & Sacchi (1987) with some modifications (Takemori *et al.* 1997). Briefly, the cells were washed with 1 ml PBS, and then dissolved in 1.6 ml guanidine isothiocyanate solution (Trizol reagent; Gibco BRL). The cell lysates were transferred into microcentrifuge tubes and mixed with 0.4 ml chloroform. After the centrifugation, an aqueous phase was taken, to which ethanol was added to precipitate total RNAs. The RNAs were washed with 70% ethanol, dried and dissolved in 200 μ l distilled water. DNA contaminant in the RNA preparation was digested by 1 U DNase I (Gibco BRL) using the buffer supplied by the manufacturer. The RNAs were again extracted by phenol–chloroform (1:1) and reprecipitated by ethanol.

Northern blot analysis

To prepare a cDNA probe for 3 β HSD II, a DNA fragment of the exon 3 was amplified by PCR using primers shown in Table 1 and ligated into the *Sma*I site of pUC18. cDNA fragments of CYP11A and CYP21 were also prepared, using RT-PCR with sets of primers listed in Table 1, and introduced into pT7R (Novagen, Madison, WI, USA). A *Nde*I–*Hind*III cDNA fragment of CYP17 was excised from pCW (Katagiri *et al.* 1995) that had been kindly provided by M Katagiri at Osaka Kyoiku University, Japan. We utilized a rat glyceraldehyde-3-phosphate dehydrogenase (G3PDH) cDNA fragment (Halder *et al.* 1998) as the control probe. Total RNAs (5 μ g) were separated by electrophoresis in 1% agarose gel in the presence of 6% formaldehyde and then transferred onto nylon membrane (Hybond-N+; Amersham, Arlington Heights, IL, USA) by capillary action in 20 × SSPE (0.2 M phosphate buffer, pH 7.7, containing 3.6 M NaCl and 0.02 M EDTA). The RNA on the membrane was hybridized with [³²P]-labeled DNA fragments in 5 × SSPE containing 50% formamide, 0.5% SDS, 0.1% polyvinylpyrrolidone, 0.1% Ficoll and 0.1% BSA at 42 °C for 16 h. After the hybridization, the membrane was washed sequentially in 2 × SSPE/0.1% SDS, 1 × SSPE/0.1% SDS, and 0.2 × SSPE/0.1% SDS at 60 °C, and then exposed to X-ray film at –80 °C with an intensifying screen. The radioactivity in hybridized signals was quantified with a phosphorimager (BAS2000; Fuji-film, Tokyo, Japan).

Determination of mRNAs for CYP11B1 and CYP11B2 by RT-PCR followed by Southern blot analysis

Total RNAs (2 μ g) were reverse-transcribed by 200 U reverse transcriptase (Super Script II; Gibco BRL) in 20 μ l reaction mixture containing 150 ng random hexamer. An aliquot (5 μ l) of the reaction mixture was used for PCR

Table 1 Sequence of oligonucleotides used in the experiments

Enzyme	Forward primers (5'→3')	Reverse primers (5'→3')
3βHSD type II	GCA CAT GGA TCT GTG CAT GTG GTT GCA G†	GAC CTG GGC TTG TGC CCC TGT TGC C†
3βHSD type I-specific	TGG TCC GCC TGT TGG TGG AA‡	CTA CCT CTA TGC TAC TGG TGT AG‡
3βHSD type II-specific	TCA TCC GCC TCT TGG TGA AG‡	(common to 3βHSD types I and II)
CYP11A	TTG CCT TTG AGT CCA TCA CT†	GAG CAG GAC TTG GGA CAG AC†
CYP21	CTC AGC TGC CTT CAT CAG TTC†	CAC CCC TTG GAG CAT GTA GT†
Common to CYP11B and CYP11B2	CAA ATG TGG CGT GTT CTT GT‡	AGT TGC TGG CTT CTA TGG‡
CYP11B1-specific		CCC AAC GCT GTG CA§
CYP11B2-specific		CCC AAG GCC GTG CA§

†Primers to generate cDNA probes for northern blot analysis.

‡Primers for RT-PCR analysis.

§Oligonucleotide probes for Southern blot analysis after RT-PCR.

amplification with 20 pmol primer sets (common primer) shown in Table 1 in 30 cycles of 94 °C for 30 s, 58 °C for 30 s and 72 °C for 30 s. The amplified cDNA fragments were separated in 1.5% agarose gel and transferred onto a nylon membrane. To distinguish the cDNA of CYP11B1 from that of CYP11B2, the membrane was subjected to Southern blot analyses using [³²P]-labeled oligonucleotide probes specific to the respective P450s (Table 1). The membrane was hybridized in 5 × SSPE containing 0.5% SDS, 0.1% polyvinylpyrrolidone, 0.1% Ficoll and 0.1% BSA at 40 °C for 16 h, and washed three times in 0.2 × SSPE containing 0.1% SDS at room temperature for 20 min.

Results

Effect of EGF and bFGF on expression of steroidogenic mRNAs in H295R cells

Northern blot analyses were performed on the growth-factor-treated H295R cells to examine the level of mRNAs of 3βHSD, CYP17, CYP11A and CYP21. To distinguish between CYP11B1 and CYP11B2 mRNAs, both were first amplified by RT-PCR using primers common to the two mRNAs and then the respective cDNAs were detected by Southern blot analysis using specific oligonucleotide probes. As shown in Fig. 1A and C, the level of mRNA of 3βHSD markedly increased in the EGF- or bFGF-treated cells, whereas that of CYP17 was decreased substantially by the EGF and bFGF treatment. The maximal stimulation of 3βHSD expression and the strongest suppression of CYP17 were found with a concentration of 10 ng/ml of either growth factor (data not shown). In contrast, the mRNA levels of the other steroidogenic P450s – CYP11A, CYP21, CYP11B1 and CYP11B2 – showed little change with the growth factor treatment (Fig. 1A, B and C). The RT-PCR analyses using the primers specific to either the type I or the type

II isoform of 3βHSD revealed that only the type II cDNA was clearly amplified from the total RNAs extracted from H295R cells (Fig. 1D).

Time courses of 3βHSD and CYP17 mRNA expression

Time courses of the EGF- and bFGF-mediated induction of 3βHSD mRNA and suppression of CYP17 mRNA were examined (Fig. 2). A distinct increase in 3βHSD mRNA was seen after the 6 h incubation with EGF, its level becoming maximal after 12 h and gradually diminishing thereafter. When bFGF instead of EGF was used as the stimulator, the mRNA level continued to increase after 12 h. In contrast, EGF or bFGF decreased the level of CYP17 mRNA, which reached the lowest point after 24 h. There was no significant change in the cell number under these growth factor treatments.

Effect of various protein kinase inhibitors on EGF- and bFGF-mediated gene expression

Numerous investigators have reported that protein kinases are involved in the signal transduction system of 3βHSD and CYP17 gene expression (McAllister & Hornsby 1987b, Brentano *et al.* 1990, Chris *et al.* 1990, Bird *et al.* 1996). Therefore, several protein inhibitors were tested to explore the signaling pathway occurring in the EGF- and bFGF-treated H295R cells. As shown in Fig. 3, protein kinase A inhibitors, H-8 (15 μM) and H-89 (20 μM), or a protein kinase C (PKC) inhibitor, bisindolylmaleimide I (10 μM), seemed to have no effect on the levels of 3βHSD and CYP17 mRNAs in both EGF- and bFGF-treated cells. A mitogen-activated protein kinase (MAPK) inhibitor, PD98059 (20 μM), in contrast, further increased the growth factor-dependently increased 3βHSD mRNA level. PD98059 appeared to restore the growth factor-mediated suppression of CYP17 mRNA, but had little effect on the basal levels of expression of both genes (data not shown).

Next, the inhibitor of Ca^{2+} /calmodulin-dependent kinase II (CaMK II), KN-93, was tested. KN-93 (5 μM) markedly decreased the level of 3βHSD mRNA in the EGF- and bFGF-treated cells. It exerted little effect on the basal mRNA expression in the non-treated cells (data not

Figure 2 Time course of 3βHSD and CYP17 mRNA expression stimulated by growth factors. Total RNAs (10 μg) were isolated from H295R cells at the indicated times after the addition of EGF (10 ng/ml: A, C) and bFGF (10 ng/ml: B, D). The levels of mRNA of 3βHSD (A, B) and CYP17 (C, D) were determined by a phosphorimager (BAS 2000) after northern blot analysis, and then normalized by those of G3PDH. Each point represents a mean value of duplicate experiments.

Figure 1 Effect of EGF and bFGF on mRNA expression of steroidogenic enzymes. (A) Total RNAs (10 μg) were prepared from H295R cells that had been incubated without or with EGF (10 ng/ml) and bFGF (10 ng/ml) for 24 h. mRNAs of 3βHSD, CYP17, CYP11A and CYP21 were subjected to northern blot analyses. mRNA of G3PDH was visualized as the internal standard. (B) The RNAs were reverse-transcribed and the cDNA fragments amplified by PCR using primers hybridizable with both CYP11B1 and CYP11B2 cDNAs. The PCR products were then subjected to Southern blot analyses using oligonucleotide probe specific to the respective enzymes. Plasmid DNAs inserted with CYP11B1 or CYP11B2 cDNA fragments (two lanes on the left) were used for the positive controls of the specific hybridization. Each blot (A and B) is representative of three independent experiments. (C) The levels of mRNAs of 3βHSD, CYP17, CYP11A and CYP21 (in A) and CYP11B1 and CYP11B2 (in B) were determined by a phosphorimager (BAS 2000) after northern blot analysis and Southern blot analysis coupled with RT-PCR, respectively. Each mRNA level was first normalized to the level of G3PDH mRNA and then expressed as the fold change (means $\pm$ S.E.M.; $n=3$). (D) The cDNA fragments were amplified with 3βHSD type I- or type II-specific primers. The DNA fragments were separated on an agarose gel, and visualized by ethidium bromide staining. Total RNAs of placenta were used as the positive control for 3βHSD type I and type II mRNAs. cont, Control; M, molecular markers.

Figure 3 Effect of various protein kinase inhibitors on the EGF and bFGF treatments. (A) H295R cells were pretreated with or without protein kinase inhibitors (bisindolylmaleimide I (10 μ M), PD98059 (20 μ M), H-8 (15 μ M), H-89 (20 μ M) and KN-93 (5 μ M)) for 1 h, and then stimulated with EGF (10 ng/ml) or bFGF (10 ng/ml) for 24 h. Total RNAs were isolated and subjected to northern blot analysis as described in Materials and Methods. The figure is representative of three independent experiments. cont, control.

shown). Another CaMK II inhibitor, KN-62 (10 μ M), or a calmodulin antagonist, W-7 (30 μ M), also inhibited the 3 β HSD expression data (not shown). It should be noted that KN-93 further decreased the growth-factor-dependent repressed mRNA level of CYP17.

Effect of CHX on EGF- and bFGF-mediated gene expression

Because the induction of 3 β HSD and suppression of CYP17 appeared to be late events of growth factor treatment, we examined whether *de novo* protein biosynthesis was required for this phenomenon. An inhibitor of protein synthesis, CHX (10 μ g/ml), was added to the culture medium together with EGF and bFGF. As shown in Fig. 4A, CHX completely suppressed the EGF- and bFGF-mediated induction of 3 β HSD mRNA, suggesting that *de novo* protein synthesis is required for the induction of 3 β HSD mRNA. The CHX treatment also completely abolished the PD98059-mediated enhancement of 3 β HSD mRNA in EGF-stimulated H295R cells (Fig. 4B). To our surprise, CHX seemed to prevent the EGF- and bFGF-mediated repression of CYP17 mRNA. The treatment with EGF in the presence of both PD98059 and CHX produced a more increased level of CYP17 mRNA than did treatment in the presence of PD98059 alone (Fig. 4B). Similar results were obtained by treatment with the two inhibitors in the bFGF-stimulated cells (data not shown).

Discussion

Previous studies of bovine (Singh & Waters 1983) and sheep (Singh *et al.* 1985) adrenal cortical cells demonstrated that the addition of EGF to culture media resulted in stimulated cortisol secretion from and cholesterol synthesis in the cells, without significant effect on the production rate of steroid intermediates. Abolition of the EGF-mediated stimulation of cortisol secretion by inhibitors of cholesterol synthesis led to the idea that HMG-CoA

Figure 4 Effect of CHX on the EGF and bFGF treatment. (A) H295R cells were stimulated with EGF (10 ng/ml) or bFGF (10 ng/ml) in the presence or absence of CHX (10 μ g/ml) for 24 h. Total RNAs were isolated and subjected to northern blot analysis as described in Materials and Methods. The panel is representative of three independent experiments. (B) H295R cells were treated with PD98059 (20 μ M), CHX (10 μ g/ml) or both PD98059 (20 μ M) and CHX (10 μ g/ml) in the presence of EGF (10 ng/ml). The panel is a representative of duplicated experiments. Cont, control.

reductase, the rate-limiting enzyme of cholesterol synthesis, may be the major target of EGF action in the cells. The presence of EGF receptor has been reported in human normal, tumor (Sasano *et al.* 1994), and fetal (Smikle *et al.* 1996) adrenocortical cells. Coulter *et al.* (1996b) reported that the administration of EGF increased the volume of the definitive zone of fetal adrenals of rhesus monkeys in late gestation. This zone-specific increase seemed to be the result of cellular hypertrophy, not of cell proliferation. It was also found that EGF treatment induced 3β HSD II protein in the definitive and transitional zones without significant effect on CYP17. Similar induction of 3β HSD II occurred in fetal rhesus monkeys in which endogenous adrenocorticotrophic hormone (ACTH) secretion was stimulated by administration of an inhibitor of CYP11B1, metyrapone (Coulter *et al.* 1996a). Coulter and colleagues were unable to determine whether the effect on adrenocortical function that was produced by the EGF administered to the animals had occurred as the result of the activation of the hypothalamic–pituitary–adrenal axis, as described by Luger *et al.* (1988) and Polk *et al.* (1987), or as the result of the direct action of EGF on adrenocortical cells. Our present study with H295R cells, however, demonstrated that treatment of the cells with EGF or bFGF stimulated transcription of the 3β HSD II gene in the absence of ACTH or cAMP, the second messenger of ACTH.

Interestingly, the EGF or bFGF treatments resulted in a decrease in CYP17 mRNA levels in H295R cells. Welsh & Hsueh (1982) also reported that gonadotropin-stimulated production of testosterone, androstenedione and 17α -hydroxyprogesterone in cultured testicular cells was inhibited by EGF. Because EGF also inhibited the conversion of exogenous 17α -hydroxyprogesterone to androstenedione, they proposed that the EGF effect occurred via inhibition of 17α -hydroxylase and $17,20$ -lyase. These results suggested that EGF and bFGF could directly modulate the gene expression of steroidogenic enzymes in steroidogenic cells.

Hornsby *et al.* (1983) reported that bFGF acted as a mitogen in human fetal adrenal cortex, and that the mitogenic effect of bFGF was greater in the definitive zone than in the fetal zone. The mRNA of bFGF in cultured human adrenocortical cells was increased by about threefold by the addition of ACTH. These results suggest that bFGF may be one of the autocrine factors of adrenocortical cells, in which secretion is regulated by ACTH. Li *et al.* (1998) reported that bFGF treatment increased the level of CYP17 mRNA in cultured bovine fasciculata cells. Our results, in contrast, showed that both bFGF and EGF decreased the level of CYP17 mRNA in H295R cells. This discrepancy may have resulted from the species difference or from the difference in the zonal origin of the cells.

Membrane-associated receptor tyrosine kinases such as EGF/TGF α receptor and bFGF receptor are believed to

stimulate several protein-kinase-mediated signaling pathways, such as those involving protein kinase C (PKC) and MAPK (Kim & Muller 1999, Rowan *et al.* 2000). Several investigators (Leers *et al.* 1997, Bird *et al.* 1998) have demonstrated that a PKC agonist, phorbol-12-myristate-13-acetate (PMA), increased 3β HSD II mRNA expression. MAPK was immunochemically detected in the rat zona glomerulosa, and MAPK seemed to be involved in gene expression of 3β HSD II, and thus activate the production of aldosterone (McNeil *et al.* 1998). These findings indicate that PKC- and MAPK-mediated signaling pathways may be involved in regulating 3β HSD II gene expression, suggesting the possibility that the effect of EGF and bFGF on 3β HSD II gene expression described here could be mediated via these signaling systems. The PKC inhibitor, bisindolylmaleimide I, however, did not block the EGF- or bFGF-mediated induction of 3β HSD II mRNA in a concentration of $10\ \mu\text{M}$ – a concentration sufficient to block the PMA-mediated induction of 3β HSD II mRNA (data not shown). Moreover, the level of 3β HSD II mRNA was further increased, not inhibited, by the addition of the MAPK inhibitor PD98059 in the presence of EGF or bFGF. These results suggest that a signaling system mediated by a member(s) other than PKC or MAPK may be important for the EGF- and bFGF-stimulated induction of 3β HSD II mRNA. A possible candidate signal would be CaMK II, because KN-93, an inhibitor of CaMK II, markedly inhibited the EGF- and bFGF-stimulated induction of 3β HSD II mRNA. However, this inhibitor also inhibited the expression of CYP17 mRNA. This indicates that the CaMK II-mediated pathway may be essential for the gene expression of 3β HSD II, but this signaling system could not account for the repressive effects of EGF and bFGF on the expression of the CYP17 gene.

It took several hours for EGF or bFGF to increase the 3β HSD II mRNA level in H295R cells (Fig. 2). As in the PKC-mediated induction of 3β HSD II mRNA (Leers *et al.* 1997), CHX blocked the effect of EGF and bFGF on 3β HSD II mRNA expression (Fig. 4), indicating that *de novo* protein synthesis is required for the EGF- and bFGF-mediated 3β HSD II gene expression. In contrast, EGF or bFGF decreased the level of CYP17 mRNA in H295R cells (Fig. 1). Interestingly also, the EGF- or bFGF-mediated suppression of CYP17 mRNA was abolished by CHX treatment (Fig. 4). These results may suggest the presence of a CHX-sensitive protein factor(s) that is induced in H295R cells by the growth factor treatment, and which stimulates the expression of the 3β HSD II gene but inhibits that of the CYP17 gene.

Treatment of the cells with EGF in the presence of both CHX and PD98059, however, revealed the complicated involvement of the MAPK signaling pathway in the expression of these genes. The incubation with both CHX and PD98059 resulted in further increase in the level of CYP17 mRNA in EGF-stimulated H295R cells, whereas

the presence of PD98059 seemed not to influence the blocking effect of CHX on β HSD II gene expression (Fig. 4B). These results indicate that more than one factor – one sensitive to the CHX-treatment and the other related to the MAPK signaling pathway – is involved in the regulation of β HSD II and CYP17 gene expression. In the EGF-mediated activation of the β HSD II gene, the CHX-sensitive factor may act more strongly than that involved in the MAPK pathway, because treatment with both CHX and PD98059 completely abolished expression of the gene. In contrast, in the EGF-mediated suppression of the CYP17 gene, the CHX-sensitive factor and the factor involved in the MAPK signaling pathway seemed to repress the gene independently. To date, we have failed in our attempts to search the database of the promoter regions of the β HSD II and CYP17 genes for sequences typical of *cis* elements of transcription factors that may act downstream of either the MAPK or CaMK II-mediated signaling pathway.

Further investigation is in progress in our laboratory to elucidate the precise mechanism underlying the enhancement of β HSD II mRNA and suppression of CYP17 mRNA by EGF and bFGF.

Acknowledgements

We thank Dr J Ian Mason at Edinburgh University for the generous gift of H295R cells and Dr Masanao Katagiri at Osaka Kyoiku University for pCW plasmid. We also thank Dr Hiromasa Tojo at Osaka University Medical School, Drs Michael R Waterman and Norio Kagawa at Vanderbilt University School of Medicine and Dr William E Rainey at University of Texas Southwestern Medical Center for their helpful discussion. A part of this work was supported by Health Sciences Research Grants from the Ministry of Health and Welfare of Japan.

References

- Basile DP & Holzwarth MA 1993 Basic fibroblast growth factor may mediate proliferation in the compensatory adrenal growth response. *American Journal of Physiology* **265** R1253–R1261.
- Basile DP & Holzwarth MA 1994 Basic fibroblast growth factor receptor in the rat adrenal cortex: effects of suramin and unilateral adrenalectomy on receptor numbers. *Endocrinology* **134** 2482–2489.
- Bird IM, Imaishi K, Pasquarette MM, Rainey WE & Mason JI 1996 Regulation of 3 beta-hydroxysteroid dehydrogenase expression in human adrenocortical H295R cells. *Journal of Endocrinology* **150** 8165–8173.
- Bird IM, Mason JI & Rainey WE 1998 Protein kinase A, protein kinase C, and Ca(2+)-regulated expression of 21-hydroxylase cytochrome P450 in H295R human adrenocortical cells. *Journal of Clinical Endocrinology and Metabolism* **83** 1592–1597.
- Brentano ST, Picado-Leonard J, Mellon SH, Moore CC & Miller WI 1990 Tissue-specific, cyclic adenosine 3',5'-monophosphate-induced, and phorbol ester-repressed transcription from the human P450c17 promoter in mouse cells. *Molecular Endocrinology* **4** 1972–1979.
- Chomczynski P & Sacchi N 1987 Single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction. *Analytical Biochemistry* **162** 156–159.
- Chris CD, Moore S, Brentano T & Miller WL 1990 Human P450 ccc gene transcription is induced by cyclic AMP and repressed by 12-O-tetradecanoylphorbol-13-acetate and A23187 through independent *cis* element. *Molecular and Cellular Biology* **10** 6013–6023.
- Coulter CL, Goldsmith PC, Mesiano S, Voytek CC, Martin MC, Mason JI & Jaffe RB 1996a Functional maturation of the primate fetal adrenal *in vivo*. II. Ontogeny of corticosteroid synthesis is dependent upon specific zonal expression of 3 beta-hydroxysteroid dehydrogenase/isomerase. *Endocrinology* **137** 4953–4959.
- Coulter CL, Read LC, Carr BR, Tarantal AF, Barry S & Styne DM 1996b A role for epidermal growth factor in the morphological and functional maturation of the adrenal gland in the fetal rhesus monkey *in vivo*. *Journal of Clinical Endocrinology and Metabolism* **81** 1254–1260.
- Crickard K, III CR & Jaffe RB 1981 Control of proliferation of human fetal adrenal cells *in vitro*. *Journal of Clinical Endocrinology and Metabolism* **53** 790–796.
- Fisher DA & Lakshmanan J 1990 Metabolism and effects of epidermal growth factor and related EGF and bFGF in mammals. *Endocrine Review* **11** 418–442.
- Gazdar AF, Oie HK, Shackleton CH, Chen TR, Triche TJ, Myers CE, Chrousos GP, Brennan MF, Stein CA & La RR 1990 Establishment and characterization of a human adrenocortical carcinoma cell line that expresses multiple pathways of steroid biosynthesis. *Cancer Research* **50** 5488–5496.
- Gospodarowicz D & Handley HH 1975 Stimulation of division of Y1 adrenal cells by a growth factor isolated from bovine pituitary glands. *Endocrinology* **97** 102–107.
- Gospodarowicz D, Baird A, Cheng J, Lui GM, Esch F & Bohlen P 1986 Isolation of fibroblast growth factor from bovine adrenal gland: physicochemical and biological characterization. *Endocrinology* **118** 82–90.
- Halder SK, Takemori H, Hatano O, Nonaka Y, Wada A & Okamoto M 1998 Cloning of a membrane-spanning protein with epidermal growth factor-like repeat motifs from adrenal glomerulosa cells. *Endocrinology* **139** 3316–3328.
- Ho MM & Vinson GP 1995 Endocrine control of the distribution of basic fibroblast growth factor, insulin-like growth factor-I and transforming growth factor-beta 1 mRNAs in adult rat adrenals using non-radioactive *in situ* hybridization. *Journal of Endocrinology* **144** 379–387.
- Hornsby PJ, Sturek M, Harris SE & Simonian MH 1983 Serum and growth factor requirements for proliferation of human adrenocortical cells in culture: comparison with bovine adrenocortical cells. *In Vitro* **19** 863–869.
- Hotta M & Baird A 1986 Differential effects of transforming growth factor type beta on the growth and function of adrenocortical cells *in vitro*. *PNAS* **83** 7795–7799.
- Katagiri M, Kagawa N & Waterman MR 1995 The role of cytochrome b5 in the biosynthesis of androgens by human P450c17. *Archives of Biochemistry and Biophysics* **317** 343–347.
- Kim H & Muller WJ 1999 The role of the epidermal growth factor receptor family in mammary tumorigenesis and metastasis. *Experimental Cell Research* **253** 78–87.
- Leers SS, Morohashi K, Mason JI & Melner MH 1997 Synergistic activation of the human type II 3 beta-hydroxysteroid dehydrogenase/delta5-delta4 isomerase promoter by the transcription factor steroidogenic factor-1/adrenal 4-binding protein and phorbol ester. *Journal of Biological Chemistry* **272** 7960–7967.
- Li X, Lee JW, Graves LM & Earp HS 1998 Angiotensin II stimulates ERK via two pathways in epithelial cells: protein kinase C suppresses a G-protein coupled receptor–EGF receptor transactivation pathway. *EMBO Journal* **17** 2574–2583.

- Luger A, Calogero AE, Kalogeras K, Gallucci WT, Gold PW, Loriaux DL & Chrousos GP 1988 Interaction of epidermal growth factor with the hypothalamic–pituitary–adrenal axis: potential physiologic relevance. *Journal of Clinical Endocrinology and Metabolism* **66** 334–337.
- McAllister JM & Hornsby PJ 1987a Improved clonal and nonclonal growth of human, rat and bovine adrenocortical cells in culture. *In Vitro Cellular and Developmental Biology* **23** 677–685.
- McAllister JM & Hornsby PJ 1987b TPA inhibits the synthesis of androgens and cortisol and enhances the synthesis non-17 α -hydroxylated steroids in cultured human adrenocortical cells. *Endocrinology* **121** 1908–1910.
- McNeil H, Puddefoot JR & Vinson GP 1998 MAP kinase in the rat adrenal gland. *Endocrine Research* **24** 373–380.
- Mesiano S & Jaffe RB 1997 Developmental and functional biology of the primate fetal adrenal cortex. *Endocrine Reviews* **18** 378–403.
- Mesiano S, Mellon SH, Gospodarowicz D, DiBlasio AM & Jaffe RB 1991 Basic fibroblast growth factor expression is regulated by corticotropin in the human fetal adrenal: a model for adrenal growth regulation. *PNAS* **88** 5428–5432.
- Mesiano S, Mellon SH & Jaffe RB 1993 Mitogenic action, regulation, and localization of insulin-like EGF and bFGF in the human fetal adrenal gland. *Journal of Clinical Endocrinology and Metabolism* **76** 968–976.
- Polk DH, Ervin MG, Padbury JF, Lam RW, Reviczky AL & Fisher DA 1987 Epidermal growth factor acts as a corticotropin-releasing factor in chronically catheterized fetal lambs. *Journal of Clinical Investigation* **79** 984–988.
- Rainey WE, Bird IM & Mason JI 1994 The NCI-H295 cell line: a pluripotent model for human adrenocortical studies. *Molecular and Cellular Endocrinology* **100** 45–50.
- Rowan BG, Weigel NL & O'Malley BW 2000 Phosphorylation of steroid receptor coactivator-1, identification of phosphorylation site and phosphorylation through the mitogen-activated protein kinase pathway. *Journal of Biological Chemistry* **275** 4475–4483.
- Sasano H, Suzuki T, Shizawa S, Kato K & Nagura H 1994 Transforming growth factor α , epidermal growth factor, and epidermal growth factor receptor expression in normal and diseased human adrenal cortex by immunohistochemistry and *in situ* hybridization. *Modern Pathology* **7** 741–746.
- Singh AP & Waters MJ 1983 Stimulation of adrenal cortisol biosynthesis by epidermal growth factor. *Molecular and Cellular Endocrinology* **30** 189–199.
- Singh AP, Waters MJ & Wilce PA 1985 A mechanism for the *in vitro* stimulation of adrenal cortisol biosynthesis by epidermal growth factor. *International Journal of Biochemistry* **17** 857–862.
- Smikle CB, Kim HS, Mesiano S & Jaffe RB 1996 Identification of the ligands for the epidermal growth factor receptor in human fetal and adult adrenal glands. *Proceedings of the 10th International Congress of Endocrinology*, San Francisco, CA 1996. Abstract OR6–3.
- Takemori H, Halder SK, Nonaka Y, Fujii T, Ohta M, Hatano O & Okamoto M 1997 Polyadenylation-mediated translational regulation of maternal P450(11 β) mRNA in frog oocytes. *European Journal of Biochemistry* **250** 197–204.
- Thomas M, Northrup SR & Hornsby PJ 1997 Adrenocortical tissue formed by transplantation of normal clones of bovine adrenocortical cells in SCID mice replaces the essential functions of the animals' adrenal glands (see Comments). *Nature Medicine* **3** 978–983.
- Welsh TH Jr & Hsueh AJ 1982 Mechanism of the inhibitory action of epidermal growth factor on testicular androgen biosynthesis *in vitro*. *Endocrinology* **110** 1498–506.

Received 3 May 2000

Revised manuscript received 3 August 2000

Accepted 31 August 2000