EUROSTERONE MEETING

Peroxisome proliferator-activated receptors in inflammation control

P Delerive, J-C Fruchart¹ and B Staels¹

U325 INSERM, Département d'Athérosclérose, Institut Pasteur de Lille, 1 rue Professeur Calmette, 59019 Lille, France

¹Faculté de Pharmacie, Université de Lille II, 59000 Lille, France

(Requests for offprints should be addressed to B Staels; Email: Bart.Staels@pasteur-lille.fr)

Abstract

Peroxisome proliferator-activated receptors (PPARs) are ligand-activated transcription factors belonging to the nuclear receptor superfamily. PPAR α is highly expressed in liver, skeletal muscle, kidney, heart and the vascular wall. PPAR γ is predominantly detected in adipose tissue, intestine and macrophages. PPARs are activated by fatty-acid derivatives and pharmacological agents such as fibrates and glitazones which are specific for PPAR α and PPAR γ respectively. PPARs regulate lipid and lipoprotein metabolism, glucose homeostasis, cell proliferation and differentiation, and apoptosis. PPAR α controls intra- and extracellular lipid metabolisms whereas PPAR γ triggers adipocyte differentiation and promotes lipid storage. In addition, PPARs also modulate the inflammatory re-

sponse. PPAR activators have been shown to exert anti-inflammatory activities in various cell types by inhibiting the expression of proinflammatory genes such as cytokines, metalloproteases and acute-phase proteins. PPARs negatively regulate the transcription of inflammatory response genes by antagonizing the AP-1, nuclear factor- κ B (NF- κ B), signal transducer and activator of transcription and nuclear factor of activated T-cells signalling pathways and by stimulating the catabolism of proinflammatory eicosanoids. These recent findings indicate a modulatory role for PPARs in inflammation with potential therapeutical applications in chronic inflammatory diseases.

Journal of Endocrinology (2001) **169**, 453–459

Introduction

Peroxisome proliferator-activated receptors (PPARs) belong to the nuclear receptor superfamily which are ligand-activated transcription factors (Issemann & Green 1990). To date, three different PPAR subtypes have been identified: PPAR α , PPAR β (NUC-1 or PPAR δ) and PPAR γ . PPARs are activated by natural ligands such as fatty acids, eicosanoids and oxidized fatty acids (Kliewer *et al.* 1995, Devchand *et al.* 1996, Forman *et al.* 1997, Nagy *et al.* 1998, Delerive *et al.* 2000a). Furthermore, the lipid-lowering fibrates and the anti-diabetic glitazones are synthetic ligands for PPAR α and PPAR γ respectively (Lehmann *et al.* 1995, Forman *et al.* 1997). PPARs regulate gene expression by binding with retinoid X receptor (RXR) as a heterodimeric partner to specific DNA sequence elements termed PPAR response elements (PPRE) (Fig. 1) (Tugwood *et al.* 1992). PPREs consist of a direct repeat of the nuclear receptor hexameric AGGTCA recognition sequence separated by one or two nucleotides (DR-1 and DR-2) (Ijpenberg *et al.* 1997, Gervois *et al.* 1999). PPARs have been reported to be involved in lipid and lipoprotein metabolism, glucose

homeostasis, cell proliferation and differentiation, and apoptosis (for review see Desvergne & Wahli 1999). It has recently been demonstrated that PPARs may also play a role in the control of the inflammatory response. In this review, we will focus on the new insights indicating the implication of PPAR α and PPAR γ in inflammation control and discuss our current understanding of the molecular mechanisms by which they regulate the expression of inflammatory response genes.

PPAR α in inflammation: from bench to bedside

The first evidence indicating a potential role for PPARs in the inflammatory response was the demonstration that leukotriene B₄ (LTB₄), a proinflammatory eicosanoid, binds to PPAR α and induces the transcription of genes involved in ω - and β -oxydation which leads to the induction of its own catabolism (Devchand *et al.* 1996). Using the mouse ear-swelling test, these authors showed that the duration of the inflammatory response is prolonged in PPAR α -deficient mice in response to LTB₄ (Devchand *et al.* 1996). Several recent studies have been

Figure 1 Mechanism of transcriptional activation by PPARs. p/CAF, p300/cyclic AMP-responsive element binding protein-associated factor.

aimed at delineating the cellular and molecular mechanisms explaining the control of the inflammatory response by PPAR α . In primary aortic smooth muscle cells which express substantial amounts of PPAR α , it was demonstrated that PPAR α ligands inhibit interleukin (IL)-1 β -induced IL-6 secretion as well as 6-keto-prostaglandin (PG) F $_{1\alpha}$ production. In addition, PPAR α agonists have been reported to decrease cytokine-induced genes, such as expression of vascular cell adhesion molecule-1 and tissue factor in endothelial cells and monocytes respectively (Marx *et al.* 1999, Neve *et al.* 2001). Subsequently, it was shown that PPAR α acts by down-regulating the transcription of these genes (Staels *et al.* 1998, Delerive *et al.* 1999a). *In vivo* evidence for an anti-inflammatory action of PPAR α in the vascular wall came with the demonstration that aortas from PPAR α -deficient mice displayed an exacerbated inflammatory response to lipopolysaccharide stimulation (Delerive *et al.* 1999a). Furthermore, fibrates did not affect LPS-induced IL-6 transcription in PPAR α -deficient mice, demonstrating that the anti-inflammatory activities of these agonists require PPAR α expression *in vivo*. In addition, Poynter & Daynes (1998) reported that PPAR α -deficient splenocytes produced, in response to lipopolysaccharide (LPS) stimulation, two to three times more IL-6 and IL-12 than splenocytes from wild-type mice. Finally, fibrates were shown to repress the expression of a number of acute-phase proteins in liver, such as fibrinogen, in a PPAR α -dependent manner (Kockx *et al.* 1999). Taken together, these observations provide evidence that PPAR α plays a role in the inflammatory response at the vascular, splenic and hepatic level.

Studies addressing the molecular mechanisms of this anti-inflammatory action demonstrated that PPAR α negatively interferes with the inflammatory response by antagonizing the nuclear factor- κ B (NF- κ B) signalling pathway (Poynter & Daynes 1998, Staels *et al.* 1998, Delerive *et al.* 1999a, Marx *et al.* 1999). In fact, a

bidirectional antagonism between the PPAR α and NF- κ B signalling pathways exists (Delerive *et al.* 1999a). PPAR α overexpression inhibits NF- κ B-driven gene transcription and co-transfection of increasing amounts of p65 led to a dose-dependent inhibition of a PPAR response element (PPRE)-driven promoter construct. Glutathion-S-transferase (GST) pull-down assays revealed that PPAR α physically interacts with p65 via its Rel homology domain which mediates homo- and heterodimerization and interaction with inhibitor of NF κ B (I κ B) (Delerive *et al.* 1999a). Since PPAR α -mediated inhibition of NF- κ B-driven gene transcription becomes more and more important upon longer exposure to PPAR α ligands, we speculated that a complementary mechanism might exist. NF- κ B activity is tightly controlled by the degradation of I κ B α which sequesters inactive NF- κ B dimers in the cytoplasm. Interestingly, PPAR α activators were found to induce I κ B α mRNA and protein expression in primary smooth muscle cells and hepatocytes (Delerive *et al.* 2000b). I κ B α induction by fibrates again requires PPAR α expression. Surprisingly, I κ B α induction did not affect p65 nuclear translocation but was associated with reduced NF- κ B DNA-binding activity (Delerive *et al.* 2000b). Western blot analysis revealed that I κ B α protein induction occurs mainly in the nucleus which may provide an explanation for the reduced NF- κ B-binding activity (Delerive *et al.* 2000b). The induction of I κ B α by fibrates in cytokine-activated cells should therefore result in an acceleration of NF- κ B nuclear deactivation. In line with this observation, the increase of I κ B α protein after treatment with PPAR α activators would lead to a halt in p65-mediated gene activation, thereby reducing the duration of the inflammatory response. This is consistent with a previous report in which PPAR α ligands were shown to affect the duration of the inflammatory response in a PPAR α -dependent manner (Devchand *et al.* 1996). In view of these results, we propose a model in which

Figure 2 Mechanisms of PPAR α -mediated inhibition of the AP-1 and NF- κ B signalling pathways. P, ; UV, ; GF, ; ET-1, endothelin-1.

PPAR α negatively interferes with NF- κ B transcription activity by forming inactive complexes with p65 and by inducing I κ B α , the major inhibitor of NF- κ B signalling (Fig. 2). Chromatin immunoprecipitation experiments revealed that the glucocorticoid receptor antagonizes NF- κ B transcriptional activity by interfering with phosphorylation of the serine-2 of the carboxy-terminal domain of the RNA polymerase II without affecting NF- κ B DNA-binding activities, although the glucocorticoid receptor strongly interacts with p65 (Nissen & Yamamoto 2000). It would be of interest to determine whether such a mechanism is also operative for PPAR α using the same technical approach. However, we cannot exclude the existence of additional mechanisms. For instance, PPAR α was reported to play a major role in the control of the cellular redox status (Poynter & Daynes 1998). Moreover, Klucis *et al.* (1984) reported that administration of PPAR α activators results in a drastic increase of the activity of catalase, an antioxidant enzyme. Finally, catalase activity and expression were

found to be increased in endothelial cells upon fibrate treatment (C Furman, E Teissier, B Staels & P Duriez, unpublished observations) (data not shown). A potential involvement of catalase in the control of NF- κ B-driven transcription by PPAR α activators is under investigation in our laboratory.

Promoter analysis revealed that PPAR α controls IL-6 transcription by negatively interfering not only with NF- κ B but also with AP-1 transcriptional activities (Delerive *et al.* 1999a). GST pull-down experiments as well as electrophoretic mobility shift assays demonstrated that PPAR α activators reduce AP-1 DNA-binding activity by physically interacting with the amino-terminal domain of c-Jun (Delerive *et al.* 1999a,b) (Fig. 2). Since most of the proinflammatory genes are under the control of the AP-1 and NF- κ B signalling pathways, it is likely that PPAR α agonists regulate a wide spectrum of genes involved in inflammatory disorders.

One of the most relevant indications regarding a role of PPAR α agonists in inflammation control comes from

Figure 3 Hypothetical model of PPAR γ -mediated inhibition of the inflammatory response genes. PPAR γ antagonizes the AP-1, NF- κ B, STAT and NFAT signalling pathways. TZDs, thiazolidinediones; JAK, Janus kinase.

clinical trials. The influence of PPAR α activators on plasma cytokine levels as well as on acute-phase proteins was determined in patients with angiographically established atherosclerosis (Staels *et al.* 1998). Fibrate treatment for 4 weeks (200 mg daily) reduced IL-6, C-reactive protein and fibrinogen levels in patients with coronary artery disease (Staels *et al.* 1998). Another group reported independently that fenofibrate treatment for 1 month resulted in a significant reduction of plasma interferon- γ (IFN γ) and tumour necrosis factor- α (TNF α) levels in patients with hyperlipoproteinaemia type IIb (Madej *et al.* 1998). These two reports demonstrate that PPAR α activators decrease inflammation in patients, thus indicating a potential use of PPAR α agonists in the treatment of chronic inflammatory diseases.

Is there a role for PPAR γ in inflammation?

A growing body of evidence suggests that PPAR γ may also play a role in inflammation. PPAR γ ligands were shown to inhibit TNF α , IL-6 and IL-1 β expression in monocytes (Jiang *et al.* 1998); inducible nitric oxide synthase (iNOS), matrix metalloproteinase-9 (MMP-9) and scavenger receptor-A expression in macrophages (Ricote *et al.* 1998); IFN-inducible protein 10, monokine induced by inteferon gamma, inteferon gamma inducible T-cell alpha chemoattractant and endothelin-1 expression in endothelial cells (Delerive *et al.* 1999b, Marx *et al.* 2000); IL-2 in T lymphocytes (Yang *et al.* 2000) and IL-8 in colonic epithelial cells (Su *et al.* 1999). Huang *et al.* (1999) demonstrated that IL-4 induces the generation of endog-

enous ligands for PPAR γ through activation of the 12/15-lipoxygenase pathway in macrophages, providing a molecular basis for IL-4-mediated down-regulation of iNOS expression. PPAR γ ligands inhibit the expression of these genes at the transcriptional level. However, one caveat with respect to the interpretation of these studies is the fact that the most pronounced effects were observed with 15-deoxy- Δ 12,14-PGJ2 which is not a very selective PPAR γ ligand. Moreover, when high affinity PPAR γ ligands such as rosiglitazone are used extremely high concentrations (>200 nM) are required to obtain anti-inflammatory activities (Jiang *et al.* 1998, Ricote *et al.* 1998). These pharmacological discrepancies suggested that 15-deoxy- Δ 12,14-PGJ2 may act through PPAR γ -independent pathways. Several recent reports indeed demonstrated that, in the absence of PPAR γ expression, 15-deoxy- Δ 12,14-PGJ2 also negatively regulates the inflammatory response (Petrova *et al.* 1999, Vaidya *et al.* 1999). Two groups demonstrated independently that 15-deoxy- Δ 12,14-PGJ2 represses NF- κ B activation by inhibiting the IKB-kinase (IKK) complex activity (Castrillo *et al.* 2000, Rossi *et al.* 2000) thereby preventing I κ B α degradation. Straus *et al.* (2000) showed that, in addition to inhibiting IKK activity, 15-deoxy- Δ 12,14-PGJ2 probably reduces NF- κ B binding by alkylating p50/p65 dimers (Straus *et al.* 2000). 15-deoxy- Δ 12,14-PGJ2 appears thus to inhibit NF- κ B activation at different levels and may thus exert its activities in both a PPAR γ -dependent and -independent manner.

The molecular mechanisms by which PPAR γ regulates inflammatory response genes are not fully understood (Fig. 3).

Using transient transfection experiments, Ricote *et al.* (1998) demonstrated that PPAR γ inhibits scavenger receptor-A, iNOS and MMP-9 expression by antagonizing the AP-1, signal transducer and activator of transcription (STAT) and NF- κ B pathways. In a recent paper, it was demonstrated that, similar to that reported for PPAR α (Delerive *et al.* 1999a), PPAR γ inhibits NF- κ B-driven transcription by physically interacting with both p65 and p50 (Chung *et al.* 2000). Using endothelin-1 promoter as a model, we demonstrated that PPAR γ inhibits AP-1 transcriptional activity by reducing AP-1 DNA binding (Delerive *et al.* 1999b). This inhibition is likely due to a direct interaction between PPAR γ and c-Jun as previously reported for PPAR α (Delerive *et al.* 1999a). Law *et al.* (1996) reported that glitazones inhibit c-Fos transcription in vascular smooth muscle cells, resulting in a reduction of cell proliferation and migration. The same group demonstrated that glitazones inhibit angiotensin II but not TNF α -mediated extra-cellular signal-regulated kinase 1/2 activation in vascular smooth muscle cells (Goetze *et al.* 1999a,b). We also demonstrated, in a model of cardiac ischemia reperfusion, that rosiglitazone significantly reduces c-Jun-NH $_2$ -terminal kinase (JNK) activation *in vivo* (N Khandoudi, P Delerive, B Staels & A Bril, unpublished observations). Previous reports suggested that nuclear receptors may regulate AP-1 activation by modulating JNK function (Caelles *et al.* 1997, Srivastava *et al.* 1999). However, little is known about the molecular mechanism of kinase inhibition by nuclear receptors.

Li *et al.* (2000b) recently proposed a model for PPAR γ -mediated inhibition of iNOS transcription. In this model, PPAR γ would inhibit STAT1, AP-1 and NF- κ B transcriptional activities by targeting CBP through direct interaction with its N-terminal domain and via SRC-1-like bridge factors (Li *et al.* 2000b). Such a model of competition for limiting amounts of co-activators to inhibit transcriptional activation has already been proposed for other nuclear receptors (Kamei *et al.* 1996, Götlicher *et al.* 1998, Sheppard *et al.* 1998). However, recent reports do not support this model of transrepression (De Bosscher *et al.* 2000, McKay & Cidlowski 2000).

Finally, Yang *et al.* (2000) showed that PPAR γ activation results in a reduction of IL-2 secretion in T lymphocytes. Using electrophoretic mobility shift assays and immunoprecipitation experiments, these authors demonstrated that this inhibition was due to a ligand-dependent interaction between the transcription factor nuclear factor of activated T-cells (NFAT) and PPAR γ (Yang *et al.* 2000). Further studies will be required to elucidate the precise molecular mechanisms of PPAR γ -mediated NF- κ B, AP-1 and STAT1 transcriptional inhibition.

Having established a role for PPAR γ in inflammation *in vitro*, a number of groups carried out *in vivo* studies to assess the potential use of glitazones as anti-inflammatory drugs (Wiesenberg *et al.* 1998, Su *et al.* 1999, Thieringer *et al.*

2000). Using a mouse model of inflammatory bowel disease, Su *et al.* (1999) demonstrated that glitazones markedly reduce colonic inflammation. In a mouse model of atherosclerosis, Li *et al.* (2000a) demonstrated that glitazones reduce TNF α and gelatinase B gene expression in the aortic root. In contrast to these data, other reports indicated that IL-6 and TNF levels were not affected by glitazone treatment in *db/db* mice in response to LPS challenge (Thieringer *et al.* 2000). These authors also showed that PPAR γ ligands, even if used at high concentrations, were ineffective in reducing cytokine levels in monocytes and macrophages (Thieringer *et al.* 2000), raising significant doubts about the potential utility of PPAR γ ligands as anti-inflammatory drugs. In a chronic autoimmune model, rosiglitazone was again inactive in reducing the development and the progression of arthritis (Wiesenberg *et al.* 1998). These apparently conflicting results may reflect the heterogeneity of the different animal models used in these various studies. Additional *in vivo* studies are necessary to determine whether glitazones possess anti-inflammatory activities *in vivo*.

Conclusion

Even though PPARs are considered as master regulators of energy homeostasis, their role no longer seems to be restricted to controlling lipid storage and usage. PPARs may mediate the modulation of the inflammatory response by nutritional and pharmacological stimuli. From a nutritional point of view, additional studies will be necessary to determine whether the beneficial effects of certain dietary fatty acids on immune response are PPAR-mediated and whether PPAR activation will result in a permanent or transient reduction of the inflammatory status. From a pharmacological point of view, clinical studies with the recently launched glitazones should allow determination as to whether these drugs, similar to PPAR α ligands, exert anti-inflammatory activities *in vivo* in humans. Our current knowledge, derived mainly from *in vitro* data, allows us to speculate that PPAR ligands may indeed be useful for the treatment of chronic inflammatory diseases such as atherosclerosis and rheumatoid arthritis.

Acknowledgements

P D is supported by a grant from the Région Nord-Pas-de-Calais. The authors thank Dr Barbier for critical reading of the manuscript.

References

- Caelles C, Gonzalez-Sancho JM & Munoz A 1997 Nuclear hormone receptor antagonism with AP-1 by inhibition of the JNK pathway. *Genes and Development* **11** 3351–3364.

- Castrillo A, Diaz-Guerra MJM, Hortelano S, Martin-Sanz P & Bosca L 2000 Inhibition of I κ B kinase and I κ B phosphorylation by 15-deoxy Δ 12,14-prostaglandin J2 in activated murine macrophages. *Molecular and Cellular Biology* **20** 1692–1698.
- Chung SW, Kang BY, Kim SH, Kim YM, Cho D, Trinchieri G & Kim TS 2000 Oxidized low density lipoprotein inhibits interleukin-12 production in lipopolysaccharide-activated mouse macrophages via direct interactions between PPAR γ and NF- κ B. *Journal of Biological Chemistry* **276** 32681–32687.
- De Bosscher K, Vanden Berghe W, Vermeulen L, Plaisance S, Boone E & Haegeman G 2000 Glucocorticoids repress NF- κ B driven genes by disturbing the interaction of p65 with the basal transcription machinery, irrespective to coactivator levels in the cells. *PNAS* **97** 3919–3924.
- Delerive P, De Bosscher K, Besnard S, Vanden Berghe W, Peters JM, Gonzalez FJ, Fruchart JC, Tedgui A, Haegeman G & Staels B 1999a PPAR α negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF- κ B and AP-1. *Journal of Biological Chemistry* **274** 32048–32054.
- Delerive P, Martin-Nizard F, Chinetti G, Trottein F, Fruchart JC, Najib J, Duriez P & Staels B 1999b PPAR activators inhibit thrombin-induced endothelin-1 production in human vascular endothelial cells by inhibiting the AP-1 signalling pathway. *Circulation Research* **85** 394–402.
- Delerive P, Furman C, Teissier C, Fruchart JC, Duriez P & Staels B 2000a Oxidized phospholipids activate PPAR α in a phospholipase A2-dependent manner. *FEBS Letters* **471** 34–38.
- Delerive P, Gervois P, Fruchart J-C & Staels B 2000b Induction of IkBa expression as a mechanism contributing to the anti-inflammatory activities of PPAR α activators. *Journal of Biological Chemistry* **275** 36703–36707.
- Desvergne B & Wahli W 1999 Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocrine Reviews* **20** 649–688.
- Devchand PR, Keller H, Peters JM, Vasquez M, Gonzalez FJ & Wahli W 1996 The PPAR α -leukotriene B4 pathway to inflammation control. *Nature* **384** 39–43.
- Forman BM, Chen J & Evans RM 1997 Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ . *PNAS* **94** 4312–4317.
- Gervois P, Chopin-Delannoy S, Fadel A, Dubois G, Kosykh V, Fruchart JC, Najib J & Staels B 1999 Fibrates increase human REV-ERB α expression in liver via a novel peroxisome proliferator-activated receptor response element. *Molecular Endocrinology* **13** 400–409.
- Goetze S, Xi XP, Graf K, Fleck E, Hsueh WA & Law RE 1999a Troglitazone inhibits angiotensin II-induced extracellular signal-regulated kinase 1/2 nuclear translocation and activation in vascular smooth muscle cells. *FEBS Letters* **452** 277–282.
- Goetze S, Xi XP, Kawano H, Fleck E, Hsueh WA & Law RE 1999b TNF- α -induced migration of vascular smooth muscle cells is MAPK dependent. *Hypertension* **33** 183–189.
- Göttlicher M, Heck S & Herrlich P 1998 Transcriptional cross-talk, the second mode of steroid hormone receptor action. *Journal of Molecular Medicine* **76** 480–489.
- Huang JT, Welch JS, Ricote M, Binder CJ, Willson TM, Kelly C, Witztum JL, Funk CD, Conrad D & Glass CK 1999 Interleukin-4-dependent production of PPAR- γ ligands in macrophages by 12/15-lipoxygenase. *Nature* **400** 378–382.
- Ijpenberg A, Jeannin E, Wahli W & Desvergne B 1997 Polarity and specific sequence requirements of PPAR:RXR heterodimer binding to DNA. *Journal of Biological Chemistry* **272** 20108–20117.
- Issemann I & Green S 1990 Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* **347** 645–650.
- Jiang C, Ting AT & Seed B 1998 PPAR- γ agonists inhibit production of monocyte inflammatory cytokines. *Nature* **391** 82–86.
- Kamei Y, Xu L, Heinzel T, Torchia J, Kurokawa R, Gloss B, Lin SC, Heyman RA, Rose DW, Glass CK & Rosenfeld MG 1996 A CBP integrator complex mediates transcriptional activation and AP-1 inhibition by nuclear receptors. *Cell* **85** 403–414.
- Kliwer SA, Lenhard JM, Wilson TM, Patel I, Morris DC & Lehmann JM 1995 A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor γ and promotes adipocyte differentiation. *Cell* **83** 803–812.
- Klucis E, Crane D & Masters C 1984 Sequential alterations in the micro-localization of catalase in mouse liver after treatment with hypolipidemic drugs. *Molecular and Cellular Biochemistry* **65** 73–82.
- Kockx M, Gervois P, Poulain P, Derudas B, Peters JM, Gonzalez FJ, Princen HMG, Kooistra T & Staels B 1999 Fibrates suppress fibrinogen gene expression in rodents via activation of the peroxisome proliferator-activated receptor- α . *Blood* **93** 2991–2998.
- Law RE, Meehan WP, Xi XP, Graf K, Wuthrich DA, Coats W, Faxon D & Hsueh WA 1996 Troglitazone inhibits vascular smooth muscle cell growth and intimal hyperplasia. *Journal of Clinical Investigation* **98** 1897–1905.
- Lehmann JM, Moore LB, Smith-Oliver TA, Wilkinson WO, Willson TM & Kliwer SA 1995 An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *Journal of Biological Chemistry* **270** 12953–12956.
- Li AC, Brown KK, Silvestre MJ, Willson TM, Palinski W & Glass CK 2000a Peroxisome proliferator-activated receptor γ ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *Journal of Clinical Investigation* **106** 523–531.
- Li M, Pascual G & Glass CK 2000b Peroxisome proliferator-activated receptor γ -dependent repression of the inducible nitric oxide synthase gene. *Molecular and Cellular Biology* **20** 4699–4707.
- McKay LI & Cidlowski JA 2000 CBP (CREB binding protein) integrates NF- κ B (nuclear factor- κ B) and glucocorticoid receptor physical interactions and antagonism. *Molecular Endocrinology* **14** 1222–1234.
- Madej A, Okopien B, Kowalski J, Zielinski M, Wysocki J, Szygula B, Kalina Z & Herman ZS 1998 Effects of fenofibrate on plasma cytokine concentrations in patients with atherosclerosis and hyperlipoproteinemia IIb. *International Journal of Clinical Pharmacology and Therapeutics* **36** 345–349.
- Marx N, Sukhova GK, Collins T, Libby P & Plutzky J 1999 PPAR α activators inhibit cytokine-induced vascular cell adhesion molecule-1 expression in human endothelial cells. *Circulation* **99** 3125–3131.
- Marx N, Mach F, Sauty A, Leung JH, Sarafi MN, Ransohoff RM, Libby P, Plutzky J & Luster AD 2000 Peroxisome proliferator-activated receptor-gamma activators inhibit IFN-gamma-induced expression of the T cell-active CXC chemokines IP-10, Mig, and I-TAC in human endothelial cells. *Journal of Immunology* **164** 6503–6508.
- Nagy L, Tontonoz P, Alvarez JGA, Chen H & Evans RM 1998 Oxidized LDL regulates macrophage gene expression through ligand activation of PPAR γ . *Cell* **93** 229–240.
- Neve BP, Corseaux D, Chinetti G, Zawadzki C, Fruchart JC, Duriez P, Staels B & Jude B 2001 PPAR α agonists inhibit tissue factor expression in human monocytes and macrophages. *Circulation* **103** 207–212.
- Nissen RM & Yamamoto KR 2000 The glucocorticoid receptor inhibits NF κ B by interfering with serine-2 phosphorylation of the RNA polymerase II carboxy-terminal domain. *Genes and Development* **14** 2314–2329.
- Petrova TV, Akama KT & Van Eldik LJ 1999 Cyclopentone prostaglandins suppress activation of microglia: down-regulation of inducible nitric-oxide synthase by 15-deoxy- Δ 12,14-prostaglandin J2. *PNAS* **96** 4668–4673.
- Poynter ME & Daynes RA 1998 Peroxisome proliferator-activated receptor α activation modulates cellular redox status, represses nuclear factor- κ B signalling, and reduces inflammatory cytokine production in aging. *Journal of Biological Chemistry* **273** 32833–32841.

- Ricote M, Li AC, Willson TM, Kelly CJ & Glass CK 1998 The peroxisome proliferator-activated receptor- γ is a negative regulator of macrophage activation. *Nature* **391** 79–82.
- Rossi A, Kapahi P, Natoli G, Takahashi T, Chen Y, Karin M & Santoro MG 2000 Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of I κ B kinase. *Nature* **403** 103–108.
- Sheppard KA, Phelps KM, Williams AJ, Thanos D, Glass CK, Rosenfeld MG, Gerritsen ME & Collins T 1998 Nuclear integration of glucocorticoid receptor and nuclear factor- κ B signalling by CREB-binding protein and steroid receptor coactivator-1. *Journal of Biological Chemistry* **273** 29291–29294.
- Srivastava S, Weitzmann MN, Cenci S, Ross FP, Adler S & Pacifici R 1999 Estrogen decreases TNF gene expression by blocking JNK activity and the resulting production of c-Jun and JunD. *Journal of Clinical Investigation* **104** 503–513.
- Staels B, Koenig W, Habib A, Merval R, Lebret M, Pineda-Torra I, Delerive P, Fadel A, Chinetti G, Fruchart JC, Najib J, Maclouf J & Tedgui A 1998 Activation of human aortic smooth-muscle cells is inhibited by PPAR α but not by PPAR γ activators. *Nature* **393** 790–793.
- Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang C-H, Sengchanthalangsy LL, Ghosh G & Glass CK 2000 15-Deoxy- Δ 12,14-prostaglandin J2 inhibits multiple steps in the NF- κ B signalling pathway. *PNAS* **97** 4844–4849.
- Su CG, Wen X, Bailey ST, Jiang W, Rangwala SM, Keilbaugh SA, Flanigan A, Murthy S, Lazar MA & Wu GD 1999 A novel therapy for colitis utilizing PPAR- γ ligands to inhibit the epithelial inflammatory response. *Journal of Clinical Investigation* **104** 383–389.
- Thieringer R, Fenyk-Melody JE, Le Grand CB, Shelton BA, Detmers PA, Somers EP, Charbin L, Moller DE, Wright SD & Berger J 2000 Activation of peroxisome proliferator-activated receptor γ does not inhibit IL-6 or TNF- α responses of macrophages to lipopolysaccharide *in vitro* or *in vivo*. *Journal of Immunology* **164** 1046–1054.
- Tugwood JD, Issemann I, Anderson RG, Bundell KR, McPheat WL & Green S 1992 The mouse peroxisome proliferator-activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl coA oxidase gene. *EMBO Journal* **11** 433–439.
- Vaidya S, Somers EP, Wright SD, Detmers PA & Bansal VS 1999 15-Deoxy- Δ 12,14-prostaglandin J2 inhibits the β 2 integrin-dependent oxidative burst: involvement of a mechanism distinct from peroxisome proliferator-activated receptor γ ligation. *Journal of Immunology* **163** 6187–6192.
- Wiesenberg I, Chiesi M, Missbach M, Spanka C, Pignat W & Carlberg C 1998 Specific activation of the nuclear receptors PPAR γ and ROR α by the antidiabetic thiazolidinedione BRL 49653 and the antiarthritic thiazolidinedione derivative CGP 52608. *Molecular Pharmacology* **53** 1131–1138.
- Yang XY, Wang LH, Chen T, Hodge DR, Resau JH, DaSilva L & Farrar WL 2000 Activation of human T lymphocytes is inhibited by peroxisome proliferator-activated receptor γ (PPAR γ) agonists. *Journal of Biological Chemistry* **275** 4541–4544.

Received 1 November 2000

Accepted 14 November 2000