Immunoelectron microscopic localization of three key steroidogenic enzymes (cytochrome P450_{scc}, 3 β -hydroxysteroid dehydrogenase and cytochrome P450_{c17}) in rat adrenal cortex and gonads

G Pelletier, S Li, V Luu-The, Y Tremblay, A Bélanger and F Labrie

Oncology and Molecular Endocrinology Research Center, Laval University Medical Center (CHUL), and Laval University, Québec, Canada G1V 4G2

(Requests for offprints should be addressed to G Pelletier, Oncology and Molecular Endocrinology Research Center, Laval University Hospital (CHUL), 2705, Laurier Boulevard, Québec, Québec, Canada G1V 4G2; Email: georges.pelletier@crchul.ulaval.ca)

Abstract

The biosynthesis of steroid hormones in endocrine steroid-secreting glands results from a series of successive steps involving both cytochrome P450 enzymes, which are mixed-function oxidases, and steroid dehydrogenases. So far, the subcellular distribution of steroidogenic enzymes has been mostly studied following subcellular fractionation, performed in placenta and adrenal cortex. In order to determine *in situ* the intracellular distribution of some steroidogenic enzymes, we have investigated the ultrastructural localization of the three key enzymes: P450 side chain cleavage (scc) which converts cholesterol to pregnenolone; 3 β -hydroxysteroid dehydrogenase (3 β -HSD) which catalyzes the conversion of 3 β -hydroxy-5-ene steroids to 3-oxo-4-ene steroids (progesterone and androstenedione); and P450_{c17} which is responsible for the transformation of C₂₁ into C₁₉ steroids (dehydroepiandrosterone and androstenedione). Immunogold labeling was used to localize the enzymes in rat adrenal cortex and gonads. The tissues were fixed in 1% glutaraldehyde and 3% paraformaldehyde and included in LR gold resin. In the adrenal cortex, both P450_{scc} and 3 β -HSD immunoreactivities were detected in the reticular, fascicular and glomerular zones. P450_{scc} was exclusively found in large

mitochondria. In contrast, 3 β -HSD antigenic sites were mostly observed in the endoplasmic reticulum (ER) with some gold particles overlying crista and outer membranes of the mitochondria. P450_{c17} could not be detected in adrenocortical cells. In the testis, the three enzymes were only found in Leydig cells. Immunolabeling for P450_{scc} and 3 β -HSD was restricted to mitochondria, while P450_{c17} immunoreactivity was exclusively observed in ER. In the ovary, P450_{scc} and 3 β -HSD immunoreactivities were found in granulosa, theca interna and corpus luteum cells. The subcellular localization of the two enzymes was very similar to that observed in adrenocortical cells. P450_{c17} could also be detected in theca interna cells of large developing and mature follicles. As observed in Leydig cells, P450_{c17} immunolabeling could only be found in the ER. These results indicate that in different endocrine steroid-secreting cells P450_{scc}, 3 β -HSD and P450_{c17} have the same association with cytoplasmic organelles (with the exception of 3 β -HSD in Leydig cells), suggesting similar intracellular pathways for biosynthesis of steroid hormones.

Journal of Endocrinology (2001) **171**, 373–383

Introduction

The biosynthesis of steroid hormones in the adrenal cortex, testis, ovary and placenta results from a series of successive steps involving both cytochrome P450 enzymes, which are mixed-function oxidases, and steroid dehydrogenases (for review see Miller 1988). The first step is the conversion of cholesterol into pregnenolone by the cholesterol side-chain cleavage P450 (P450_{scc}) enzyme. Pregnenolone can be either directly hydroxylated at the 17 α -position or dehydrogenated at the 3 β -position by the enzyme 3 β -hydroxysteroid dehydrogenase (3 β -HSD) to yield progesterone. The hydroxylation of pregnenolone and progesterone on the 17 α -position and the scission of the

C17,20-carbon bond of the 17 α -hydroxylated steroids are catalyzed by the steroid 17 α -hydroxylase/17,20-lyase P450 (P450_{c17}). This enzyme is a key branch point in the pathways of biosynthesis of steroid hormones, such as glucocorticoids and sex steroids. All these enzymes are distributed throughout different cellular compartments.

So far, most of the data on the subcellular localization of steroidogenic enzymes have been obtained following subcellular fractionation studies performed in placenta and adrenal cortex (Katagiri *et al.* 1976, Kominami *et al.* 1980, Pudney *et al.* 1985, Shinzawa *et al.* 1988, Luu-The *et al.* 1989, 1990, 1991, Cherradi *et al.* 1993, Sauer *et al.* 1994, Thomas *et al.* 1999). The ultrastructural localization of steroidogenic enzymes has, so far, not been extensively

Figure 1 Localization of P450_{sc} in fascicular zone of the adrenal cortex. Colloidal gold particles are located in mitochondria (M), overlying vesicular structures (→). Note the absence of labeling over the endoplasmic reticulum and lipid inclusions (L). Magnification $\times 64\,000$.

investigated and the few reports on the subject are restricted to the adrenal cortex (Ishimura *et al.* 1988, Shinzawa *et al.* 1988, Whitnall *et al.* 1993, Cherradi *et al.* 1997, Ishimura & Fujita 1997).

From the previous biochemical and morphological studies it appears that P450_{sc} is almost exclusively associated with mitochondria and that 3 β -HSD is predominant in microsomal fractions (endoplasmic reticulum) and has also been found in mitochondrial preparations (Katagiri *et al.* 1976, Luu-The *et al.* 1989, 1990, 1991, Cherradi *et al.* 1994, 1997, Ishimura & Fujita 1997, Thomas *et al.* 1999). P450_{c17} has been found to be associated only with microsomal fractions (Kominami *et al.* 1980, Miller 1988, Ishimura & Fujita 1997).

In order to clarify the subcellular distribution of these three enzymes in different steroid-secreting cell types, we felt it was of interest to study the comparative *in situ* localization of the enzymes in adult rat adrenal cortex and gonads. The studies were performed at the electron microscopic level using specific antibodies for each enzyme.

Materials and Methods

Animals

Four adult male (225–250 g) and female (175–200 g) Sprague-Dawley rats were housed under constant

temperature ($21 \pm 1^\circ\text{C}$) and light (lights on from 0600 to 2000 h) regimens. Purina Chow (Ralston-Purina, St Louis, MO, USA) and tap water were available *ad libitum*. The rats were all perfused between 0900 and 1000 h for histological procedures, as described below. The females were at random stages of the estrous cycle.

Histological procedures

The animals were perfused transcardially with 200 ml 1% glutaraldehyde and 3% (w/v) paraformaldehyde in 0.1 M phosphate buffer (pH 7.4). The different tissues, namely testes, ovaries and adrenals were excised and post-fixed in the same fixative for 24 h at 4°C .

Preparation of the tissue blocks in LR gold resin (London Resin Co., Reading, Berks, UK) was performed as previously described (Thorpe 1999). Briefly, following fixation, tissue fragments were rinsed with phosphate buffer and dehydrated in graded series of up to 100% ethanol. The tissue fragments were then infiltrated with LR gold resin and placed in gelatin capsules for polymerization under UV lamps at 4°C for 24 h.

Immunocytochemistry

Semithin sections were first performed for selection of the areas of interest in each gland. Ultrathin sections were then cut with a diamond knife and collected on nickel grids.

Figure 2 Localization of 3 β -HSD in the adrenal cortex (fascicular zone). The colloidal gold particles are mostly associated with the endoplasmic reticulum (ER). In the mitochondria (M), the gold particles are overlying vesicular cristae membranes as well as the outer membranes (arrows). Magnification $\times 64\,000$.

They were immunostained using protein A-gold complex (10 nm; British Biocell Int., Cardiff, UK), as described (Roth *et al.* 1978). The antisera to human P450_{scc} (supplied by Dr Miller; Black *et al.* 1993), human type 1 3 β -HSD (Luu-The *et al.* 1989, 1990, Pelletier *et al.* 1992) and human P450_{c17} (Tremblay *et al.* 1994) were all used at dilutions ranging from 1:500 to 1:1000. Control experiments were performed by substituting non-immunized rabbit serum (1:500) or the antiserum (1:500) absorbed with an excess of their respective antigen (10^{-6} M). Following immunostaining procedures, the sections were counterstained with 1% uranyl acetate and lead citrate.

Results

In all the tissues examined, the fixation with aldehydes and embedding in LR gold did not always provide a clear distinction between the smooth and rough endoplasmic reticulum (ER). Since the steroid-secreting cells contain large amounts of smooth ER, especially in close proximity of lipid inclusions and mitochondria, we consider that in general the labeling occurring in ER was mostly associated with smooth ER.

Adrenal cortex

In the adrenal cortex, immunolabeling for P450_{scc} and 3 β -HSD was observed in secretory cells of reticular,

fascicular and glomerular zones. The adrenocortical cells are characterized by the presence of large amounts of smooth ER, numerous lipid inclusions and large mitochondria with tubular or vesicular cristae. P450_{scc} immunogold labeling was restricted to mitochondria, with the majority of colloidal gold particles overlying vesicular structures (Fig. 1). Neither ER nor lipid inclusions were labeled. In contrast, 3 β -HSD immunolabeling was mostly observed over ER (Fig. 2; Table 1). Labeling also occurred in mitochondria, with gold particles being located over the vesicular crista membranes of mitochondria as well as over the outer membranes. No staining was obtained with the antibodies to P450_{c17}.

Testis

In the testis, the three enzymes were immunolocalized exclusively in Leydig cells. The cytoplasm of these cells is characterized by abundant smooth ER, numerous mitochondria and a few lipid inclusions. As observed in the adrenal cortex, P450_{scc} immunoreactivity was exclusively observed in mitochondria, although the intensity of labeling was less than that observed in the adrenocortical cells (Fig. 3). Colloidal gold particles were observed over the limiting and crista membranes of the mitochondria. In contrast to what has been observed in the adrenal cortex, 3 β -HSD immunostaining was exclusively detected in mitochondria, with no specific labeling occurring in the

Table 1 Subcellular distribution of 3β-HSD (percent of gold particles). Data were calculated from the analysis of 820-1310 gold particles per cell type

Subcellular compartments	Cell types				
	Adrenocortical cells	Leydig cells	Ovarian cells		
			Granulosa cells	Thecal cells	Luteal cells
Mitochondria	25.5	100	25.8	30.6	26.1
Endoplasmic reticulum	74.5	0	74.2	69.4	73.9

Figure 3 Localization of P450_{scc} in a Leydig cell of the testis. The labeling is seen in mitochondria (M). Very few particles, which can be associated to background, are observed over the endoplasmic reticulum. Magnification × 64 000.

ER (Fig. 4; Table 1). As shown in Fig. 5, P450_{c17} immunoreactivity was only observed in ER. The very few gold particles which were occasionally seen overlying mitochondria could possibly be attributed to background.

Ovary

In the ovary, the ultrastructural characteristics of the steroid secreting cells are similar to those observed in the Leydig cells with abundant smooth ER and lipid inclusions varying in number and size. The lipid inclusions appeared larger and more numerous in luteal and thecal cells than in granulosa cells. P450_{scc} could be immunolocalized in

granulosa cells and theca interna cells in large antral and preovulatory follicles (Fig. 6). In small developing follicles, the labeling was weak and inconsistent in both granulosa and thecal cells. Interstitial and corpus luteum cells were also immunolabeled (Fig. 7). As observed in the Leydig and adrenocortical cells, in all the positive cells, P450_{scc}-associated gold particles were only observed over mitochondria, mostly in association with crista membranes.

3β-HSD immunoreactivity was detected in thecal and granulosa cells of growing and preovulatory follicles. Primary follicles remained unstained. Interstitial and luteal cells were also strongly labeled. (Figs 8 and 9; Table 1). As observed in adrenocortical cells, in all the reactive cells,

Figure 4 Localization of 3β -HSD in a Leydig cell of the testis. Immunolabeling is restricted to mitochondria (M). Magnification $\times 64\,000$.

Figure 5 Localization of $P450_{c17}$ in a Leydig cell of the testis. Labeling is exclusively detected in the endoplasmic reticulum (ER). Very few particles, which correspond to background level, are present over mitochondria (M). Magnification $\times 64\,000$.

most of the immunoreactivity had an extra-mitochondrial localization, most of the gold particles being associated with the ER. In the mitochondria, the labeling was associated with the crista membranes.

$P450_{c17}$ was detected in the theca interna cells of large antral and preovulatory follicles (Fig. 10) and interstitial cells while granulosa cells remained unlabeled. Few luteal and interstitial cells were observed to exhibit

Figure 6 Localization of $P450_{\text{scc}}$ in the ovary: theca interna cell. Colloidal gold particles are almost exclusively associated with the tubular-vesicular cristae membranes of mitochondria (M). Very few particles, corresponding to the background, are present outside mitochondria. RER: rough endoplasmic reticulum. Magnification $\times 64\,000$.

immunolabeling. The subcellular localization of the enzyme was identical to that observed in the Leydig cells of the testis with labeling exclusively associated with ER.

When non-immunized rabbit serum or antisera immunoadsorbed with their respective antigens were used, no association of gold particles with any organelle in the reactive cell types could be observed. Only a few dispersed gold particles could be detected throughout the sections (Fig. 11).

Discussion

The present data obtained at the ultrastructural level provides new information about the *in situ* localization of three key steroidogenic enzymes. Although due to the fixation and embedding methods used it was not always possible to clearly identify the smooth ER, the abundance of smooth ER in the steroid-secreting cells suggest that most of the extra-mitochondrial immunolabeling was in fact associated with smooth ER. Moreover, the rough ER was not seen to be consistently labeled in any of the $P450_{\text{scc}}$ -, $3\beta\text{-HSD}$ - or $P450_{\text{c17}}$ -positive cells.

$P450_{\text{scc}}$ immunoreactivity was detected in the three layers of the adrenal cortex, the Leydig cells in the testis as

well as in thecal, granulosa, interstitial and luteal cells in the ovary. These results are in good agreement with previous localization studies performed in several species including the rat (Miller 1988, Sasano *et al.* 1989, LeGoascogne *et al.* 1991, Ishimura & Fujita 1997, Sanders & Stouffer 1997). In all the reactive cell types, immunolabeling was restricted to mitochondria, with gold particles overlying vesicular structures. These results are in agreement with a previous immunoelectron microscopic study indicating that $P450_{\text{scc}}$ immunoactivity was associated with the matrix side of the inner mitochondrial membranes in rat adrenocortical cells as well in granulosa, theca interna and interstitial cells in the ovary (Farkash *et al.* 1986). They are also in agreement with previous subcellular fractionation studies indicating that in rat and bovine adrenal glands $P450_{\text{scc}}$ activity was associated with mitochondrial fractions (Hanukoglu *et al.* 1981, Kramer *et al.* 1984, Cherradi *et al.* 1994). Submitochondrial fractionation studies combined with immunolabeling have shown that immunoreactive $P450_{\text{scc}}$ was associated with inner mitochondrial membranes (Cherradi *et al.* 1997). These biomedical and morphological studies thus suggest that in steroid-secreting cells cholesterol is transferred from the outer to the inner membranes (crista membranes facing matrix) where $P450_{\text{scc}}$ is present to initiate the steroid synthesis.

Figure 7 Localization of $P450_{scc}$ in the ovary: corpus luteum cell. Immunolabeling is observed only over cristae membranes of mitochondria (M). Neither the endoplasmic reticulum nor the lipid inclusions (L) are labeled. Magnification $\times 64\,000$.

Figure 8 Localization of 3β -HSD in the ovary: granulosa cell. Most of the colloidal gold particles are associated with the endoplasmic reticulum, with a few particles overlying mitochondria (M). L: lipid inclusion. Magnification $\times 64\,000$.

With antibodies which have been extensively used to localize 3β -HSD in tissues of several species including rat (Dupont *et al.* 1990a,b, Pelletier *et al.* 1992), we have shown that, in steroid-secreting cells of the adrenal cortex,

3β -HSD is mostly observed in association with membranes of the ER, with consistent labeling of mitochondria. In bovine and rat adrenal cortex, 3β -HSD has been shown to be associated with both the mitochondrial and

Figure 9 Localization of 3β -HSD in the ovary: corpus luteum cell. Gold particles are detected over the endoplasmic reticulum with a high density in membranes surrounding the lipid inclusions (L). Immunolabeling can also be observed inside the mitochondria (M) and their outer membranes. LY: lysosome. Magnification $\times 64\,000$.

Figure 10 Localization of $P450_{c17}$ in a theca interna cell of the ovary. Strong immunolabeling of endoplasmic reticulum (ER) can be observed. Neither mitochondria (M) nor lipid inclusions (L) appear to be significantly labeled. Magnification $\times 64\,000$.

microsomal fractions (Cherradi *et al.* 1993, 1994, Sauer *et al.* 1994). By immunoelectron microscopy, 3β -HSD immunoreaction was found in the ER in bovine adrenal

cortex (Ishimura *et al.* 1988, Ishimura & Fujita 1997). Recently, Cherradi *et al.* (1997) using immunoelectron microscopy have reported that 3β -HSD immunoreactivity

Figure 11 Control adrenal cortex (fascicular zone) section. Immunoabsorption of the antiserum to 3 β -HSD with the antigen has completely prevented any staining (compare with Fig. 2). Very few dispersed gold particles can be detected. M: mitochondria. Magnification $\times 64\,000$.

could be detected in both ER and mitochondria, the highest density being observed in the ER, as shown in the present study. In mitochondria, we observed that 3 β -HSD labeling was mostly associated with crista membranes and was also found over the outer membranes. Cherradi *et al.* (1997) using immunogold-labeling also observed that 60% of 3 β -HSD immunoreactivity was associated with the crista membranes, while the rest of antigenic sites were distributed between the outer membranes, the mitochondrial matrix and the intermembrane spaces.

In the ovary, we confirm previous studies obtained in the rat which indicated that 3 β -HSD was expressed in all ovarian steroid-secreting cells including granulosa cells in growing follicles (Dupont *et al.* 1990b). At the ultrastructural level, we clearly establish that the subcellular localization of 3 β -HSD is identical in all the positive ovarian cells. As in adrenocortical cells, 3 β -HSD was mainly found in the ER with consistent labeling of mitochondrial crista membranes. This is the first report on the *in situ* ultrastructural localization of 3 β -HSD in ovarian tissue. The present data strongly suggest that the steroid metabolism involving 3 β -HSD follows an intracellular pathway similar to that observed in adrenocortical cells.

In the testis, immunoreactive material was only observed in Leydig cells, in agreement with previous results obtained at the light microscopic level (Dupont *et al.* 1990a,b, Pelletier *et al.* 1992). Contrary to what has

been observed in the adrenal cortex and ovary, 3 β -HSD immunoreactivity was confined to the mitochondria. Using immunoblot analysis, Cherradi *et al.* (1997) had previously shown the presence of immunoreactive 3 β -HSD in MH-10 mouse Leydig cell mitochondrial fractions. Such a localization suggests that the type of 3 β -HSD expressed in the rat testis is different, although immunologically related, from that found in the adrenal cortex and ovary. The exact reason for such a differential localization is unclear.

No P450_{C17} immunoreactivity could be detected in the adrenal cortex, as previously shown by light microscope immunostaining studies (LeGoascogne *et al.* 1991). This is in complete agreement with the biochemical data indicating the absence of 17 α -hydroxylase/17,20 lyase activity in the rat adrenal cortex (Miller 1988). In the testis, P450_{C17} immunoreactivity was only detected in the Leydig cells, confirming previous light microscope studies performed in the rat and guinea pig (LeGoascogne *et al.* 1991, Suzuki *et al.* 1992). Strong immunolabeling was only observed in the ER. In the ovary, the immunostaining was mostly restricted to theca interna cells, with a few interstitial and luteal cells being weakly labeled. At the light microscopic level, immunoreactive P450_{C17} was only found in theca interna of rat ovaries (LeGoascogne *et al.* 1991). In the bovine ovary, both theca interna and interstitial cells were shown to express P450_{C17} mRNA and the

immunoreactive protein (Suzuki *et al.* 1992). In the human ovary, immunoreactive P450_{c17} was confined to theca interna cells and a few luteinized theca cells in corpora lutea (Sasano *et al.* 1989). As observed in the Leydig cells, gold particles were only detected over ER. Up until now, there had been no report on the subcellular localization of P450_{c17} in testis and ovary. The exclusive detection of immunoreactive material in the ER is in agreement with previous biochemical studies indicating that 17 α -hydroxylase and 17,20-lyase activities were associated with bovine and guinea pig adrenocortical microsomes (Inano *et al.* 1969, Kominami *et al.* 1982, Shinzawa *et al.* 1988). By immunoelectron microscopy involving the use of the direct peroxidase-labeled antibody technique, Shinzawa *et al.* (1988) reported that staining for P450_{c17} occurred on smooth-surface ER in the guinea pig adrenal cortex. In their studies, due to the diffusion of the reaction product, it was not possible to identify clearly all the labeled structures. From the present data obtained in the rat gonads, it appears that the subcellular distribution of the P450_{c17} is identical in all steroidogenic endocrine cells, thus extending previous results obtained by subcellular fractionation and immunoelectron microscopy in the adrenal cortex.

In summary, it clearly appears that in steroid-secreting cells of endocrine glands three key enzymes involved in steroid biosynthesis are associated with the same organelles. The only exception is the exclusive localization of 3 β -HSD to mitochondria in Leydig cells. The present data indicate that, in general, steroid metabolism follows similar intracellular pathways in different steroid-secreting cell types.

References

- Black S, Syklarz G, Harikrishna J, Lin D, Wolf C & Miller WL 1993 Regulation of proteins in the cholesterol side-chain cleavage system in JEG-3 and Y-1 cells. *Endocrinology* **135** 539–545.
- Cherradi N, Defaye G & Chambaz EM 1993 Dual subcellular localization of the 3 β -hydroxysteroid dehydrogenase isomerase: characterization of three key steroidogenic enzymes in the bovine adrenal cortex. *Journal of Steroid Biochemistry and Molecular Biology* **46** 773–779.
- Cherradi N, Defaye G & Chambaz EM 1994 Characterization of the 3 β -hydroxysteroid dehydrogenase activity associated with bovine adrenocortical mitochondria. *Endocrinology* **134** 1358–1364.
- Cherradi N, Rossier MF, Vallotton MB, Timberg R, Friedberg I, Orly J, Wang XJ, Stocco DM & Capponi AM 1997 Submitochondrial distribution of three key steroidogenic proteins (steroidogenic acute regulatory protein and cytochrome P450_{scc} and 3 β -hydroxysteroid dehydrogenase isomerase enzymes) upon stimulation by intracellular calcium in adrenal glomerulosa cells. *Journal of Biological Chemistry* **272** 7899–7907.
- Dupont É, Luu-The V, Labrie F & Pelletier G 1990a Light microscopic immunocytochemical localization of 3 β -hydroxy-5-ene-steroid dehydrogenase/ Δ 5- Δ 4 isomerase (3 β -HSD) in the gonads and adrenal glands of the guinea pig. *Endocrinology* **126** 2906–2909.
- Dupont É, Zhao HF, Rhéaume E, Simard J, Luu-The V, Labrie F & Pelletier G 1990b Localization of 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4-isomerase in rat gonads and adrenal glands by immunocytochemistry and *in situ* hybridization. *Endocrinology* **127** 1394–1403.
- Farkash Y, Timberg R & Orly J 1986 Preparation of antiserum to rat cytochrome P-450 cholesterol side chain cleavage, and its use for ultrastructural localization of the immunoreactive enzyme by protein A-gold technique. *Endocrinology* **118** 1353–1365.
- Hanukoglu J, Spitzberg V, Bumpus JA, Dus KM & Jefcoate CR 1981 Adrenal mitochondrial cytochrome P450 scc. *Journal of Biological Chemistry* **256** 4301–4329.
- Inano H, Inano A & Tamaoki B 1969 Submicrosomal distribution of adrenal enzymes and cytochrome P450-related to corticoidogenesis. *Biochimica et Biophysica Acta* **191** 257–271.
- Ishimura K & Fujita H 1997 Light and electron microscopic immunohistochemistry of the localization of adrenal steroidogenic enzymes. *Microscopy Research and Technique* **36** 445–453.
- Ishimura K, Yoshinaga-Hirabayashi T, Fujita H, Ishii-Ohba H, Inano H & Tamaoki B 1988 Light and electron microscopic immunocytochemistry on the localization of 3 β -hydroxysteroid dehydrogenase/isomerase in the bovine adrenal cortical cells. *Histochemistry* **89** 35–39.
- Katagiri M, Takemori S, Stajeki E, Suhara K, Gomi T & Sato H 1976 Characterization of purified cytochrome P450 scc and P450-11 β from bovine adrenocortical mitochondria. *Advances in Experimental Medicine and Biology* **74** 281–292.
- Kominami S, Ochi H, Kobayashi Y & Takemori S 1980 Studies on the steroid hydroxylation system in adrenal cortex microsomes. Purification and characterization of cytochrome P-450 specific for steroid C-21 hydroxylation. *Journal of Biological Chemistry* **255** 3386–3394.
- Kominami S, Shinzawa K & Takemori S 1982 Purification and some properties of cytochrome P450-specific for steroid 17 α -hydroxylation and C₁₇-C₂₀ bond cleavage from guinea pig adrenal microsome. *Biochemical and Biophysical Research Communications* **109** 916–921.
- Kramer RE, Rainey WE, Furkenstein B, Du A, Simpson ER & Waterman ER 1984 Induction of synthesis of mitochondrial steroidogenic enzymes of bovine adrenocortical cells by analogs of cyclic AMP. *Journal of Biological Chemistry* **259** 707–712.
- LeGoascogne C, Savaris N, Goviezan M, Takemori S, Kuminami S, Beaulieu EE & Robel P 1991 Immunoreactive cytochrome P-450 17 α in rat and guinea-pig gonads, adrenal glands and brain. *Journal of Reproduction and Fertility* **93** 609–622.
- Luu-The V, Lachance Y, Labrie C, Leblanc G, Thomas JL, Strickle RC & Labrie F 1989 Full length cDNA structure and deduced amino acid sequence of human 3 β -hydroxy-5-ene steroid dehydrogenase. *Molecular Endocrinology* **3** 1310–1312.
- Luu-The V, Takahashi M & Labrie F 1990 Purification of microsomal and mitochondrial 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4 isomerase from human placenta. *Annals of the New York Academy of Sciences* **595** 386–388.
- Luu-The V, Takahashi M, de Launoit Y, Dumont M, Lachance Y & Labrie F 1991 Evidence for distinct dehydrogenase and isomerase sites within a single 3 β -hydroxysteroid dehydrogenase/5-ene-4-ene isomerase protein. *Biochemistry* **30** 8861–8865.
- Miller WL 1988 Molecular biology of steroid hormone synthesis. *Endocrine Reviews* **9** 295–318.
- Pelletier G, Dupont E, Simard J, Luu-The V, Bélanger A & Labrie F 1992 Ontogeny and subcellular localization of 3 β -hydroxysteroid dehydrogenase (3 β -HSD) in the human and rat adrenal, ovary and testis. *Journal of Steroid Biochemistry and Molecular Biology* **43** 451–467.
- Pudney J, Canick JA, Clifford NM, Knapp JB & Callar GV 1985 Location of enzymes of androgen and estrogen biosynthesis in the testis of the ground squirrel. *Biology of Reproduction* **33** 971–980.
- Roth J, Bendayan M & Orci L 1978 Ultrastructural localization of intracellular antigens by the use of protein-A gold complex. *Journal of Histochemistry and Cytochemistry* **26** 1074–1081.

- Sanders SL & Stouffer RL 1997 Localization of steroidogenic enzymes in macaque luteal tissue during the menstrual cycle and stimulated early pregnancy: immunocytochemical evidence supporting the two-cell model for estrogen production in the primate corpus luteum. *Biology of Reproduction* **56** 1077–1087.
- Sasano H, Okamoto M, Mason JD, Simpson ER, Mendelson CR, Sasano N & Silverberg SG 1989 Immunolocalization of aromatase, 17 α -hydroxylase and side-chain-cleavage cytochromes P-430 in the human ovary. *Journal of Reproduction and Fertility* **85** 163–169.
- Sauer LA, Chapman JC & Dauchy RT 1994 Topology of 3 β -hydroxy-5-ene-steroid dehydrogenase/ Δ 5- Δ 4-isomerase in adrenal cortex mitochondria and microsomes. *Endocrinology* **134** 751–759.
- Shinzawa K, Ishibashi S, Murakoshi M, Watanabe K, Kominami S, Karvohara A & Takemori S 1988 Relationship between zonal distribution of microsomal P450s (P-450 17 α , lyase and P-450c21) and steroidogenic activities in guinea-pig adrenal cortex. *Journal of Endocrinology* **119** 191–200.
- Suzuki T, Sasano H, Suwai T, Mason JL & Nagura H 1992 Immunohistochemistry and *in situ* hybridization of P450 17 α (17 α -hydroxyl/17-20-lyase). *Journal of Histochemistry and Cytochemistry* **40** 903–908.
- Thomas JL, Evans BW, Blanco G, Mason JI & Stickler RC 1999 Creation of a fully active, cytosolic form of human type 3 β -hydroxysteroid dehydrogenase/isomerase by the deletion of a membrane spanning domain. *Journal of Molecular Endocrinology* **23** 231–239.
- Thorpe JR 1999 The application of LR gold resin for immunogold labeling. *Methods in Molecular Biology* **117** 99–110.
- Tremblay Y, Fleury A, Beaudoin C, Vallee M & Belanger A 1994 Molecular cloning and expression of guinea pig cytochrome P450c17 cDNA (steroid 17 α -hydroxylase/17,20 lyase): tissue distribution, regulation, and substrate specificity of the expressed enzyme. *DNA and Cell Biology* **13** 1199–1212.
- Whitnall MH, Driscoll WJ, Lu YC & Strott CA 1993 Estrogen and hydroxysteroid sulfatransferases in guinea pig adrenal cortex: cellular and subcellular distribution. *Endocrinology* **133** 2284–2291.

Received 21 May 2001

Accepted 9 July 2001