Genetic variability in melatonin secretion originates in the number of pinealocytes in sheep

A Gómez Brunet^{1,2}, B Malpaux¹, A Daveau¹, C Taragnat¹
and P Chemineau¹

¹Neuroendocrinologie, Physiologie Reproduction et Comportements-INRA-CNRS-Univ. F. Rabelais UMR6073, 37380 Nouzilly, France

²Departamento Reproducción Animal y Conservación de Recursos Zoogenéticos, INIA, 28040 Madrid, Spain

(Requests for offprints should be addressed to P Chemineau, PRC-INRA-CNRS-Univ. UMR6073, 37380 Nouzilly, France; Email: Philippe.Chemineau@tours.inra.fr)

Abstract

Genetic variability in plasma melatonin concentrations in ewes results from variations in pineal weight. This study investigated whether it is due to a difference in the number of pinealocytes, or in their size. Two groups of lambs were assigned before birth to being extremes (18 High and 21 Low) by calculating their genetic value on the basis of the melatonin concentrations of their parents. Lambs were bled from 1 week of age until 14 weeks of age. Pineal gland, brain and pituitary weights, length and width of the brain, and length of the hypothalamus were recorded. A significant effect (ANOVA) of genetic group ($P<0.05$) and age ($P<0.05$) was detected on mean nocturnal plasma melatonin concentrations, as soon as the first week after birth (mean $\pm$ S.E.M.; High: 51.7 ± 10.7 vs

Low: 31.9 ± 3.2 pg/ml). There was no difference between the two genetic groups in any of the brain parameters measured, but the pineal glands of the High group were heavier and contained significantly more pinealocytes (High: 27.8 ± 2.4 vs Low: $21.0 \pm 2.4 \times 10^6$; $P<0.05$) than those in the Low group. The mean size of pinealocytes did not differ between the two genetic groups. Thus, the genetic variability in nocturnal plasma melatonin concentrations in sheep is expressed by 1 week of age and higher levels of secretion are the consequence of larger pineal glands containing a greater number of pinealocytes.

Journal of Endocrinology (2002) **172**, 397–404

Introduction

In mammals, the pineal gland secretes melatonin, which transduces the neural photoperiodic information received by the retina into a hormonal message which is read by a large variety of tissues in the whole organism (Arendt 1995). Melatonin is synthesized and released into the general circulation with a marked day–night rhythm characterized by low or undetectable concentrations during the day and increases many fold at night (Bittman *et al.* 1983). A large variability in nocturnal plasma melatonin concentrations among individuals has been described in several mammalian species, including humans (Arendt 1995) and sheep (Malpaux *et al.* 1987, 1988).

In sheep, it has been clearly shown that nocturnal plasma melatonin concentrations is a very stable and highly repeatable characteristic within each ewe (Chemineau *et al.* 1996), and estimation of the heritability coefficient of this trait showed that this between-individual variation in melatonin secretion is under strong genetic control (Zarazaga *et al.* 1998a). Similar findings were obtained in humans concerning melatonin in urine (Wetterberg *et al.*

1983). While the physiological significance of this genetic variability in melatonin secretion is unknown, investigations conducted regarding their physiological origin have demonstrated that such a variability comes from differences in the synthesis of melatonin from the pineal gland (Zarazaga *et al.* 1998b). More specifically, this ability of the pineal gland is not related to enzymatic activity per mg of tissue, but to the size of the pineal gland, which can vary from 30 mg to more than 200 mg (Coon *et al.* 1999). However, despite a well-established characterization of ‘high secretors’ and ‘low secretors’ of melatonin in adult sheep, we do not know whether this is due to differences in the number, or in the size, of pinealocytes in the pineal gland.

In the literature, results are scarce about the existence of a relationship between melatonin secretion and pineal size and/or tissue composition of the pineal gland. In rats, no correlation was detected between serum melatonin and pineal size (Vollrath & Welker 1984), and ‘classical’ books on the pineal gland detailing the anatomy of the gland (Kappers & Pévet 1979, Vollrath 1981) do not raise the question of such a relationship. In other endocrine organs,

there is evidence that an increase in secretory activity may be due to an increase in the number of secretory cells (Shimokawa *et al.* 1996, Francis *et al.* 2000), or to an increase in cell size (Fouquet *et al.* 1983, Couillard *et al.* 2000), or to changes in secretion rate without changes in cell size or number (Heath *et al.* 1996).

Considering that the sheep pineal gland is composed primarily of pinealocytes (>80%; Arendt 1995), it is possible that the genetic differences in plasma melatonin concentration among ewes occur either because the larger pineal glands of the animals secreting more melatonin contains a greater pinealocyte number, or because pinealocytes are larger, or both.

Regarding the onset of melatonin secretion in life, although a night-time increase in plasma melatonin concentrations has been reported to occur in sheep fetuses (Yellon & Longo 1987, Zemdegs *et al.* 1988) as a direct consequence of melatonin secretion from the maternal pineal gland (Yellon & Longo 1988, McMillen & Nowak 1989) and in the newborn lambs by 1–6 weeks of age (Rodway *et al.* 1985, Claypool *et al.* 1999), it is not known whether the genetic difference in plasma melatonin concentrations previously described in adult ewes is already present soon after birth, or whether it is progressively acquired with age.

Thus, in the present study, we used two extreme groups of lambs genetically selected on the basis of their parent's plasma melatonin concentrations to: (1) determine how early in life the genetic variability in plasma melatonin concentrations is established and (2) investigate whether the genetic difference in melatonin secretion is related to a difference in the number of pinealocytes or in their size.

Materials and Methods

The experimental procedure reported in this study was carried out at the Institut National de la Recherche Agronomique, Research Center of Nouzilly, France, in accordance with the authorization for animal experimentation no. A37801 of the French Ministry of Agriculture.

Animals, blood sampling and slaughtering

The study was conducted with 39 spring-born male lambs (mean $\pm$ s.e.m birth date, 7 March $\pm$ 4 days; range, 3–14 March). The Ile-de-France breed used in this study results from crossbreedings performed in France between 1830 and 1900 between the English Dishley (Leicester) and Merinos Rambouillet (imported from Spain in the 18th century) breeds (Perret 1986). The Ile-de-France breed was chosen because it was known that the large variability in the nocturnal plasma melatonin concentrations among ewes (Chemineau *et al.* 1996) is under a strong genetic control (Zarazaga *et al.* 1998a,b). It is therefore a good model to study the genetic plasma melatonin variability in

mammals (Chemineau *et al.* 2001). The whole Ile-de-France flock from which the experimental lambs were obtained is a large flock of about 2500 animals which is divided into six different families. At regular intervals sires are purchased from various private external flocks and are introduced to prevent inbreeding and maintain genetic connections with the national French scheme of genetic improvement of the Ile-de-France breed.

The experimental lambs were chosen from about 400 lambs born at the same time of the year (mid March), in which the genetic value – defined as the sum of the average effects of the genes an individual carries (Falconer 1989) – was calculated on the basis of the endogenous nocturnal plasma melatonin concentration of their parents, previously determined at the June and December solstices (Zarazaga *et al.* 1998a). The 39 lambs were chosen and assigned before birth to two extreme groups referred to as Low ($n=21$) and High ($n=18$) groups, on the basis of differences in their genetic value. Lambs of the Low group were the progeny of nine sires and lambs of the High group were the progeny of 11 sires. Immediately after birth, lambs were removed from their mothers, maintained in artificial suckling with reconstituted milk until weaning at 7 weeks of age, and kept under a lighting regime of 16 h light : 8 h darkness (lights on from 0600 to 2200 h) throughout the experiment. Beginning at birth, and until lambs were slaughtered at an average of 14 weeks of age, lambs were weighed weekly to monitor their growth.

Blood (3 ml) was collected from all lambs at weekly intervals from 1 to 7 weeks of age, then at weeks 9, 11 and finally during the week of slaughtering at week 14 of age. At each sampling session, four blood samples were taken at hourly intervals during the night (from 2300 to 0200 h), and then during the following day (from 1000 to 1300 h). At night, blood samples were collected under a dim-red light (<1 lx at 20 cm) with care taken to avoid any direct illumination of the animal's eyes (using a rag to cover animal's eyes during sampling). All blood samples were obtained by venepuncture of jugular vein and plasma was immediately separated by centrifugation at 3000 r.p.m. for 20 min and stored at -20°C until assayed for melatonin.

Lambs were slaughtered at 14 weeks of age, during the day, and the brain, pineal and pituitary glands were removed and weighed rapidly. Length and width of the brain and length of hypothalamus were also measured. Carcass weight was recorded 24 h after slaughter. For each lamb in the two genetic groups, pineal gland was used for the morphometric study in order to determine the number and size of their pinealocytes.

Tissue preparation and morphometric analysis

Pineal glands were fixed in paraformaldehyde (4%) for 24 h, dehydrated in a graded series of alcoholic solutions, cleared in butanol, and embedded in paraffin wax. The

tissues were cut into 8 µm thick sections and stained with 0.1% cresyl violet.

Pineal volumes (V_0) were estimated by the Cavalieri principle (Gundersen 1986) according to the formula $V_0 = \sum ah$, where a is the area in µm² of every 10th section (8 µm thickness) and h is the distance in µm between the sections used to determine a .

Analysis of the number and individual area of pinealocytes was performed using the SAMBA 2005 Image Analyser (System for Analytical Microscopy in Biological Application, ALCATEL TITN Co., Massy, France). Light microscope images were input into the analyser through an objective with a $\times 40$ magnification, and a camera. A preliminary experiment, performed on a single pineal gland, demonstrated that the number and surface of pinealocytes were homogeneous between sections performed at various levels of the gland (same means and same variances). Thus, within each pineal, two sections, randomly selected at the level of the middle of the gland, were used in the morphometric analysis. On each section, 5350 µm² fields, selected according a grid covering the entire section, were analysed. Threshold cut-off for detection of pinealocytes was 10.4 µm². Accordingly, a total of approximately 1150 pinealocytes per pineal gland were counted and their surfaces measured. The total number of pinealocytes (n) from each pineal was estimated from the formula: $n = (\text{Number of pinealocytes per field} / \text{Volume of the field (µm}^3\text{)}) \times \text{Volume of the gland (µm}^3\text{)}$. The mean volume (µm³) for the pinealocytes from each pineal gland was determined, assuming that the shape of the pinealocyte was spherical.

Melatonin assay

Melatonin concentrations were quantified in duplicate aliquots of 100 µl of plasma by the RIA of Fraser *et al.* (1983) using an antibody first raised by Tillet *et al.* (1986). The sensitivity of the assay was 4 pg/ml. Mean intra-assay coefficient of variation, estimated by assaying three plasma pools (low, medium and high concentrations of melatonin) in duplicate every 100 unknown samples, was 5.8%. Mean inter-assay coefficient of variation for the same three plasma pools was 8.7% (two assays).

Statistical analysis

Statistical analyses of the data were performed by ANOVA (Super Anova, Statistical Software, Inc., Berkeley, CA, USA). For melatonin data, an effect of the genetic group and an effect of age was included in the analysis. All melatonin measurements were log-transformed, to correct for heterogeneity of variance, and plasma samples with values below the sensitivity of the RIA were arbitrarily assigned the limit of detection (4 pg/ml of plasma) for statistical analysis. The Pearson correlation coefficient R^2 was used to assess correlations among pineal parameters

and between pineal parameters and plasma melatonin concentrations, after combining the two groups of lambs. Multiple and step-to-step regressions (STATVIEW, Abacus Concept, Berkeley, CA, USA) were used for the descriptions of relationships among pineal weight and number and/or volume of pinealocytes. A value of 0.05 for P was taken as significant. Group data are presented as means $\pm$ S.E.M.s.

Results

Melatonin concentrations

A clear mean day/night (D/N) difference in plasma melatonin concentrations with higher values at night was evident by 1 week of age in the two genetic groups (mean $\pm$ S.E.M.; High group, D: 5.0 ± 0.4 vs N: 51.7 ± 10.7 pg/ml, $P < 0.01$; Low group, D: 4.9 ± 0.3 vs N: 31.9 ± 3.2 pg/ml, $P < 0.01$). From 1 week to slaughtering, no difference was found between the two genetic groups in mean daytime plasma melatonin (mean $\pm$ S.E.M.; 5.0 ± 0.2 pg/ml). However, ANOVA revealed a significant effect of the genetic group ($P < 0.05$) and of age ($P < 0.05$) on the mean night-time plasma melatonin concentrations, which were consistently lower in group Low compared with group High (Fig. 1). This difference was detected as soon as the first week after birth (Low: 31.9 ± 3.2 vs High: 51.7 ± 10.7 pg/ml) and was maintained until lambs were slaughtered at 14 weeks of age (Low: 125.0 ± 15.8 vs High: 162.9 ± 14.0 pg/ml).

Body growth and slaughtering measurements

No significant difference was observed on body growth between the two genetic groups. Body weight increased ($P < 0.01$) with age. The average weekly liveweight gain from birth to slaughtering was 2.5 ± 0.2 kg/week.

Among the brain parameters measured at slaughter (Table 1), only pineal weight was significantly different between the two genetic groups, with the mean pineal weight higher in the High group than in the Low group whereas no differences were detected between the High and Low groups for total brain weight, pituitary weight, length and width of the brain, and length of hypothalamus. Carcass weight of lambs did not differ between groups (21.5 ± 0.3 kg).

Pineal parameters

Morphometric analysis of pineal glands for the two genetic groups revealed that lambs in the High group had a significantly larger gland volume (High: 55.3 ± 5.5 vs Low: 39.9 ± 4.6 mm³; $P < 0.05$) and a significantly higher total number of pinealocytes (High: 27.8 ± 2.4 vs Low: $21.0 \pm 2.4 \times 10^6$; $P < 0.05$) compared with those in the

Figure 1 Mean $\pm$ S.E.M. nocturnal plasma melatonin concentrations from genetically extreme lambs (High, filled bars, $n=18$ and Low, open bars, $n=21$), sampled four times during the night, at weekly intervals, from 1 to 7 weeks of age, and at 9, 11 and 14 weeks of age. * indicates significant difference between groups ($P<0.05$).

Low group (Fig. 2). In contrast, no significant difference was found on mean individual volume of pinealocytes between the High and Low groups (185.5 ± 9.9 and $166.3 \pm 6.8 \mu\text{m}^3$ respectively). An illustration of the pinealocytes is given in Fig. 3.

Pineal weight was highly correlated with pineal volume ($R^2=0.86$, $P<0.001$), with total number of pinealocytes ($R^2=0.79$, $P<0.001$), and with volume of pinealocyte ($R^2=0.24$, $P<0.01$; Fig. 4). Multiple regression indicated that the number of pinealocytes ($P<0.0001$) and the volume of pinealocytes ($P<0.05$), when combined, significantly contributed to explain more than 80% of the total variance of the pineal weight. Step-to-step regression analysis revealed that about 63% was due to pineal weight and 17% to the volume of pinealocytes. Mean nocturnal plasma melatonin concentrations at slaughtering, at 14 weeks of age, were also significantly correlated with the number of pinealocytes ($R^2=0.29$, $P<0.001$), with pineal volume ($R^2=0.34$, $P<0.001$) and with the weight of

the pineal gland ($R^2=0.24$, $P<0.01$). No significant correlation was detected between plasma melatonin concentrations and pinealocyte size.

Discussion

The present study clearly shows that the two groups of extreme lambs differed by the number of pinealocytes that were counted within their pineal gland, rather than by the size of their pinealocytes. At slaughter, at 14 weeks of age, the lambs of the High group, selected on the basis of the plasma melatonin concentration of their parents, had 32% more pinealocytes in their pineal gland than the lambs of the Low group. Conversely, the size of the pinealocytes did not differ between the two groups.

However, when pooling the two genetic groups, a significant correlation between volume of pinealocytes and pineal weight existed. This apparent contradiction with

Table 1 Mean values ($\pm$ S.E.M.) of brain parameters measured at slaughtering at 14 weeks of age in the Low ($n=21$) and High ($n=18$) Ile-de-France lamb groups

Parameter	Low group (n=21)	Difference between groups	High group (n=18)
Pineal weight (mg)	88.1 $\pm$ 7.7	P<0.05	118.1 $\pm$ 9.9
Pituitary weight (mg)	471.9 $\pm$ 27.3	N.S.	443.7 $\pm$ 31.6
Brain			
Total weight (g)	75.0 $\pm$ 0.9	N.S.	74.4 $\pm$ 1.1
Length (mm)	52.4 $\pm$ 0.8	N.S.	52.3 $\pm$ 0.5
Width (mm)	42.1 $\pm$ 0.6	N.S.	42.2 $\pm$ 0.4
Hypothalamus length (mm)	11.1 $\pm$ 0.2	N.S.	11.5 $\pm$ 0.3

Figure 2 Pineal weight (a), pineal volume (b), number of pinealocytes (c) and volume of pinealocytes (d) from genetically extreme lambs (High, filled bars, $n=18$ and Low, open bars, $n=21$) slaughtered at 14 weeks of age. Data are means $\pm$ S.E.M.s.

the results obtained when comparing the two genetic groups may indicate that pineal weight of lambs also partly depends on the volume of pinealocytes. By referring to the multiple and step-to-step regression analyses, we can assume that the variation in the number of pinealocytes

Figure 3 Illustrative examples of pinealocytes from genetically extreme lambs (High, $n=18$ and Low, $n=21$) slaughtered at 14 weeks of age. Bar=20 μm .

may explain about 80%, and the volume of pinealocytes about 20%, of the total variation in pineal weight.

Thus, the difference between the two genetic groups of extreme lambs regarding their pineal weight at slaughter, already found in a previous experiment (Coon *et al.* 1999), originated from a higher multiplication rate of pinealocytes rather than an enlargement of the same number of cells. This difference in number of pinealocytes was observed at slaughter, at 14 weeks old. As attested by the substantial increase in mean melatonin plasma concentration from 1 to 14 weeks of age in the present study, already observed in previous publications (Claypool *et al.* 1989), it is highly probable that pineal glands of young-born lambs continue their development in the first weeks of life. We can wonder if the difference between the two groups regarding the number of pinealocytes was already present at birth, or if the difference is acquired progressively during the first weeks of life. The existence, in the present experiment, of a difference in plasma melatonin concentrations as early as less than 1 week after birth, strongly suggests that the lambs of the High group had a higher

Figure 4 Individual relationship between number of pinealocytes and pineal weight (upper graph) and volume of pinealocytes and pineal weight (lower graph) in genetically extreme lambs (both groups of lambs were combined) slaughtered at 14 weeks of age.

number of pinealocytes at birth, compared with the number present in the Low group.

As the difference between these two groups is due to the effects of alleles of specific genes (because animals were selected on the performances of their parents), we can suggest that the genes controlling the multiplication of pinealocytes probably act during embryonic or fetal development, when the pineal gland is at the beginning of its development. In vertebrates, the pineal develops from the diencephalon, and in the human species has been identified as soon as the second month of gestation, when the embryo is 6–9 mm long (Binkley 1988). Unfortunately, the identification of these genes will probably not be easy because the specific alleles producing the difference between groups may be expressed and act during a limited period of time during embryonic or fetal development. This was demonstrated

for homeobox genes (Gehring 1993), for example, by modifying the yield of multiplication of the stem cells at the beginning of pineal differentiation.

Interestingly, in our study, only pineal weight was different between the two extreme groups of lambs, whereas the other simple brain measurements, such as brain weight, pituitary weight, or brain size, did not differ between the two groups of animals. This may suggest that the specific genes controlling multiplication of pinealocytes could be specific to the pineal gland or the diencephalon, and may not be involved in the general development of the whole brain. However, as these structures are not linked embryologically with the development of the pineal gland, we cannot fully exclude that the genetic differences in pineal size are not associated with other changes in brain characteristics; more detailed measurements on structures linked to the diencephalon would be necessary to ascertain this conclusion.

The absence of observed difference in body growth between the two groups of lambs raises the question of the biological consequence(s) of this genetic variability in pineal weight/melatonin secretion. To our knowledge, only a few studies have addressed this question. Regarding seasonal breeding, a trait known to be under melatonin control, in a limited number of ewes it was suggested that the relative day/night melatonin ratio could be related to the date of onset of the ovulatory activity (Chemineau *et al.* 1993, Zarazaga *et al.* 1996), but this relationship needs to be confirmed on a larger set of ewes. In other species, studies were more focused on duration of melatonin secretion than on amplitude, with the exception of the relationship between melatonin amplitude and ageing in humans (Touitou *et al.* 1981, Sack *et al.* 1986, Wetterberg *et al.* 1993, Kripke *et al.* 1998, Touitou 2001) and primates (Aujard *et al.* 1998, Roth *et al.* 2001). In this latter case, melatonin amplitude was considered as a good marker of ageing processes. In this regard, sheep may constitute a good model for the genetic control of plasma melatonin in mammals.

In conclusion, these results show for the first time (a) that the genetic differences previously observed in adult sheep are detected in young lambs as soon as the first week of life, and (b) that a high secretion of melatonin is due directly to a hyperplasia of the pineal gland that is associated with an increase in pinealocyte number. The identification of genes controlling pineal size (which is in progress in our laboratory) could be of interest for other mammalian species, including humans.

Acknowledgements

A.G.B. was supported by a postdoctoral grant from INIA and INRA-DSPA. The authors thank Loys Bodin (Saga Toulouse) for the estimation of the genetic values of lambs, Jean Voisin and Christian Moussu for slaughtering the

lambs, the breeders of the INRA Research Center of Nouzilly for the supply and care of experimental animals, and the Laboratory of Anatomopathology of Veterinary Faculty of Complutense University in Madrid.

References

- Arendt J 1995 *Melatonin and the Mammalian Pineal Gland*, London: Chapman & Hall.
- Aujard F, Boissy I & Claustrat B 1998 Melatonin secretion in a nocturnal prosimian primate: effect of photoperiod and aging. In *Biological Clocks. Mechanisms and Applications*, pp 337–340. Ed. Y Touitou. Paris: Elsevier Science BV.
- Binkley S 1988 *The Pineal. Endocrine and Nonendocrine Function*, p. 304. Englewood Cliffs, NJ: Prentice Hall.
- Bittman EL, Dempsey RL & Karsch FJ 1983 Pineal melatonin secretion drives the reproductive response to daylength in the ewe. *Endocrinology* **113** 2276–2283.
- Chemineau P, Maurice F & Daveau A 1993 Re-initiation of ovulatory activity by melatonin given as a constant-release implant in long-day treated Ile-de-France ewes, depends on endogenous secretion of melatonin. In *Melatonin and the Pineal Gland – from Basic Science to Clinical Application*, pp 247–250. Eds Y Touitou, J Arendt & P Pévet. Paris: Elsevier Science Publishers BV.
- Chemineau P, Beltran De Heredia I, Daveau A & Bodin L 1996 High repeatability of the amplitude and duration of the nycthemeral rhythm of the plasma melatonin concentrations in the Ile-de-France ewes. *Journal of Pineal Research* **21** 1–6.
- Chemineau P, Daveau A, Bodin L, Zarazaga L, Gomez-Brunet A & Malpoux B 2001 The sheep as a mammalian model of genetic plasma melatonin variability. *Reproduction* **59** (Suppl) (In Press).
- Claypool LE, Wood RI, Yellon SM & Foster DL 1989 The ontogeny of melatonin secretion in the lamb. *Endocrinology* **124** 2135–2143.
- Coon SL, Zarazaga LA, Malpoux B, Ravault JP, Bodin L, Voisin P, Weller JL, Klein DC & Chemineau P 1999 Genetic variability in plasma melatonin in sheep is due to pineal weight, not to variations in enzyme activities. *American Journal of Physiology* **277** (Endocrinology and Metabolism 40) E792–E797.
- Couillard C, Mauriege P, Imbeault P, Prud'homme D, Nadeau A, Tremblay A, Bouchard C & Despres JP 2000 Hyperleptinemia is more closely associated with adipose tissue cell hypertrophy than with adipose tissue hyperplasia. *International Journal of Obesity and Related Metabolic Disorders* **24** 782–788.
- Falconer DS 1989 *Introduction to Quantitative Genetics*, edn 3, p. 438. Harlow, UK: Longman.
- Fouquet JP, Meusy-Dessolle N & Dang DC 1983 Morphometry of fetal Leydig cells in the monkey (*Macaca fascicularis*), correlation with plasma testosterone. *Biology of the Cell* **49** 267–272.
- Francis SM, Venters SJ, Duxson MJ & Suttie JM 2000 Differences in pituitary cell number but not cell type between genetically lean and fat Coopworth sheep. *Domestic Animal Endocrinology* **18** 229–239.
- Fraser S, Cowen P, Franklin M, Franey C & Arendt J 1983 Direct radio-immunoassay for melatonin in plasma. *Clinical Chemistry* **20** 396–397.
- Gehring WJ 1993 Exploring the homeobox. *Gene* **135** 215–221.
- Gundersen HJG 1986 Stereology of arbitrary particles: a review of unbiased number and size estimators and the presentation of some new ones. *Journal of Microscopy* **143** 3–45.
- Heath DA, Caldani M & McNatty KP 1996 Relationship between the number of immunostaining gonadotropes and the plasma concentrations of gonadotrophins in ewes with and without the FecBB gene. *Journal of Reproduction and Fertility* **106** 73–78.
- Kappers JA & Pevet P 1979 The pineal gland of vertebrates including man. In *Progress in Brain Research*, vol 52, p. 562. Eds JA Kappers & P Pevet. Amsterdam: Elsevier.
- Kripke DF, Elliot JA, Youngstedt SD & Smith JS 1998 Melatonin: marvel or marker? *Annals of Medicine* **30** 81–87.
- Malpoux B, Robinson J, Brown MB & Karsch FJ 1987 Reproductive refractoriness of the ewe to inductive photoperiod is not caused by inappropriate secretion of melatonin. *Biology of Reproduction* **36** 1333–1341.
- Malpoux B, Moenter SM, Wayne NL, Woodfill CJI & Karsch FJ 1988 Reproductive refractoriness of the ewe to inhibitory photoperiod is not caused by alteration of the circadian secretion of melatonin. *Neuroendocrinology* **48** 264–270.
- Mcmillen IC & Nowak R 1989 Maternal pinealectomy abolishes the diurnal rhythm in plasma melatonin concentrations in the fetal sheep and pregnant ewe during late gestation. *Journal of Endocrinology* **120** 459–464.
- Perret G 1986 *Races Ovines*, p. 440. Paris: SPEOC.
- Rodway RG, Swift AD, Nowak R, Smith TA & Padwick D 1985 Plasma concentrations of melatonin and the onset of puberty in female lambs. *Animal Reproduction Science* **8** 241–246.
- Roth GS, Lesnikov V, Lesnikov M, Ingram DK & Lane MA 2001 Dietary caloric restriction prevents the age-related decline in plasma melatonin levels of rhesus monkeys. *Journal of Clinical Endocrinology and Metabolism* **86** 3292–3295.
- Sack RL, Lewy DL, Erb DL, Vollmer WM & Singer CM 1986 Human melatonin production decreases with age. *Journal of Pineal Research* **3** 379–388.
- Shimokawa I, Yu BP, Higami Y & Ikeda T 1996 Morphometric analysis of somatotrophs: effects of age and dietary restriction. *Neurobiology of Aging* **17** 79–86.
- Tillet Y, Ravault JP, Selve C, Evain G, Castro B & Dubois MP 1986 Conditions d'utilisation d'anticorps spécifiques pour la visualisation immunohistochimique de la sérotonine et de la mélatonine dans la glande pinéale du mouton. *Compte Rendus de l'Académie des Sciences, Paris (III)* **303** 77–82.
- Touitou Y 2001 Human aging and melatonin. Clinical relevance. *Experimental Gerontology* **36** 1083–1100.
- Touitou Y, Ferve M, Lagouey M, Carayon A, Bogdan A, Reinberg A, Beck H, Cesselin F & Touitou C 1981 Age- and mental-health related circadian rhythms of plasma levels of melatonin, prolactin, luteinizing hormone and follicle-stimulating hormone in man. *Journal of Endocrinology* **91** 467–475.
- Vollrath L 1981 *The Pineal Organ*, pp 665. Ed. L Vollrath. Berlin: Springer-Verlag.
- Vollrath L & Welker HA 1984 Pineal gland size and serum melatonin levels in male rats are not correlated. *Neuroscience Letters* **49** 25–28.
- Wetterberg L, Iselius L & Lindsten J 1983 Genetic regulation of melatonin excretion in urine. *Clinical Genetics* **24** 399–402.
- Wetterberg LG, Eberhard L, Von Knorring MA, Kohan E & Lippa Y 1993 The influence of age, sex, height, urine volume and latitude on melatonin concentrations in urine; multinational study. In *Light and Biological Rhythms in Man*, pp 275–286. Ed. L Wetterberg. Oxford: Pergamon Press.
- Yellon SM & Longo LD 1987 Melatonin rhythms in fetal and maternal circulation during pregnancy in the sheep. *American Journal of Physiology* **252** E799–E802.
- Yellon SM & Longo LD 1988 Effect of pinealectomy and reverse photoperiod on the circadian melatonin rhythm in the sheep and the fetus during the late trimester of pregnancy. *Biology of Reproduction* **39** 1093–1099.
- Zarazaga LA, Forcada F, Abecia JA & Lozano JM 1996 Date of reinitiation of the breeding season could be related with relative changes in plasma melatonin amplitude in ewes. In *Pineal Update – from Molecular Mechanisms to Clinical Implications*, pp 295–300. Eds SM Webb, M Puig-Domingo, M Möller & P Pévet. New York: PJD Publications Ltd.

Zarazaga LA, Malpaux B, Bodin L & Chemineau P 1998a The large variability in melatonin blood levels in ewes is under strong genetic influence. *American Journal of Physiology* **274** (*Endocrinology & Metabolism* 37) E607–E610.

Zarazaga LA, Malpaux B, Guillaume D, Bodin L & Chemineau P 1998b Genetic variability in melatonin concentrations in ewes originates in its synthesis, not in its catabolism. *American Journal of Physiology* **274** (*Endocrinology & Metabolism* 37) E1086–E1090.

Zemdegs IZ, Mcmillen IC, Walker DW, Thorburn GD & Nowak R 1988 Diurnal rhythms in plasma melatonin concentrations in the fetal sheep and pregnant ewe during late gestation. *Endocrinology* **123** 284–289.

Received 27 July 2001

Accepted 9 October 2001