Calcium balance in sea bream (*Sparus aurata*): the effect of oestradiol-17 β

P M Guerreiro, J Fuentes, A V M Canario and D M Power

Centre of Marine Sciences, University of Algarve, 8000–810 Faro, Portugal

(Requests for offprints should be addressed to J Fuentes, Centro de Ciências do Mar (CCMar), Universidade do Algarve, Campus de Gambelas, 8000–810 Faro, Portugal; Email: jfuentes@ualg.pt)

(P M Guerreiro and J Fuentes contributed equally to this work)

Abstract

In all teleost fishes vitellogenesis is triggered and maintained by oestradiol-17 β (E₂) and is accompanied by an increase of blood plasma calcium and phosphate. The action of this hormone on calcium metabolism was investigated by treating fast-growing immature juvenile sea bream (*Sparus aurata*) with coconut butter implants alone (control) or implants containing 10 $\mu\text{g/g}$ E₂. Treatment with E₂ induced the production of circulating vitellogenin, a 2.5-fold increase in plasma ionic Ca²⁺ and a 10-fold increase in plasma total calcium, largely bound to protein. In contrast to freshwater species, which obtain most of their calcium from the environment directly through the gills, the intestinal component of calcium uptake of the salt water-living sea bream represented up to 60–70% of the total uptake. The whole body calcium uptake, expressed as the sum of calcium obtained via intestinal and

extra-intestinal (likely branchial) routes increased significantly in response to E₂. Combined influx and unchanged efflux rates resulted in a significant 31% increase in net calcium uptake. There was no evidence for an effect of E₂ on the calcium and phosphate content of the scales or the tartrate-resistant acid phosphatase activity (an index for bone/scale osteoclast activity). While most freshwater fish appear to rely on internal stores of calcium, i.e. bone and/or scales to increase calcium availability, the marine sea bream accommodates calcium-transporting mechanisms to obtain calcium from the environment and preserve internal stores. These observations suggest that a fundamental difference may exist in the E₂-dependent calcium regulation between freshwater and marine teleosts.

Journal of Endocrinology (2002) **173**, 377–385

Introduction

Calcium balance in fish has been extensively studied in freshwater species, mostly in salmonids and cichlids (for review see Flik *et al.* 1995). The majority of the calcium transport mechanisms are proposed to take place actively at the level of the gills (Flik *et al.* 1995) and, at least in some species, partly through the skin (McCormick *et al.* 1992, Marshall *et al.* 1992). In fresh water, fish drink very little and as a consequence virtually all the calcium incorporated via the intestine is of dietary origin. Thus, if dietary calcium is low, calcium requirements can be compensated by uptake via the gills from the surrounding medium (Ichii & Mugiya 1983). Seawater fish, in contrast to freshwater fish, ingest large amounts of seawater to compensate for the osmotic water loss. Seawater contains around 10 mMolar calcium and thus drinking of seawater may provide a significant additional mode of calcium uptake. Although the contribution of the intestine to whole body calcium balance remains little studied in seawater species, there is some evidence that a varying proportion of the

calcium imbibed may be absorbed by the intestine and this proportion may be species-specific (Schoenmakers *et al.* 1993, Sundell *et al.* 1993, Larsson *et al.* 1995).

Oestrogen treatment, administered usually in the form of oestradiol-17 β (E₂), is known to induce an increase in circulating levels of vitellogenin in fish (Bradley & Grizzle 1989, Burzawa-Gerard & Dumas-Vidal 1991, Mosconi *et al.* 1998), which is paralleled by an increase in blood plasma calcium (Persson *et al.* 1994, 1995, 1997). This increase in calcium demand during the reproductive period in freshwater fish is met not only from external sources but also by mobilisation of internal calcium stores, such as bone and scales which, in tilapia, account for up to 95% of the total body calcium (Flik *et al.* 1986). In goldfish, killifish (Mugiya & Watabe 1977) and rainbow trout (Carragher & Sumpter 1991, Persson *et al.* 1994, 1995, 1997), bone is initially spared and calcium mobilisation takes place predominantly from the scales. Moreover, in juvenile rainbow trout the effect of oestrogen at the level of the scales appears to be via high-affinity, low-capacity E₂-binding sites (Persson *et al.* 2000) and

oestrogen receptor mRNA is expressed in this tissue (Armour *et al.* 1997).

In the present study, the effect of E_2 on plasma calcium in a marine teleost, *Sparus aurata*, was investigated and the mechanism which increases internal calcium availability was also determined. The specific aims of the present study were: first, to establish the intestinal and extra-intestinal contribution to whole body calcium uptake in seawater, secondly, to determine the changes in calcium metabolism in response to E_2 and, thirdly, to evaluate the relative importance of internal and external calcium sources during periods of high demand such as vitellogenesis.

Materials and Methods

Fish

Immature sea bream (*Sparus aurata*) juveniles (1+ year old, body mass 31.0 ± 0.9 g) were obtained from a stock raised from fertilised eggs at Ramalhete Marine Station (University of Algarve) and maintained under natural annual conditions of water temperature ($17\text{--}25^\circ\text{C}$; 18°C during the experiments), photoperiod and salinity (36–40‰; 37‰ during the experiments).

E_2 treatment and sampling

In April, juvenile sea bream ($n=60$) were randomly distributed between two tanks and acclimated for 2 weeks prior to the start of the experiment. At the start of the experiment fish were anaesthetised with 2-phenoxyethanol (1:10 000 2-phenoxyethanol:seawater; Sigma-Aldrich, Madrid, Spain), one group ($n=30$) received a peritoneal implant of coconut butter (10 $\mu\text{l/g}$ body weight; Sigma-Aldrich) containing 10 $\mu\text{g/g}$ body weight E_2 (Sigma-Aldrich) and the other group ($n=30$) received the coconut butter alone (control). This E_2 dose induces a chronic increase in the circulating levels of plasma E_2 and vitellogenin in the sea bream and other teleosts (Persson *et al.* 1997, Mosconi *et al.* 1998). Control and E_2 -treated fish were then distributed between two flow-through seawater tanks per treatment, each with 15 fish, and left undisturbed for 15 days. During this period, fish were fed daily at a rate of 2% of body weight/day on commercial sea bream pellets (Provimi, Faro, Portugal) between 1000 and 1100 h, except that food was withheld 24 h prior to and on the day of the experiments and sampling. No deaths occurred during the experiment. At day 15, fish ($n=7$) from both groups were sampled under anaesthesia (1:10 000 2-phenoxyethanol:seawater). Each fish was weighed to the nearest 0.1 g and subsequently a 0.3 ml blood sample was collected by puncture of the caudal vessels into heparinised (ammonium heparin, 30 units/ml; Sigma-Aldrich) 2 ml syringes fitted with 25

gauge needles. The puncture area was cleaned with tissue paper before blood was withdrawn to avoid contamination of the blood samples with seawater. Plasma was obtained by centrifugation of whole blood (10 000 r.p.m. for 5 min), aliquoted into four (1.5 ml) vials, snap-frozen in liquid N_2 and stored at -80°C for later analysis.

After collection of blood samples, fish were killed by decapitation and immediately blotted dry, the remaining mucus adhering to the scales was removed with a scalpel and the scales covering the lateral line collected, snap-frozen in liquid N_2 and stored at -80°C for subsequent analysis. Livers were removed and weighed for calculation of the hepato-somatic index: $\text{HSI} = (\text{liver weight/body weight}) \times 100$.

Whole body calcium uptake

For determinations of whole body calcium uptake, control ($n=7$) and E_2 -treated ($n=7$) fish were transferred to individual seawater tanks (1.5 litres/fish) set up with closed circuit water flow with aeration. Fish were left for 30 min and then 37 kBq/l ^{45}Ca as $^{45}\text{CaCl}_2$ (specific activity 14.8 MBq/ μmol ; NEN, Life Sciences Products, Boston, USA) and 37 kBq/l ^{51}Cr as $^{51}\text{Cr-EDTA}$ (specific activity 96.2 MBq/ μmol ; NEN, Life Sciences Products) were added to the tank water and allowed to mix. The tanks were partially darkened with a black lid and fish were left to swim undisturbed. Exactly 2 h after the addition of the tracers, triplicate 2 ml water samples were collected for counting and the radioactive solution was immediately replaced by tracer-free seawater. The fish were left to swim in this solution for 10 min to prevent any radioactivity from adhering to the body surfaces and then killed with an overdose of 2-phenoxyethanol (1:250 2-phenoxyethanol:seawater), blotted dry, quickly weighed to the nearest 0.1 g and frozen at -20°C . The whole frozen intestine, from oesophagus to rectum, from individual fish was subsequently dissected out. The intestine and carcass were then transferred separately to pre-weighed glass tubes and digested with nitric acid (1:10 of the sample weight) at 30°C for 1 week. After digestion, samples were neutralised with equal volumes of 2 M NaOH, and a 2 ml aliquot transferred to a scintillation counting vial. All samples, including tank water, digested intestines and carcasses were bleached with 2 ml 35% hydrogen peroxide (Fluka; Sigma-Aldrich), to prevent colour quenching, incubated overnight at room temperature and then counted for 15 min in a Beckman LS6000IC scintillation counter after the addition of 20 ml scintillation cocktail (OptiPhase HiSafe II, Wallac; Amersham Pharmacia Biotech). Total ^{51}Cr was measured in 2 ml aliquots of tank water and in digested intestines by counting for 15 min in a Wallac 1470 Wizard gamma counter (Amersham Pharmacia Biotech).

Whole body calcium uptake (i.e. calcium influx) was calculated according to the following formula: $\text{CI} =$

$(Af \times Cw)/(Aw \times t \times w)$, where Af is the total ^{45}Ca activity in the fish carcass (i.e. excluding the activity in the whole intestine which can be considered to be outside of the fish), Cw is the total calcium concentration in the water, Aw is total activity in the water, t is duration of exposure (h) and w is fish wet weight (g). Whole body calcium influx rate is expressed as nmol/h per g. The total water calcium content was measured in triplicate (Sigma-Aldrich procedure no. 587).

Intestinal calcium uptake

The combination of whole body calcium influx and the amount of water imbibed (i.e. drinking rate) allows for estimation of the contribution of intestinal and extra-intestinal uptake routes to whole body calcium uptake. Drinking rates (DR) were calculated as: $DR = Af/(Aw \times t \times w)$, where Af is the total activity of ^{51}Cr -EDTA in the intestine, Aw is the tracer concentration (c.p.m./ml) in water, t is the duration of exposure (h) and w is fish wet weight (g). Results are expressed as $\mu\text{l/h}$ per g. Using the same expression for ^{45}Ca , another estimate, the 'drinking rate', can be calculated. However, there is a crucial difference in the behaviour of the two tracers; while ^{51}Cr -EDTA is not absorbed, ^{45}Ca crosses the intestinal wall. The difference in the estimated drinking rates using both tracers provides the amount of calcium imbibed which is actually absorbed by the intestine. The intestinal contribution to whole body calcium uptake (Int_{Ca}) can be calculated as: $\text{Int}_{\text{Ca}} = (\text{DR}_{\text{Cr}} - \text{DR}_{\text{Ca}}) \times W_{\text{Ca}}$, where DR_{Cr} is the drinking rate calculated from ^{51}Cr -EDTA values ($\mu\text{l/h}$ per g), DR_{Ca} is the drinking rate calculated from ^{45}Ca values ($\mu\text{l/h}$ per g) and W_{Ca} is the water calcium concentration (nmol/ μl). Results are expressed as nmol/g per h. The difference between Int_{Ca} and the value for whole body calcium uptake estimates the contribution of the intestinal and extra-intestinal routes to the whole body calcium uptake.

Whole body calcium efflux

Control ($n=7$) and E_2 -treated fish ($n=7$) were loaded with ^{45}Ca by intraperitoneal injection of 37 kBq $^{45}\text{CaCl}_2$ dissolved in 500 μl 0.9% NaCl and transferred to separate tanks with flowing seawater for 48 h for recovery and to achieve a constant specific activity of the readily available calcium pool in the whole fish. Fish were then randomly and individually housed in 0.75 litre tanks and 1 ml water samples were taken after 30 min and every 20 min thereafter for 2 h. Fish were then killed by an overdose of anaesthetic (1:250 2-phenoxyethanol), weighed, a blood sample collected and plasma separated as described earlier. Both water and plasma samples were measured for ^{45}Ca radioactive decay as previously described. Calcium efflux rate (ER) was calculated as follows: $ER = (Aw \times Ct)/(At \times t \times w)$, where Aw is the final specific activity in

water, Ct is the total plasma calcium (mMol), At is the specific activity of ^{45}Ca in the plasma, t is time (h) and w is fish wet weight (g). Results are expressed as nmol/h per g.

E_2 assay

Individual plasma samples (50 μl) were extracted twice with diethyl ether (80% E_2 recovery), resuspended in phosphate buffer containing 0.5 g/l gelatine, pH 7.6 and radioimmunoassay was carried out with specific antibodies (Research Diagnostics, USA) following the general methods described by Scott *et al.* (1982). Cross-reactivity of the antibody (expressed as %) was as follows: 4-pregnene-3,20-dione <0.2; 11 β ,17,21-trihydroxy-4-pregnene-3,20-dione <0.2; 4-androstene-3,17-dione <0.2; 17 β -hydroxy-4-androsten-3-one <0.2; 3 β -hydroxy-5-pregnen-20-one <0.2; 3 β -hydroxy-5-androsten-17-one <0.2; 3-hydroxy-1,3,5(10)-estratrien-17-one = 15; 3,17 β -dihydroxy-1,3,5(10)-estratrien-16-one = 8; 3,16 α ,17 β -trihydroxy-1,3,5(10)-estratrien-3-one = 0.7; 3,16 α -dihydroxy-1,3,5(10)-estratrien-17-one <0.2. All samples were assayed in duplicate in a single assay. Inter- and intra-assay coefficients of variation determined were 8.6% ($n=3$) and 4.01% ($n=5$) respectively.

Protein analysis

Plasma protein Total plasma protein was measured in diluted samples (1:100) using the method of Lowry *et al.* (1951) with minor modifications for a micro plate reader (Benchmark; BioRad, Hercules, CA, USA). Bovine serum albumin (Sigma-Aldrich) was used as the standard. Results are given as mg/ml plasma.

Vitellogenin Western blotting was used to confirm the identity of the plasma protein putatively designated as vitellogenin using a sea bream-specific vitellogenin polyclonal antibody. Diluted plasma (12.5 μg protein per sample) was mixed with an equal volume of sample buffer (6% (w/v) SDS, 6% (v/v) 2-mercaptoethanol, 40% (w/v) sucrose, 0.02% bromophenol blue in 0.125 M Tris-HCl, pH 6.8) boiled for 5 min, spun (30 s, 12 000 r.p.m.) and fractionated by SDS-PAGE according to Laemmli (1970). Molecular weight markers (Sigma-Aldrich) were run on all the gels to assist protein size identification. Gels were lightly stained with Coomassie blue (0.025%) and images captured with an image analysis system (Image Master VDS system; Amersham Pharmacia Biotech). Fractionated plasma proteins were transferred to a nitrocellulose membrane (NCP, Amersham Pharmacia Biotech) by electroblotting, using Tris-lysine (pH 8.2) as the transfer buffer. Immunoblotting of membranes was carried out by blocking non-specific binding in MTw (0.1 M Tris, 2% milk powder and 0.05% Tween 20) for 3 h, rinsing in Tris (0.1 M), followed by incubation overnight at 4 °C with

anti-sea bream vitellogenin serum (1:7000). Membranes were then washed with MTw and incubated with anti-rabbit IgG complexed with peroxidase (1:2000) for 1 h at room temperature with continuous agitation. After rinsing in Tris (0.1 M), membranes were developed using 4-chloro-1-naphthol as the chromogen.

Plasma calcium analysis

Individual total plasma calcium (bound and free) and plasma calcium activity were measured in triplicate ($n=14$ controls; $n=14$ E_2 -implanted). Total plasma calcium was measured by colorimetric assay (Sigma-Aldrich procedure no. 587). Free calcium was measured by means of calcium-sensitive electrodes (MI-600; Microelectrodes Inc., Bedford, MD, USA) using an Ag-AgCl electrode (MI-409; Microelectrodes Inc.) as an external reference. Plasma phosphorus was determined in triplicate in pre-diluted plasma using an endpoint colorimetric assay (Sigma-Aldrich procedure no. 360).

Analysis of scales

Calcium and phosphate scale content Frozen scales (20–30 mg) collected from individual fish were weighed, placed in glass tubes and digested with 1 ml nitric acid for 24 h at 30 °C. The resulting digest was neutralised with equal volumes of 2 M NaOH and the volume in each tube increased to 5 ml with double-deionised water. Total calcium and phosphate in the digests were measured in triplicate using the colorimetric methods described in the previous sections. Results are expressed as μmol calcium or phosphate/mg scale tissue.

Tartrate-resistant acid phosphatase (TRAP) To examine the changes in TRAP activity, 30–40 mg frozen scales from individual fish ($n=7$ per treatment) were homogenised at 4 °C using a glass–glass manual homogeniser in 1 ml of a 0.1 M sodium acetate buffer (pH 5.0). Homogenates were kept on ice and aliquots incubated in triplicate for 30 min at 22 °C, in acetate buffer (0.1 M, pH 5.3) in the presence of 10 mM para-nitrophenyl phosphate (Sigma-Aldrich) and 20 mM tartrate. Para-nitro phenol (pNP; Sigma-Aldrich) was used as the standard. Enzyme activity measurements were performed according to the optimal assay conditions for TRAP measurements for fish scales (Persson *et al.* 1995) and shown to be similar to those reported for mammalian bone tissue (Lau *et al.* 1985).

Statistics

Values are shown as means $\pm$ S.E.M. unless otherwise stated. Differences between groups were established by one-way

analysis of variance (ANOVA) after assessing normality and homogeneity of variances. Groups were considered significantly different at $P<0.05$.

Results

Effect of E_2 implants on plasma E_2 , calcium and phosphate

Fifteen days after implantation, control sea bream had circulating plasma E_2 levels of 0.49 ± 0.18 ng/ml while in the E_2 -implanted group, levels had increased 14-fold to 7.18 ± 1.89 ng/ml (Fig. 1A).

Total plasma calcium increased tenfold in response to E_2 treatment, reaching 29.83 ± 3.09 mmol/l compared with 2.84 ± 0.23 mmol/l in control fish. Levels of ionic calcium were 1.46 ± 0.20 mmol/l in control fish (51.4% of the total calcium) while, in E_2 -treated fish, ionic calcium was 3.71 ± 0.21 mmol/l (decreasing to only 12.4% of total). Despite the significant increase in both total and ionic calcium in the E_2 -treated group compared with the control group, only a relatively small proportion of the calcium remained in an ionic form and the majority (87.5%) was present in the protein-bound fraction of the total circulating calcium (Fig. 1B).

In parallel with increases in plasma calcium, E_2 implants also caused a significant increase in plasma phosphate levels ($P<0.01$, Fig. 1C) from 1.33 ± 0.24 mmol/l in controls to 8.01 ± 1.56 mmol/l in E_2 -treated fish.

Drinking and intestinal calcium absorption

The drinking rate measured by ^{51}Cr -EDTA intake was unaffected by E_2 treatment (Table 1) and was $5.2\text{--}5.3$ $\mu\text{l/h}$ per g in both groups. However, the estimates of drinking based on ^{45}Ca accumulation in the intestine were lower, ranging from 0.36 to 0.49 $\mu\text{l/h}$ per g. The difference between the two measurements was used to determine the proportion of calcium in the drinking water that was absorbed by the intestine, $89.3 \pm 1.4\%$ for controls and $92.8 \pm 0.8\%$ for E_2 -implanted fish (Table 1).

Whole body calcium uptake

Whole body calcium uptake increased significantly in response to E_2 (Fig. 2A) from 64.1 ± 1.8 nmol/h per g in control fish to 72.5 ± 1.6 nmol/h per g in E_2 -implanted fish ($P<0.01$). The intestinal component of the whole body calcium uptake was $64.9 \pm 2.1\%$ in the control group and $69.5 \pm 3.3\%$ in the E_2 -implanted sea bream group (Fig. 2A).

Calcium efflux and net calcium balance

No differences in efflux were found between consecutive 20-min periods in control ($P=0.837$) or E_2 -treated

Figure 1 (A) Circulating plasma levels (ng/ml) of E_2 in sea bream juveniles 15 days after implantation of $10 \mu\text{g/g}$ E_2 in coconut butter implants (E_2) or coconut butter alone (control). Each bar is the mean $\pm$ S.E.M. of seven fish. $*P < 0.05$ (one-way ANOVA). (B) Total and ionic plasma calcium (mmol/l) and (C) total plasma phosphate levels (mmol/l) in sea bream juveniles 15 days after treatment with $10 \mu\text{g/g}$ E_2 in coconut butter implants (E_2) or coconut butter implants alone (control). Values are given as means $\pm$ S.E.M. of 14 fish per treatment. $*P < 0.001$ compared with corresponding controls (one-way ANOVA).

Table 1 Drinking rates based on ^{51}Cr uptake (DR_{Cr}) and ^{45}Ca uptake (DR_{Ca}), the calculated percentage of absorbed calcium (% absorbed), the intestinal calcium uptake (Int_{Ca}) and the percentage contribution of intestinal calcium uptake to whole body calcium (% intestinal calcium uptake) in sea bream juveniles 15 days after treatment with $10 \mu\text{g/g}$ E_2 in coconut butter implants (E_2) or implants only (control). Values are the means $\pm$ S.E.M. of $n=7$

	Control	E_2	<i>P</i> value
DR_{Cr} ($\mu\text{l/h per g}$)	5.26 ± 0.74	5.27 ± 0.60	0.961
DR_{Ca} ($\mu\text{l/h per g}$)	0.49 ± 0.05	0.36 ± 0.07	0.146
% Absorbed	89.3 ± 1.4	92.8 ± 0.8	0.056
Int_{Ca} (nmol/h per g)	41.6 ± 7.9	50.4 ± 5.6	0.374
% Intestinal calcium uptake	64.9 ± 2.1	69.5 ± 3.3	0.360

P values of one-way ANOVA are shown.

($P=0.557$) groups. Efflux data were therefore pooled for subsequent analysis and the average value of the 2-h efflux period in each group is shown in Fig. 2B. Calcium efflux remained unchanged in response to E_2 treatment. Both groups were in positive calcium balance, i.e. there was an average net uptake of calcium of $18.33 \text{ nmol/h per g}$ in control and $24.03 \text{ nmol/h per g}$ in E_2 -treated fish (Fig. 2C).

Scale calcium, phosphate and TRAP

The total calcium and phosphate content of the scales remained unaffected after 15 days of E_2 treatment (Table 2). The calcium content of the scales in control fish was $1.25 \pm 0.07 \mu\text{mol/mg}$ vs $1.23 \pm 0.05 \mu\text{mol/mg}$ for E_2 -treated fish, while phosphate scale content was $0.96 \pm 0.07 \mu\text{mol/mg}$ in controls and $1.00 \pm 0.05 \mu\text{mol/mg}$ in E_2 -treated fish (Table 2). Furthermore, TRAP activity, a marker of osteoclastic activity, in the scales was not altered by E_2 treatment (Table 2).

Plasma protein analysis

Circulating levels of plasma protein were increased nearly threefold in response to E_2 treatment ($28.3 \pm 6.2 \text{ mg/ml}$ plasma vs $95.2 \pm 13.0 \text{ mg/ml}$ plasma, $P < 0.001$). SDS-PAGE and Western blot analysis of plasma protein confirmed that E_2 treatment induced a significant increase in circulating levels of vitellogenin; the latter protein was undetectable in the plasma of control fish. Associated with the increased vitellogenin production by the liver was a significant increase in the HSI of E_2 -treated fish compared with the control group, $3.3 \pm 0.4\%$ plasma vs $2.27 \pm 0.2\%$ plasma ($P < 0.05$) respectively.

Discussion

In the present study, the increase in circulating levels of E_2 in response to E_2 treatment was associated with

Figure 2 Whole body (A) calcium uptake, (B) calcium efflux and (C) net calcium flux in sea bream juveniles 15 days after treatment with 10 µg/g E₂ in coconut butter implants (E₂) or coconut butter implants alone (control). Each bar represents the mean ± S.E.M. of seven fish. **P*<0.001 compared with corresponding controls (one-way ANOVA).

significantly elevated plasma levels of both protein bound and ionic calcium. The experimental evidence indicated that there are at least two major components involved in the elevation of plasma calcium levels: an increase in calcium uptake from the water and an increase in intestinal calcium absorption. Thus, in the sea bream, E₂ appears to participate directly or indirectly in the regulation of these components.

Table 2 Calcium, phosphate and TRAP activity (means ± S.E.M., *n*=7) in the scales of sea bream juveniles 15 days after treatment with 10 µg/g E₂ in coconut butter implants (E₂) or implants only (control)

	Control	E ₂	P value
Calcium (µmol/mg)	1.26 ± 0.07	1.24 ± 0.05	0.838
Phosphate (µmol/mg)	0.96 ± 0.07	1.00 ± 0.05	0.697
TRAP (µmol pNP/h per mg)	11.9 ± 1.2	14.9 ± 1.9	0.235

P values of one-way ANOVA are shown.

Calcium exchange in fish takes place mostly at the level of the gills, with contributions from the skin (i.e. opercular membrane; McCormick *et al.* 1992, Marshall *et al.* 1992) and from the diet via the intestine. In the present study, we aimed to determine the contribution of the intestine to calcium regulation since marine fish drink large amounts of seawater (containing 10 mmol/l calcium in our experiments) and the intestine is known to play an important role in osmoregulation. Previous studies have shown that the proportion of the calcium taken up from water in the intestine of seawater fish is very variable. For example, the cod (*Gadus morhua*) absorbs about 30% of the calcium from drinking water (Sundell *et al.* 1993), while the flounder (*Paralichthys lethostigma*) absorbs about 70% (Hickman 1968). In tilapia (*Oreochromis mossambicus*), it has been shown that the net calcium transport in the intestine of seawater-adapted fish is significantly lower than in their freshwater counterparts, in which the uptake of calcium by the intestine represents a high proportion of whole body calcium uptake (Schoenmakers *et al.* 1993). In the present study, when the various routes of calcium entry into the body are considered, the high absorption rate in the intestine of sea bream translates into an intestinal component of the whole body calcium uptake in the range of 60–70% (Table 1). Moreover, since the branchial component of calcium uptake is in negative balance, it is the intestinal route that compensates for the calcium loss and maintains the fish in a positive calcium balance (Fig. 2).

The drinking rate of control and E₂-treated fish was the same (5.2 µl/h per g) and is similar to that previously observed in other marine species (Fuentes & Eddy 1997). However, one of the effects of E₂ treatment appears to be an increase in the relative proportion of the calcium taken up as drinking water that is absorbed (Table 1). An increase of calcium uptake at the intestinal level, 89% in controls versus 93% in the E₂-treated fish (Fig. 2A), coupled with an unchanged rate of calcium efflux (Fig. 2B) results in a 31% increase of the net whole body calcium uptake (Fig. 2C). However, this 31% increase cannot account for the final plasma calcium differences found between control and E₂-treated fish at the end of the experiment. Intriguingly, when plasma ionic calcium

(the physiologically important fraction) instead of total calcium is used to estimate calcium efflux rates, a threefold reduction was observed in E_2 -treated animals, which results in a nearly 40% calculated increment in the net influx. This change would be sufficient to explain the increase in total (and free) plasma calcium in the E_2 -treated group, assuming that the E_2 starts to have an effect soon after implantation.

In the E_2 -treated sea bream, vitellogenin was very abundant in plasma but was not detected in control fish. It was not determined whether the tenfold increase in plasma calcium in the E_2 -treated fish was exclusively associated with higher levels of circulating vitellogenin or whether there may have been additional calcium-binding proteins in the plasma. However, previous reports suggest that virtually all of the calcium in oestrogen-treated freshwater trout is bound to vitellogenin (Persson *et al.* 1994, 1995).

That the source of plasma calcium appears to be mainly the environment was supported by the fact that the phosphate and calcium content of the scales in E_2 -implanted juveniles remained unchanged, while the plasma calcium levels were increased to a level higher than those described in salmonids (Persson *et al.* 1997). This is in marked contrast to studies with rainbow trout in which considerable mobilisation of calcium from the scales was reported (Persson *et al.* 1997). The lack of calcium mobilisation from the scales in sea bream was further confirmed by the absence of E_2 -stimulated TRAP activity (Table 2) and contrasts with the effects of E_2 seen in rainbow trout (Persson *et al.* 1994, 1997). In salmonids, it has been suggested that increased calcium demand responds to a need for bone remodelling during reproductive riverine upstream migration. This increased calcium demand for bone remodelling is obtained not only from the environment but also from internal stores. The hypothesis of a direct effect of oestrogen on bone and/or scale remodelling in salmonids is supported by detection of oestrogen receptor α (ER α) mRNA expression in these tissues in the trout (Armour *et al.* 1997). Also, Persson *et al.* (2000) recently characterised a high-affinity, low-capacity binding of E_2 to trout scales. In the sea bream, ER α mRNA is not detected in bone and ER β is expressed at low levels (only detectable by RT-PCR; Socorro *et al.* 2000). The expression of ER in sea bream scales has yet to be determined, but the failure of E_2 to stimulate calcium mobilisation from scales in the sea bream suggests that the receptor is probably absent.

Whether the effect of oestradiol on calcium transport mechanisms in the sea bream occurs by a direct action on the gills, intestine and/or kidney, or is mediated by its action via the hypothalamic–pituitary axis and/or other endocrine systems, remains to be established. However, it has recently been shown (Socorro *et al.* 2000) that neither ER α nor ER β are expressed by the gills of sea bream, making it unlikely that oestradiol has a direct effect on calcium transport via ER in this tissue but mediates its

effects, if any, via a calcitropic factor. In contrast, there is high expression of ER β in the sea bream intestine and kidney (Socorro *et al.* 2000). The increase in net calcium influx (Fig. 2C) in response to E_2 in sea bream may be explained, at least in part, by a direct action via ER β in these tissues. In some tetrapods, a similar effect of E_2 on the intestine and kidney, which can be blocked by oestrogen antagonists (e.g. Nordin *et al.* 1991, ten Bolscher *et al.* 1999), has been observed.

Teleosts possess two hormones with hypocalcaemic action, but so far no hypercalcaemic factor has been identified. The hypocalcaemic factors include stannio-calcin which was originally isolated, characterised and its biological activity determined in fish (for review see Wendelaar Bonga & Pang 1991, Wagner *et al.* 1998) and calcitonin which has hypocalcaemic effects in goldfish (Sasayama *et al.* 1993) and salmonids (Wagner *et al.* 1997). The principal hypercalcaemic factor identified in tetrapods, parathyroid hormone (PTH), has not been identified in fish. However, both puffer fish (Power *et al.* 2000) and the sea bream (Flanagan *et al.* 2000) have recently been shown to express the gene for PTH-related protein (PTHrP), and a peptide corresponding to the amino acid residues 1–38 of the N-terminal region of PTHrP was able to cause an elevation in whole body calcium uptake in sea bream larvae, by increasing uptake and decreasing efflux of calcium (Guerreiro *et al.* 2001). Although the involvement of PTHrP in the hypercalcaemic effect of E_2 in the present study was beyond the experimental objectives, there is supporting evidence from mammalian studies that E_2 increases renal expression of PTHrP mRNA without modifying PTH/PTHrP receptor level of expression (Cros *et al.* 1998), thus preventing the renal leak of calcium in osteoporotic women (Nordin *et al.* 1991, Cros *et al.* 1998).

In fish, several other candidate hypercalcaemic hormones have been identified and include prolactin (Flik *et al.* 1994), cortisol and growth hormone (Flik & Perry 1989, Takagi *et al.* 1992). The link between calcium balance, oestrogen and pituitary hormone expression, i.e. prolactin, growth hormone and somatolactin in sea bream, is currently under study and preliminary results show a down-regulation of pituitary gene expression of these hormones in response to E_2 (J Fuentes, unpublished observations). However, the relative importance of calcium and E_2 on this effect has yet to be established.

In conclusion, our data suggest that the sea bream, in common with freshwater teleosts, respond to increases in circulating E_2 with an increase in circulating plasma levels of vitellogenin, total calcium and an accommodation of the calcium-transporting mechanisms. However, the source of calcium and mechanisms by which internal circulating levels are increased appear to be different between freshwater and marine teleosts. While freshwater fish may rely on the internal stores of calcium, i.e. bone and/or scales to increase calcium availability, the sea bream, a marine fish, fulfils the extra calcium demand by utilising the high

environmental calcium and, by doing so, protects internal calcified structures. It remains to be seen if the profound effects of E₂ on calcium balance in sea bream are direct, or are mediated by other endocrine factors.

Acknowledgements

This study was funded by the project PRAXIS/P/BIA/13174/98. P M G and J F were funded by the Ministry of Science, FCT, Portugal, through grants BD/9207/96 and BPD/22033/99 respectively. We thank Dr Patricia M Ingleton for helpful discussions during the study and for reading and commenting on the manuscript.

References

- Armour KJ, Lehan DB, Pakdel F, Valotaire Y, Graham R, Russell RG & Henderson IW 1997 Estrogen receptor mRNA in mineralized tissues of rainbow trout: calcium mobilization by estrogen. *FEBS Letters* **411** 145–148.
- ten Bolscher M, Netelenbos JC, Barto R, Van Buuren LM & Van der Vijgh WJF 1999 Estrogen regulation of intestinal calcium absorption in the intact and ovariectomized adult rat. *Journal of Bone and Mineral Research* **14** 1197–1202.
- Bradley JT & Grizzle JM 1989 Vitellogenin induction by estradiol in channel catfish, *Ictalurus punctatus*. *General and Comparative Endocrinology* **73** 28–39.
- Burzawa-Gerard E & Dumas-Vidal A 1991 Effects of 17 β -estradiol and carp gonadotropin on vitellogenesis in normal and hypophysectomized European silver female eel (*Anguilla anguilla* L.) employing a homologous radioimmunoassay for vitellogenin. *General and Comparative Endocrinology* **84** 264–276.
- Carragher JF & Sumpter JP 1991 The mobilization of calcium from calcified tissues of rainbow trout (*Oncorhynchus mykiss*) induced to synthesize vitellogenin. *Comparative Biochemistry and Physiology* **99A** 169–172.
- Cros M, Silve C, Graulet AM, Morieux C, Urena P, de Vernejoul MC & Bouzard 1998 Estrogen stimulates PTHrP but not PTH/PTHrP receptor gene expression in the kidney of ovariectomized rat. *Journal of Cellular Biochemistry* **70** 84–93.
- Flanagan JA, Power DM, Bendall LA, Guerreiro PM, Fuentes J, Clark MS, Canario AVM, Danks JA, Brown BL & Ingleton PM 2000 Cloning of the cDNA for sea bream (*Sparus aurata*) parathyroid hormone-related protein. *General and Comparative Endocrinology* **118** 373–382.
- Flik G & Perry SF 1989 Cortisol stimulates whole body calcium uptake and the branchial calcium pump in freshwater rainbow trout. *Journal of Endocrinology* **120** 75–82.
- Flik G, Fenwick JC, Kolar Z, Mayer-Gostan N & Wendelaar Bonga SE 1986 Effects of low ambient calcium levels on Ca²⁺ flux rates and internal calcium pools in the freshwater cichlid teleost, *Oreochromis mossambicus*. *Journal of Experimental Biology* **120** 249–264.
- Flik G, Rentier-Delrue F & Wendelaar Bonga SE 1994 Calcitropic effects of recombinant prolactins in *Oreochromis mossambicus*. *American Journal of Physiology* **266** R1302–R1308.
- Flik G, Verboost PM & Wendelaar Bonga SE 1995 Calcium transport processes in fishes. In *Fish Physiology*, vol. 14: *Cellular and Molecular Approaches to Fish Ionic Regulation*, pp 317–342. Eds CM Wood & T Shuttleworth. San Diego: Academic Press.
- Fuentes J & Eddy FB 1997 Drinking in freshwater, euryhaline and marine teleosts. In *Ionic Regulation in Animals*, pp 135–149. Eds N Hazon, FB Eddy & G Flik. Berlin, Heidelberg: Springer-Verlag.
- Guerreiro PM, Fuentes J, Power DM, Ingleton PM, Flik G & Canario AVM 2001 Parathyroid hormone-related protein: a calcium regulatory factor in sea bream (*Sparus aurata* L.) larvae. *American Journal of Physiology* **281** R855–R860.
- Hickman CPJ 1968 Ingestion, intestinal absorption and elimination of sea water and salts in the southern flounder, *Paralichthys lethostigma*. *Canadian Journal of Zoology* **46** 457–466.
- Ichii T & Mugiya Y 1983 Effects of dietary deficiency in calcium on growth and calcium uptake from the aquatic environment in the goldfish, *Carassius auratus*. *Comparative Biochemistry and Physiology* **74A** 259–262.
- Laemmli UK 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227** 680–685.
- Larsson D, Björnsson BT & Sundell K 1995 Physiological concentrations of 24,25-dihydroxyvitamin D₃ rapidly decrease the *in vitro* intestinal calcium uptake in the Atlantic cod, *Gadus morhua*. *General and Comparative Endocrinology* **100** 211–217.
- Lau K-WH, Farley JR & Baylink DJ 1985 Phosphotyrosyl-specific protein phosphatase activity of a bovine skeletal acid phosphatase isoenzyme. *Journal of Biological Chemistry* **260** 4653–4660.
- Lowry OH, Rosebrough NJ, Farr AL & Randall RJ 1951 Protein measurement with the folin-phenol reagents. *Journal of Biological Chemistry* **193** 265–275.
- McCormick SD, Hasegawa S & Hirano T 1992 Calcium uptake in the skin of a freshwater teleost. *PNAS* **89** 3635–3638.
- Marshall WS, Bryson SE & Wood CM 1992 Calcium transport by isolated skin of rainbow trout. *Journal of Experimental Biology* **166** 297–316.
- Mosconi G, Carnevali O, Carletta R, Nabissi M & Polzonetti-Magni AM 1998 Gilthead sea bream (*Sparus aurata*) vitellogenin: purification, partial characterization, and validation of an enzyme-linked immunosorbent assay (ELISA). *General and Comparative Endocrinology* **110** 252–261.
- Mugiya Y & Watabe N 1977 Studies on fish scale formation and resorption. II. Effect of estradiol on calcium homeostasis and skeletal tissue in the goldfish, *Carassius auratus* and the killifish, *Fundulus heteroclitus*. *Comparative Biochemistry and Physiology* **57A** 197–202.
- Nordin BEC, Need AG, Morris HA, Horowitz M & Robertson WG 1991 Evidence for a renal calcium leak in postmenopausal women. *Journal of Clinical Endocrinology and Metabolism* **72** 401–407.
- Persson P, Sundell K & Björnsson BT 1994 Estradiol-17 β induced calcium uptake and resorption in juvenile rainbow trout, *Oncorhynchus mykiss*. *Fish Physiology and Biochemistry* **13** 379–386.
- Persson P, Takagi Y & Björnsson BT 1995 Tartrate resistant acid phosphatase as a marker for scale resorption in rainbow trout, *Oncorhynchus mykiss*: effects of estradiol-17 β treatment and refeeding. *Fish Physiology and Biochemistry* **14** 329–339.
- Persson P, Johannsson SH, Takagi Y & Björnsson BT 1997 Estradiol-17 β and nutritional status affect calcium balance, scale and bone resorption, and bone formation in rainbow trout, *Oncorhynchus mykiss*. *Journal of Comparative Physiology B* **167** 468–473.
- Persson P, Shrimpton JM, McCormick SD & Björnsson BT 2000 The presence of high-affinity, low-capacity estradiol-17 β binding in rainbow trout scale indicates a possible route for the regulation of scale resorption. *General and Comparative Endocrinology* **120** 35–43.
- Power DM, Ingleton PM, Flanagan J, Canario AVM, Danks JA, Elgar G & Clark MS 2000 Genomic structure and expression of parathyroid hormone-related protein gene (PTHrP) in a teleost, *Fugu rubripes*. *Gene* **250** 67–76.
- Sasayama Y, Ukawa K, Kai-Ya H, Oguro C, Takei Y, Watanabe TX, Nakajima K & Sakakibara S 1993 Goldfish calcitonin: purification, characterization, and hypocalcemic potency. *General and Comparative Endocrinology* **89** 189–194.
- Schoenmakers TJM, Verboost PM, Flik G & Wendelaar Bonga SE 1993 Transcellular intestinal calcium transport in freshwater and seawater fish and its dependence on sodium/calcium exchange. *Journal of Experimental Biology* **176** 195–206.

- Scott AP, Sheldrick EL & Flint APF 1982 Measurement of 17 alpha, 20 beta-dihydroxy-4-pregnen-3-one in plasma of trout (*Salmo gairdneri* Richardson): seasonal changes and response to salmon pituitary extract. *General and Comparative Endocrinology* **46** 444–451.
- Socorro S, Power DM, Olsson P-E & Canario AVM 2000 Two estrogen receptors expressed in the teleost fish, *Sparus aurata*: cDNA cloning, characterization and tissue distribution. *Journal of Endocrinology* **166** 293–306.
- Sundell K, Norman AW & Björnsson BT 1993 1,25(OH)₂ vitamin D₃ increases ionized plasma calcium concentrations in the immature Atlantic cod, *Gadus morhua*. *General and Comparative Endocrinology* **91** 344–351.
- Takagi Y, Moriyama S, Hirano T & Yamada J 1992 Effects of growth hormones on bone formation and resorption in rainbow trout (*Oncorhynchus mykiss*), as examined by histomorphometry of the pharyngeal bone. *General and Comparative Endocrinology* **86** 90–95.
- Wagner GF, Jaworski EM & Radman DP 1997 Salmon calcitonin inhibits whole body Ca²⁺ uptake in young rainbow trout. *Journal of Endocrinology* **155** 459–465.
- Wagner GF, Jaworski EM & Haddad M 1998 Stanniocalcin in the seawater salmon: structure, function, and regulation. *American Journal of Physiology* **274** R1177–R1185.
- Wendelaar Bonga SE & Pang PKT 1991 Control of calcium regulating hormones in the vertebrates: parathyroid hormone, calcitonin, prolactin and stanniocalcin. *International Review of Cytology* **128** 139–213.

Received in final form 2 January 2002

Accepted 22 January 2002