Cloning of *Octopus* cephalotocin receptor, a member of the oxytocin/vasopressin superfamily

Atsuhiko Kanda, Kyoko Takuwa-Kuroda, Eiko Iwakoshi-Ukena, Yasuo Furukawa¹, Osamu Matsushima¹ and Hiroyuki Minakata

Suntory Institute for Bioorganic Research, 1-1-1 Wakayamadai, Shimamoto, Mishima, Osaka 618-8503, Japan

¹Department of Biological Science, Faculty of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan

(Requests for offprints should be addressed to H Minakata; Email: minakata@sunbor.or.jp)

Abstract

We reported that the common octopus, *Octopus vulgaris*, in common with vertebrates, possesses two members of the oxytocin/vasopressin superfamily: octopressin (OP) and cephalotocin (CT). This was the first observation of its kind in invertebrates. As OP and CT have different biological activities, the presence of specific receptors has been proposed. We cloned the cDNA of an orphan receptor from *Octopus* brain and found it to encode a polypeptide of 397 amino acids that displays sequences characteristic of G-protein coupled receptors. The orphan receptor showed high homology to receptors of the oxytocin/vasopressin superfamily and seemed to conserve the agonist-binding pocket common to the oxytocin and vasopressin receptors. *Xenopus* oocytes that express the orphan receptor responded to the application of CT by an

induction of membrane Cl^- currents coupled to the inositol phosphate/ Ca^{2+} pathway. OP and the other members of the oxytocin/vasopressin superfamily did not activate this receptor. HPLC fractionation of the *Octopus* brain extract combined with an oocyte assay yielded a single substance that was identical to CT. On the basis of these results, we conclude that the cloned receptor is the CT receptor (CTR). Expression of CTR mRNA in *Octopus* was detected in the central and the peripheral nervous systems, the pancreas, the oviduct and the ovary. This receptor may mediate physiological functions of CT in *Octopus* such as neurotransmission, reproduction and metabolism.

Journal of Endocrinology (2003) **179**, 281–291

Introduction

Oxytocin (OT) and vasopressin (VP) are nonapeptides that are present in all placental mammals. Although they are similar in structure, they serve different functions. OT has uterotonic and milk-ejection activities and regulates aspects of reproductive behaviour, whereas VP has antidiuretic activities and is involved in hydromineral regulation.

Members of the OT/VP superfamily are widely distributed in both invertebrates and vertebrates, and have key roles as signalling molecules in the regulation of reproduction and osmotic balance (reviewed by Gimpl & Fahrenholz 2001). In invertebrates, they include VP-like diuretic hormone from the locust *Locusta migratoria*, annetocin from the earthworm *Eisenia foetida*, Lys- and Arg-conopressins from the gastropod molluscs *Conus geographus*, *Conus striatus* and *Lymnaea stagnalis* (reviewed by Hoyle 1998) and cephalotocin (CT) from the cephalopod mollusc, *Octopus vulgaris* (Reich 1992). Although all vertebrate species except for cyclostomes have peptides from both the OT and the VP families, invertebrate species examined to date have usually possessed only a

single peptide belonging to the OT/VP superfamily. We showed previously that, in addition to CT, *Octopus* possesses another member of the OT/VP superfamily, octopressin (OP) (Takuwa-Kuroda *et al.* 2003). This was the first report of the co-occurrence of two members of the superfamily in a single invertebrate species. Because OP, but not CT, induced contractile activity in smooth muscles of several *Octopus* tissues, we proposed that there are more than two distinct receptors in the octopus for members of the oxytocin/vasopressin superfamily (Takuwa-Kuroda *et al.* 2003).

Receptors of the OT/VP superfamily are generally G-protein coupled receptors (GPCRs), a superfamily of proteins characterized by seven transmembrane-spanning domains. GPCRs mediate the actions of numerous neurotransmitters and hormones (Probst *et al.* 1992). The V2 receptor mediates the antidiuretic effects of VP and couples positively to adenylate cyclase (Lolait *et al.* 1992, Gorbulev *et al.* 1993). The OT receptor (Kimura *et al.* 1992, Gorbulev *et al.* 1993) mediates various central and peripheral functions in reproduction in mammals. V1a (Thibonnier *et al.* 1994) and V1b (Sugimoto *et al.* 1994)

Table 1 Sequences of primers

Primer	Sequence
OP/CT-R TM6	5'-TA(C/T)(A/G)TI(T/G/C)TITG(C/T)TGG(A/T/G)CITT(C/T)TT-3'
OP/CT-R TM7a	5'-TA(A/T/G)ATCCAIGG(A/G)TT(A/G)(T/G/C)(A/T)(A/G)CAIGA(A/G)TT-3'
OP/CT-R 3'-1F	5'- ¹¹²⁸ GCAGCTTGGGATGAAAATGC ¹¹⁴⁷ -3'
OP/CT-R 3'-2F	5'- ¹¹⁴⁶ GCTCCATATTCAAGTGCAATC ¹¹⁶⁶ -3'
OP/CT-R 5'-1R	5'- ¹⁴¹¹ AAGCCGGATTTTAGTAGTGG ¹³⁹² -3'
OP/CT-R 5'-2R	5'- ¹²⁶¹ CTAGTCTTTGCTCTGTGTTGG ¹²⁴¹ -3'
OP/CT-R 5'-3R	5'- ¹²⁰⁹ TAGTACAGCTGTTGAGGGAC ¹¹⁹⁰ -3'
OP/CT-R 5'-4R	5'- ⁷⁸³ TCCATTCTGGGTCAAAGGTC ⁷⁶⁴ -3'
OP/CT-R 5'-5R	5'- ⁷¹¹ CGGAGTACTAAAGAGAAATG ⁶⁹¹ -3'
OP/CT-R 5'-6R	5'- ⁶⁵² GCAGTTGTCCACTTGTGAC ⁶³⁴ -3'
OP/CT-R 5'-7R	5'- ²⁵⁴ ACTCCATATTTTCGGTCGATGAC ²³³ -3'
OP/CT-R 5'-8R	5'- ²¹⁶ GCGTCGATATTTTCAATTTGGATG ¹⁹⁵ -3'
OP/CT-R 5'-9R	5'- ¹⁹⁰ GTAATGTACCTCATTGCTGC ¹⁷¹ -3'
OP/CT-R FwA	5'- ⁵² GTTGTGGTTTATTTTCATCTGGC ⁷³ -3'
OP/CT-R FwB	5'- ⁸⁰ CTGTTTACAACACTACAAAAGCG ¹⁰² -3'
OP/CT-R Rev	5'- ¹⁴⁰⁸ CCGGATTTTAGTAGTGGTATTC ¹³⁸⁷ -3'

I, inosine.

receptors mediate the effects of VP on liver glycogenolysis and the release of adrenocorticotropin, respectively. Similar effects of OT on the liver are also mediated by the OT receptor. These receptors couple to the inositol phosphate/ Ca^{2+} signal transduction pathway.

In invertebrates, the cloning of two Lys-conopressin receptors, LSCPR1 and LSCPR2, from *Lymnaea* has been reported (van Kesteren *et al.* 1995, 1996). These receptors are expressed in different tissues and mediate both functions of OT-like reproduction and VP-like glycogenolysis of Lys-conopressin in *Lymnaea*.

We report here the cloning of an octopus homologue of the OT/VP receptor from the brain of *Octopus*. The receptor, named CT receptor (CTR), is a GPCR that is specific to CT. Because CTR mRNA was detected in the central and the peripheral nervous systems, the pancreas, the oviduct and the ovary, CTR may mediate physiological functions such as neurotransmission, reproduction and metabolism.

Material and Methods

Animals

Adult octopuses (*Octopus vulgaris*, ca. 2 kg), were purchased from a local fish shop, and kept in artificial seawater at 18 °C.

Preparation of total RNA

Total RNA was prepared using Sepasol-RNA I Super (Nacalai Tesque, Kyoto, Japan) according to the manufacturer's instructions.

Oligonucleotide primers

Oligonucleotide primers were purchased from Qiagen and Kiko-tech (Osaka, Japan). Degenerate primers, OP/

CT-R TM6 and OP/CT-R TM7a, were designed that were based on sequences in the sixth and seventh transmembrane domains, respectively, both of which are conserved among the receptors of the OT/VP superfamily in vertebrates and Lys-conopressin receptors. All primers used are listed in Table 1.

Cloning of partial-length cDNA

All PCR amplifications were carried out in a reaction mixture containing Ex Taq polymerase (Takara Shuzo, Kyoto, Japan) and 200 μM dNTP in a thermal cycler (GeneAmp PCR System 9700; Applied Biosystems Japan, Tokyo, Japan). To confirm the fidelity of Taq polymerase, several sets of clones (at least four clones) were sequenced in every PCR experiment. First-strand cDNA was synthesized with the oligo(dT)-anchor primer supplied in the 5'/3'-RACE kit (Roche Diagnostics, Basel, Switzerland). PCR was performed using OP/CT-R TM6 and OP/CT-R TM7a under the following conditions: 5 min at 94 °C; 50 cycles of 30 s at 94 °C, 30 s at 50 °C, 1 min at 72 °C (7 min for the last cycle). PCR products separated by 1.5% agarose gel electrophoresis were purified using Qiaquick Gel Extraction kit (Qiagen), and subcloned into the pDrive Cloning vector using a Qiagen PCR cloning kit and transformed into *Escherichia coli* DH5 α (TOYOBO, Tokyo, Japan). Subcloned inserts were sequenced on an ABI Prism 310 Genetic Analyser (Applied Biosystems Japan) using a Big-Dye sequencing kit version 3 (Applied Biosystems Japan) and analysed on Genetyx-Mac software (Software Development, Tokyo, Japan). Universal M13 primers or gene-specific primers were used to sequence both strands.

3'-Rapid amplification of cDNA ends

First-strand cDNA was synthesized with the oligo(dT)-anchor primer supplied in the 5'/3'-RACE kit (Roche Diagnostics). The first PCR was performed using the PCR anchor primer and OP/CT-R 3'-1F under the following conditions: 5 min at 94 °C; 30 cycles of 30 s at 94 °C, 30 s at 50 °C, 2 min at 72 °C (7 min for the last cycle). The second PCR was performed using the PCR anchor primer and OP/CT-R 3'-2F under the following conditions: 94 °C for 5 min; 30 cycles of 30 s at 94 °C, 30 s at 51 °C, 2 min at 72 °C (7 min for the last cycle). The second PCR products were subcloned and sequenced as described above.

5'-Rapid amplification of cDNA ends

Three sets of template cDNAs were synthesized using as primers OP/CT-R 5'-1R, OP/CT-R 5'-4R and OP/CT-R 5'-7R, and tailed by dA using dATP and terminal transferase (Roche Diagnostics). The first template was amplified using the oligo(dT)-anchor primer and OP/CT-R 5'-2R. The remaining two templates were also amplified between the anchor primer and OP/CT-R 5'-5R, and between the anchor and OP/CT-R 5'-8R, respectively. Each of the first PCR products was re-amplified using the anchor primer and OP/CT-R 5'-3R, OP/CT-R 5'-6R or OP/CT-R 5'-9R. Both first and second PCR were performed for 5 min at 94 °C; 30 cycles for 30 s at 94 °C, 30 s at 52 °C, and 2 min at 72 °C (7 min for the last cycle). The second PCR products were subcloned and sequenced as described in the previous section. Full-length cDNA was obtained using primers, OP/CT-R FwA and OP/CT-R Rev, and the following PCR conditions: 5 min at 94 °C; 50 cycles for 30 s at 94 °C, 30 s at 50 °C, and 2 min at 72 °C (7 min for the last cycle). Thus a 1706 bp-long cDNA encoding an orphan receptor was obtained.

RT-PCR Southern blot analysis

Each of total RNAs (1 µg) extracted from various tissues was reverse-transcribed by Superscript II (Invitrogen, Groningen, The Netherlands) using oligo dT₁₂₋₁₈ primer. The first PCR was performed using OP/CT-R FwA and OP/CT-R 5'-1R under the following conditions: 5 min at 94 °C; 50 cycles of 30 s at 94 °C, 30 s at 50 °C, 1 min at 72 °C (7 min for the last cycle). The second PCR was performed using OP/CT-R FwB and OP/CT-R Rev under the following conditions: 94 °C for 5 min; 20 cycles of 30 s at 94 °C, 30 s at 50 °C, 1 min at 72 °C (7 min for the last cycle). The second PCR products were separated by 1.5% agarose gel electrophoresis, and then transferred onto Hybond-N⁺ membrane (Amersham) and crosslinked by UV irradiation. Hybridization and detection were processed according to the manufacturer's

procedure (Roche Diagnostics). The cDNA of the orphan receptor was labelled with the DIG DNA labelling kit (Roche Diagnostics) and used as a probe for Southern blotting according to the manufacturer's instructions. The partial cDNA of *Octopus* actin (AB053937) amplified by the primer set (position 25–44 and position 595–614) was hybridized and detected by the DIG-labelled oligonucleotide (position 347–366) as an internal control. As a negative control, the extracted total RNAs, which were not reverse transcribed, were used as templates for PCR, and it was confirmed that there was no amplification of traces of the genomic DNA.

Expression of the cloned orphan receptor in *Xenopus* oocytes

The open reading frame region of the orphan receptor cDNA was amplified and inserted into a pSP64 poly(A) vector (Promega). The plasmid was linearized with *EcoRI*, and cRNA was prepared using SP6 RNA polymerase (Ambion, Austin, TX, USA). Fifty nanolitres of the cRNA solution (0.05 µg/µl) was injected into *Xenopus* oocytes. The oocytes were incubated for 2–4 days at 17 °C in ND96 buffer [96 mM NaCl, 2 mM KCl, 1.8 mM CaCl₂, 1 mM MgCl₂ and 5 mM HEPES (pH 7.6)]. The oocytes were voltage-clamped at –80 mV. Peptides diluted with ND96 buffer were applied to the oocytes every 15 min to prevent desensitization of the receptor at concentrations between 10^{–10} and 10^{–6} M. Four oocytes were tested at each concentration to obtain a dose–response relationship. OP and CT were synthesized with a solid-phase peptide synthesizer using standard Fmoc chemistry (Model 433A, Applied Biosystems Japan). Peptides of the OT/VP superfamily and antagonists for OT, V1 and V2 receptors were purchased from Bachem (Bubendorf, Switzerland). A mixture of an antagonist and a peptide was dissolved in the buffer at 10^{–5} M and 10^{–7} M. The dose–response data were analysed using ORIGIN 6.1 software (Microcal Software Inc., Northampton, MA, USA) to estimate the EC₅₀ value.

Isolation and structure determination of an endogenous ligand of the orphan receptor from *Octopus* brain

The methods for dissection, extraction and purification were the same as those previously described (Iwakoshi *et al.* 2000). Briefly, the suboesophageal brain (10 g) was boiled in water and extracted with 3% acetic acid. The extract was applied to a solid-phase extraction column (Sep-pak Vac C18, Waters, Milford, MA, USA). The retained material was eluted with 60% acetonitrile in 0.1% trifluoroacetic acid (TFA), and condensed *in vacuo*. The residue was dissolved in 0.1% TFA and fractionated by a linear gradient elution of 0–36% acetonitrile in 0.1% TFA at a flow rate of 1.5 ml/min on a reversed-phase HPLC with a C18 column (Capcell Pak C18 UG80 10 mm × 250 mm, Shiseido, Tokyo, Japan). Fractions were

collected every 2 min. A 1/100 aliquot of each fraction was condensed to dryness *in vacuo*, dissolved in ND96 buffer and then assayed in the *Xenopus* oocyte that expressed the orphan receptor. The active fractions were further purified by using the following columns: a cation-exchange column (TSKgel SP-5PW, 7.5 mm × 75 mm, Tosoh, Tokyo, Japan; 0–0.6 M NaCl/60-min linear gradient in 20 mM phosphate buffer, pH 7.0, flow rate of 1.5 ml/min, fraction/2 min), a C18 column (Capcell Pak C18, MG80, 4.6 mm × 250 mm, Shiseido; 18–30% acetonitrile in 0.1% TFA/40-min linear gradient, flow rate of 1.0 ml/min, fraction/1 min). In the assay of cation-exchange HPLC fractions, 1/100 aliquot of each fraction was diluted in ND96 buffer, and applied to the assay chamber.

The active peak from the third C18 HPLC was subjected to amino acid sequencing (PSQ-1; Shimadzu, Kyoto, Japan). The molecular weight was determined by nanoflow electrospray ionization time-of-flight mass spectrometry (nano ESI-TOF-MS) (Q-TOF; Micromass UK, Wythenshawe, UK). The molecular mass peak was further examined in nano ESI-TOF-MS/MS analysis. The capillary voltage was optimized at 1200 V and the cone voltage was set at 50 V. Argon was used as the collision gas and the energy was set at 25 V.

Results

Cloning and structure of an Octopus receptor of the OT/VP superfamily

Within the amino acid sequences of the known receptors of the OT/VP superfamily, in both vertebrates and invertebrates, those of the sixth and seventh transmembrane domains are highly conserved. To identify receptors for OP and CT in *Octopus*, two degenerate primers were designed that were based on the conserved region, and PCR was performed between these primers. PCR products with the expected size of 134 bp were cloned and sequenced. The encoded protein fragment showed the greatest homology with structures for receptors of the OT/VP superfamily found in a database, DDBJ. The full-length cDNA sequence (1706 bp) encoding a putative OP/CT receptor was determined using the 5′- and 3′-RACE methods (Fig. 1A). It has an open reading frame of 397 amino acids and contains 176 bp of 5′-untranslated and 336 bp of 3′-untranslated sequences with two polyadenylation signals (AATAAA) 113 bp and 244 bp upstream from the poly(A) tail. The nucleotide sequence was deposited into the DDBJ/EMBL/GeneBank databases under the accession number AB090945.

Several sets of clones in every PCR were analysed and gave exactly the same nucleotide sequence in each experiment, thus confirming the fidelity of Taq polymerase.

Hydrophobicity analysis of the predicted amino acid sequence revealed seven hydrophobic domains character-

istic of GPCR (Fig. 1B). The orphan receptor has two Cys residues at positions 118 and 194 in the first and second extracellular loops. These residues are highly conserved within GPCR and are probably connected by a disulphide bridge. As shown in Fig. 1A, the sequence displays several other characteristics of GPCR such as N-linked glycosylation sites (Asn-X-Ser/Thr; Asn^{18,27,33}) in the N-terminal extracellular domain, consensus sequences for phosphorylation by protein kinase A (Arg/Lys-X-(X)-Ser/Thr; Ser^{240,251,258,271,287,362,386}, Thr^{78,241,259,361}), by protein kinase C (Ser/Thr-X-Arg/Lys; Ser^{153,244,258,377,386}, Thr^{158,254,361,394}), and by casein kinase II (Ser/Thr-X-X-Asp/Glu; Ser^{271,378}). The orphan receptor has an Asp⁹¹ in transmembrane domain II and a tripeptide (Asp-Arg-Tyr) at the interface of transmembrane III and the first intracellular loop. These residues are believed to be important for receptor activation (Bockaert & Pin 1999). These results suggest that the cloned orphan receptor is a member of the GPCR family.

An amino acid sequence alignment of the cloned receptor and receptors of the OT/VP superfamily is presented in Fig. 2. The sequence homologies of the cloned receptor were estimated as follows: 42.5% human V1a, 39.9% human V1b, 35.5% human V2, 42.2% chicken vasotocin, 41.1% white sucker vasotocin, 39% human OT, 42.2% toad mesotocin, 39.4% white sucker isotocin, 43.2% *Lymnaea* LSCPR1, and 37.9% *Lymnaea* LSCPR 2. Several amino acid sequences (Phe-Gln-Val-Leu-Pro-Gln-Leu at the C-terminal end of transmembrane domain II, Gly-Pro-Asp in the first and Asp-Cys-Trp and Pro-Trp-Gly in the second extracellular loop) that are conserved among receptors of the OT/VP superfamily in vertebrates are considered to be important for ligand binding (Sharif & Hanley 1992) (Fig. 2). In the cloned receptor, these sequences are somewhat homologous, namely, Phe-Asn-Asn-Leu-Pro-Gln-Met, Gly-Thr-Asp, Thr-Cys-Gln, and Glu-Trp-Thr. Such amino acid substitutions also occurred in LSCPR1 and LSCPR2. On the basis of these results, we suggest that the cloned receptor is a homologue of receptors of the OT/VP superfamily.

Functional expression of the cloned receptor in Xenopus oocyte

If the cloned receptor can couple to Gq protein of *Xenopus* oocyte, active ligands will exhibit Cl[−] current mediated by the inositol phosphate/Ca²⁺-signalling pathway. The voltage-clamped oocytes expressing the cloned receptor displayed inward membrane currents after the application of CT at 10^{−7} M (Fig. 3A). In contrast, OP did not trigger the current even at 10^{−6} M (Fig. 3B). We also tested both vertebrate and invertebrate peptides belonging to the OT/VP superfamily (VP, vasotocin, anetocin, Lys-conopressin, or Arg-conopressin), but they had no effects even at 10^{−5} M (data not shown). Among the tested peptides, only OT caused a weak effect at 10^{−6} M

(A)

CACACACACACACACACACACACACACACAGATTCTGTACATAGTTGTGGTTTTCATCTGGCAATTCTGTTTACAACACTCAAAAGCGATTGTGAGAATTGAAAA 120
TAACAAATTTTGTAACTGATCAACAGAGAATTATTTTAAATAAAATTAAGACGCAATGAGGTACATTACTACACATCCAAATGAAATATCGACGCAAAATTGGAATTAATTTCTCATCGA 1240
M R Y I T T H P N E I S T Q I W N N F S S
10 20

CGAAATATGGAGTAACTTTCCGCAGCAAAAAATGAACTCAGCCCAATAGACGTAATCAAGACCTGGCTAATGCTGAAGTTATTACGTTGGCTGTTGTTATAAATACAGTTATAG 360
T E I W S N F S A A K N E T Q P I R R N Q D L A N A E V I T L A V V I I I T V I
30 40 50 60

GTAATGATATTGTTTAAATCAACTGTTCCAGCGTAGAAAAGAACTGACACGAATGCACCTGTTTATTTTACACCTGTCTGTAACCTGACCTCTTCTGCTGCCCTCTTCTTAATTAATTTGCCGC 480
G N S I V L I T I L F Q R R K K L T R M H L F L H L S V T D L F V A F F N N L P
70 80 90 100

AGATGATTTGGACATTACATTTTATTTCTTGGAACTGATTACTGTGTCTGCTGTTACTTACTTACAAAGCTAGCTATGTATGCATCGAGCTATGCTCTTGTGGCAACAGCTATTG 600
Q M I W D I T F L F L G T D L L C R L V T Y L Q S V A M Y A S S Y V L V A T A T I
110 120 130 140

ACCGTTATTTGCTATATGCTACCCCTTTATCAAGTCACAAGTGGACAACTGCCAGAGTACAGCTCATGGTTTTCATGCTGGAGTTATCATTTCTCTCAGTACTCCGCAGCTCTTCA 720
D R Y F A I C H P L S S H K W T T A R V H V M V F I A W M L S F L F S T P Q L F
150 160 170 180

TATGGAGCATGCAGTTCTCAAATATTGGAATGACTTGTGAGCGACCTTTGACCCAGAATGGACTCTGAAATTTTATACCTTGGTTAAACAGTCGCGATTGGATTTTACCAACAATAG 840
I W S M Q F S N I G L T C Q A T F D P E W T L K F Y I T W L T V A I W I L P T I
190 200 210 220

CAGTGACTCTGTTTATGGAATGATGCTGTTGCTGTATGGAAAAGAGGACGTTCTACTTTGGCGAGTCCCGAACACGAAATAGATCTTTCTCAGCAATCGTGTCTACACGATTG 960
A L T L F Y G M M C F A V A W K R G R S T L G S S R T R N R S F L T N R V S T R I
230 240 # 250 # 260 #

GTCAAAGCCATTAGCTCGAGGGTTCCTCCGAAGGAGTATGGAAGGACAAGGCTCAATTACAACAGAGGAATTTCTCGAGCCAAAGTCGGGTCTGGCTCTAACTTAAAGTGGTTG 1080
G Q S H L A R G F S E E D M E G Q S V N Y N R G I S R A K V R S V A L T L S V V
270 # 280 # 290 300

CCTGTTACTTTATATGTTGGTCACCGTTTTTGTGTGTCAGATGTGGGCAGCTTGGGATGAAATGCTCCATATTCAGGTGCAATCTATACGATATATTGCTACTCTGTCCTCAACA 1200
A C Y F I C W S P F F V C Q M W A A W D E N A P Y S L N A I Y T I L L L L S S L N
310 320 330 340

GCTGTACTAATCCCTGGATTACATGATCTTTAGTGATTCCAAACAGACGAAAGACTAGTCGCTTGTGTAATGACGAAGAACGACCTCAGTGACTGTGCTGTCATCAGGAATGATA 1320
S C T N P W I Y M I F S V F Q H R A K T S R F V N D E E T T S V T V L S S R N D
350 360 # 370 380

TCGATTTGATGCTCAATGAAGAAGAAATGGAAACAAACAGCCAGAAACTGAAAAAATTCAGAAAATTAATACCACCTACTAAAAATCCGGCTTCCATTATGTTGTAATTTGAGTGACATGT 1440
I R L M S M K K K L E Q T A R N *
390 397

#

TAATAACCTGGAAGAACACAGAAATTTAGAACAAATTTAAGTGCAACGTAATTCATTTCTGTGACGTATATGATATGGATGATGCATATAAGTACTTGCATGTGTACTGA 1560
TATATATATGTAATAATATATAGAAATCAGCATGATCAAAAGCAGCGATACGCAACACGCCAGATATACCTGTAACCTACTTATACGTATACATGCTTACAGTCCACACTT 1680
CAGCATCATCAAAAAA 1706

(B)

Figure 1 Primary structure of the precursor of the cloned orphan receptor. (A) Nucleotide sequence and deduced amino acid sequence of the precursor of the orphan receptor cloned from the *Octopus* brain. The transmembrane regions are underlined. *, N-Linked glycosylation sites. Potential phosphorylation sites: ●, protein kinase C; #, protein kinase A; ◆, casein kinase II. The polyadenylation signals (AATAAA) are in boxes. (B) Hydrophobicity plot of the orphan receptor according to Kyte & Doolittle (1982).

Figure 2 Sequence alignment of the orphan receptor (CTR) and receptors of the OT/VP superfamily. The amino acid sequence of the orphan receptor is aligned with those of *Lymanaea* conopressin receptors (LSCPR) 1 and 2 (van Kesteren *et al.* 1995, 1996), the human OT receptor (Kimura *et al.* 1992), the human V1a receptor (Thibonnier *et al.* 1994), the human V1b receptor (Sugimoto *et al.* 1994) and the human V2 receptor (Lolait *et al.* 1992). Amino acid residues that are identical in all receptors are indicated by an asterisk. Amino acid residues in boxes are highly conserved in the vertebrate receptors of the OT/VP superfamily, but not in other GPCRs, and have been suggested to be important in receptor–ligand interactions (Sharif & Hanley 1992). Residues in the agonist-binding pocket described in the text are shown by arrows. Bars indicate the seven putative transmembrane domains.

(Fig. 3D). Thus the cloned receptor seems to be specific to CT. The EC_{50} for CT was estimated from the dose–response curve to be 14.0 ± 1.7 nM (Fig. 3C). This value is comparable to those for OT at the human OT receptor and Lys–conopressin at the LSCPR1, which are also expressed in *Xenopus* oocytes (Kimura *et al.* 1992, van Kesteren *et al.* 1995, 1996).

Isolation of an endogenous ligand from Octopus brain

To search for a genuine endogenous ligand for the cloned receptor, we assayed HPLC fractions of the *Octopus* subesophageal brain extract in oocytes expressing the cloned receptor. Three steps of HPLC yielded a single material that showed potent activity in the oocyte assay (Fig. 4). Apart from this material, no fractions exhibited active signals in the oocyte assay. The amino acid sequence

of the purified material was X-Try-Phe-Arg-Asn-X-Pro-Ile-Gly (X is an unidentified residue). Sequencing profiles at the first and the sixth cycles in a protein sequencer were similar to those of CT: the first half-cysteine at position 1 did not appear on the chromatogram, whereas the second half-cysteine at position 6 was detected as two specific peaks: the dithiothreitol adduct of PTH-dehydroalanine and PTH-Cys (Nokihara *et al.* 1990, Takuwa–Kuroda *et al.* 2003). The observed mass number, $535.22 [M+2H]^{2+}$ in the nano ESI-TOF-MS analysis was very close to the mass number ($535.25 [M+2H]^{2+}$) calculated for CT. Moreover, the fragmentation pattern of the isolated peptide in the TOF-MS/MS analysis was superimposable on that of CT. Taking these results together, we conclude that the cloned *Octopus* homologue of receptors of the OT/VP superfamily is the CTR.

Figure 3 Functional expression of the cloned orphan receptor in *Xenopus* oocytes. (A) Membrane current induced by CT at 10^{-7} M in an oocyte expressing the receptor. CT was applied during the period indicated by the black bar. (B) No current was triggered by application of 10^{-6} M OP, also indicated by a black bar. (C) Dose-response curve from the concentration range of 10^{-10} M to 10^{-6} M CT. Maximum membrane currents elicited by CT are plotted. The current caused at 10^{-6} M CT was taken as 100%. The mean of four independent experiments at each dose is shown with S.E.M. (error bars). (D) Membrane current elicited by 10^{-6} M OT.

Localization of CTR mRNA

As CTR mRNA was not detected by northern blot analysis (data not shown), RT-PCR Southern blot analysis was performed to determine possible expression of the CTR genes in the central and peripheral nervous systems and in several peripheral tissues of *Octopus*. Figure 5 shows that CTR transcripts are present in the brain, the buccal ganglion, the gastric ganglion, the olfactory lobe and the peduncle lobe, the optic lobe, the pancreas, the oviduct and the ovary.

Antagonists

Many subtype-specific peptide or non-peptide antagonists of VP and OT have been developed to meet clinical demand. In this study, four peptide antagonists were tested to determine whether the activity of CT at the CTR expressed in oocytes was inhibited or not. An OT antag-

onist [d(CH₂)₅¹, Tyr(Me)², Orn⁸]-OT at 10^{-5} M showed no effects on the response to 10^{-7} M CT (Fig. 6A). The following antagonists, tested at 10^{-5} M, completely inhibited the response to 10^{-7} M CT (Fig. 6B, C, D): an OT antagonist [d(CH₂)₅¹, Tyr(Me)², Thr⁴, Orn⁸, Tyr-NH₂⁹]-vasotocin; a V1 antagonist [d(CH₂)₅¹, Tyr(Me)², Arg⁸]-VP; a V2 antagonist [d(CH₂)₅¹, D-Ile², Ile⁴, Arg⁸]-VP.

Discussion

We have previously shown that, in common with vertebrates, the common octopus, *Octopus vulgaris*, possesses two members of the OT/VP superfamily, OP and CT – the first time this was observed in invertebrates (Takuwa-Kuroda *et al.* 2003). As OP and CT have different biological activities, the presence of specific receptors has been proposed. We have cloned receptors of

Figure 4 Purification of the endogenous ligand of the CTR from *Octopus* suboesophageal brain. (A) HPLC fractionation on a Capcell Pak UG80 column. Fraction 26 showed a Cl^- current in the oocyte assay. (B) Fraction 26 in (A) was condensed *in vacuo*, dissolved in 20 mM phosphate buffer, and injected into a cation-exchange HPLC with TSKgel SP-5PW column. Fraction 14 was active in the oocyte assay. (C) Fraction 14 in (B) was separated by a reversed-phase HPLC with a Capcell Pak C18 MG80 column. Fractions 21 and 22 both exhibited Cl^- currents. The peak was outside the range of detection in the chromatogram and collected in two fractions, but a single material was eluted at the peak.

Figure 5 Southern blot analysis of RT-PCR products for CTR (upper panel) and actin (lower panel) transcripts isolated from the *Octopus* tissues. To exclude the possibility that a signal was caused by contamination with genomic DNA during RNA preparation, the PCR-Southern experiment was also performed without preincubation with reverse transcriptase (middle panel). L1: brain; L2: buccal ganglion; L3: gastric ganglion; L4: oesophagus; L5: crop; L6: stomach; L7: rectum; L8: aorta; L9: systemic heart; L10: salivary gland; L11: vas deferens; L12: testis; L13: optic gland; L14: olfactory lobe and peduncle lobe; L15: optic lobe; L16: kidney; L17: branchia; L18: liver; L19: pancreas; L20: white body; L21: egg; L22: branchial vessel; L23: oviductal ball; L24: oviduct; L25: ovary.

Figure 6 Effects of OT and VP antagonists on the response of oocyte-expressed CTR to CT. A mixture of CT (10⁻⁷ M) and an antagonist (10⁻⁵ M) was applied during the period indicated by the black bar. (A) OT antagonist [d(CH₂)₅¹, Tyr(Me)², Orn⁸]-OT. (B) OT antagonist [d(CH₂)₅¹, Tyr(Me)², Thr⁴, Orn⁸, Tyr-NH₂⁹]-vasotocin. (C) V1 antagonist [d(CH₂)₅¹, Tyr(Me)², Arg⁸]-VP. (D) V2 antagonist [d(CH₂)₅¹, D-Ile², Ile⁴, Arg⁸]-VP.

OP and CT from the *Octopus* brain by PCR amplification, using degenerate primers based on the amino acid sequences in the conserved transmembrane regions of the mammalian OT and VP receptors and Lys-conopressin receptors. The cloned orphan receptor is characteristic of a GPCR. Moreover, on the basis of amino acid sequence alignment and homologies in the transmembrane domains, the cloned receptor is suggested to be a homologue of receptors of the OT/VP superfamily.

The agonist-binding site for the OT and VP receptors has been suggested to be located in a narrow cleft delimited by the ringlike arrangement of the transmembrane domains (Mouillac *et al.* 1995). Five Gln residues in the transmembrane II, III, IV and VI and a Lys residue localized in transmembrane III are well conserved in all the OT and VP receptors (Mouillac *et al.* 1995) and are important for the interaction of OT and VP with their receptors (see Fig. 2). Four of them are also conserved in the cloned receptor (Gln^{102,125,179,315}). The remaining one, the Gln¹⁰⁴ residue in the V1a receptor, is substituted by the analogous Asn⁹⁸ residue. The Lys¹²⁸ residue in the V1a receptor, which is proposed to interact with the Gln residue at position 4 in VP, is substituted by a Thr¹²² residue. The residue at position 4 in CT is a basic one, Arg. It may be unfavourable for interaction if a basic side chain is present at position 122. However, the ligand of the cloned receptor was not clarified at this stage. The residues at position 8 in OT and VP are known to be necessary for their high-affinity and selective binding to their receptors. Molecular modelling of VP interaction with the V1a receptor supported the view that the side

chain of Arg⁸ in VP projects outside the transmembrane core of the receptor and could interact with Tyr¹¹⁵ in the first extracellular loop (Chini *et al.* 1995). When the Tyr¹¹⁵ residue in the V1a receptor is replaced by Asp or Phe, the residue naturally occurring in the human V2 and OT receptors (see Fig. 2), a potent increase in V2 or OT agonist-binding affinities is induced (Chini *et al.* 1995). The Phe¹⁰⁹ residue homologous to Phe¹⁰³ in the human OT receptor is found in the cloned receptor. The residue at position 8 in CT and OP is Ile. Thus the cloned receptor seems to conserve the agonist-binding pocket common to OT and VP receptors.

Two adjacent Cys residues, which are found 15 residues downstream of transmembrane domain VII in the vertebrate OT/VP superfamily of receptors and LSCPR2 (van Kesteren *et al.* 1996), may be responsible for anchoring the C-terminus to the plasma membrane through palmitoylation (see Fig. 2). These Cys residues have also been observed for the β -adrenergic receptor (O'Dowd *et al.* 1989). However, palmitoylation does not seem to be important for receptor binding and signalling at the human V2 receptor (Sadeghi *et al.* 1995, 1997). That may also be the case in the cloned receptor, as only a Cys residue is present at the corresponding position in transmembrane domain VII.

Xenopus oocytes that express the orphan receptor responded to the application of CT by an induction of membrane Cl⁻ currents coupled to the inositol phosphate/Ca²⁺ pathway. OP and the other members of the OT/VP superfamily did not activate the expressed receptor. Only OT at 10⁻⁶ M had any effect, and it was a weak one (see Fig. 3), which might be attributable to the conserved Phe¹⁰⁹ in the receptor. HPLC fractionation of the *Octopus* brain extract, combined with the oocyte assay, yielded a single substance identical to CT (see Fig. 4). Thus it was revealed that the cloned receptor is the CTR.

Expression of CTR mRNA

Expression of CTR mRNA was detected in *Octopus* in the central and peripheral nervous systems, the pancreas, the oviduct and the ovary (see Fig. 5).

In rats, OT receptors are abundantly present in several brain regions, namely some cortical areas, the olfactory system, the basal ganglia, the limbic system, the thalamus, the hypothalamus, the brain stem and the spinal cord. OT and VP are considered to have pivotal roles in various aspects of sexual, social, feeding and learned behaviour (reviewed by Gimpl & Fahrenholz 2001). The expression of the CTR gene in the brain may suggest the involvement of CT in neurotransmission, neuromodulation, or both, in the *Octopus* central nervous system. The expression of CTR in the optic lobe, the olfactory lobe and the peduncle lobe may suggest the role of CT in sensory systems such as vision and olfaction, and its expression in the

buccal ganglion and the gastric ganglion may suggest a contribution of CT to feeding behaviour.

In the pancreas of mouse and rat, OT and VP increased the release of somatostatin, glucagon and insulin. OT and VP amplified glucose-induced release of insulin, probably via stimulation of phosphoinositide metabolism (Gao *et al.* 1991, Yibchok-Anun *et al.* 1999). In *Lymnaea*, insulin-related peptides are released from neuroendocrine cells, light-green cells, which are located in the cerebral ganglion, and control carbohydrate metabolism. Lys-conopressin induced a depolarizing response in the light-green cells, which express LSCPR1 (van Kesteren *et al.* 1995). Although insulin-related peptides have not been identified, and the presence of endocrine cells like the islets of Langerhans has not been investigated, in the *Octopus* pancreas, the expression of the *CTR* gene may be interesting from a comparative endocrinological point of view.

The ovary has been shown to contain OT, and may be a site of local production of OT in mammals (Ivell & Richter 1984). Functional OT receptors have been detected in bovine granulosa cells, suggesting that OT may be an autocrine factor during follicular growth (Okuda *et al.* 1997). OT also increased the rate of mouse blastocyst development and might therefore play some part in the early stage of development of fertilized oocytes (Furuya *et al.* 1995). Although the levels of expression in the ovary are very low, *CTR* may mediate some of the processes in fertilization and early embryonic development in *Octopus*. The oviduct expressed the *CTR* gene, but CT had no contractile effects on the oviduct (Takuwa-Kuroda *et al.* 2003). The *Octopus* ovary sac contains various stages of oocytes. The mature eggs travel down the proximal oviduct into the oviductal gland, where they become encapsulated (Villanueva 1992). It is not unlikely that one of the activities in which CT may be involved in the oviduct might be maturation and encapsulation of eggs.

CT was also expressed in neurons in the ventral median vasomotor lobe. Some of these send out axons and form an extensive neuropile inside the vena cava in direct contact with the circulating blood, making a neurosecretory system (Takuwa-Kuroda *et al.* 2003). It is suggested that CT may be carried to peripheral tissues that express the *CTR* gene, not only by nerve fibres, but also by the bloodstream after being secreted into the vena cava.

Response of the *CTR* to CT

The response of the *CTR* to CT was inhibited by both OT and VP antagonists (see Fig. 6). V1 and V2 receptors are coupled to different signalling transduction systems and V1 and V2 antagonists used in this experiments show good selectivity for each of receptors. However, both antagonists completely inhibited the response to CT. This result means that the *CTR* does not conserve properties of V1 and V2 selectivity. Nevertheless, the *CTR* may not

belong to any of the known subtypes of OT and VP receptors. The differences in amino acid residues, in the aromaticity at position 3, and in the polarity at position 8 are believed to enable peptides of the OT and VP families to discriminate and interact with their respective receptors in vertebrates (Barberis *et al.* 1998). Such discrimination based on the residue at position 8 does not make sense in invertebrates (Satake *et al.* 1999, Takuwa-Kuroda *et al.* 2003). Both Lys-conopressin and its Ile-analog at position 8 could activate LSCPR2 (van Kesteren *et al.* 1996). Both annetocin (Thr at position 8) and Lys-conopressin induced egg-laying-like behaviour in the leech (Fujino *et al.* 1999). Our findings in the present study suggest that peptides and receptors of the OT/VP superfamily in invertebrates do not follow the rule for vertebrates, but have their own alternative one.

In conclusion, we have cloned a novel receptor of the OT/VP superfamily that may mediate physiological functions such as neurotransmission, reproduction and metabolism in *Octopus*. Functional experiments show that the endogenous ligand for the cloned receptor is CT, and that the receptor can be slightly activated by OT, but not by other members of the OT/VP superfamily, including OP. As *Octopus* enjoys a highly developed brain and endocrine system, somewhat like vertebrates, it may have gained two OT/VP superfamily genes and specific receptors during the evolution of the molluscs.

Acknowledgements

The authors gratefully acknowledge Dr Terumi Nakajima (Professor emeritus, The University of Tokyo) for his encouragement of this work.

Funding

A part of this work was funded by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science to H M (12640669, 13024274).

References

- Barberis C, Mouillac B & Durroux T 1998 Structural bases of vasopressin/oxytocin receptor function. *Journal of Endocrinology* **156** 223–229.
- Bockaert J & Pin JP 1999 Molecular tinkering of G protein-coupled receptors: an evolutionary success. *EMBO Journal* **18** 1723–1729.
- Chini B, Mouillac B, Ala Y, Balestre MN, Trumpp-Kallmeyer S, Hoflack J, Elands J, Hibert M, Manning M, Jard S *et al.* 1995 Tyr115 is the key residue for determining agonist selectivity in the V1a vasopressin receptor. *EMBO Journal* **14** 2176–2182.
- Fujino Y, Nagahama T, Oumi T, Ukena K, Morishita F, Furukawa Y, Matsushima O, Ando M, Takahama H, Satake H *et al.* 1999 Possible functions of oxytocin/vasopressin-superfamily peptides in annelids with special reference to reproduction and osmoregulation. *Journal of Experimental Zoology* **284** 401–406.

- Furuya K, Mizumoto Y, Makimura N, Mitsui C, Murakami M, Tokuoaka S, Ishikawa N, Nagata I, Kimura T & Ivell R 1995 A novel biological aspect of ovarian oxytocin: gene expression of oxytocin and oxytocin receptor in cumulus/luteal cells and the effect of oxytocin on embryogenesis in fertilized oocytes. *Advances in Experimental Medicine and Biology* **395** 523–528.
- Gao ZY, Drews G & Henquin JC 1991 Mechanisms of the stimulation of insulin release by oxytocin in normal mouse islets. *Biochemical Journal* **276** 169–174.
- Gimpl G & Fahrenholz F 2001 The oxytocin receptor system: structure, function, and regulation. *Physiological Review* **81** 629–683.
- Gorbulev V, Buchner H, Akhundova A & Fahrenholz F 1993 Molecular cloning and functional characterization of V2 [8-lysine] vasopressin and oxytocin receptors from a pig kidney cell line. *European Journal of Biochemistry* **215** 1–7.
- Hoyle CH 1998 Neuropeptide families: evolutionary perspectives. *Regulatory Peptides* **73** 1–33.
- Ivell R & Richter D 1984 The gene for the hypothalamic peptide hormone oxytocin is highly expressed in the bovine corpus luteum: biosynthesis, structure and sequence analysis. *EMBO Journal* **3** 2351–2354.
- Iwakoshi E, Hisada M & Minakata H 2000 Cardioactive peptides isolated from the brain of a Japanese octopus, *Octopus minor*. *Peptides* **21** 623–630.
- van Kesteren RE, Tensen CP, Smit AB, van Minnen J, van Soest PF, Kits KS, Meyerhof W, Richter D, van Heerikhuizen H, Vreugdenhil E *et al.* 1995 A novel G protein-coupled receptor mediating both vasopressin- and oxytocin-like functions of Lys-conopressin in *Lymnaea stagnalis*. *Neuron* **4** 897–908.
- van Kesteren RE, Tensen CP, Smit AB, van Minnen J, Kolakowski LF, Meyerhof W, Richter D, van Heerikhuizen H, Vreugdenhil E & Geraerts WP. 1996 Co-evolution of ligand-receptor pairs in the vasopressin/oxytocin superfamily of bioactive peptides. *Journal of Biological Chemistry* **271** 3619–3626.
- Kimura T, Tanizawa O, Mori K, Brownstein MJ & Okayama H 1992 Structure and expression of a human oxytocin receptor. *Nature* **356** 526–529.
- Kyte J & Doolittle RF 1982 A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology* **157** 105–132.
- Lolait SJ, O'Carroll AM, McBride OW, Konig M, Morel A & Brownstein MJ 1992 Cloning and characterization of a vasopressin V2 receptor and possible link to nephrogenic diabetes insipidus. *Nature* **357** 336–339.
- Mouillac B, Chini B, Balestre MN, Elands J, Trumpp-Kallmeyer S, Hoflack J, Hibert M, Jard S & Barberis C 1995 The binding site of neuropeptide vasopressin V1a receptor. Evidence for a major localization within transmembrane regions. *Journal of Biological Chemistry* **270** 25771–25777.
- Nokihara K, Morita N, Yamaguchi M & Watanabe T 1990 Routine determination of cystine containing peptides and proteins using protein sequencer, Shimadzu PSQ-1, with isocratic separation of PTH-amino acids. *Proceedings of the 28th Symposium on Peptide Chemistry*, Osaka, Japan. 159–164.
- O'Dowd BF, Hnatowich M, Caron MG, Lefkowitz RJ & Bouvier M 1989 Palmitoylation of the human beta 2-adrenergic receptor. Mutation of Cys341 in the carboxyl tail leads to an uncoupled nonpalmitoylated form of the receptor. *Journal of Biological Chemistry* **264** 7564–7569.
- Okuda K, Uenoyama Y, Fujita Y, Iga K, Sakamoto K & Kimura T 1997 Functional oxytocin receptors in bovine granulosa cells. *Biology of Reproduction* **56** 625–631.
- Probst WC, Snyder LA, Schuster DI, Brosius J & Sealfon SC 1992 Sequence alignment of the G-protein coupled receptor superfamily. *DNA and Cell Biology* **11** 1–20.
- Reich G 1992 A new peptide of the oxytocin/vasopressin family isolated from nerves of the cephalopod *Octopus vulgaris*. *Neuroscience Letters* **134** 191–194.
- Sadeghi HM, Innamorati G & Birnbaumer M 1995 Functional properties of mutant V2 receptors. In *Neurohypophysis: Recent Progress of Vasopressin and Oxytocin Research*, pp 445–451. Eds T Saito, K Kurokawa & S Yoshida. Amsterdam: Elsevier.
- Sadeghi HM, Innamorati G, Dagarag M & Birnbaumer M 1997 Palmitoylation of the V2 vasopressin receptor. *Molecular Pharmacology* **52** 21–29.
- Satake H, Takuwa K, Minakata H & Matsushima O 1991 Evidence for conservation of the vasopressin/oxytocin superfamily in Annelida. *Journal of Biological Chemistry* **274** 5605–5611.
- Sharif M & Hanley MR 1992 Peptide receptors. Stepping up the pressure. *Nature* **357** 279–280.
- Sugimoto T, Saito M, Mochizuki S, Watanabe Y, Hashimoto S & Kawashima H 1994 Molecular cloning and functional expression of a cDNA encoding the human V1b vasopressin receptor. *Journal of Biological Chemistry* **269** 27088–27092.
- Takuwa-Kuroda K, Iwakoshi-Ukena E, Kanda A & Minakata H 2003 *Octopus*, which owns the most advanced brain in invertebrates, has two members of vasopressin/oxytocin superfamily as in vertebrates. *Regulatory Peptides* **115** 139–149.
- Thibonnier M, Auzan C, Madhun Z, Wilkins P, Berti-Mattera L & Clauser E 1994 Molecular cloning, sequencing, and functional expression of a cDNA encoding the human V1a vasopressin receptor. *Journal of Biological Chemistry* **269** 3304–3310.
- Villanueva R 1992 Continuous spawning in the cirrate octopods *Opisthoteuthis agassizi* and *O. vossi*: features of sexual maturation defining a reproductive strategy in cephalopods. *Marine Biology* **114** 265–275.
- Yibchok-Anun S, Cheng H, Heine PA & Hsu WH 1999 Characterization of receptors mediating AVP- and OT-induced glucagon release from the rat pancreas. *American Journal of Physiology* **277** E56–E62.

Received 3 July 2003

Accepted 1 August 2003

Made available online as an

Accepted Preprint 8 August 2003