RAPID COMMUNICATION

Steroidogenic factor-1 enhances basal and forskolin-stimulated transcription of the human glycoprotein hormone α -subunit gene in GH₃ cells

Robert C Fowkes and Jacky M Burrin

Department of Endocrinology, Barts and the Royal London School of Medicine and Dentistry, West Smithfield, London EC1A 7BE, UK

(Requests for offprints should be addressed to R C Fowkes who is now at Department of Physiology, University of California San Francisco, 513 Parnassus Avenue, San Francisco, CA 94143-0444, USA; Email: rfow0187@itsa.ucsf.edu)

Abstract

Steroidogenic factor-1 (SF-1) is a key regulator of endocrine development, and mediates expression of gonadotrophin-specific genes in the pituitary. Basal and hormone stimulated transcription of the human glycoprotein hormone α -subunit gene (α GSU) in gonadotrophs involves SF-1 and its cognate binding site, the gonadotroph-specific element (GSE). In this study, we demonstrate that SF-1 significantly enhances basal and forskolin-stimulated transcription of the human α GSU promoter in GH₃ cells. Mutation of the GSE abolished the

SF-1-mediated transactivation of basal α GSU promoter activity, and significantly attenuated the forskolin effect by 50%. Mutation of the Ser203 residue in SF-1 to Ala blocked basal transactivation of α GSU promoter activity, and halved the forskolin effect. These data collectively reveal a direct role for SF-1 and the GSE in mediating basal and forskolin-stimulated transcription of the human α GSU promoter in GH₃ cells. The phosphorylation site at Ser203 appears to be required for these effects.

Journal of Endocrinology (2003) **179**, R1–R6

Introduction

The orphan nuclear receptor, steroidogenic factor-1 (SF-1, NR5A1) is expressed in gonadal and adrenal tissues, where it plays an important role in endocrine development and maintenance of steroidogenic gene expression (Parker *et al.* 2002). However, SF-1 is also expressed in the ventromedial hypothalamus and in gonadotrophs of the anterior pituitary (Ingraham *et al.* 1994). In gonadotrophs, SF-1 regulates the expression of the gonadotrophin subunit genes, glycoprotein hormone α -subunit (α GSU), luteinising hormone- β (LH β), follicle-stimulating hormone- β (FSH- β), as well as other important genes such as the gonadotrophin releasing hormone receptor (GnRH-R) and neuronal nitric oxide synthase (nNOS) (for review see Fowkes and Burrin 2003).

The human α GSU promoter is regulated by several response elements which reside in the proximal 435 bp of the promoter (Heckert *et al.* 1996). We have recently shown that SF-1 and its cognate binding site in the α GSU promoter (the gonadotroph-specific element, GSE) contribute to basal and hormone-stimulated α GSU transcription in α T3–1 and L β T2 gonadotroph cell lines (Fowkes *et al.* 2002, Fowkes *et al.* 2003). As these cell lines express

high levels of endogenous SF-1, previous investigators have utilised the rat somatolactotroph GH₃ cell line as a surrogate gonadotroph model to examine the role of exogenous SF-1 on gonadotrophin gene expression (Halvorson *et al.* 1999, Kaiser *et al.* 2000). The regulation of α GSU by SF-1 in GH₃ cells has been poorly elucidated.

SF-1 transcriptional activity is regulated by post-translational modification, such as acetylation and phosphorylation (for review see Fowkes and Burrin 2003). The Serine 203 residue of SF-1 resides in a consensus mitogen activated protein kinase (MAPK) sequence, and has been shown to be phosphorylated by these kinases *in vitro* (Hammer *et al.* 1999, Gyles *et al.* 2001, Desclozeaux *et al.* 2002, Fowkes *et al.* 2003). Mutation of Ser203 to Ala (S203A) reduces SF-1 transcriptional activity and co-activator recruitment (Hammer *et al.* 1999), and S203A-SF-1 does not transactivate the α GSU promoter in α T3–1 cells (Fowkes *et al.* 2003). However, it is unclear whether this residue is important for mediating the effects of SF-1 in heterologous expression models such as GH₃ cells.

Our recent observations show that SF-1 can mediate pituitary adenylate cyclase-activating polypeptide (PACAP)-stimulated α GSU transcription in gonadotrophs (Fowkes *et al.* 2003). In this study, we show that SF-1 and

the GSE mediate basal and forskolin-stimulated α GSU transcription in GH₃ cells, an effect which requires an intact Ser 203 residue in SF-1 for maximal activation.

Materials and Methods

Materials

All chemicals were purchased from Sigma (Sigma Chemicals, Poole, Dorset, UK) unless otherwise stated. Forskolin (FSK) was stored as a stock solution of 100 mM in DMSO at 4 °C, before dilution in culture medium.

Plasmids used in transfection studies

An expression vector encoding mouse SF-1 was prepared by sub-cloning the mouse full-length SF-1 cDNA sequence (from Prof. K L Parker, UT SouthWestern, TX, USA) into pCIneo (Promega Corporation, Southampton, UK), an expression vector containing a cytomegalovirus promoter for high levels of constitutive expression and imparting neomycin resistance. Restriction endonuclease digests verified correct orientation of the SF-1 coding sequence, and the vector was termed pCI-SF-1 neo. The reporter construct -517 α LUC contains 517 bp of the 5' flanking sequence and 44 bp of exon 1 of the human α GSU gene, linked to the luciferase (LUC) reporter gene in the plasmid pA3 LUC. The promoter-less LUC expression vector pA3 LUC was used as an internal control plasmid for basal luciferase expression. The internal control plasmid Bos β Gal contains the promoter of the human elongation factor 1 gene driving expression of β -galactosidase, and was used as an internal control to normalize transfection efficiencies. The expression vector encoding the mutant SF-1, S203A-SF-1 in pCIneo plasmid, and the prolactin promoter containing 5000 bp of the human PRL promoter (PRL-LUC) have been described previously (Hammer *et al.* 1999, Takasuka *et al.* 1998). All constructs were verified for orientation and correct sequence by restriction endonuclease digests and the dideoxy-DNA sequencing method. Large-scale preparation and purification of plasmids were performed by alkaline lysis and resin purification (Qiagen).

Site-directed Mutagenesis

The -517 α LUC construct was sub-cloned in to the HindIII site in pBluescript and site-directed mutagenesis using the QuikChange kit (Stratagene, Cambridge, UK) was conducted to introduce a 2 bp mutation within the GSE region. Mutagenesis was performed according to manufacturer's instructions and using the following primers:

forward: 5'-CTCTCTTTTCATGGGCTGATTTTGTCGTACCATCACCTG-3', reverse: 5'-CAGGTG

ATGGTGACGACAAAATCAGCCCATGAAAAGAGAG-3', with the mutation highlighted. Following sequence analysis, the mutated -517 α LUC (named -517 α MUT) was cloned back in to the HindIII site of pA3 LUC.

Generation of stable cell lines

Sub-confluent GH₃ cells were transfected with pCIneo or pCI-SF-1 neo by electroporation. Briefly, 1×10^6 cells were placed in 0.8 ml of ice-cold DMEM containing 10 μ g of either pCIneo (GH₃neo cells) or pCI-SF-1neo (GH₃SF-1 cells). The cell suspension containing the DNA to be transfected was transferred to an electroporation cuvette (Biorad Laboratories Ltd, Herts., UK). The cells received a single electrical pulse of 250 V from a total capacitance of 1080 μ F with a 1000 msec discharge interval using the PG200 Progenitor II electroporator (Hoefer Scientific Instruments, Staffs., UK). Transfected cells were then selected with the Neomycin analogue Geneticin G418 (Invitrogen, Paisley, UK) at a concentration (250 μ g/ml) shown in preliminary experiments to kill all untransfected GH₃ cells (data not shown). Continuous G418 exposure selected cells carrying the stably integrated Neomycin resistance gene and surviving colonies were propagated as a mixed population of cells and maintained in tissue culture medium containing G418.

Transient expression assays

GH₃SF-1 and GH₃NEO cells (3×10^6 cells/cuvette) were transfected by electroporation as described above with 10 μ g of the expression vectors pA3 LUC, -517 α LUC, -517 α MUT or PRL-LUC in the presence of 5 μ g of the control plasmid Bos β Gal. Forty hours after electroporation, medium was changed and cells were treated with or without FSK (10 mM) for a further 8 h. Cells were harvested after washing the cell monolayer with ice-cold phosphate buffered saline (PBS), and the cellular extracts were assayed for luciferase and β -galactosidase activity as previously described (Holdstock *et al.* 1996). In some experiments, 5 μ g of SF-1, S203A-SF-1 or pCIneo were co-transfected with the -517 α LUC and Bos β Gal. Luciferase data from separate experiments were pooled by normalizing the data to the level of β -galactosidase activity.

Statistical analysis

All graphical data were prepared using GraphPad Prism 3.0 (GraphPad, San Diego, USA) and analyzed using pre-programmed analysis equations within Prism. Transfection data are presented as normalized data pooled from multiple experiments (each in triplicate, and performed at least twice). Where appropriate, an ANOVA was

performed on data followed by Student's *t*-test or Tukey's multiple comparisons test, accepting $P < 0.05$ as significant.

Results

SF-1 regulation of α GSU promoter activity in transfected GH₃ cells requires the GSE

To examine the role of SF-1 and the GSE in GH₃ cells, we created a stable GH₃ cell-line that expressed SF-1. Use of an SF-1 deficient cell line allowed us to investigate FSK effects on basal as well as SF-1 stimulated α GSU promoter activity. Our initial studies compared basal luciferase expression following transient transfection of the GH₃SF-1 and GH₃NEO cells with -517α LUC and Bos β Gal, before harvesting 48 h later. Basal α GSU promoter activity was 2.6 ± 0.6 -fold greater in GH₃SF-1 cells compared with GH₃NEO cells ($P < 0.05$, Fig. 1A). To establish whether the presence of SF-1 could alter the transcriptional response to FSK, the same transfection was performed but with 0 or 10 μ M FSK present for the last 8 h. In the absence of SF-1, FSK stimulated α GSU promoter activity by 4.3 ± 0.8 -fold over basal (Fig. 1B, $P < 0.001$). However, the FSK effect in GH₃SF-1 cells was still significantly greater, by 2.2 ± 0.2 -fold ($P < 0.001$). To determine whether this effect was promoter specific, the FSK effect on the prolactin promoter (PRL-LUC) was compared between cell lines, and showed no significant difference in the absence or presence of SF-1 (Fig. 1C).

To examine the role of the GSE in mediating the observed effects of SF-1 in GH₃ cells, we transiently transfected -517α LUC or -517α MUT constructs into both GH₃SF-1 and GH₃NEO cells. Basal expression of -517α LUC was 3.4 ± 1.0 -fold greater in the stably transfected GH₃SF-1 cell line as compared with GH₃NEO ($P < 0.05$), but mutation of the GSE completely abolished this increase in basal luciferase activity (Fig. 2A). Thus, the basal enhancement of α GSU expression achieved by stable transfection of SF-1 is mediated by the GSE. The expression of the promoter-less control pA3 LUC was not significantly different in either cell line (data not shown).

Following transient transfection of GH₃SF-1 or GH₃NEO cells with either -517α LUC or -517α MUT, cells were stimulated in the presence or absence of FSK (10 μ M) for 8 h. As expected, -517α LUC promoter activity following FSK stimulation was significantly greater in GH₃SF-1 cells compared with GH₃NEO cells (13.3 ± 2.2 -fold cf 6.3 ± 1.5 -fold, $P < 0.05$). However, in the SF-1 expressing cells, mutation of the GSE reduced FSK-stimulated α GSU promoter activity to 2.8 ± 1.2 -fold when using -517α MUT, further indicating that SF-1 acts at the GSE to mediate the response to cAMP ($P < 0.01$, Fig. 2B). In GH₃NEO cells, a similar FSK response was observed with the wild-type (6.3 ± 1.5 -fold) and

Figure 1 Effect of SF-1 on basal and FSK-stimulated α GSU promoter activity. A, GH₃SF-1 and GH₃NEO cells were transiently transfected with -517α LUC and Bos β Gal and harvested 48 h later. B, GH₃SF-1 and GH₃NEO cells were transfected as above, and stimulated with 0 or 10 μ M FSK for 8 h. C, GH₃SF-1 and GH₃NEO cells were transfected with PRL-LUC and Bos β Gal, and stimulated with 0 or 10 μ M FSK for 8 h. All data represents the mean $\pm$ S.E.M. of at least 3 independent experiments each performed in triplicate. *, $P < 0.05$, *** $P < 0.001$, significantly different as indicated.

mutated -517α MUT constructs (5.8 ± 1.8 -fold). These data suggest that the enhanced basal and FSK-responsiveness seen in GH₃SF-1 cells was specific to SF-1 binding at the GSE.

Figure 2 Role of the GSE in mediating the SF-1 effect on basal and FSK-stimulated α GSU promoter activity. A, GH₃SF-1 and GH₃NEO were transiently transfected with either -517 α LUC or -517 α MUT and harvested 48 h post-transfection. The data are normalized to basal -517 α LUC promoter activity in GH₃NEO cells. Mean fold increase over basal is indicated above the relevant bar. *, $P < 0.05$, significantly different from GH₃NEO. B, GH₃SF-1 and GH₃NEO cells were transfected as above, and stimulated with 0 or 10 mM FSK for 8 h. The data are normalized to basal -517 α LUC or -517 α MUT promoter activity. Mean fold increase over basal is indicated above the relevant bar. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, significantly different from the relevant basal, except where indicated by brackets.

The Ser203 residue in SF-1 is required for full basal and FSK-stimulated transcriptional activity

The MAPK phosphorylation site on Ser203 in SF-1 has been shown to regulate SF-1 activity *in vitro* (Hammer *et al.* 1999, Desclozeaux *et al.* 2002, Fowkes *et al.* 2003). To assess the effect of mutating this residue in SF-1 on the regulation of the human α GSU promoter, we performed a series of transient co-transfections in wild-type GH₃ cells. We co-transfected 5 μ g of the wild-type -517 α LUC promoter with 5 μ g of a wild-type SF-1 expression vector (the maximally effective concentration as determined in preliminary experiments, data not shown), 5 μ g of empty expression vector (pCIneo) or 5 μ g of the S203A-SF-1

Figure 3 Role of Ser 203 in SF-1 on basal and FSK-stimulated α GSU promoter activity in GH₃ cells. A, GH₃ cells were transiently co-transfected with -517 α LUC and either wild-type SF-1, pCIneo or S203A-SF-1 and harvested 48 h after transfection. The data are normalized to -517 α LUC activity with pCIneo. *, $P < 0.05$, significantly different from pCIneo. B, Transiently co-transfected GH₃ cells were cultured for 48 h post-transfection and stimulated with 0 or 10 μ M FSK for the last 8 h before harvesting. The data are normalized to the basal expression of each promoter construct to account for differences in basal activity. Mean fold increase over basal is indicated above the bar. *, $P < 0.05$, ***, $P < 0.001$, significantly different from pCIneo or as indicated.

expression vector in which the Ser203 residue had been converted to alanine to prevent MAPK-mediated phosphorylation. As shown (Fig. 3A), wild-type SF-1 transactivated the α GSU promoter by 1.9 ± 0.2 -fold ($P < 0.05$) compared with empty vector. This effect was absent when S203A-SF-1 was co-transfected with the α GSU promoter (1.0 ± 0.3 -fold). When these co-transfected GH₃ cells were stimulated for 8 h with 10 μ M FSK, the 12.7 ± 2.9 -fold increase seen in the presence of wild-type SF-1 was reduced to 5.2 ± 2.0 -fold in the presence of mutant SF-1 ($P < 0.05$, Fig. 3B). Collectively, these data suggest that an intact MAPK phosphorylation site at Ser203 is required for the SF-1 enhancement of basal and FSK-stimulated α GSU promoter activity in GH₃ cells.

Discussion

Investigation of the role of SF-1 in regulating gonadotrophin gene transcription is often hampered by the high levels of endogenous SF-1 expressed in gonadotroph cell lines. In this study, we have used GH₃ cells that have previously been reported as lacking SF-1 expression but have been used by others as a useful heterologous model to study regulation of expression of the gonadotrophin sub-unit genes (Halvorson *et al.* 1999, Kaiser *et al.* 2000). We used two approaches to examine the role of SF-1 in regulating α GSU promoter activity; generation of a stably transfected GH₃ cell lines (GH₃SF-1 and GH₃NEO), and transient co-transfection of wild-type and mutant SF-1 and α GSU-LUC in native GH₃ cells. Both these approaches yielded comparable results. We observed a 50% decrease in activity with the mutant GSE in GH₃SF-1 cells, similar to that reduction seen in α T3-1 gonadotrophs (Fowkes *et al.* 2003). Furthermore, when a mutant form of SF-1 (S203A-SF-1) was co-transfected with -517 α LUC in GH₃ cells, the observed increases in basal and FSK-stimulated promoter activity were blocked (basal) or reduced by at least 50% (FSK-stimulated).

The failure to completely abrogate the effects of FSK on α GSU promoter activity by either mutating the GSE or SF-1 are not surprising given that multiple response elements can be found within the proximal -435 bp of the human α GSU promoter (Heckert *et al.* 1996). Indeed, tandem cAMP response elements (CRE) reside proximally of the GSE at -146 bp. Our earlier studies revealed that mutation of these elements could significantly blunt basal and PACAP or 8-Br-cAMP-stimulated α GSU promoter activity in α T3-1 cells (Burrin *et al.* 1998) similar to our more recent observations with GSE mutations (Fowkes *et al.* 2002, Fowkes *et al.* 2003). However, mutation of the CRE was not sufficient to completely block the effect of PACAP on the α GSU promoter (Burrin *et al.* 1998). Instead, PACAP responsiveness was only lost when the promoter was deleted at -195 bp, removing the GSE-containing region, even though the CRE's remained intact. It is likely that in addition to the GSE and CREs, other factors are required for maximal hormonal regulation of pituitary α GSU expression.

The GSE has previously been implicated in the basal regulation of α GSU promoter activity (Barnhart and Mellon 1994, Ingraham *et al.* 1994), and similar CC to TT mutations have displayed a comparable 50% inhibition in basal promoter activity. We have recently showed that this same mutation abolishes binding of SF-1 in α T3-1 and L β T2 gonadotrophs by electrophoretic mobility shift assays (Fowkes *et al.* 2002, Fowkes *et al.* 2003), and inhibits both GnRH and PACAP-stimulated α GSU promoter activity when introduced within the context of the -517 α LUC promoter. Our current observations that FSK-stimulated α GSU transcription is partially mediated by SF-1 and the GSE in GH₃ cells adds to a growing list of target gene

promoters where hormone-responsiveness maps to the SF-1 site. For example, FSK and cAMP stimulation in adrenocortical Y-1 cells increases StAR transcription via a MAPK and SF-1-dependent mechanism (Gyles *et al.* 2001). Furthermore, in α T3-1 cells, PACAP and protein kinase A (PKA)-stimulation of the GnRH-R promoter requires an intact SF-1 response element, similar to the GSE (Pincas *et al.* 2001). Thus, the propagation of peptide hormone signalling to gene transcription via SF-1 is likely to involve a number of intracellular signalling pathways that ultimately impinge upon mechanisms of post-translational modification.

Phosphorylation of SF-1 has previously been shown to occur on Ser203 *in vitro*, and is mediated via the activity of extracellular signal-regulated kinase (ERK) members of the MAPK family (Hammer *et al.* 1999, Desclozeaux *et al.* 2001). We recently have shown that PACAP utilises the same ERK/MAPK pathway to enhance phosphorylation of SF-1 in α T3-1 cells (Fowkes *et al.* 2003). Our current observations that basal and FSK-stimulated α GSU promoter activity are impaired in the presence of the S203A mutant further support the requirement for this MAPK phosphorylation site. Furthermore, we have observed that FSK stimulates ERK and CREB phosphorylation, and ERK and CREB-mediated transcription with similar kinetics and magnitude in GH₃ cells (data not shown) compared with PACAP effects in α T3-1 cells (Fowkes *et al.* 2001, Fowkes *et al.* 2003b). It remains to be seen whether S203A is as influential *in vivo* as it clearly is *in vitro*.

The regulation of orphan nuclear receptor activity by peptide hormone signalling is becoming a commonplace occurrence in endocrine tissues (Hammer *et al.* 1999, Desclozeaux *et al.* 2002). In the absence of identified ligand, it is possible that a class of nuclear receptors, similar to SF-1, are predominantly mediated by the propagation of signalling mechanisms from the cell membrane to the nucleus. Our current observations suggest that this is an important mechanism in the regulation of SF-1-mediated gene transcription in pituitary cells.

Acknowledgements

The authors thank Prof. H A Ingraham (UCSF, USA) for providing the S203A SF-1 expression vector, Prof. K L Parker for the wild-type SF-1 expression vector (SWMED, USA), Prof. J R E Davis (University of Manchester, UK) for the PRL-LUC, and Drs M Lee and M Desclozeaux for reading this manuscript. This project was funded by a Wellcome Trust Project grant (to JMB).

References

- Barnhart KM & Mellon PL 1994 The orphan nuclear receptor, steroidogenic factor-1, regulates the glycoprotein hormone α -subunit gene in pituitary gonadotrophs. *Molecular Endocrinology* **8** 878-885

- Burrin JM, Aylwin SJB, Holdstock JG & Sahye U 1998 Mechanism of action of pituitary adenylate cyclase-activating polypeptide on human glycoprotein hormone α -subunit transcription in α T3-1 gonadotropes. *Endocrinology* **139** 1731–1737
- Desclozeaux M, Krylova IN, Horn F, Fletterick RJ & Ingraham HA 2002 Phosphorylation and intramolecular stabilization of the ligand binding domain in the nuclear receptor steroidogenic factor 1. *Molecular and Cellular Biology* **22** 7193–203
- Fowkes RC & Burrin JM 2003 Steroidogenic factor-1 (SF-1): a pivotal regulator of gonadotroph gene expression. *Journal of Endocrinology* **177** 345–350
- Fowkes RC, Burch J & Burrin JM 2001 Stimulation of extracellular-regulated kinase by pituitary adenylate cyclase-activating polypeptide in α T3-1 gonadotrophs. *Journal of Endocrinology* **171** R5–R10
- Fowkes RC, King P & Burrin JM 2002 Regulation of human glycoprotein hormone α -subunit (α GSU) transcription in L β T2 gonadotropes by protein kinase C (PKC) and extracellular signal-regulated kinase 1/2 (ERK1/2). *Biology of Reproduction* **67** 725–734
- Fowkes RC, Desclozeaux M, Patel MV, Aylwin SJ, King P, Ingraham HA & Burrin JM 2003a Steroidogenic factor-1 (SF-1) and the gonadotrope-specific element (GSE) enhance basal and pituitary adenylate cyclase-activating polypeptide (PACAP)-stimulated transcription of the human glycoprotein hormone α -subunit gene (α GSU) in gonadotropes. *Molecular Endocrinology* **17** (In Press)
- Fowkes RC, Sidhu KK, Sosabowski JK, King P & Burrin JM 2003b Absence of pituitary adenylate cyclase activating polypeptide (PACAP)-stimulated transcription of the human glycoprotein α -subunit (α GSU) gene in L β T2 gonadotrophs reveals disrupted cyclic 3'5'-adenosine monophosphate (cAMP)-mediated gene transcription. *Journal of Molecular Endocrinology*, **31** 263–278.
- Gyles SL, Burns CJ, Whitehouse BJ, Sugden D, Marsh PJ, Persaud SJ & Jones PM 2001 ERKs regulate cyclic AMP-induced steroid synthesis through transcription of the steroidogenic acute regulatory (StAR) gene. *Journal of Biological Chemistry* **276** 34888–34895
- Halvorson LM, Kaiser UB & Chin WW 1999 The protein kinase C system acts through the early growth response protein 1 to increase LH β gene expression in synergy with steroidogenic factor-1. *Molecular Endocrinology* **13** 106–116
- Hammer GD, Krylova I, Zhang Y, Darimont BD, Simpson K, Weigel NL & Ingraham HA 1999 Phosphorylation of the nuclear receptor SF-1 modulates cofactor recruitment: integration of hormone signaling in reproduction and stress. *Molecular Cell* **3** 521–526
- Heckert LL, Schultz K & Nilson JH 1996 The cAMP response elements of the α -subunit gene bind similar proteins in trophoblasts and gonadotropes but have distinct functional sequence requirements. *Journal of Biology Chemistry* **271** 31650–31656
- Holdstock JG, Aylwin SJB & Burrin JM 1996 Calcium and glycoprotein hormone α -subunit expression and secretion in α T3-1 gonadotropes. *Molecular Endocrinology* **10** 1308–1317
- Ingraham HA, Lala DS, Ikeda Y, Luo X, Shen WH, Nachtigal MW, Abbud R, Nilson JH & Parker KL 1994 The nuclear receptor steroidogenic factor 1 acts at multiple levels of the reproductive axis. *Genes and Development* **8** 2302–12
- Kaiser UB, Halvorson LM and Chen MT 2000 Sp1, steroidogenic factor 1 (SF-1), and early growth response protein 1 (EGR-1) binding sites form a tripartite gonadotropin-releasing hormone response element in the rat luteinizing hormone- β gene promoter: an integral role for SF-1. *Molecular Endocrinology* **14** 1235–1245
- Parker KL, Rice DA, Lala DS, Ikeda Y, Luo X, Wong M, Bakke M, Zhao L, Frigeri C, Hanley NA *et al.* 2002 Steroidogenic factor 1: an essential mediator of endocrine development. *Recent Progress in Hormone Research* **57** 19–36
- Pincas H, Laverriere JN & Counis R 2001 Pituitary adenylate cyclase-activating polypeptide and cyclic adenosine 3',5'-monophosphate stimulate the promoter activity of the rat gonadotropin-releasing hormone receptor gene via a bipartite response element in gonadotrope-derived cells. *Journal of Biology Chemistry* **276** 23562–71
- Takasuka N, White MR, Wood CD, Robertson WR & Davis JR 1998 Dynamic changes in prolactin promoter activation in individual living lactotrophic cell. *Endocrinology* **139** 1361–1368