COMMENTARY

Photoperiodic regulation of prolactin secretion: changes in intra-pituitary signalling and lactotroph heterogeneity

J D Johnston

School of Biological Sciences, University of Aberdeen, Zoology Building, Tillydrone Avenue, Aberdeen AB24 2TZ, UK
(Requests for offprints should be addressed to J D Johnston; Email: j.johnston@abdn.ac.uk)

Abstract

Many mammalian species utilise day-length (photoperiod) to adapt their physiology to seasonal changes in environmental conditions, via secretion of pineal melatonin. Photoperiodic regulation of prolactin secretion is believed to occur via melatonin-mediated changes in the secretion of a putative prolactin secretagogue, tuberulin, from the pituitary pars tuberalis. Despite the *in vivo* and *in vitro* evidence in support of this intra-pituitary signalling mech-

anism, the identity of tuberulin has yet to be elucidated. This paper reviews recent advances in the characterisation of tuberulin and the regulation of its secretion. Furthermore, the hypothesis that pituitary lactotroph cells display heterogeneity in their response to changing photoperiod and tuberulin secretion is examined.

Journal of Endocrinology (2004) **180**, 351–356

Introduction

Regions of extreme latitude are subject to marked changes in environmental conditions. In order to maximise survival, many mammalian species utilise changing annual day-length (photoperiod) to adapt their physiology in anticipation of these environmental changes. The master circadian clock, located in the suprachiasmatic nuclei (SCN) of the hypothalamus, integrates photoperiodic information (reviewed in Schwartz *et al.* 2001) and continues to reflect ambient day-length even after many months of exposure to constant photoperiod (Carr *et al.* 2003). The SCN innervates the pineal gland via a polysynaptic neural pathway to drive rhythmic, nocturnal secretion of melatonin, the duration of which varies in proportion to the length of the night. Altered melatonin signal duration is then interpreted by target tissues to drive downstream changes in physiology.

In seasonal mammals, photoperiod drives robust rhythms of prolactin secretion from the pars distalis (PD) region of the adenohypophysis. Prolactin secretion peaks during the summer months and reaches a nadir during the winter. Circulating prolactin concentration primarily regulates seasonal changes in pelage, but also has important effects on reproductive status, along with other aspects of physiology and behaviour (reviewed in Lincoln 1989). It was initially thought that melatonin might regulate prolactin secretion via altered release of hypothalamic factors (e.g. dopamine, noradrenaline). However, this mechanism

is now not believed to be a major contributory factor (Steger & Bartke 1991, Badura & Goldman 1992, Lincoln & Clarke 2002).

Compelling *in vivo* evidence for a novel, intra-pituitary mechanism of prolactin regulation came from the hypothalamo-pituitary disconnected (HPD) rat, which bears a surgical lesion of the neural connection between pituitary and hypothalamus, leaving the pituitary circulation intact. Photoperiodic prolactin regulation persists in these animals, suggesting that melatonin acts directly at the level of the pituitary gland to drive changes in prolactin secretion (Lincoln & Clarke 1994). It was subsequently demonstrated that hamsters bearing hypothalamic lesions exhibited normal prolactin rhythms, despite loss of their reproductive response to photoperiod (Maywood & Hastings 1995). These key studies formed the basis of the 'dual site' hypothesis, whereby melatonin acts at the level of the hypothalamus to drive reproductive rhythms, but directly within the pituitary to regulate changes in prolactin secretion (Fig. 1).

The pars tuberalis (PT) and tuberulin

The pituitary PT is a thin layer of the adenohypophysis that surrounds the pituitary stalk and extends rostrally along the ventral surface of the median eminence. It therefore lies in close proximity to the primary plexus of pituitary portal vasculature, an ideal position from which

Figure 1 A model of intra-pituitary mechanisms driving photoperiodic prolactin secretion. Melatonin binds to MT_1 receptors expressed on PT-specific thyrotrophs. The duration of the nocturnal melatonin signal is decoded, through the regulation of circadian clock genes and/or sensitisation, to modulate the stimulation of PT by an endogenous agonist (e.g. adenosine or PACAP). This mechanism regulates the secretion of tuberalin, which stimulates prolactin promoter activity in a subpopulation of lactotrophs, leading to an increase in mRNA expression and hormone secretion. PRL, prolactin.

to modulate the activity of cells in the distal pituitary gland. The PT consists of three main cell types; PD-like endocrine cells, follicular cells and 'PT-specific' cells, which are thyrotrophs morphologically distinct from those found in the PD (reviewed in Wittkowski *et al.* 1999). Seasonal changes in the physiology of PT-specific cells (e.g. Merks *et al.* 1993) are also induced by melatonin administration (Bockers *et al.* 1995), suggesting that the function of the PT is dependent on photoperiod-driven melatonin signal duration (additional references in Wittkowski *et al.* 1999).

High affinity binding sites for the radioligand [125 I]iodomelatonin have been detected in the PT of most species examined (reviewed in Morgan *et al.* 1994), suggesting that it is an important melatonin target tissue. There are two major subtypes of mammalian melatonin receptor, termed MT_1 and MT_2 (previously Mel_{1a} and Mel_{1b} ; reviewed in Reppert *et al.* (1996)), of which MT_1 is the dominant pituitary subtype (Weaver *et al.* 1996,

von Gall *et al.* 2002). Despite transient expression of MT_1 receptors in neonatal pituitary gonadotrophs (Johnston *et al.* 2003b), co-localisation studies have shown that MT_1 is only expressed in the PT-specific thyrotroph cells in adult rodents (Klosen *et al.* 2002, Dardente *et al.* 2003).

The restriction of melatonin receptor expression to the PT in the adult pituitary (Morgan *et al.* 1994), together with the inability of melatonin to directly regulate either prolactin gene expression or secretion in PD cultures (Stirland *et al.* 2001), is consistent with the hypothesis that melatonin drives seasonal changes in prolactin secretion via altered secretion of a PT-derived prolactin secretagogue(s). This hypothesis was supported by the demonstration that ovine PT cells secrete a peptide(s), termed tuberalin, that stimulates prolactin secretion from primary cultures of PD cells (Hazlerigg *et al.* 1996, Morgan *et al.* 1996). Tuberalin secretion was subsequently demonstrated from bovine (Lafarque *et al.* 1998) and hamster (Stirland *et al.* 2001) PT cells, indicating conservation of this PT function between

species. Two key challenges that arose from these observations were to identify tuberalin and also understand the cellular and molecular mechanisms that regulate its secretion.

The extensive list of secretory molecules expressed in PT-specific cells (reviewed in Wittkowski *et al.* 1999) has recently been extended (D'Este *et al.* 2000, Guerra & Rodriguez 2001). However, the ability of many of these molecules to fulfil the putative physiological role of tuberalin has not been rigorously examined. In order to partially characterise the chemical nature of tuberalin, different groups have utilised size fractionation of PT-conditioned medium. It has been suggested that bovine tuberalin is >30 kDa, based on Percoll gradient separation (Lafarque *et al.* 1998). By contrast, exclusion filtration studies indicate that ovine tuberalin is <1 kDa in size (Graham *et al.* 2002). This discrepancy may be due to interspecies differences in tuberalin identity, or contamination of PT-conditioned medium by factors of hypothalamic or serum origin present in primary PT cell cultures. Irrespective of this, the identity of tuberalin remains elusive and to date no candidate molecules have been identified that are both regulated by melatonin and able to stimulate prolactin secretion.

Regulation of tuberalin secretion

In the PT, melatonin acutely inhibits the stimulation of cAMP-dependent signal transduction (reviewed in Hazlerigg *et al.* 2001). Furthermore, melatonin also acutely inhibits the ability of forskolin, a pharmacological stimulator of adenylyl cyclase activity, to increase tuberalin secretion *in vitro* (Morgan *et al.* 1996, Stirland *et al.* 2001, Graham *et al.* 2002). Together, these data suggest that cAMP-mediated pathways play an important role in regulating tuberalin secretion.

Critical to the model described in Fig. 1, tuberalin secretion from the hamster PT is dependent upon prior photoperiod exposure (Stirland *et al.* 2001, Johnston *et al.* 2003a). However, the mechanisms by which melatonin signal duration drives PT physiology are poorly understood. Prolonged (6–24 h) incubation of PT cell cultures with melatonin sensitises subsequent stimulation of cAMP (Hazlerigg *et al.* 1993, von Gall *et al.* 2002), probably resulting from changes in tyrosine phosphorylation of adenylyl cyclase (Barrett *et al.* 2000). This has fuelled speculation that melatonin signal duration may modulate the induction of cAMP by stimulatory factor(s), such as pituitary adenylyl cyclase-activating polypeptide (PACAP) or adenosine (Barrett *et al.* 2002, von Gall *et al.* 2002). Over a range of melatonin signal duration that mimics change from long to short photoperiod, the sensitisation of PT cells *in vitro* increases in proportion to the duration of melatonin incubation (Hazlerigg *et al.* 1993). However, decreased tuberalin secretion and amplitude of

cAMP-responsive gene expression on short days suggests that longer melatonin signals result in decreased cAMP levels *in vivo*. This apparent paradox has yet to be resolved.

A separate avenue of research has provided evidence that altered phasing and/or amplitude of circadian clock gene expression within the PT decodes melatonin signal duration (Messenger *et al.* 1999, Lincoln *et al.* 2002, Johnston *et al.* 2003a). This topic is the subject of a recent review (Lincoln *et al.* 2003).

Unmasking the enigma: how might tuberalin be identified?

Despite these recent advances in the understanding of PT physiology, the identity of tuberalin remains unresolved. Due to the small size of the rodent PT, previous attempts to identify this enigmatic factor(s) have relied upon studies of secretory products from the PT of larger ungulate species. However, even in these animals, PT explants contain median eminence nerve terminals, and therefore it is likely that PT-conditioned medium will be contaminated by hypothalamic factors (Graham *et al.* 2002). In order to progress with this line of research, it would thus be preferable to first generate a population of PT cells that is independent of hypothalamic input. One solution to this problem would be to generate a PT-derived cell line by one of a number of established *in vivo* and *in vitro* immortalisation techniques (e.g. Mellon *et al.* 1991, Katakura *et al.* 1998). PT secretion from these cells could then be compared in various experimental conditions, such as the presence of forskolin with or without melatonin. However, a major drawback of this approach is the uncertainty that the immortalised cells will retain the same physiology as PT cells *in vivo*. An alternative solution would be to compare PT secretory products from primary PT cells derived from HPD rams exposed to appropriate photoperiodic conditions. However, the generation of HPD animals requires complex surgery and therefore can only generate a finite number of PT cells for experimentation.

Rather than analysis of secreted products from the PT, future experiments could take advantage of more recent molecular techniques, such as differential display and microarray technology, to identify genes expressed in the PT in a photoperiod-dependent manner. To be successful, these experiments would rely upon photoperiodic tuberalin regulation occurring at the level of mRNA expression and not merely secretion. Furthermore, if tuberalin is a low molecular weight molecule, as predicted from recent studies of the ovine PT (Graham *et al.* 2002), it is possible that such an approach would not identify mRNA for tuberalin itself. In this instance, one might hypothesise that the gene(s) identified would instead encode enzyme(s) involved in tuberalin biosynthesis.

Figure 2 Schematic representation of putative heterogeneity in prolactin promoter stimulation by tuberlin. It is hypothesised that some lactotroph cells will be unresponsive to tuberlin, while others will show altered steady-state or phasic prolactin promoter activity.

The identification of tuberlin will provide the opportunity for a number of additional studies. Within the field of seasonal endocrinology, it would be of interest to determine whether the identity of tuberlin is conserved amongst photoperiodic mammals. Since the photoperiodic regulation of prolactin secretion is regarded as an 'ancient' aspect of physiology (Lincoln 1999), it might be predicted that the identity of tuberlin would be conserved between species. Additionally, a specific assay for tuberlin synthesis/secretion would provide an important endocrine endpoint to aid studies attempting to understand the decoding of melatonin signal duration by the PT. Such experiments would enable direct examination of the hypothesis that the photoperiodic changes in circadian clock gene expression are directly associated with the endocrine activity of the PT. This line of experimental analysis is currently under active consideration. Finally, in a wider endocrine context, one might also determine if tuberlin is secreted from, or acts upon, any other regions of the neuroendocrine system, distinct from the intrapituitary regulation of prolactin secretion. The ability of PT-conditioned medium to stimulate *c-fos* expression in non-lactotroph cells (Morgan *et al.* 1996) suggests that

tuberlin may play a role in multiple neuroendocrine pathways.

Photoperiodic changes in prolactin transcription and lactotroph heterogeneity

Photoperiodic changes in prolactin secretion are accompanied by robust changes in pituitary prolactin mRNA expression (Hegarty *et al.* 1990, Stirland *et al.* 2001), which may result from changes in either gene transcription or mRNA degradation (Khodursky & Bernstein 2003). Recent studies have shown that both ovine and hamster PT-conditioned media stimulate a prolactin-luciferase reporter construct transfected into lactotroph cells (Stirland *et al.* 2001, 2003, Johnston *et al.* 2003a), suggesting that transcriptional regulation may be a key factor driving photoperiodic prolactin mRNA expression.

Heterogeneity of lactotroph physiology is well established (reviewed in Lamberts & MacLeod 1990). However, the cellular mechanisms underlying the regulation of individual endocrine cells by photoperiod are poorly

understood. A study of freshly dispersed hamster PD cells revealed heterogeneity of prolactin mRNA expression in animals housed under both long and short photoperiod (Johnston *et al.* 2003c). Despite a large overall change in the frequency distribution of prolactin mRNA expression per cell, a similar number of low-expressing cells were observed in both photoperiods suggesting that there may be heterogeneity in the response of lactotrophs to photoperiod. This hypothesis is consistent with the observation that tuberalin is only able to activate *c-fos* expression in a subpopulation of ovine lactotroph cells (Morgan *et al.* 1996). Additional heterogeneity may result from regional differences in tuberalin secretion into the portal vasculature, as has been previously reported for hypothalamic dopamine (Reymond *et al.* 1983). Together, these findings suggest a novel dimension to lactotroph heterogeneity in seasonal mammals.

Individual lactotroph cells exhibit phasic, temporal variation in prolactin gene expression (Takasuka *et al.* 1998, Villalobos *et al.* 1999, Shorte *et al.* 2002, Stirland *et al.* 2003). Individual cells exhibited heterogeneous temporal profiles of promoter activity and also qualitative differences in promoter regulation by prolactin secretagogues; some cells exhibit gradual changes in promoter activity whereas others respond with bursts of transient promoter activity (Takasuka *et al.* 1998, Villalobos *et al.* 1999, Stirland *et al.* 2003). Given these findings, it is therefore plausible that some hamster pituitary cells reported as expressing low levels of prolactin mRNA (Johnston *et al.* 2003c) are actually in a stimulated state, but responded to changing photoperiod with altered temporal patterns of gene expression. In this instance, the timing of pituitary collection and dispersion would influence the mRNA expression profiles obtained. Use of viral transfection technology now permits the efficient transfection of reporter constructs into primary cells (Stirland *et al.* 2003). Pituitary cell transfection, coupled with real-time promoter analysis, could therefore greatly extend our understanding of the regulation of lactotroph physiology by photoperiod and more accurately define the heterogeneity of response of lactotroph cells to changing photoperiod and tuberalin secretion (Fig. 2). Such studies, coupled with the identification of tuberalin and its regulatory mechanism(s), would fully substantiate the 'dual site' hypothesis and also reveal how a novel intra-pituitary signalling mechanism can alter lactotroph cell heterogeneity.

Acknowledgements

The author thanks Dr Gerald A Lincoln for critical reading of an early draft of the manuscript.

Funding

The author is supported by a Biotechnology and Biological Sciences Research Council grant awarded to Dr David G

Hazlerigg. There is no conflict of interest that would prejudice the impartiality of this manuscript.

References

- Badura LL & Goldman BD 1992 Prolactin-dependent seasonal changes in pelage: role of the pineal gland and dopamine. *Journal of Experimental Zoology* **261** 27–33.
- Barrett P, Choi W-S, Morris M & Morgan P 2000 A role for tyrosine phosphorylation in the regulation and sensitisation of adenylate cyclase by melatonin. *FASEB Journal* **14** 1619–1628.
- Barrett P, Messenger S, Schuster C, Moar KM, Mercer JG & Morgan PJ 2002 Pituitary adenylate cyclase-activating polypeptide acts as a paracrine regulator of melatonin-responsive cells of the ovine pars tuberalis. *Endocrinology* **143** 2366–2375.
- Bockers TM, Niklowitz P, Bockmann J, Fauteck J-D, Wittkowski W & Kreutz MR 1995 Daily melatonin injections induce cytological changes in pars tuberalis-specific cells similar to short photoperiod. *Journal of Neuroendocrinology* **7** 607–613.
- Carr A-JF, Johnston JD, Semikhodskii AG, Nolan T, Cagampang FRA, Stirland JA & Loudon ASI 2003 Photoperiod differentially regulates circadian oscillators in central and peripheral tissues of the Syrian hamster. *Current Biology* **13** 1543–1548.
- Dardente H, Klosin P, Pevet P & Masson-Pevet M 2003 MT1 melatonin receptor mRNA expressing cells in the pars tuberalis of the European hamster: effect of photoperiod. *Journal of Neuroendocrinology* **15** 778–786.
- D'Este L, Kulaksiz H, Rausch U, Vaccaro R, Wenger T, Tokunaga Y, Renda TG & Cetin Y 2000 Expression of guanylin in 'pars tuberalis-specific cells' and gonadotrophs of rat adenohypophysis. *PNAS* **97** 1131–1136.
- Graham ES, Webster CA, Hazlerigg DG & Morgan PJ 2002 Evidence for the biosynthesis of a prolactin-releasing factor from the ovine pars tuberalis, which is distinct from thyrotropin-releasing hormone. *Journal of Neuroendocrinology* **14** 945–954.
- Guerra M & Rodriguez EM 2001 Identification, cellular and subcellular distribution of 21 and 72 kDa proteins (tuberalins?) secreted by specific cells of the pars tuberalis. *Journal of Endocrinology* **168** 363–379.
- Hazlerigg DG, Gonzalez-Brito A, Lawson W, Hastings MH & Morgan PJ 1993 Prolonged exposure to melatonin leads to time-dependent sensitization of adenylate cyclase and down-regulates melatonin receptors in pars tuberalis cells from ovine pituitary. *Endocrinology* **132** 285–292.
- Hazlerigg DG, Hastings MH & Morgan PJ 1996 Production of a prolactin releasing factor by the ovine *pars tuberalis*. *Journal of Neuroendocrinology* **8** 489–492.
- Hazlerigg DG, Morgan PJ & Messenger S 2001 Decoding photoperiodic time and melatonin in mammals: what can we learn from the pars tuberalis? *Journal of Biological Rhythms* **16** 326–335.
- Hegarty CM, Jonassen JA & Bittman EL 1990 Pituitary hormone gene expression in male golden hamsters: interactions between photoperiod and testosterone. *Journal of Neuroendocrinology* **2** 567–573.
- Johnston JD, Cagampang FRA, Stirland JA, Carr A-JF, White MRH, Davis JRE & Loudon ASI 2003a Evidence for an endogenous *per1*- and *ICER*-independent seasonal timer in the hamster pituitary gland. *FASEB Journal* **17** 810–815.
- Johnston JD, Messenger S, Ebling FJP, Williams LM, Barrett P & Hazlerigg DG 2003b Gonadotrophin-releasing hormone drives melatonin receptor down-regulation in the developing pituitary gland. *PNAS* **100** 2831–2835.
- Johnston JD, Stirland JA, White MRH, Davis JRE & Loudon ASI 2003c Heterogeneous regulation of individual lactotroph cells by photoperiod in the Syrian hamster (*Mesocricetus auratus*). *General and Comparative Endocrinology* **134** 182–186.

- Katakura Y, Alam S & Shirahata S 1998 Immortalization by gene transfection. *Methods in Cell Biology* **57** 69–91.
- Khodursky AB & Bernstein JA 2003 Life after transcription – revisiting the fate of messenger RNA. *Trends in Genetics* **19** 113–115.
- Klosen P, Bienvenu C, Demarteau O, Dardente H, Guerrero H, Pevet P & Masson-Pevet M 2002 The MT1 melatonin receptor and ROR β receptor are co-localised in specific TSH-immunoreactive cells in the pars tuberalis of the rat pituitary. *Journal of Histochemistry and Cytochemistry* **50** 1647–1657.
- Lafarque M, Oliveros L & Aguado L 1998 Efecto de secreciones de pars tuberalis de la adenohipofisis sobre la liberación de prolactina desde pars distalis. *Medicina* **58** 36–40.
- Lamberts SWJ & MacLeod RM 1990 Regulation of prolactin secretion at the level of the lactotroph. *Physiological Reviews* **70** 279–318.
- Lincoln GA 1989 Significance of seasonal cycles in prolactin secretion in male mammals. In *Perspectives in Andrology*, vol 3, pp 299–306. Ed. M Serio. New York: Raven Press.
- Lincoln GA 1999 Melatonin modulation of prolactin and gonadotrophin secretion. Systems ancient and modern. *Advances in Experimental Medicine and Biology* **460** 137–153.
- Lincoln GA & Clarke IJ 1994 Photoperiodically-induced cycles in the secretion of prolactin in hypothalamo–pituitary disconnected rams: evidence for translation of the melatonin signal in the pituitary gland. *Journal of Neuroendocrinology* **6** 251–260.
- Lincoln GA & Clarke IJ 2002 Noradrenaline and dopamine regulation of prolactin secretion in sheep: role in prolactin homeostasis but not photoperiodism. *Journal of Neuroendocrinology* **14** 36–44.
- Lincoln GA, Messenger S, Andersson H & Hazlerigg DG 2002 Temporal expression of seven clock genes in the suprachiasmatic nucleus and the pars tuberalis of the sheep: evidence for an internal coincidence timer. *PNAS* **99** 13890–13895.
- Lincoln GA, Andersson H & Loudon ASI 2003 Clock genes in calendar cells as the basis of annual timekeeping in mammals – a unifying hypothesis. *Journal of Endocrinology* **179** 1–13.
- Maywood ES & Hastings MH 1995 Lesions of the iodomelatonin-binding sites of the mediobasal hypothalamus spare the lactotropic, but not the gonadotropic response of male Syrian hamsters to short photoperiod and to melatonin. *Endocrinology* **136** 144–153.
- Mellon PL, Windle JJ & Weiner RI 1991 Immortalization of neuroendocrine cells by targeted oncogenesis. *Recent Progress in Hormone Research* **47** 69–93.
- Merks T, Schulze-Bonhage A & Wittkowski W 1993 Photoperiod-dependent changes in exocytotic activity in the hypophyseal pars tuberalis of the Djungarian hamster, *Phodopus sungorus*. *Cell and Tissue Research* **273** 287–291.
- Messenger S, Ross AW, Barrett P & Morgan PJ 1999 Decoding photoperiodic time through *Per1* and *ICER* gene amplitude. *PNAS* **96** 9938–9943.
- Morgan PJ, Barrett P, Howell HE & Helliwell R 1994 Melatonin receptors: localization, molecular pharmacology and physiological significance. *Neurochemistry International* **24** 101–146.
- Morgan PJ, Webster CA, Mercer JG, Ross AW, Hazlerigg DG, MacLean A & Barrett P 1996 The ovine pars tuberalis secretes a factor(s) that regulates gene expression in both lactotropic and nonlactotropic pituitary cells. *Endocrinology* **137** 4018–4026.
- Reppert SM, Weaver DR & Godson C 1996 Melatonin receptors step into the light: cloning and classification of subtypes. *Trends in Pharmacological Sciences* **17** 100–102.
- Reymond MJ, Speciale SG & Porter JC 1983 Dopamine in plasma of lateral and medial hypophysial portal vessels: evidence for regional variation in the release of hypothalamic dopamine into hypophysial portal blood. *Endocrinology* **112** 1958–1963.
- Schwartz WJ, de la Iglesia HO, Zlomanczuk P & Illnerova H 2001 Encoding *Le Quattro Stagioni* within the mammalian brain: photoperiodic orchestration through the suprachiasmatic nucleus. *Journal of Biological Rhythms* **16** 302–311.
- Shorte SL, Leclerc GM, Vazquez-Martinez R, Leamont DC, Faught WJ, Frawley LS & Boockfor FR 2002 PRL gene expression in individual living mammatropes displays distinct functional pulses that oscillate in a noncircadian temporal pattern. *Endocrinology* **143** 1126–1133.
- Steger RW & Bartke A 1991 Temporal sequence of neuroendocrine events associated with the transfer of male golden hamsters from a stimulatory to a nonstimulatory photoperiod. *Biology of Reproduction* **44** 76–82.
- Stirland JA, Johnston JD, Cagampang FRA, Morgan PJ, Castro MG, White MRH, Davis JRE & Loudon ASI 2001 Photoperiodic regulation of prolactin gene expression in the Syrian hamster by a pars tuberalis-derived factor. *Journal of Neuroendocrinology* **13** 147–157.
- Stirland JA, Seymour ZC, Windeatt S, Norris AJ, Stanley P, Castro MG, Loudon ASI, White MRH & Davis JRE 2003 Real-time imaging of gene promoter activity using an adenoviral reporter construct demonstrates transcriptional dynamics in normal anterior pituitary cells. *Journal of Endocrinology* **178** 61–69.
- Takasuka N, White MRH, Wood CD, Robertson WR & Davis JRE 1998 Dynamic changes in prolactin promoter activation in individual living lactotrophic cells. *Endocrinology* **139** 1361–1368.
- Villalobos C, Faught WJ & Frawley LS 1999 Dynamics of stimulus–expression coupling as revealed by monitoring of prolactin promoter-driven reporter activity in individual, living mammatropes. *Molecular Endocrinology* **13** 1718–1727.
- von Gall C, Garabette ML, Kell CA, Frenzel S, Dehghani F, Schumm-Draeger P-M, Weaver DR, Korf H-W, Hastings MH & Stehle JH 2002 Rhythmic gene expression in pituitary depends on heterologous sensitisation by the neurohormone melatonin. *Nature Neuroscience* **5** 234–238.
- Weaver DR, Liu C & Reppert SM 1996 Nature's knockout: the Mel_{1b} receptor is not necessary for reproductive and circadian responses to melatonin in Siberian hamsters. *Molecular Endocrinology* **10** 1478–1487.
- Wittkowski W, Bockmann J, Kreutz MR & Bockers TM 1999 Cell and molecular biology of the pars tuberalis of the pituitary. *International Review of Cytology* **185** 157–194.

Received in final form 17 October 2003

Accepted 20 October 2003

Made available online as an

Accepted Preprint on 22 October 2003