Human growth hormone and insulin-like growth factor-I inhibit erythropoietin secretion from the kidneys of adult rats

Motoi Sohmiya and Yuzuru Kato

Department of Endocrinology, Metabolism and Hematological Oncology, Shimane University School of Medicine, Izumo 693-8501, Japan
(Requests for offprints should be addressed to M Sohmiya; Email: motoi@med.shimane-u.ac.jp)

Abstract

Growth hormone (GH) and insulin-like growth factor-I (IGF-I) play important roles in erythropoiesis and erythropoietin (EPO) secretion. We examined the effects of GH and IGF-I on EPO production in adult rat kidney and liver *in vivo* and *in vitro*. Male Wistar rats aged 8–10 weeks were used. Recombinant human GH (hGH) was continuously infused (20 µg/kg per h) subcutaneously for 48 h using a micro-osmotic infusion pump. Octreotide (10 µg/kg) was subcutaneously injected every 12 h beginning 12 h before the hGH treatment. GH increased plasma EPO levels earlier than it increased plasma IGF-I levels. At 24 h, the IGF-I content in the liver and kidney was increased from 172.8 ± 14.6 to 232.6 ± 17.8 ng/g tissue (means $\pm$ S.E.) and from 53.8 ± 3.1 to 112.8 ± 7.2 ng/g tissue, respectively. The EPO content in the liver was increased from

7.5 ± 1.2 to 15.1 ± 1.4 mIU/g tissue at 48 h, whereas the EPO content in the kidney was decreased at 12, 24, and 48 h after the start of hGH treatment. When the kidneys were organ-cultured, hGH considerably decreased EPO levels in the culture medium in a dose-related manner. The addition of anti-hGH IgG blunted the GH-induced inhibition of EPO secretion from the kidneys. IGF-I also decreased EPO levels in the medium in a dose-related manner. The addition of anti-IGF-I IgG blunted the IGF-I-induced inhibition of EPO secretion from the kidneys, whereas the GH-induced inhibition of EPO secretion was not affected. These findings suggest that both hGH and IGF-I have direct inhibitory effects on EPO secretion from adult rat kidneys.

Journal of Endocrinology (2005) **184**, 199–207

Introduction

The relationship between the endocrine system and hemopoiesis has been known for more than half a century (Watkinson *et al.* 1947, Gordon 1954, Crafts & Meineke 1959, Jepson *et al.* 1968, Nagy & Berczi 1989). Hypopituitarism is often accompanied by normochromic and normocytic anemia. Androgen and thyroid hormones are known to stimulate erythropoiesis (Alexanian 1969). Growth hormone (GH) has a stimulatory effect on erythropoiesis (Fisher *et al.* 1964, Meineke & Crafts 1968, Peschle *et al.* 1972, Golde *et al.* 1977). GH directly stimulated the proliferation of erythroid progenitor cells *in vitro* (Golde *et al.* 1977). Erythropoiesis was stimulated in rat bone marrow perfused with GH (Meineke & Crafts 1968) and in nephrectomized and hypophysectomized rats (Fisher *et al.* 1964) *in vivo*. Hypoplastic bone marrow was restored by GH treatment, but not by adrenocorticotrophic hormone, follicle stimulating hormone, luteinizing hormone, or thyroid-stimulating hormone (Jepson & Lowenstein 1964, Nagy & Berczi 1989). GH administration improved anemia in GH-deficient animals (Jepson & McGarry 1972, Ten Have *et al.* 1997).

Erythropoietin (EPO), a potent regulator of erythropoiesis, is a glycoprotein with a molecular mass of 34 000 Da. EPO secretion was strongly stimulated by hypoxia (Goldberg *et al.* 1989). EPO secretion is regulated endocrinologically. Thyroid hormones and adrenocortical hormone enhance EPO secretion (Peschle *et al.* 1978, Nagy & Berczi 1989, Brenner *et al.* 1994, Fandrey *et al.* 1994). Although GH has a stimulatory effect on erythropoiesis, there have been few reports on the role of GH in EPO secretion. We previously reported that GH treatment increased plasma EPO levels in anemic patients with diabetic nephropathy (Sohmiya *et al.* 1998) and in patients with GH deficiency (Sohmiya & Kato 2001). EPO is secreted from various organs depending on various conditions. EPO is secreted mainly from the peritubular endothelia and interstitial fibroblasts of renal tubules. It is also secreted partly from the liver in fetal rats (Bondurant & Koury 1986, Clemons *et al.* 1986, Koury *et al.* 1988, 1989, Kurtz *et al.* 1989) and sheep (Zanjani *et al.* 1977, 1981), placenta (Davis *et al.* 2003), and astrocytes in the brain (Masuda *et al.* 1994). The liver is the primary site of EPO production in the fetus (Gruber *et al.* 1977, Zanjani *et al.* 1977, 1981), and extrarenal EPO production was

increased in anemic rats (Erslev *et al.* 1980). In adult rats, EPO is secreted mainly from the kidney, suggesting that some mechanism switches EPO secretion from the liver to the kidney (Zanjani *et al.* 1981). This functional switching mechanism has yet to be fully elucidated. Fetal hypoxemia experiments showed that the EPO secretion from the placenta was increased by hypoxic stimuli (Davis *et al.* 2003). Physiological conditions might induce EPO secretion from the various organs. In particular, as the liver is the main target organ of GH, GH administration affected EPO secretion from the liver. However, the roles of GH on EPO secretion and intervention of the kidney and liver *in vivo* and *in vitro* have not been elucidated fully. In the present study, we investigated the effects of GH and insulin-like growth factor-I (IGF-I) on EPO production in adult rat kidney and liver *in vivo* and *in vitro*.

Materials and Methods

Reagents

Recombinant human GH (rhGH) of the 22 kDa form (rhGH-22k; Genotropin; Sumitomo Pharmaceutical Co., Tokyo, Japan) was kindly provided by the Sumitomo Pharmaceutical Co. Recombinant human IGF-I was purchased from GroPep Co. (Adelaide, SA, Australia). Octreotide (Sandostatin) was kindly provided by Sandoz Co. (Basel, Switzerland). Anti-hGH rabbit plasma and anti-IGF-I rabbit plasma were produced by multiple subcutaneous injection of hGH or recombinant IGF-I with Freund's complete adjuvant to rabbits. The anti-hGH IgG and anti-IGF-I IgG, both purified to the IgG fraction by Protein A column (Pharmacia, Uppsala, Sweden), were used in organ-culture experiments. Normal rabbit IgG was obtained by using the same method from normal rabbit plasma.

Animal experiments

Male Wistar rats (Japan Crea Co., Tokyo, Japan) aged 8–10 weeks were used throughout the experiments. The animals were maintained on a constant light/dark cycle (lights on 08:00–20:00 h) in a room controlled for temperature ($24 \pm 1^\circ\text{C}$) and humidity (50–60%). Tap water and laboratory chow pellets (Japan Crea Co.) were given *ad libitum*.

Octreotide (10 $\mu\text{g/kg}$) was injected subcutaneously every 12 h beginning 12 h before hGH administration. rhGH was infused subcutaneously for 48 h (20 $\mu\text{g/kg}$ per h) using a micro-osmotic infusion pump (model 1003D; Alzet, Cupertino, CA, USA). Rats were divided into seven groups of eight rats each, for analysis at before (0 h) or after (3, 6, 12, 24, or 48 h) the start of GH infusion with octreotide treatment, or no treatment. At the end of each time point, rats were anesthetized with pentobarbital.

Each was placed on its back, and the abdominal cavity was opened. Blood samples were obtained from the abdominal aorta and centrifuged immediately. Plasma samples were stored at -20°C until assayed. The liver and kidney were perfused with 0.1 M ice-cold PBS, pH 7.4. The kidneys were perfused via the abdominal aorta after the blood sampling. The liver was perfused via the hepatic portal vein using the once-through procedure (Waynforth & Flecknell 1992). After the blood was completely displaced from the liver by massaging the organ gently between the fingers, the liver was excised and frozen at -80°C until extraction of EPO.

Extraction of EPO

EPO was extracted from the kidneys and liver using a modified extraction method (Sherwood & Goldwasser 1978). The tissues of rat kidney and liver were homogenized with a Polytron homogenizer in 0.1 M ice-cold PBS, pH 7.4, at a ratio of 5 ml buffer to 1 g tissue. The homogenates were centrifuged at 15 000 *g* for 30 min, and the supernatants were applied to an ultracentrifugation device (Centricon 100; Amicon, Beverly, MA, USA) to remove the large proteins ($>100\,000\text{ Da}$). The filtrated fractions were kept frozen at -20°C until assayed.

Organ culture of the kidneys

Organ culture of the kidney was performed by a modified method of Sherwood *et al.* (1972). Briefly, the kidneys were excised by an aseptic method and placed in ice-cold modified Tröwell's T8 medium supplemented with 10% fetal calf serum, with 50 U/ml penicillin and 50 $\mu\text{g/ml}$ streptomycin. The kidneys were cut into pieces (approximately 1 mm^3 in volume), and between seven and 10 pieces were placed on a membrane filter (0.65 μm ; Durapore; Millipore Co., Bedford, MA, USA) in an organ-culture dish (Falcon; Becton Dickinson Co., Franklin Lakes, NJ, USA). The level of the culture medium was adjusted so that only the membrane filter would be wetted. The kidneys were incubated with 1 ml modified Tröwell's T8 medium in each culture dish. Kidney explants were cultured in serum-free Tröwell's T8 medium for up to 5 days. After preincubation for 24 h, different concentrations of rhGH (10^{-6} , 10^{-7} , and 10^{-8} M) and IGF-I (10^{-6} , 10^{-7} , and 10^{-8} M) with or without anti-hGH rabbit IgG (100 $\mu\text{g/ml}$), anti-IGF-I rabbit IgG (100 $\mu\text{g/ml}$), or normal rabbit IgG (100 $\mu\text{g/ml}$) were added to the medium, and the explants were further incubated for 72 h in a humidified atmosphere of 95% air/5% CO_2 at 37°C . The medium was frozen immediately at -20°C until assay. After the organ culture, the kidney fragment was fixed with formaldehyde and stained with hematoxylin/eosin, and the histological changes

Figure 1 Effects of rhGH treatment on plasma hGH (left upper panel), rat GH (left lower panel), IGF-I (right upper panel), and EPO level (right lower panel) in rats pretreated with octreotide (SMS). ND, not detectable. SMS(-), not treated with octreotide. Means $\pm$ S.E. are shown. * $P < 0.005$ versus SMS(-); # $P < 0.05$ versus time 0; ## $P < 0.005$ versus time 0.

were assessed. Little central necrosis was recognized in any of the fragments.

Assays

EPO levels in plasma, extracted solution, and culture medium were measured by a sensitive enzyme immunoassay (EIA) as previously described (Sohmiya & Kato 2000). The minimal detectable quantity was 0.15 mIU/ml using 100 μL . The intra- and inter-assay coefficients of variation were 6.8% and 8.2%, respectively.

Plasma hGH levels were measured by highly sensitive EIA as previously described (Sohmiya & Kato 1992, Sohmiya *et al.* 1993). Plasma rat GH levels were measured by a highly sensitive EIA as previously described (Sohmiya & Kato 1994). There is no cross-reactivity of rat GH with hGH in the EIA. IGF-I levels in plasma and in the extracts

of kidney and liver were measured by specific RIA after acid/ethanol extraction as described previously (Yamamoto *et al.* 1991).

Statistical analysis

The data are expressed as means $\pm$ S.D. In the *in vivo* experiments, the experimental data were analyzed by repeated-measures ANOVA followed by Student's *t*-test. In the organ-culture experiments, the data were handled using Fisher's method as a *post-hoc* analysis. $P < 0.05$ was considered significant.

Results

As shown in Fig. 1, plasma hGH levels increased from undetectable at 0 h to $23.8 \pm 1.8 \mu\text{g/L}$ (means $\pm$ S.E.) at 6 h

Figure 2 Effects of rhGH treatment on the content of IGF-I (upper panels) and EPO (lower panels) in liver (right-hand panels) and kidney (left-hand panels) pretreated with octreotide. SMS(-) shows not treated with octreotide. Means $\pm$ S.E. are shown. # P <0.05 versus time 0; ## P <0.01 versus time 0; ### P <0.005 versus time 0.

after the start of hGH infusion. Octreotide administration suppressed plasma rat GH levels. Plasma IGF-I levels increased from 625.8 ± 8.7 to 725 ± 8.7 μ g/l at 12 h and peaked at 962.0 ± 12.2 μ g/l at 48 h. The octreotide treatment made no difference in plasma IGF-I levels. Plasma EPO levels increased from 8.0 ± 1.9 to 12.4 ± 1.8 IU/l at 3 h and peaked at 36.5 ± 1.8 IU/l at 48 h. Plasma EPO levels increased earlier than plasma IGF-I levels. There was a significant difference in plasma EPO levels with and without octreotide treatment.

The contents of IGF-I and EPO in the kidney and liver are shown in Fig. 2. The IGF-I content in the kidney increased from 172.8 ± 14.6 to 232.6 ± 17.8 ng/g tissue at 12 h. The IGF-I content in the liver increased from 53.8 ± 3.1 to 112.8 ± 7.2 ng/g tissue at 24 h. The EPO content in the liver increased from 7.5 ± 1.2 to 15.1 ± 1.4 mIU/g tissue at 48 h, whereas the EPO content in the kidney was decreased at 12, 24, and 48 h after the start of hGH treatment.

As shown in Fig. 3, hGH considerably decreased medium EPO levels in organ-cultured kidney in a dose-related manner (10^{-7} M GH, 1.22 ± 0.20 , and 10^{-6} M GH, 0.65 ± 0.18 , versus control, 1.81 ± 0.13 mIU/ml; P <0.05 and P <0.005, respectively). The addition of anti-hGH IgG (100 μ g/ml) blunted the GH-induced inhibition of EPO release from the organ-cultured kidneys, whereas normal rabbit IgG did not affect the GH-induced inhibition of EPO release. As shown in Fig. 4, IGF-I also suppressed medium EPO levels in a dose-related manner (10^{-7} M IGF-I, 1.36 ± 0.07 , and 10^{-6} M IGF-I, 0.99 ± 0.09 , versus control, 1.85 ± 0.18 mIU/ml; P <0.01 and P <0.0001, respectively). The addition of anti-IGF-I IgG (100 μ g/ml) blunted the IGF-I-induced inhibition of EPO release from the organ-cultured kidneys. The addition of normal rabbit IgG did not affect the suppressive effect of IGF-I on EPO release. As shown in Fig. 5, we also found that the addition of anti-IGF-I IgG (100 μ g/ml) did not affect the

Figure 3 Effects of hGH on medium EPO levels in organ-cultured kidney. The EPO levels decreased by hGH in a dose-related manner (10^{-7} M hGH, 1.22 ± 0.20 , and 10^{-6} M hGH, 0.65 ± 0.18 , versus control, 1.81 ± 0.13 mIU/ml; $P < 0.05$ and $P < 0.005$, respectively; left-hand panel). Addition of anti-hGH IgG (100 µg/ml) blunted the GH-induced inhibition of EPO release from organ-cultured kidney. Addition of normal rabbit IgG did not affect the GH-induced inhibition of EPO release (right-hand panel). Means $\pm$ S.E. are shown.

GH-induced inhibition of EPO secretion from the organ-cultured kidneys.

Discussion

EPO is a major humoral regulator of erythropoiesis. In the rat, it is secreted mainly from the endothelia and interstitial fibroblasts of renal tubules and partly from the liver (Kurtz *et al.* 1989, Koury *et al.* 1988, 1989). EPO secretion is controlled mainly by oxygen pressure (Koury *et al.* 1988, Eckardt *et al.* 1989). We previously reported that plasma EPO levels increased after the start of rhGH infusion in anemic patients with chronic renal failure (Sohmiya *et al.* 1998) and in adult patients with GH deficiency (Sohmiya & Kato 2001). Human plasma EPO levels increased within 6 h after the start of rhGH administration, suggesting that GH directly stimulates EPO production. In the present study, rat plasma EPO levels were increased earlier than the increase of plasma IGF-I, suggesting that GH directly stimulates EPO secretion *in vivo*. On the other hand, it was reported that erythroid and myeloid progenitor precursor cells increased after GH replacement in adult patients with GH deficiency, whereas plasma EPO levels remained almost unchanged (Kotzmann *et al.* 1996). In the present study, rhGH and octreotide administration increased circulating EPO levels in adult rats. The octreotide administration dosage was sufficient to suppress endogenous rat

GH secretion. The decreased plasma EPO levels due to octreotide administration suggest that endogenous rat GH plays an important stimulatory role in EPO secretion *in vivo*.

EPO content decreased in the kidney, whereas it increased in the liver. These findings suggest that hGH's effects on EPO secretion differ depending on whether it is secreted from the kidney or the liver. The kidney could be the main site of EPO production in adult rats, and increased circulating EPO should originate from the kidney. However, our data suggest that the increased circulating EPO levels might originate from the liver instead. It is known that the liver is the major site of EPO production in the ovine fetus, whereas the kidney takes a major role in EPO production in adult sheep (Zanjani *et al.* 1981). In the neonatal rat, EPO was found to originate from the liver rather than the kidney, and hypoxic stimuli induced EPO production in both organs (Clemons *et al.* 1986). However, the results do not clarify the mechanism by which EPO is secreted from the liver. The present data suggest that GH suppresses EPO production from the kidney and that GH stimulates EPO secretion from the liver in a different manner than that from the kidney.

The organ-culture method was suitable for evaluating EPO secretion from kidney tissue (Sherwood *et al.* 1972), since EPO was secreted from some cell populations in the kidney, such as interstitial cells and peritubular endothelial cells. Sherwood *et al.* reported EPO production using organ culture of the rat kidney (Sherwood *et al.* 1972).

Figure 4 Effects of IGF-I on medium EPO levels in organ-cultured kidney. The EPO levels decreased by IGF-I in a dose-related manner (10^{-7} M IGF-I, 1.36 ± 0.07 , and 10^{-6} M IGF-I, 0.99 ± 0.09 , versus control, 1.85 ± 0.18 mIU/ml; $P < 0.01$ and $P < 0.0001$, respectively; left-hand panel). Addition of anti-IGF-I IgG (100 µg/ml) blunted the IGF-I-induced inhibition of EPO secretion from organ-cultured kidney. Addition of normal rabbit IgG did not affect the suppressive effect of IGF-I on EPO secretion (right-hand panel). Means $\pm$ S.E. are shown.

They used a bioassay to measure EPO levels. We measured medium EPO levels using a highly sensitive EIA with 0.15 mIU/ml as the detectable quantity; this EIA was able to measure all of the EPO levels in the medium.

In the present study, hGH suppressed EPO release from organ-cultured kidneys into the medium in a dose-dependent manner. Anti-IGF-I IgG administration with hGH could not inhibit the suppression of EPO secretion by hGH, suggesting that hGH inhibited EPO secretion directly.

IGF-I stimulates erythropoiesis *in vitro* (Claustres *et al.* 1987, Merchav *et al.* 1988) and *in vivo* (Kurtz *et al.* 1988). However, there have been few reports on the relationship between IGF-I and EPO (Claustres *et al.* 1987, Kurtz *et al.* 1988, Merchav *et al.* 1988). IGF-I stimulates EPO secretion in patients with insulin resistance (Quin *et al.* 1994). On the other hand, it exerts a negative control function on oxygen-regulated EPO production in a human hepatoma cell line (Hep G2; Scholz *et al.* 1992). In the present study, IGF-I suppressed EPO secretion from the organ-cultured kidney. It decreased EPO content in the kidney, but increased it in the liver. IGF-I suppressed EPO secretion from the kidney organ culture, and the addition of anti-IGF-I IgG with IGF-I inhibited the suppression of EPO secretion, suggesting that IGF-I directly stimulated

EPO secretion from the kidney. IGF-I content was increased in the kidney and the liver. The minimum concentration of IGF-I that suppressed EPO secretion was 10^{-7} M. The effect was weaker than that of the same concentration of hGH. IGF-I might affect EPO secretion directly as well as by a paracrine mechanism. In the present study, the kidney fragments used in the organ culture consisted of heterogeneous cell populations. Therefore, the interaction between different cell types might be considered. Further investigation in prepared individual normal cells should be required.

The cellular mechanism of EPO secretion from the kidney is not fully elucidated. EPO is secreted mainly from interstitial cells and peritubular endothelial cells in normal kidneys of adult rats (Goldberg *et al.* 1989, Kurtz *et al.* 1989). Although there have been many perfusion experiments, there has been no report on culture experiments using interstitial cells and peritubular endothelial cells. This is because it is very complicated to prepare a primary culture of interstitial cells and endothelial cells from normal kidney tissue. Calcium regulates EPO secretion (Sherwood *et al.* 1987, Fisher 1988). An increase in intracellular calcium leads to the inhibition of EPO biosynthesis and/or of secretion, and a decrease in intracellular calcium increases EPO production in an established human renal cell

Figure 5 Effects of anti-IGF-I IgG on hGH-induced inhibition of EPO secretion. Addition of anti-IGF-I IgG (100 µg/ml) did not affect the GH-induced inhibition of EPO secretion from organ-cultured kidney. Means $\pm$ S.E. are shown.

carcinoma cell line (Sherwood *et al.* 1987). Primary culture experiments using a human renal cell carcinoma cell line indicate that a decreased Ca^{2+} concentration and the addition of a calcium antagonist induce EPO secretion in culture medium. Increased cAMP induces EPO secretion from an intracellular EPO-storage pool (Sherwood *et al.* 1987). Increased intracellular calcium activates a calcium/calmodulin kinase and produces a phosphoprotein that inhibits EPO production and secretion (Fisher 1988). However, for a mechanism based on a carcinoma cell line, it is unknown whether or not the mechanism acts in normal cells and in different tissues from different organs.

Both GH receptor and GH-binding protein mRNAs are present in the liver, kidney, adrenal gland, heart, muscle, ovary, mammary gland, gastrointestinal tract, and adipose tissue (Tiong & Herington 1991). IGF-I receptor is also present in many tissues such as liver, kidney, adipose tissue, muscle, neuron, cartilage, and others, and it plays a role in the regulation of metabolism, growth, and differentiation (De Meyts *et al.* 1994, Van Obberghen 1994). Therefore, the post-receptor pathway of GH and/or IGF-I might be related to EPO secretion. GH increases the intracellular free-calcium concentration in a variety of cell types (Schwartz & Goodman 1990, Schwartz *et al.* 1992, Gaur *et al.* 1996, Udy *et al.* 1997, Zhu *et al.* 2001). An

increase in intracellular calcium can be achieved by an influx of Ca^{2+} through voltage-dependent Ca^{2+} channels located in the plasma membrane or by mobilization of Ca^{2+} from intracellular stores (Cruzalegui & Bading 2000). IGF-I also increases intracellular Ca^{2+} (Kojima *et al.* 1988, Poiraudau *et al.* 1997). An increase in the intracellular calcium concentration decreased EPO secretion. Therefore, the increase in intracellular Ca^{2+} by GH and/or by IGF-I might at least partly affect the suppressive mechanism of EPO secretion.

GH and IGF-I levels were superphysiological levels in the organ-culture experiments. Stimulatory effects of GH and IGF-I on EPO secretion were not observed at concentrations lower than 10 µM for GH and IGF-I. This might be attributed to the characteristics of the cell or tissue culture. Therefore, it was unknown whether EPO secretion from the kidney was suppressed by GH or IGF-I administration under physiological conditions *in vivo* or not.

Primate GH binds to both prolactin and GH receptors. Primate GH might partly have stimulatory effect on EPO secretion mediated by prolactin receptor. Our experiment using anti-GH antibody did not clarify whether the stimulatory effect of GH on EPO secretion was mediated by prolactin receptor and/or GH receptor. Further studies using GH receptor or prolactin receptor blocker are required.

The most interesting thing is the origin of increased circulating EPO level. In the next set of preliminary experiments, we found that GH stimulates EPO secretion from interstitial cells containing Kupffer cells, fibroblasts, and vascular endothelial cells whereas GH rather inhibits EPO release from parenchymal cells in the rat liver, indicating that an increase in plasma EPO levels after GH treatment might be derived from interstitial cells in the liver (data not shown).

In conclusion, the continuous subcutaneous infusion of hGH stimulated EPO secretion in rats. Plasma EPO levels were increased earlier than plasma IGF-I levels. The EPO content decreased in the kidney but increased in the liver, and EPO secretion decreased in organ-cultured kidneys. In organ culture of rat kidneys, EPO secretion was inhibited by the addition of hGH and IGF-I. The inhibitory effect of EPO secretion by hGH was not blunted by anti-IGF-I IgG. These findings suggest that both GH and IGF-I have direct inhibitory effects on EPO secretion from rat kidneys.

Acknowledgments

This work was supported in part by grants from the Ministry of Education, Science and Culture, Japan, the Ministry of Health and Welfare, Japan, the Growth Science Foundation, Japan, and the Foundation for Renal Anemia Therapy, Japan. We are indebted to Mrs Akiko

Kawakami and Mrs Akiko Kanayama for their secretarial assistance. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Alexanian R 1969 Erythropoietin and erythropoiesis in anemic man following androgen. *Blood* **33** 564–572.
- Bondurant MC & Koury MJ 1986 Anemia induces accumulation of erythropoietin mRNA in the kidney and liver. *Molecular Cell Biology* **6** 2731–2733.
- Brenner B, Fandrey J & Jelkmann W 1994 Serum immunoreactive erythropoietin in hyper- and hypothyroidism: clinical observations related to cell culture studies. *European Journal of Haematology* **53** 6–10.
- Claustres M, Chatelain P & Sultan C 1987 Insulin-like growth factor I stimulates human erythroid colony formation in vitro. *Journal of Clinical Endocrinology and Metabolism* **65** 78–82.
- Clemons GK, Fitzsimmons SL & Pemanincor D 1986 Immunoreactive erythropoietin concentration in fetal and neonatal rats and the effects of hypoxia. *Blood* **68** 892–899.
- Crafts SG & Meineke HA 1959 The anemia of hypophysectomized animals. *Annals of New York Academy of Science* **77** 501–517.
- Cruzalegui FH & Bading H 2000 Calcium-regulated protein kinase cascades and their transcription factor targets. *Cellular and Molecular Life Sciences* **57** 402–410.
- Davis LE, Widness JA & Brace RA 2003 Renal and placental secretion of erythropoietin during anemia or hypoxia in the ovine fetus. *American Journal of Obstetrics and Gynecology* **189** 1764–1770.
- De Meyts P, Wallach B, Christoffersen CT, Urso B, Gronskov K, Latus LJ, Yakushiji F, Ilondo MM & Shymko RM 1994 The insulin-like growth factor-I receptor. Structure, ligand-binding mechanism and signal transduction. *Hormone Research* **42** 152–169.
- Eckardt K, Kurtz A, Scholz H & Bauer C 1989 Regulation of erythropoietin formation in vivo. *Contribution to Nephrology* **76** 33–38.
- Erslev AJ, Caro J, Kansu E & Silver R 1980 Renal and extrarenal erythropoietin production in anaemic rats. *British Journal of Haematology* **45** 65–72.
- Fandrey J, Pagel H, Frede S, Wolff M & Jelkmann W 1994 Thyroid hormones enhance hypoxia-induced erythropoietin production in vitro. *Experimental Hematology* **22** 272–277.
- Fisher JW 1988 Pharmacologic modulation of erythropoietin production. *Annual Review of Pharmacology and Toxicology* **28** 101–122.
- Fisher JW, Roh BL, Couch C & Nightingale WO 1964 Influence of cobalt, sheep erythropoietin and several hormones on erythropoiesis in bone marrows of isolated perfused hind limbs of dogs. *Blood* **23** 87–98.
- Gaur S, Yamaguchi H & Goodman HM 1996 Growth hormone increases calcium uptake in rat fat cells by a mechanism dependent on protein kinase C. *American Journal of Physiology Cell Physiology* **270** C1485–C1492.
- Goldberg MA, Imagawa S, Dunning PS & Bunn HF 1989 Oxygen sensing and erythropoietin gene regulation. *Contribution to Nephrology* **70** 39–56.
- Golde DW, Bersch N & Li CH 1977 Growth hormone: species-specific stimulation of erythropoiesis in vitro. *Science* **196** 1112–1113.
- Gordon AS 1954 Endocrine influences upon the formed elements of blood and blood forming organs. *Recent Progress in Hormone Research* **10** 339–394.
- Gruber DF, Zucali JR & Mirand EA 1977 Identification of erythropoietin producing cells in fetal mouse liver cultures. *Experimental Hematology* **5** 392–398.
- Jepson JH & Lowenstein L 1964 Effect of prolactin on erythropoiesis in the mouse. *Blood* **24** 726–738.
- Jepson JH & McGarry EE 1972 Hemopoiesis in pituitary dwarfs treated with human growth hormone and testosterone. *Blood* **39** 238–248.
- Jepson JH, McGarry EE & Lowenstein LL 1968 Erythropoietin excretion in a hypopituitary patient. Effects of testosterone and vasopressin. *Archives of Internal Medicine* **122** 265–270.
- Kojima I, Matsunaga H, Kurokawa K, Ogata E & Nishimoto I 1988 Calcium influx: an intracellular message of the mitogenic action of insulin-like growth factor-I. *Journal of Biological Chemistry* **263** 16561–16567.
- Kotzmarn H, Riedl M, Clodi M, Barnas U, Kaider A, Hocker P & Luger A 1996 The influence of growth hormone substitution therapy on erythroid and myeloid progenitor cells and on peripheral blood cells in adult patients with growth hormone deficiency. *European Journal of Clinical Investigation* **26** 1175–1181.
- Koury ST, Bondurant MC & Koury MJ 1988 Localization of erythropoietin synthesizing cells in murine kidneys by in situ hybridization. *Blood* **71** 524–527.
- Koury ST, Koury MJ, Bondurant MC, Caro J & Graber SE 1989 Quantitation of erythropoietin-producing cells in kidneys of mice by in situ hybridization: correlation with hematocrit, renal erythropoietin mRNA, and serum erythropoietin concentration. *Blood* **74** 645–651.
- Kurtz A, Zapf J, Eckardt KU, Clemons G, Froesch ER & Bauer C 1988 Insulin-like growth factor I stimulates erythropoiesis in hypophysectomized rats. *PNAS* **85** 7825–7829.
- Kurtz A, Eckardt KU, Neumann R, Kaissling B, Le Hir M & Bauer C 1989 Site of erythropoietin formation. *Contribution to Nephrology* **76** 14–23.
- Masuda S, Okano M, Yamagishi K, Nagao M, Ueda M & Sasaki R 1994 A novel site of erythropoietin production. Oxygen-dependent production in cultured rat astrocytes. *Journal of Biological Chemistry* **269** 19488–19493.
- Meineke HA & Crafts RC 1968 Further observations on the mechanism by which androgens and growth hormone influence erythropoiesis. *Annals of the New York Academy of Sciences* **149** 298–307.
- Merchav S, Tatarksky I & Hochberg Z 1988 Enhancement of erythropoiesis in vitro by human growth hormone is mediated by insulin-like growth factor I. *British Journal of Haematology* **70** 267–271.
- Nagy E & Berczi I 1989 Pituitary dependence of bone marrow function. *British Journal of Haematology* **71** 457–462.
- Peschle C, Rappaport IA, Sasso GF, Gordon AS & Condorelli M 1972 Mechanism of growth hormone (GH) action on erythropoiesis. *Endocrinology* **91** 511–517.
- Peschle C, Rappaport IA, Magli MC, Marone G, Lettieri F, Cillo C & Gordon AS 1978 Role of the hypophysis in erythropoietin production during hypoxia. *Blood* **51** 1117–1124.
- Poiraudau S, Lieberherr M, Kergosie N & Corvol MT 1997 Different mechanisms are involved in intracellular calcium increase by insulin-like growth factors 1 and 2 in articular chondrocytes: voltage-gated calcium channels, and/or phospholipase C coupled to a pertussis-sensitive G-protein. *Journal of Cellular Biochemistry* **64** 414–422.
- Quin J, Miell J, Smith K, Gordon D, Strachan J, Dick JB & MacCuish AC 1994 Effect of insulin-like growth factor-I therapy on erythropoietin concentrations in extreme insulin resistance. *Diabetologia* **37** 439.
- Scholz H, Baier W, Ratcliffe P, Eckardt K, Zapf J, Kurtz A & Bauer C 1992 Insulin-like growth factors decrease oxygen-regulated erythropoietin production by human hepatoma cells (Hep G2). *American Journal of Physiology Cell Physiology* **263** C474–C479.
- Schwartz Y & Goodman HM 1990 Refractoriness to the insulin-like effects of growth hormone depends upon calcium. *Endocrinology* **127** 170–176.

- Schwartz Y, Yamaguchi H & Goodman HM 1992 Growth hormone increases intracellular free calcium in rat adipocytes: correlation with actions on carbohydrate metabolism. *Endocrinology* **131** 772–778.
- Sherwood JB & Goldwasser E 1978 Extraction of erythropoietin from normal kidneys. *Endocrinology* **103** 866–870.
- Sherwood JB, Robinson SH, Bassan LR, Rosen S & Gordon AS 1972 Production of erythropoietin by organ cultures of rat kidney. *Blood* **40** 189–197.
- Sherwood JB, Burns ER & Shouval D 1987 Stimulation by cAMP of erythropoietin secretion by an established human renal carcinoma cell line. *Blood* **69** 1053–1057.
- Sohmiya M & Kato Y 1992 Renal clearance, metabolic clearance rate and half life of human growth hormone in young and aged subjects. *Journal of Clinical Endocrinology and Metabolism* **75** 1487–1490.
- Sohmiya M & Kato Y 1994 A highly sensitive enzyme immunoassay of rat growth hormone (GH) and the inhibitory effect of GH on GH secretion in the rat. *Biomedical Research* **5** 207–214.
- Sohmiya M & Kato Y 2000 Molecular and electrical heterogeneity of circulating human erythropoietin measured by sensitive enzyme immunoassay. *European Journal of Clinical Investigation* **30** 344–349.
- Sohmiya M & Kato Y 2001 Effect of long-term administration of recombinant human growth hormone (rhGH) on plasma erythropoietin (EPO) and haemoglobin levels in anaemic patients with adult GH deficiency. *Clinical Endocrinology* **55** 749–754.
- Sohmiya M, Yamamoto H & Kato Y 1993 Age and sex related changes in urinary growth hormone (GH) levels in normal adults. *Biomedical Research* **4**:227–238.
- Sohmiya M, Ishikawa K & Kato Y 1998 Stimulation of erythropoietin secretion by continuous subcutaneous infusion of recombinant human GH in anemic patients with chronic renal failure. *European Journal of Endocrinology* **138** 302–306.
- Ten Have SMTH, van der Lely AJ & Lamberts SWJ 1997 Increase in haemoglobin concentrations in growth hormone deficient adults during human recombinant growth hormone replacement therapy. *Clinical Endocrinology* **47** 565–570.
- Tiong TS & Herington AC 1991 Tissue distribution, characterization, and regulation of messenger ribonucleic acid for growth hormone receptor and serum binding protein in the rat. *Endocrinology* **129** 1628–1634.
- Udy GB, Towers RP, Snell RG, Wilkins RJ, Park SH, Ram PA, Waxman DJ & Davey HW 1997 Requirement of STAT5b for sexual dimorphism of body growth rates and liver gene expression. *PNAS* **94** 7239–7244.
- Van Obberghen E 1944 Signalling through the insulin receptor and the insulin-like growth factor-I receptor. *Diabetologia* **37** S125–S134.
- Watkinson G, McMenemey WH & Evans G 1947 Hypopituitarism, hypogonadism, and anemia treated with testosterone. *Lancet* **252** 631–634.
- Waynforth HB & Flecknell PA 1992 In *Experimental and Surgical Technique in the Rat*, pp 316–322. San Diego: Academic Press.
- Yamamoto H, Sohmiya M, Oka N & Kato Y 1991 Effects of aging and sex on plasma insulin-like growth factor I (IGF-I) levels in normal adults. *Acta Endocrinologica* **124** 497–500.
- Zanjani ED, Foster J, Burlington H, Mann LJ & Wasserman LR 1977 Liver as the primary site of erythropoietin formation in the fetus. *Journal of Laboratory and Clinical Medicine* **89** 640–644.
- Zanjani ED, Ascensao JL, McGlave PB, Banisadre M & Ash RC 1981 Studies of the liver to kidney switch of erythropoietin production. *Journal of Clinical Investigation* **67** 1183–1188.
- Zhu T, Goh EL, Graichen R, Ling L & Lobie PE 2001 Signal transduction via the growth hormone receptor. *Cellular Signaling* **13** 599–616.

Received 6 October 2004
Accepted 15 October 2004