Gene expression of glucose transporter (GLUT) 1, 3 and 4 in bovine follicle and corpus luteum

H Nishimoto, R Matsutani, S Yamamoto¹, T Takahashi, K-G Hayashi, A Miyamoto², S Hamano¹ and M Tetsuka

Department of Agricultural and Life Science, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido 080-8555, Japan

¹Animal Biotechnology Center, Livestock Improvement Association of Japan Inc, Tokyo, Japan

²Graduate School of Animal and Food Hygiene, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido 080-8555, Japan

(Requests for offprints should be addressed to M Tetsuka; Email: mtetsuka@obihiro.ac.jp)

Abstract

Glucose is the main energy substrate in the bovine ovary, and a sufficient supply of it is necessary to sustain the ovarian activity. Glucose cannot permeate the plasma membrane, and its uptake is mediated by a number of glucose transporters (GLUT). In the present study, we investigated the gene expression of GLUT1, 3 and 4 in the bovine follicle and corpus luteum (CL). Ovaries were obtained from Holstein × Japanese Black F1 heifers. Granulosa cells and theca interna layers were harvested from follicles classified into five categories by their physiologic status: follicular size (≥ 8.5 mm: dominant; < 8.5 mm: subordinate), ratio of estradiol (E_2) to progesterone in follicular fluid (≥ 1 : E_2 active; < 1 : E_2 inactive), and stage of estrous cycle (luteal phase, follicular phase). CL were also classified by the stage of estrous cycle. Expression levels of GLUT1, 3 and 4 mRNA were quantified by a real-time PCR. The mRNA for GLUT1 and 3 were detected in the bovine follicle and CL at comparable levels to those in classic GLUT-expressing

organs such as brain and heart. Much lower but appreciable levels of GLUT4 were also detected in these tissues. The gene expression of these GLUT showed tissue- and stage-specific patterns. Despite considerable differences in physiologic conditions, similar levels of GLUT1, 3 and 4 mRNA were expressed in subordinate follicles as well as dominant E_2 -active follicles in both luteal and follicular phases, whereas a notable increase in the gene expression of these GLUT was observed in dominant E_2 -inactive follicles undergoing the atretic process. In these follicles, highly significant negative correlations were observed between the concentrations of glucose in follicular fluid and the levels of GLUT1 and 3 mRNA in granulosa cells, implying that the local glucose environment affects glucose uptake of follicles. These results indicate that GLUT1 and 3 act as major transporters of glucose while GLUT4 may play a supporting role in the bovine follicle and CL.

Journal of Endocrinology (2006) **188**, 111–119

Introduction

The ovary is one of the most dynamic endocrine organs, and it nurtures the cyclic development of several follicles, maturation and ovulation of a selected follicle, and subsequent formation and function of the corpus luteum (CL). To sustain its activity, a substantial energy source is necessary. In the bovine ovary, the predominant energy substrate appears to be glucose, and the whole ovarian respiratory exchange ratio was calculated to be 0.95 (Rabiee *et al.* 1997).

Glucose, a hydrophilic molecule, cannot permeate the plasma membrane, and its uptake is mediated by a number of facilitative sugar transporters. To date, 13 facilitative sugar transporters (GLUT1–12 and HMIT) that exhibit different substrate specificities, kinetic properties and tissue expression profiles have been identified along with

six Na^+ -dependent glucose transporters (SGLT1–6) (Wood & Trayhurn 2003), of which GLUT1, 3 and 4 have been so far reported to be expressed in the ovary of sheep (Williams *et al.* 2001), rat (Kol *et al.* 1997, Kodaman & Behrman 1999) and mouse (Zhou *et al.* 2000) with considerable species differences in the expression profiles. The authors of these reports also demonstrated that the expression of these GLUT is regulated by intraovarian factors involved in follicular development, maturation and ovulation, such as estradiol (E_2), insulin growth factor (IGF)-I and interleukin-1 β (Kol *et al.* 1997, Zhou *et al.* 2000) as well as the classic regulator of ovarian function, gonadotropin (Kodaman & Behrman 1999). These results imply that the ovary equips a regulatory mechanism for controlling its glucose uptake. Nevertheless, there is no report that studies the expression of GLUT in bovine ovary. In the present study, therefore, we investigated the

Table 1 Follicular classification

Follicular type	Follicular diameter (mm)	E ₂ /P ₄	CL stage	Accompanied dominant follicle
POD	≥ 8.5	≥ 1	IV	–
EAD	≥ 8.5	≥ 1	I–III	–
EID	≥ 8.5	<1	I–IV	–
ASF	<8.5	–	I–IV	EID
SSF	<8.5	–	I–IV	POD or EAD

POD: preovulatory dominant follicle; EAD: E₂-active dominant follicle; EID: E₂-inactive dominant follicle; ASF: active subordinate follicle; SSF: suppressed subordinate follicle.

expression levels of GLUT1, 3 and 4 mRNA in the bovine follicle and CL in various physiologic conditions, and correlated them with concentrations of glucose in the follicular fluid.

Materials and Methods

Sample collection and storage

The tissue samples (liver, kidney, heart, pancreas, spleen, adrenal, muscle, fat, brain, lung and mammary gland) were collected from Holstein cows at the university farm. Because of the government regulation that restricts collection of fresh organ samples, only two sets of organ samples were used in the present study. They were sliced into small pieces and soaked in RNAlater (Ambion, Austin, TX, USA) and then kept at –30 °C until RNA extraction.

Paired ovaries were obtained from 21–26-month-old Holstein × Japanese Black F1 heifers at a local slaughterhouse. Only healthy ovarian pairs with a CL and follicles were used in the present study. CL were dissected and weighed, and their stages were assessed. They were cut in half vertically, and a 2 mm slice was taken from the central plane. Follicular fluid (FF) was aspirated from follicles by a syringe fitted with a 20 G needle and kept at –30 °C. Follicular walls were carefully peeled off from the ovaries, and individually placed in RNAlater and frozen at –30 °C. At the laboratory, the follicular walls were cut in half, and granulosa cells (GC) were harvested by gently scraping the walls with a spatula under a dissection microscope. The GC preparations obtained by this method were essentially free from contaminating thecal tissues. The remaining follicular walls were further scraped and washed several times with PBS to remove as much GC as possible. Surrounding stroma and theca externa were also removed from the follicular walls, and the cleaned follicular walls were used as theca interna layers (TL). The TL preparations obtained by this method were estimated to contain up to 5% of GC by determining the levels of aromatase mRNA arisen from contaminating GC. The tissue samples

were then placed in RNAlater and kept at –30 °C until RNA extraction.

Classification of CL and follicles

CL were macroscopically assessed by their color, vascularity and consistency by the criteria of Ireland *et al.* (1980), and were classified into four stages (I: days 1–4 after ovulation; II: days 5–10; III: days 11–17; and IV: days 18–20).

Follicles were classified into five groups by diameter (≥ 8.5 mm: dominant; <8.5 mm: subordinate), relative levels of E₂ and progesterone (P₄) in FF (E₂/P₄ ≥ 1: E₂ active; E₂/P₄ <1: E₂ inactive), the stage of CL (luteal phase: stages I–III; follicular phase: stage IV), and the relationship between dominant and subordinate follicles (subordinate follicles with either E₂-active or -inactive dominant follicles): that is, preovulatory dominant follicle (POD); E₂-active dominant follicle (EAD); E₂-inactive dominant follicle (EID); active subordinate follicle (ASF); and suppressed subordinate follicle (SSF). The details of these follicles are summarized in Table 1.

RNA extraction

Tissue samples were homogenized in a denaturing solution containing 4 M guanidium thiocyanate (Wako Pure Chemical Industries, Osaka, Japan), 25 mM sodium citrate (Wako), 0.5% sarkosyl (Sigma) and 0.1 M β-mercaptoethanol (Kanto Chemical, Tokyo Japan). Total RNA was extracted with phenol-chloroform (Chomczynski & Sacchi 1987) and then further purified and treated with DNase with a commercial kit (SV Total RNA Isolation System; Promega, Madison, WI, USA), and were frozen at –30 °C in THE RNA Storage Solution (Ambion).

Reverse-transcription and quantitative PCR

Single-strand cDNA was generated from total RNA (0.5–1.0 µg) with the 1st Strand cDNA Synthesis Kit for

Table 2 Primers for quantitative PCR

	Primer	Sequence	GenBank no. ^a	Position ^b
Gene (size: bp)				
GLUT1: standard (702)	Sense	caatgctgatgatgaacctgct	M60448	454–474
	Anti-sense	ccaccacaaatagcgacacga	M60448	1135–1155
GLUT1: real-time (134)	Sense	atcctcattgccgtgggtct	M60448	984–1003
	Anti-sense	acgatgccagagccgatggt	M60448	1098–1117
GLUT3: standard (290)	Sense	cgccttggcactctcaac	NM_174603	537–555
	Anti-sense	tccgcatactctcatcctca	NM_174603	806–826
GLUT3: real-time (142)	Sense	cgccttggcactctcaac	NM_174603	537–555
	Anti-sense	gcactggatgatggctgtaa	NM_174603	658–678
GLUT4: standard (720)	Sense	gcgtattctcgcgtgttctgg	D63150	273–294
	Anti-sense	gcttcgccttctcctctca	D63150	971–992
GLUT4: real-time (113)	Sense	ccaccaggcacactaccaca	D63150	410–430
	Anti-sense	ctcttcttcccagcactga	D63150	502–522
β -actin: standard (278)	Sense	gttcaacactcctgccatgat	K00622	149–161
	Anti-sense	gtagcagagcttctcctgatg	K00622	405–426
β -actin: real-time (228)	Sense	tgccatgatgtggccatcc	K00622	162–181
	Anti-sense	cgctcggctgtggtgtaa	K00622	371–389

^aGenBank (www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=Nucleotide). ^bNucleotide position in the reported sequence.

RT-PCR (AMV) (Roche) and a random primer. The RT cycle comprises 10-min annealing at 25 °C, 60-min cDNA synthesis at 42 °C, and 5-min inactivation at 99 °C.

Genes for GLUT1, 3 and 4 and β -actin were quantified by real-time PCR (LightCycler; Roche) with a commercial kit (QuantiTect SYBR Green PCR; Qiagen). The primers were designed with Primer3 (Rozen & Skaletsky 2004: <http://frodo.wi.mit.edu/cgi-bin/primer3/primer3-www.cgi>) by bovine sequences (Table 2). The amplification program consists of initial 15-min activation at 95 °C followed by 40 cycles of PCR steps (15-s denaturation at 94 °C, 30-s annealing at 65 °C, and 20-s extension at 72 °C). For the quantification of the target genes, a series of standards was constructed by amplifying a fragment of DNA (278–720 bp) that contains the target sequence for real-time PCR (113–228 bp). The PCR products were subjected to electrophoresis, and the target band was cut out and purified with a DNA purification kit (SUPREC-01, TaKaRa Bio, Otsu, Japan). Five to eight stepwise-diluted DNA standards were included in every PCR run. The intra- and interassay coefficients of variation were 15.0% and 21.8% for GLUT1, 15.3% and 10.1% for GLUT3, 19.4% and 26.6% for GLUT4, and 11.2% and 16.4% for β -actin. The values were normalized with β -actin as the internal standard.

Steroid hormone extraction and assay

Concentrations of E_2 and P_4 in the FF samples were determined by an enzyme immunoassay (EIA), as detailed previously (Miyamoto *et al.* 1992). Steroid hormones were extracted with diethyl ether. The extraction efficiency was 85%. Ranges of the standard curves were 2–2000 pg/ml for E_2 and 0.05–50 ng/ml for P_4 .

The intra- and interassay coefficients of variation were 6.2% and 8.5% for E_2 , and 4.5% and 7.4% for P_4 .

Glucose assay

Concentrations of glucose in the FF samples were determined by the colorimetric method with a commercial kit (Glucose CII Test, Wako). The intra- and interassay coefficients of variation were 1.4% and 1.1% respectively.

Statistical analysis

The data were subjected to one-way analysis of variance (ANOVA) followed by the Steel–Dwass multiple comparison test. The relationships among the levels of GLUT mRNA and concentrations of glucose in FF were determined by regression analysis.

Results

Characteristics of follicles

A total of 59 follicles (1–5 follicles/animal) and 32 CL from 32 animals were used in the present study. Concentrations of steroids and glucose in follicular fluid and diameter of follicles in each group are shown in Table 3. There were significant differences in steroid concentrations among large follicles as well as between large and small follicles, while no clear difference was observed in any factor between ASF and SSF. Concentrations of glucose in FF were comparable to the levels previously reported (30–70 mg/dl) and slightly lower than those

Table 3 Characteristics of each follicular category: diameter, concentrations of E₂, P₄ and glucose in follicular fluid

	Follicle diameter (mm)	E ₂ (ng/ml)	P ₄ (ng/ml)	E ₂ /P ₄	Glucose (mg/dl)
Group (n)					
POD (8)	16.0 ± 1.3 ^a	81.3 ± 12.6 ^a	23.6 ± 2.6 ^a	3.6 ± 0.6 ^a	66.5 ± 3.2 ^a
EAD (13)	11.5 ± 0.7 ^a	27.7 ± 2.2 ^b	16.5 ± 2.0 ^{ac}	2.0 ± 0.3 ^a	63.1 ± 3.5 ^a
EID (13)	13.4 ± 0.6 ^a	3.2 ± 1.3 ^c	251.3 ± 67.0 ^c	0.0 ± 0.0 ^b	50.6 ± 5.5 ^{ab}
ASF (15)	7.2 ± 0.2 ^b	2.0 ± 0.6 ^c	9.8 ± 1.9 ^{cd}	0.3 ± 0.1 ^c	37.6 ± 4.6 ^b
SSF (10)	7.3 ± 0.3 ^b	1.7 ± 0.5 ^c	7.7 ± 2.0 ^d	0.2 ± 0.1 ^c	54.4 ± 3.0 ^{ab}

Mean ± S.E.M. (range). POD: preovulatory dominant follicle; EAD: E₂-active dominant follicle; EID: E₂-inactive dominant follicle; ASF: active subordinate follicle; SSF: suppressed subordinate follicle. ^{a,b,c,d}P<0.05, significantly different from each other.

in serum (80 mg/dl) in dairy cows (Leroy *et al.* 2004). Glucose concentration in ASF was significantly lower than that in POF and EAD.

Gene expression of GLUT1, 3 and 4 in tissues

The mRNA encoding GLUT1, 3 and 4 were detected in various tissues (Table 4). The gene expression pattern of these GLUT was consistent with that reported in bovine (Zhao *et al.* 1993) and other species (Mueckler 1994, Hocquette & Abe 2000). While GLUT1 and 3 mRNA were expressed in all tissues examined at substantial levels, GLUT4 mRNA was preferentially expressed in muscle, heart and subcutaneous fat.

Gene expression of GLUT1, 3 and 4 in follicles

To obtain an approximate idea of GLUT expression levels in relation to known GLUT-expressing organs, levels of GLUT1, 3 and 4 mRNA in ovarian tissues were normalized with values obtained from brain, heart and muscle respectively. The mRNA encoding GLUT1, 3 and 4 were all detected in bovine follicles (Fig. 1). The expression levels of

GLUT1 and 3 in GC and TL appeared to be as high as those in brain and heart, whereas the expression level of GLUT4 was less than 1% of that in muscle. The expression patterns of these GLUT appear to be different between GC and TL and among physiologic statuses of follicles. Approximately similar levels of GLUT1 mRNA were detected in GC and TL. On the other hand, the expression of GLUT3 mRNA was higher in GC than TL, while that of GLUT4 mRNA was much higher in TL, and negligible levels of expression were detected in GC (Fig. 1). The expressions of GLUT1 and 3 mRNA in GC and TL, and GLUT4 mRNA in TL were significantly higher or tended to be higher in EID follicles than in the other follicular classes (Fig. 1). The expressions of these GLUT among E₂-active dominant follicles (POD and EAD) and subordinate follicles (ASF and SSF) were virtually the same except for GLUT3 in GC of SSF, where the expression level was significantly lower than in the other groups (Fig. 1).

Gene expression of GLUT1, 3 and 4 in corpus luteum

The bovine CL expresses mRNA for GLUT1, 3 and 4 at comparable levels to the follicle (Fig. 2). The expression of

Table 4 Relative expression of GLUT mRNA in tissues

	GLUT1	GLUT3	GLUT4
Tissue (n=2)			
Liver	27.7 ± 0.5	22.7 ± 0.8	1.8 ± 0.9
Kidney (cortex and medulla)	26.8 ± 17.5	9.0 ± 7.3	0.5 ± 0.4
Adrenal (cortex and medulla)	36.0 ± 31.1	37.0 ± 33.0	0.6 ± 0.3
Pancreas	19.6 ± 14.7	45.4 ± 7.7	3.9 ± 2.0
Spleen	9.7 ± 2.5	33.5 ± 13.1	0.6 ± 0.3
Muscle	7.9 ± 0.5	25.3 ± 4.9	100.0 ± 41.2
Heart	13.1 ± 2.7	100.0 ± 29.5	40.4 ± 10.2
Mammary gland	26.0 ± 20.4	14.2 ± 4.3	0.3 ± 0.0
Subcutaneous fat	16.8 ± 13.7	28.1 ± 6.4	14.6 ± 11.2
Uterine endometrium	20.6 ± 11.1	11.8 ± 8.1	0.9 ± 0.8
Oviduct (ampulla)	54.0 ± 53.6	13.6 ± 8.9	1.5 ± 1.7
Lung*	6.4	49.3	0.4
Brain (cerebral cortex)*	100.0	42.9	0.1

Values are expressed as percentage relative to the values in the highest expressing organs (mean ± S.D.): mean of two samples was used for GLUT3 (heart) and GLUT4 (muscle), while a single sample was used for GLUT1 (brain). *Only one sample was obtained for lung and brain.

Figure 1 Gene expression of GLUT1 (a), GLUT3 (b) and GLUT4 (c) in bovine granulosa cells (GC) and theca interna layer (TL). Follicles were harvested from Holstein × Japanese Black F1 heifers and classified into five categories by physiologic status (Table 3). The levels of mRNA encoding GLUT1, 3 and 4 were quantified by real-time PCR and normalized with β -actin as the internal standard. Values (mean $\pm$ S.E.M.) are expressed as percentage relative to brain (GLUT1), heart (GLUT3) and muscle (GLUT4). ^{a,b} $P < 0.05$ and ^{*} $P < 0.05$, significantly different from each other.

Figure 2 Gene expression of GLUT1 (a), GLUT3 (b) and GLUT4 (c) in bovine corpora lutea (CL). CL were harvested from Holstein × Japanese Black F1 heifers and classified into four luteal stages (I–IV), as described in Materials and Methods. The levels of mRNA encoding GLUT1, 3 and 4 were quantified by a real-time PCR and normalized with β -actin as the internal standard. Values (mean $\pm$ S.E.M.) are expressed as percentage relative to brain (GLUT1), heart (GLUT3) and muscle (GLUT4). ^{a,b} $P < 0.05$, significantly different from each other.

Table 5 Correlations between GLUT mRNA levels in granulosa cells and theca interna and glucose concentrations in follicular fluid

	GLUT1		GLUT3		GLUT4	
	GC	TL	GC	TL	GC	TL
Group (n)						
POD (8)	-0.25	0.27	-0.40	0.69	-0.18	0.70
EAD (13)	-0.09	-0.35	-0.18	0.31	0.09	0.46
EID (13)	-0.86***	-0.47	-0.87***	-0.41	-0.50	-0.51
ASF (15)	-0.38	-0.31	-0.50	0.55*	-0.05	0.65**
SSF (10)	0.40	-0.20	-0.23	0.33	0.62	0.80**

POD: preovulatory dominant follicle; EAD: E₂-active dominant follicle; EID: E₂-inactive dominant follicle; ASF: active subordinate follicle; SSF: suppressed subordinate follicle. * $P<0.05$; ** $P<0.01$; *** $P<0.001$.

GLUT1 mRNA was stable throughout the luteal phase (i.e., stages I–III) and then decreased significantly as CL regressed in stage IV. The expression of GLUT3, on the other hand, increased sharply in stage IV. The expression of GLUT4 followed the pattern of the luteal progesterone secretion, with the highest levels attained during the midluteal phase (stages II and III).

Relationship between concentrations of glucose in follicular fluid and levels of GLUT mRNA

There were highly significant negative correlations between the concentration of glucose in FF and the mRNA levels of GLUT1 ($r=-0.86$, $P<0.0001$) and GLUT3 ($r=-0.87$, $P<0.0001$) in GC for EID (Table 5). Significant positive correlations were also found between the concentration of glucose and the mRNA level of GLUT4 in TL for ASF ($r=0.65$, $P<0.01$) and SSF ($r=0.80$, $P<0.01$).

Discussion

The present study demonstrated that the mRNA encoding three members of the sugar transporter family, GLUT1, 3 and 4, is expressed in the bovine ovary. The expression levels of GLUT1 and 3 were comparable to the known GLUT-expressing organs, such as brain (Wood & Trayhurn 2003), indicating that they also play important roles in the bovine ovary. However, the expression level of GLUT4 was much lower than that in the classic GLUT4-expressing organs, muscle and adipose tissues, and this expression suggests a possible supporting role in the ovary.

The gene expression pattern of these GLUT appeared to be different between GC and TL. A ubiquitous transporter GLUT1 (Mueckler 1994, Hocquette & Abe 2000, Wood & Trayhurn 2003) is expressed in both GC and TL at equal levels. On the other hand, high-affinity transporter GLUT3 (Wood & Trayhurn 2003) is expressed predominantly in GC, while the insulin-responsive transporter GLUT4 (Wood & Trayhurn 2003) is expressed

almost exclusively in TL. These results indicate that glucose transport is regulated in a tissue-specific manner in the bovine ovary.

The expression of these GLUT has been demonstrated in the ovary of sheep (Williams *et al.* 2001), rat (Kol *et al.* 1997, Kodaman & Behrman 1999) and mouse (Zhou *et al.* 2000), but the expression pattern appears to be different among species. In the ovine ovary, GLUT1 and 4 were shown to be expressed in both GC and TL (Williams *et al.* 2001), while in the rat ovary both GLUT1 and 3 were localized only in GC (Kol *et al.* 1997). Our results add another proof of the considerable species difference in the ovarian expression of GLUT.

Despite the substantial differences in physiologic condition, there was no difference in the gene expression level of GLUT among follicular groups except EID, where a notable increase in the expression of all three GLUT was observed. In the present study, we were unable to define clearly small growing and atretic follicles and thus ASF and SSF groups likely to contain both healthy and atretic follicles. Nevertheless, the results indicate that GLUT1, 3 and 4 are equally expressed in follicles over various physiologic conditions, at least on the transcriptional level per cell basis (i.e., actin normalized value).

The substantial increase in the gene expression of these GLUT in EID follicles was unexpected. Highly significant negative correlations found between levels of GLUT mRNA and concentrations of glucose in FF, however, may be a clue to understand this phenomenon. Glucose is a known regulator of GLUT expression both *in vivo* and *in vitro*. For example, it was demonstrated that glucose deprivation causes upregulation of GLUT1 and 3 in cultured bovine chromaffin cells (Fladeby *et al.* 2003). Long-term hypoglycemia, likewise, was shown to increase GLUT3 density in rat brain (Duelli *et al.* 1999). Taken together, these results imply the presence of a local regulatory mechanism of glucose uptake, which compensates for glucose deprivation and thus maintains vital energy supply to follicular cells. During the luteal phase, dominant follicles (i.e., EAD) lose their functional dominance approximately 3 days after the initiation of

dominance and undergo the atretic process (i.e., EID) (Ali *et al.* 2001). One of the first signs of atresia is the gradual loss of thecal vascularity (Moor *et al.* 1978, Jiang *et al.* 2003). In consequence, the supply of glucose, as well as other nutrients and oxygen to follicles, is diminished. Meanwhile, FF acts as a nutritional reservoir for the avascular follicles. Without replenishment, however, glucose in FF is gradually used up, leading to a condition of hypoglycemia. In the present study, follicular concentrations of glucose in the EID varied widely (6.9–71.3 mg/dl), whereas those in the POD and EAD were largely stable (47.4–96.2 mg/dl). These results imply that a severe glucose shortage occurs as the atretic process progresses. Thus, in the face of glucose deprivation, GC might over-express GLUT mRNA to maintain glucose homeostasis. The exact mechanism that regulates GLUT expression in the ovary, however, needs further investigation.

The mRNA encoding GLUT1, 3 and 4 were all expressed in CL. The expression level of GLUT1 mRNA appeared to be much higher in CL than in follicles whereas the levels of GLUT3 and 4 mRNA in CL were comparable to those in follicles. The expression patterns of GLUT1, 3 and 4 differed substantially among the genes, indicating that these GLUT are differentially regulated in bovine CL. To our knowledge, gene expression of GLUT in the CL has not been reported other than in rat, where GLUT1 and 3 were shown to be expressed in the newly formed CL (Kol *et al.* 1997). However, the presence of GLUT has been indicated in CL. GLUT1 is suggested to be responsible for luteal uptake of dehydroascorbic acid, which serves as the substrate of ascorbic acid in rat (Kodaman *et al.* 1998). The bovine CL was also shown to take up large amounts of glucose during the early and midluteal phase (Chase *et al.* 1992). The glucose uptake decreases in late or regressing CL (Chase *et al.* 1992). Our results are consistent with these results and suggest that GLUT1 is the major glucose transporter in the bovine CL. GLUT3 and 4 may play supporting roles in the CL in a stage-specific manner.

In summary, the bovine follicle and CL express mRNA encoding GLUT1 and 3 at levels comparable to the classic GLUT expressing organs such as brain. Much less but appreciable levels of GLUT4 were also detected in these tissues. The gene expression of these GLUT showed a stage- and tissue-specific pattern and is likely to be regulated by glucose. Thus, it is concluded that GLUT1 and 3 act as major glucose transporters, while GLUT4 may play a supporting role in the bovine follicle and CL. Together, these GLUT maintain a vital glucose supply to the ovary to sustain follicular development, ovulation, CL formation and maintenance.

Acknowledgements

This study was supported by a Grant-in-Aid for Scientific Research (16580237) from the Japan Society for the

Promotion of Science and a grant from the 21st Century COE Program (A-1), Ministry of Education, Culture, Sports, Science, and Technology, Japan. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Ali A, Lange A, Gilles M & Glatzel PS 2001 Morphological and functional characteristics of the dominant follicle and corpus luteum in cattle and their influence on ovarian function. *Theriogenology* **56** 569–576.
- Chase CC Jr, Del Vecchio RP, Smith SB & Randel RD 1992 In vitro metabolism of glucose by bovine reproductive tissues obtained during the estrous cycle and after calving. *Journal of Animal Science* **70** 1498–1508.
- Chomczynski P & Sacchi N 1987 Single step method of RNA isolation by acid guanidinium thiocyanate phenol chloroform extraction. *Annals of Biochemistry* **162** 156–159.
- Duelli R, Staudt R, Duembgen L & Kuschinsky W 1999 Increase in glucose transporter densities of Glut3 and decrease of glucose utilization in rat brain after one week of hypoglycemia. *Brain Research* **831** 254–262.
- Fladeby C, Skar R & Serck-Hanssen G 2003 Distinct regulation of glucose transport and GLUT1/GLUT3 transporters by glucose deprivation and IGF-I in chromaffin cells. *Biochimica et Biophysica Acta* **1593** 201–208.
- Hocquette JF & Abe H 2000 Facilitative glucose transporters in livestock species. *Reproduction, Nutrition, Development* **40** 517–533.
- Ireland JJ, Murphee RL & Coulson PB 1980 Accuracy of predicting stages of bovine estrous cycle by gross appearance of the corpus luteum. *Journal of Dairy Science* **63** 155–160.
- Jiang JY, Macchiarelli G, Tsang BK & Sato E 2003 Capillary angiogenesis and degeneration in bovine ovarian antral follicles. *Reproduction* **125** 211–223.
- Kodaman PH & Behrman HR 1999 Hormone-regulated and glucose-sensitive transport of dehydroascorbic acid in immature rat granulosa cells. *Endocrinology* **140** 3659–3665.
- Kodaman PH, Aten RF & Behrman HR 1998 Accumulation of ascorbate by endocrine-regulated and glucose-sensitive transport of dehydroascorbic acid in luteinized rat ovarian cells. *Biology of Reproduction* **58** 407–413.
- Kol S, Ben-Shlomo I, Ruutinen K, Ando M, Davies-Hill TM, Rohan RM, Simpson IA & Adashi EY 1997 The midcycle increase in ovarian glucose uptake is associated with enhanced expression of glucose transporter 3. Possible role for interleukin-1, a putative intermediary in the ovulatory process. *Journal of Clinical Investigation* **99** 2274–2283.
- Leroy JLMR, Vanholder T, Delanghe JR, Opsomer G, Van Soom A, Bols PEJ & de Kruif A 2004 Metabolite and ionic composition of follicular fluid from different-sized follicles and their relationship to serum concentrations in dairy cows. *Animal Reproduction Science* **80** 201–211.
- Miyamoto A, Okuda K, Schweigert FJ & Schams D 1992 Effects of basic fibroblast growth factor, transforming growth factor-beta and nerve growth factor on the secretory function of the bovine corpus luteum in vitro. *Journal of Endocrinology* **135** 103–114.
- Moor RM, Hay MF, Dott HM & Cran DG 1978 Macroscopic identification and steroidogenic function of atretic follicles in sheep. *Journal of Endocrinology* **77** 309–318.
- Mueckler M 1994 Facilitative glucose transporters. *European Journal of Biochemistry* **219** 713–725.
- Rabiee AR, Lean IJ, Gooden JM, Miller BG & Scaramuzzi RJ 1997 An evaluation of transovarian uptake of metabolites using arterio-venous difference methods in dairy cattle. *Animal Reproduction Science* **48** 9–25.

- Rozen S & Skaletsky HJ 2000 Primer3 on the WWW for general users and for biologist programmers. In *Bioinformatics Methods and Protocols: Methods in Molecular Biology*, pp 365–386. Eds S Krawetz & S Misener. Totowa, New Jersey: Humana Press.
(http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi).
- Williams SA, Blache D, Martin GB, Foot R, Blackberry MA & Scaramuzzi RJ 2001 Effect of nutritional supplementation on quantities of glucose transporters 1 and 4 in sheep granulosa and theca cells. *Reproduction* **122** 947–956.
- Wood IS & Trayhurn P 2003 Glucose transporters (GLUT and SGLT): expanded families of sugar transport proteins. *British Journal of Nutrition* **89** 3–9.
- Zhao FQ, Glimm DR & Kennelly JJ 1993 Distribution of mammalian facilitative glucose transporter messenger RNA in bovine tissues. *International Journal of Biochemistry* **25** 1897–1903.
- Zhou J, Bievre M & Bondy CA 2000 Reduced GLUT1 expression in Igfl^{-/-} null oocytes and follicles. *Growth Hormone and IGF Research* **10** 111–117.

Received in final form 30 August 2005

Accepted 17 October 2005

Made available online as an Accepted Preprint
9 November 2005