Modulation of ACTH-induced cortisol release by polyunsaturated fatty acids in interrenal cells from gilthead seabream, *Sparus aurata*

R Ganga, L Tort¹, L Acerete¹, D Montero and M S Izquierdo

Grupo de Investigación en Acuicultura, Instituto Canario de Ciencias Marinas, PO Box 56, 35200-Telde, Las Palmas, Canary Islands, Spain

¹Department of Cell Biology, Physiology and Immunology, Universitat Autònoma de Barcelona, Bellaterra 08193, Spain

(Requests for offprints should be addressed to R Ganga; Email: rgaqua2003@yahoo.fr)

Abstract

Highly unsaturated fatty acids are essential components of cellular membranes of vertebrates and can modulate physiological processes, including membrane transport, receptor function and enzymatic activities. In gilthead sea bream, dietary deficiencies of essential fatty acids of marine fish raise the basal cortisol levels and alter the pattern of cortisol release after stress. The aim of the present study was to clarify the effect of different essential fatty acids on adrenocorticotrophic hormone (ACTH)-induced cortisol production and release in fish, through *in vitro* studies of sea bream interrenal cells maintained in superfusion and incubated with different types of fatty acids and eicosanoid production inhibitors. Results showed the first evidence of the effect of certain fatty acids on cortisol production by

ACTH-stimulated interrenal cells in fish. Both arachidonic acid (ARA) and particularly eicosapentaenoic acid (EPA) promoted cortisol production in sea bream interrenal cells. Moreover, incubation with indometacin (INDO) reduced the increased cortisol production induced by EPA and ARA, suggesting mediation by their cyclooxygenase-derived products. Docosahexaenoic acid stimulated cortisol production to a lesser extent than that caused by EPA or ARA, but the inhibitory effect of INDO was not as marked as it was for the other fatty acids. In contrast, supplementation with dihomo-gammalinoleic acid reduced cortisol production, denoting the inhibitor effect of this fatty acid in cortisol secretion.

Journal of Endocrinology (2006) **190**, 39–45

Introduction

Highly unsaturated fatty acids with 20 or more carbon atoms and three or more double bonds (HUFA) are essential components of cellular membranes and can modulate physiological processes, including membrane transport, receptor function and enzymatic activities. Hence, dietary fatty acids have been shown to have marked effects on a variety of immunological and haemostatic parameters (Balfry *et al.* 2001, Montero *et al.* 2001). HUFA possess a wide range of cellular functions. One of the most important functions is to supply precursors for the synthesis of eicosanoids, which are produced in response to various extracellular stimuli by two main types of dioxygenase enzymes: cyclooxygenases (COX) and lipoxygenases (Horrobin 1983). Following cell stimulation, both arachidonic acid (ARA; 20:4n-6) and eicosapentaenoic acid (EPA; 20:5n-3) are released from the membrane by the action of phospholipase A₂. Later these fatty acids are transformed by a range of lipoxygenases and cyclooxygenases to yield prostaglandins (PG), leukotrienes, lipoxins and other compounds, which can modulate several immune functions (Uhing *et al.* 1990).

Eicosanoids have been found in a large range of freshwater and marine fish (Matsumoto *et al.* 1989, Mustafa & Srivastava

1989) and in many tissues (Henderson & Tocher 1987, Bell *et al.* 1994a, Tocher 1995). In fish, a preferred eicosanoid precursor for cyclooxygenase seems to be ARA (20:4n-6) (Tocher & Sargent 1987, Bell *et al.* 1994a, 1994b, 1998), but EPA (20:5n-3) and dihomo- γ -linolenic acid (DHGLA; 20:3n-6) are also important eicosanoid precursors which can modulate production and biological efficacy of ARA-derived eicosanoids (Horrobin 1983, Bell *et al.* 1994a, Ganga *et al.* 2005). In addition, the high content of docosahexaenoic acid (DHA; 22:6n-3) in cellular membranes affects eicosanoid production (Nablone *et al.* 1990). This fatty acid is also recognised as a precursor of certain biologically active trioxilated derivatives (German *et al.* 1983, Hong *et al.* 2005). Therefore, the supply of precursor polyunsaturated fatty acids with 18 or more carbon atoms and two or more double bonds (PUFA) for eicosanoid synthesis is directly related to the fatty acid composition of membrane phospholipids, which in turn is influenced by dietary PUFA intake and metabolism (Lands 1989).

In gilthead sea bream, dietary deficiencies on n-3 HUFA, essential fatty acids for marine fish (Izquierdo 1996), raised the basal plasma cortisol levels and altered the pattern of cortisol release after stress (Montero *et al.* 1998). Cortisol is a key corticosteroid hormone for homeostatic response to stress

in all vertebrates, through its effects on metabolism and immune function (Hontela 1997, Wendelaar Bonga 1997) as well as the osmoregulation process (Wendelaar Bonga 1997). Thus, the increase in plasma cortisol levels is regarded as the most reliable method for differentiating between stressed and non-stressed fish (Thompson *et al.* 1993, Yin *et al.* 1995, Rotllant & Tort 1997). Moreover, feeding relatively low levels of n-3 HUFA, although not affecting growth and feed efficiency, significantly raised plasma cortisol levels (Montero *et al.* 2003).

However, the physiological mechanisms by which these HUFA regulate the hormone-induced plasma cortisol levels are not clear. In fish, several studies have suggested that ARA is involved in the release of cortisol, although the actual mechanisms have not been investigated (Gupta *et al.* 1985, Bessonart *et al.* 1999, Harel *et al.* 2001, Koven *et al.* 2003, Van Anholt *et al.* 2004). In mammals, certain studies suggest that PG play an important role in mediating the corticosteroidogenic action of adrenocorticotrophic hormone (ACTH) (Kocsis *et al.* 1999), and thus the role of fatty acids in stress response seems to be mediated by the production of eicosanoids.

The present study aims to clarify the effect of different HUFA on ACTH-induced cortisol production and release by gilthead sea bream interrenal cells.

Material and Methods

Animals

Sexually immature gilthead sea bream (*Sparus aurata*) of body weight 54.7 ± 11.2 g supplied by a Spanish fish farm (Masnou, Barcelona, Spain) were kept for 2 weeks in two fibreglass tanks of 1000 l held in a semi-closed seawater circulation system equipped with physical and biological filters. Water temperature was maintained at 16–18 °C, the salinity at 35–40‰ and photoperiod at 12 h light:12 h darkness. Fish were fed once a day with a commercial feed until 24 h before the *in vitro* trials to avoid feed interference. A total number of 30 fish were employed in the experiments.

Superfusion trials

After 2 weeks of acclimatisation, fish were randomly taken from the tanks in less than 1 min, immediately anaesthetised with 2-phenoxyethanol (1:1000 v/v) and blood collected with a hypodermic syringe from the caudal vein to minimise the haemorrhage. Head kidney tissue was removed from two fish in each superfusion trial and cut into very small fragments in Hepes Ringer medium, which was used as the superfusion medium. Afterwards, head kidney homogenates were pooled and distributed in eight superfusion chambers (volume: 0.2 ml) in order to obtain a homogeneous aliquot from each of them. Tissues were superfused with a Hepes (pH 7.4) Ringer's solution containing 171 mM NaCl, 2 mM KCl, 2 mM CaCl₂·H₂O, 0.25% (w/v) glucose and 0.03% (w/v)

bovine serum albumin (Rotllant *et al.* 2001). The system was temperature-controlled at 15 °C and superfusion medium was pumped through the chamber at a rate of 75 µl/min by a Masterplex L/S^R multichannel peristaltic pump (Cole Parmer Instrument Co. Vernon Hills, IL, USA).

Trials were started after 3 h of superfusion when cortisol reached a stable baseline level (Rotllant *et al.* 2000a, 2000b) due to several factors such as the different dispersion of interrenal cells in the perfusion preparation, individual differences and the pre-stress level of each fish. After the stabilisation period of 3 h, tissues were subsequently incubated with different fatty acids. A series of preliminary tests were performed in quadruplicate, to determine the adequate fatty acid concentration (50, 150 or 300 µM) and incubation time (1 or 3 h) for any of the three fatty acids assayed (ARA, EPA and DHA). Best cortisol stimulation was found with fatty acid concentrations of 50 µM and an incubation time of 1 h (Table 1) and these conditions were used afterwards in all the research experiments. Both in these preliminary tests and in the research experiments, perfusion medium was supplemented with the corresponding concentration of different fatty acids ARA, EPA, DHA and DHGLA (diluted in less than 0.5% of ethanol/medium v/v) prior to tissue incubation. In a second series of experiments to clarify the action mechanisms of these fatty acids, tissues were incubated with a COX inhibitor indometacin (INDO) for 20 min at a concentration of 25 µM diluted in superfusion medium. After incubation with the fatty acids, the perfused tissues were stimulated with ACTH at a concentration of 5 nM hACTH_{1–39} (Sigma) for 20 min. Subsequently, perfusion was maintained for another 170 min, fraction samples being collected every 20 min during this period. Cortisol stimulation factor was calculated by the comparison of maximum cortisol released after ACTH stimulation with baseline cortisol released (maximum release—baseline release)/(baseline release) (Rotllant *et al.* 2001). In all the series of experiments, each treatment was assayed in quadruplicate.

Cortisol measurements

Cortisol concentration in the perfused fluid was determined by RIA (Rotllant *et al.* 2001). The antibody used for the assay was purchased from Biolink, S.L. (Costa Mesa, CA, USA) in a

Table 1 Effect of two fatty acid concentrations (50 and 150 µM) and two incubation times (1 h and 3 h) for three polyunsaturated fatty acids on cortisol secretion stimulation factor

	1 h	3 h	
Treatment			
Control	14.71 ± 2.41	13.28	
EPA			
50 µM	29.63 ± 2.59	7.79 ± 3.29	
150 µM	7.79 ± 3.29	—	
ARA			
50 µM	22.26 ± 6.29	11.75 ± 4.16	
150 µM	12.25 ± 1.86	—	
DHA			
50 µM	35.72 ± 9.28	2.60 ± 1.16	
150 µM	4.47 ± 0.28	—	

final dilution of 1:6000. This antibody cross reactivity is 100% with cortisol, 11.40% with 21-desoxycorticosterone, 8.90% with 11-desoxycortisol and 1.60% with 17 α -hydroxyprogesterone. The radioactivity was quantified using a liquid scintillation counter. Cortisol levels are given as ng/g/h.

Statistical analysis

Significance of difference ($P < 0.05$) between dietary treatments was determined by ANOVA, followed by Duncan's multiple comparison test (Sokal & Rolf 1995). Analyses were performed using SPSS software (SPSS for Windows 11.5; SPSS Inc., Chicago, IL, USA).

Results

The different incubation times and fatty acid concentrations assayed showed that 1 h of incubation time and a concentration of 50 μ M of fatty acid were the best conditions to obtain the highest effect of fatty acid on cortisol secretion stimulation factor (Table 1). As expected, after the stabilisation period of 3 h, cortisol values remained at basal levels for these fish species and no significant differences were found among basal values for the different superfused tissues (Fig. 1).

The effects of supplementation with different HUFA on cortisol secretion are illustrated in Fig. 1. The production of cortisol by interrenal cells was modified when the medium was supplemented with HUFA in comparison with the control. Addition of n-3 fatty acids, DHA and EPA induced a higher and earlier cortisol response to ACTH than the control without fatty acid incubation. Addition of n-6 fatty acids did not modify the time of cortisol response in comparison

to the control, but induced a higher response. Cortisol response was higher when ARA, EPA or DHA was added to fatty acid and lower when DHGLA was used. Such response expressed as stimulation factor was significantly ($P < 0.05$) higher with EPA (33.71 ± 4.5 basal secretion) and ARA (28.7 ± 4.47) incubation than control and DHGLA treatment groups (Fig. 2). With DHA incubation, no significant differences were found in the stimulation factor. By contrast, DHGLA showed the lowest ($P < 0.05$) stimulation factor with an increase of only 8.95 ± 2.17 .

Supplementation of INDO, a COX inhibitor, induced the stimulation of cortisol production by EPA and ARA observed in the former set of experiments, with cortisol absolute values not being different from those of the control (Fig. 3). However, a significantly higher ($P < 0.05$) cortisol peak was obtained when the tissue was incubated with DHA, despite the addition of INDO (Fig. 3).

Comparison of cortisol stimulation factors when INDO was added showed a significantly higher ($P < 0.05$) cortisol secretion in the tissue supplemented with EPA, ARA and DHA (Fig. 4). Thus, the stimulation factor of cortisol was 7.83 ± 3.31 when tissue was supplemented with EPA, 6.97 ± 4.56 with ARA, 13.67 ± 2.66 with DHA and only 1.58 ± 0.45 for control.

In addition, the comparison of cortisol stimulation factors between experiments with or without INDO showed that the addition of INDO significantly decreased ACTH-stimulated cortisol secretion in all the treatment use of this COX inhibitor (Fig. 5). However, this impaired stimulation of cortisol production was lower in the DHA-supplemented group in which INDO caused a 40.84% reduction in cortisol secretion, giving values that were significantly different ($P < 0.05$) compared to EPA treatment where INDO caused

Figure 1 Absolute cortisol secretion (ng/g Hk/h) by sea bream head kidney (Hk) after ACTH stimulation following incubation with HUFA (different letters for a given time indicate significant difference, $P < 0.05$).

Figure 2 Cortisol stimulation factor in sea bream head kidney after ACTH stimulation following incubation with different HUFA (different letters for different treatments indicate significant difference, $P < 0.05$ ANOVA).

a 76.76% reduction in cortisol secretion and a 75.71% reduction with ARA treatment.

Discussion

The present study showed the first evidence of the effect of HUFA on cortisol production by ACTH-stimulated interrenal cells in fish. These results are in agreement with the observed modulating effect of dietary fatty acids in sea bream plasma cortisol levels

(Montero *et al.* 1998, 2001) and confirm the hypothesis of these authors about the effective action of these fatty acids on the cortisol secretion by the interrenal cells in gilthead sea bream.

Both ARA and EPA promoted ACTH-induced cortisol production in sea bream interrenal cells used in the present experiment. Dietary EPA has been shown to affect fish stress resistance in several species. Although it promoted the growth and survival of the red sea bream (Watanabe *et al.* 1989), gilthead sea bream (Liu *et al.* 2002) and Japanese flounder (Furuita *et al.* 1998), its effects on larval stress resistance seem to depend on species and dietary levels. For instance, elevation of dietary EPA increased red sea bream handling stress resistance (Watanabe *et al.* 1989) and gilthead sea bream resistance to air exposure and temperature shock, but not to salinity stress (Liu *et al.* 2002). On the contrary, too high EPA levels reduced stress resistance to air exposure in Japanese flounder (Furuita *et al.* 1998). ARA has also been shown to affect stress resistance in several fish species. Dietary ARA levels of about 1% dry weight feed are necessary not only for optimum growth and survival of sea bream larvae (Bessonart *et al.* 1999), but also for improved stress resistance after handling (Koven *et al.* 2003, Van Anholt *et al.* 2004). Dietary ARA levels close to those used by these authors did not affect the handling of stress resistance in Japanese flounder, whereas higher ones reduced larval stress resistance (Furuita *et al.* 1998).

These differences of the effects of dietary EPA or ARA on stress resistance in different species may also be related to different ratios among these fatty acids, since both are competing substrates for cyclooxygenase enzymes (Izquierdo *et al.* 2001). For instance, in Atlantic salmon, alteration in the dietary ratio of n-3/n-6 fatty acids has been shown to prevent stress susceptibility to transport (Bell *et al.* 1991). The present

Figure 3 Absolute cortisol secretion (ng/g Hk/h) by sea bream head kidney (Hk) after ACTH stimulation following incubation with different HUFA and INDO supplementation (different letters for a given time indicate significant difference, $P < 0.05$).

Figure 4 Cortisol stimulation factor in sea bream head kidney after ACTH stimulation following incubation with different HUFA and INDO supplementation (different letters for indicate significant differences among treatment).

study shows the first evidence found in fish that cyclooxygenase-derived metabolites are involved in ACTH-induced cortisol release by interrenal cells. The strong reduction of cortisol release caused by INDO addition in EPA and ARA supplemented groups suggested that the effect of these fatty acids was, at least partly, mediated by their cyclooxygenase-derived metabolites. Both fatty acids have been found to be good precursors of cyclooxygenase-derived PG in fish (Bell *et al.* 1994a, Ganga *et al.* 2005). In turn, cyclooxygenase-derived PG have been shown to increase *in vitro* cortisol release in interrenal tissue of female frogs during

Figure 5 Comparison of cortisol stimulation factor in sea bream head kidney after ACTH stimulation following incubation with different HUFA and including (filled bars) or not (dotted bars) INDO supplementation (different letters for control or each fatty acid incubation indicates significant differences by INDO addition).

ovulation (Gobbetti & Zerani 1993) and in human adrenal cells as well (Vakharia & Hinson 2005).

Interestingly, DHA stimulation of ACTH-induced cortisol production was lower than that caused by EPA or ARA. Besides, the inhibitory effect of INDO in the DHA-supplemented group was not so marked as in the other treatments, suggesting that the action of DHA in cortisol release from ACTH-stimulated interrenal cells is less dependent on COX metabolites in gilthead sea bream. Indeed, this fatty acid is a poorer substrate for COX than EPA or DHA. The action of DHA on interrenal cells, whether it is direct or mediated by its lipoygenase derivatives, still has to be elucidated since lipoygenase metabolites have been shown to modify the hormone-induced release of cortisol in mammal adrenal tissues (Wang *et al.* 2000, Yamazaki *et al.* 2001). Using nordihydroguaiaretic acid, a lipoygenase inhibitor, cortisol secretion was inhibited in response to ACTH in bovine adrenocortical cells (Wang *et al.* 2000).

DHA has long been known for its high value as an essential fatty acid for marine fish (Watanabe 1982), particularly during larval stages (Izquierdo *et al.* 1989) when it invariably promotes growth, survival and stress resistance to a higher extent than EPA or ARA in all the studied species (Watanabe *et al.* 1989, Kanazawa 1997, Rodríguez *et al.* 1997, Furuita *et al.* 1999, Izquierdo *et al.* 2005). In gilthead sea bream, dietary deficiencies of n-3 HUFA and especially DHA have been shown to increase plasma cortisol levels after both acute (net chasing) and chronic (high stocking density) stress (Montero *et al.* 1998, 2001). Besides, imbalances in the dietary n-3/n-6 fatty acids ratio induced by the inclusion of vegetable oils in the diet have been shown to alter the release of cortisol after stress in this species (Montero *et al.* 2003) and in other species such as chinook salmon (Welker & Congleton 2003). The role of dietary oils on stress response in fish remains unclear, but results indicate that dietary fatty acids could be regulating the *in vivo* stress response through the mechanisms discussed above. Moreover, vegetable oils in fish diets have been shown to regulate COX-derived eicosanoids directly (Ganga *et al.* 2005). Dietary supplementation of other fatty acids such as ARA seems to be affecting plasma cortisol levels after stress (Van Anholt *et al.* 2004), although the effect on cortisol release *in vivo* is dose dependent, since high levels of ARA in diet seem to be detrimental to chronic stress resistance in larval gilthead sea bream (Koven *et al.* 2003).

Concentrations of fatty acids used in the present study were those providing the maximum cortisol stimulation factor (50 μ M). However, higher concentrations reduced and even inhibited cortisol secretion (Acerete L, Ganga R, Tort L & Izquierdo MS. unpublished results), suggesting a concentration dependency in the type of effect of these fatty acids. This is the case in other organs such as testicles where medium concentrations of ARA (3–30 μ M) induced testosterone production in testicular cells of male sea bass, whereas high concentrations (300 μ M) inhibited it (Asturiano 1999). Despite the fact that no previous data has been published on *in vitro* exposures of fish interrenal cells to fatty acids, cytotoxic

effects at the membrane level have been found in mammalian tissues, including renal cells (Zager *et al.* 1997). Particularly, excess of ARA and its derived eicosanoids has been found to cause apoptosis associated with oxidative stress in human leukocytes (Pompeia *et al.* 2002). Studies are being conducted at present to elucidate the effect of low and high physiological concentrations of fatty acids on ACTH-induced cortisol release by fish interrenal cells. Although this superfusion method is widely used for studies of interrenal tissue in fish (Rotllant *et al.* 2001), it prevents the possible control of blood flow by COX-derived prostanoids. Nevertheless, new results have shown the vasoregulatory function of COX-derived products in fish (Stenslkken *et al.* 2002), none of them relate though to interrenal tissues.

In summary, the role of ARA and EPA in the ACTH-induced release of cortisol from gilthead sea bream interrenal cells seems to be partly related to COX-derived metabolites dependent on ARA and EPA, whereas the role of DHA seems to be dependent on other factors. Both EPA and ARA as single supplemented fatty acids increased the ACTH-induced cortisol release from gilthead sea bream interrenal cells whereas the effect of DHA was weaker. A well-balanced supplementation of these three fatty acids could be necessary to regulate cortisol release from interrenal cells in gilthead sea bream.

Acknowledgements

The authors want to acknowledge the provision of fish by the Granja Marina del Masnou. This study was supported by grants AGL-2003-03112 (Ministry of Education) and Centre de Referencia en Aqicultura (Generalitat Catalunya). The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Asturiano JF 1999 El proceso reproductivo de la lubina europea (*Dicentrarchus labrax* L.), Efectos de los ácidos grasos de la dieta: estudios *in vivo* e *in vitro* PhD. Thesis. Valencia University, Spain, p. 251.
- Balfry SK, Maule AG & Iwama GK 2001 Coho salmon *Onchorhynchus kisutch* strain differences in disease resistance and non-specific immunity, following immersion challenges with *Vibrio anguillarum*. *Diseases of Aquatic Organisms* **47** 39–48.
- Bell JG, McVicar AH, Park MT & Sargent JR 1991 High dietary linoleic acid affects the fatty acid compositions of individual phospholipids from tissues of Atlantic salmon (*Salmo salar*): association with stress susceptibility and cardiac lesion. *Journal of Nutrition* **121** 1163–1211.
- Bell JG, Tocher DR & Sargent JR 1994a Effect of supplementation with 20:3(n-6), 20:4(n-6) and 20:5(n-3) on the production of prostaglandins E and F of the 1-, 2- and 3-series in turbot (*Scophthalmus maximus*) brain astroglial cells in primary culture. *Biochimica et Biophysica Acta* **1211** 335–342.
- Bell JG, Tocher DR, MacDonald FM & Sargent JR 1994b Effects of diets rich in linoleic (18:2n-6) and α -linolenic acids on the growth, lipid class and fatty acid compositions and eicosanoid production in juvenile turbot (*Scophthalmus maximus* L.). *Fish Physiology and Biochemistry* **13** 105–118.
- Bell JG, Tocher DR, Farndale BM & Sargent JR 1998 Growth, mortality, tissue histopathology and fatty acid composition, eicosanoid production and response to stress, in juvenile turbot fed diets rich in γ -linolenic acid in combination with eicosapentaenoic acid or docosahexaenoic acid. *Prostaglandins, Leukotrienes and Essential Fatty Acids* **58** 353–364.
- Bessonart M, Izquierdo MS, Salhi M, Hernandez-Cruz CM, Gonzalez MM & Fernandez-Palacios H 1999 Effect of dietary arachidonic acid levels on growth and survival of gilthead sea bream (*Sparus aurata* L.) larvae. *Aquaculture* **179** 265–275.
- Furuuta H, Takeuchi T & Uematsu K 1998 Effects of eicosapentaenoic and docosahexaenoic acids on growth, survival and brain development of larval Japanese flounder (*Paralichthys olivaceus*). *Aquaculture* **161** 59–69.
- Furuuta H, Kooichi K & Takeuchi T 1999 Effect of different levels of eicosapentaenoic acid and docosahexaenoic acid in *Artemia* nauplii on growth, survival and salinity tolerance of larvae of the Japanese flounder, *Paralichthys olivaceus*. *Aquaculture* **170** 59–69.
- Ganga R, Bell JG, Montero D, Robaina L, Caballero MJ & Izquierdo MS 2005 Effect of dietary lipids on plasma fatty acid profiles and prostaglandin and leptin production in gilthead seabream (*Sparus aurata*). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* **142** 410–418.
- German B, Bruckner G & Kinsella J 1983 Evidence against a PGF₄ prostaglandin structure in trout tissue—a correction. *Prostaglandins* **6** 207–210.
- Gobbetti A & Zerani M 1993 Prostaglandin E₂ and prostaglandin F₂ alpha involvement in the corticosterone and cortisol release by the female frog, *Rana esculenta*, during ovulation. *Journal of Experimental Zoology* **267** 164–170.
- Gupta OP, Lahlou B, Botella J & Porthé-Nibelle J 1985 *In vivo* and *in vitro* studies on the release of cortisol from interrenal tissue in trout. I. Effects of ACTH and prostaglandins. *Experimental Biology* **43** 201–212.
- Harel M, Gavasso S, Leshin J, Gubernatis A & Place AR 2001 The effect of tissue docosahexaenoic and arachidonic acids levels on hypersaline tolerance and leucocyte composition in striped bass (*Morone saxatilis*) larvae. *Fish Physiology and Biochemistry* **24** 113–123.
- Henderson RJ & Tocher DR 1987 The lipid composition and biochemistry of freshwater fish. *Progress in Lipid Research* **26** 281–347.
- Hong S, Tjonahen E, Morgan EL, Lu Y, Serhan CN & Rowley AF 2005 Rainbow trout (*Oncorhynchus mykiss*) brain cells biosynthesize novel docosahexaenoic acid-derived resolvins and protectins—mediator lipidomic analysis. *Prostaglandins, other lipid mediators* **78** 107–116.
- Hontela A 1997 Endocrine and physiological responses of fish xenobiotics: Role of glucocorticosteroid hormones. In *Reviews in Toxicology. Environmental Toxicology*, pp 1–46. The Netherlands: ISO Press.
- Horrobin DF 1983 The regulation of prostaglandin biosynthesis by the manipulation of essential fatty acid metabolism. *Revision of Pure Applied Science* **4** 339–383.
- Izquierdo MS 1996 Essential fatty acid requirements of cultured marine fish larvae. *Aquaculture Nutrition* **2** 183–191.
- Izquierdo MS, Watanabe T, Takeuchi T, Arakawa T & Kitajima C 1989 Requirement of larval red seabream *Pagrus major* for essential fatty acids (Nippon Suisan Gakkaishi). *Bulletin of Japan Social Science Fish* **55** 859–867.
- Izquierdo MS, Tandler A, Salhi M & Kolkovski S 2001 Influence of dietary polar lipids quantity and quality on ingestion and assimilation of labelled fatty acids by larval gilthead sea bream. *Aquaculture Nutrition* **6** 153–160.
- Izquierdo MS, Montero D, Robaina L, Caballero MJ, Rosenlund G & Ginés R 2005 Alterations in fillet fatty acid profile and flesh quality in gilthead seabream (*Sparus aurata*) fed vegetable oils for a long term period, recovery of fatty acid profiles by fish oil feeding. *Aquaculture* **250** 431–444.
- Kanazawa A 1997 Effects of docosahexaenoic acid and phospholipids on stress tolerance of fish. *Aquaculture* **155** 129–134.
- Kocsis JF, Rinkardt NE, Satterlee DG, Weber H & Carsia RV 1999 Concentration-dependent, biphasic effect of prostaglandins on avian corticosteroidogenesis *in vitro*. *General and Comparative Endocrinology* **115** 132–142.
- Koven WM, Van Anholt RD, Lutzky S, Ben Atia I, Nixon O, Ron B & Tandler A 2003 The effect of dietary arachidonic acid on growth, survival,

- and cortisol levels in different-age gilthead seabream larvae (*Sparus auratus*) exposed to handling or daily salinity change. *Aquaculture* **228** 307–320.
- Lands WEM 1989 Differences in n-3 and n-6 eicosanoid precursors. In *Advances in Prostaglandin Thromboxane and Leukotriene Research*, Vol 9, pp 602–605. Eds B Samuelsson & PYK Wong. New York, NY, USA: Raven Press.
- Liu J, Caballero MJ, El-Sayed Ali T, Izquierdo MS, Hernández Cruz CM, Valencia A & Fernández-Palacios H 2002 Necessity of dietary lecithin and eicosapentaenoic acid for growth, survival, stress resistance and lipoprotein formation in gilthead sea bream (*Sparus aurata*). *Fish Science* **68** 1165–1172.
- Matsumoto H, Iijima N & Kayama M 1989 The prostaglandin synthesis in marine fish thrombocytes. *Comparative Biochemistry and Physiology* **93B** 397–402.
- Montero D, Tort L, Izquierdo MS, Robaina L & Vergara JM 1998 Depletion of serum alternative complement pathway activity in gilthead seabream caused by alpha-tocopherol and n-3 HUFA dietary deficiencies. *Fish Physiology and Biochemistry* **18** 399–407.
- Montero D, Tort L, Robaina L, Vergara JM & Izquierdo MS 2001 Low vitamin E in diet reduces stress resistance of gilthead seabream (*Sparus aurata*) juveniles. *Fish and Shellfish Immunology* **11** 473–490.
- Montero D, Kalinowski T, Obach A, Robaina L, Tort L, Caballero MJ & Izquierdo MS 2003 Vegetable lipid sources for gilthead seabream (*Sparus aurata*): effects on fish health. *Aquaculture* **225** 353–370.
- Mustafa T & Srivastava KC 1989 Prostaglandins (eicosanoids) and their role in ectothermic organisms. *Advance Comparative Environmental Physiology* **5** 157–207.
- Nablon G, Grynberg A, Chevalier A, Leonardi J, Termine E & Lafont H 1990 Phospholipase A activity of cultures rat ventricular myocytes is affected by the nature of cellular polyunsaturated fatty acids. *Lipids* **25** 301–306.
- Pompeia C, Freitas JJS, Kim JS, Zyngier SB & Curi R 2002 Arachidonic acid cytotoxicity in leukocytes: implications of oxidative stress and eicosanoids synthesis. *Biology of Cells* **94** 251–265.
- Rodríguez C, Perez JA, Diaz M, Izquierdo MS, Fernandez-Palacios H & Lorenzo A 1997 Influence of the EPA/DHA ratio in rotifers on gilthead seabream (*Sparus aurata*) larval development. *Aquaculture* **150** 77–89.
- Rotllant J & Tort L 1997 Cortisol and glucose responses after acute stress by net handling in the spard red porgy previously subjected to crowding stress. *Journal of Fish Biology* **51** 21–28.
- Rotllant J, Balm PHM, Ruane NM, Pérez-Sánchez J, Wendelaar Bonga SE & Tort L 2000a Pituitary proopiomelanocortin-derived peptides and hypothalamic-pituitary-interrenal axis activity in gilthead sea bream (*Sparus aurata*) during prolonged crowding stress: differential regulation of adrenocorticotropin hormone and melanocyte-stimulating hormone release by corticotrophin-releasing hormone and thyrotropin-releasing hormone. *General and Comparative Endocrinology* **119** 152–163.
- Rotllant J, Balm PHM, Wendelaar Bonga SE, Pérez-Sánchez J & Tort L 2000b A drop in ambient temperature results in a transient reduction of interrenal ACTH responsiveness in the gilthead sea bream (*Sparus aurata* L.). *Fish Physiology and Biochemistry* **23** 265–273.
- Rotllant J, Balm PHM, Pérez-Sánchez J, Wendelaar Bonga SE & Tort L 2001 Pituitary and interrenal function in gilthead sea bream (*Sparus aurata* L., Teleostei) after handling and confinement stress. *General and Comparative Endocrinology* **121** 333–342.
- Sokal RR & Rolf SJ 1995 Biometry. In *The Principles and Practice of Statistics in Biological Research*. 3rd edn, p 419. New York, NY, USA: WH Freeman and Company.
- Stenslokken KO, Sundin L & Nilsson GE 2002 Cardiovascular effects of prostaglandin F_{2α} and prostaglandin E₂ in Atlantic cod (*Gadus morhua*). *Journal of Comparative Physiology B Biochemical, Systemic, and Environmental Physiology* **172** 363–369.
- Thompson I, White A, Fletcher TC, Houlihan DF & Secombes CJ 1993 The effect of stress on the immune system of Atlantic salmon (*Salmo salar* L.) fed diets containing different amounts of vitamin C. *Aquaculture* **114** 1–18.
- Tocher DR 1995 Glycerophospholipid metabolism. In *Biochemistry and Molecular Biology of Fishes*, vol. 4, *Metabolic and Adaptational Biochemistry*, pp 119–1957. Eds W Hochachka & TP Mommsen. Amsterdam: Elsevier Press.
- Tocher DR & Sargent JR 1987 The effects of calcium ionophore A23187 on the metabolism of arachidonic and eicosapentaenoic acids in neutrophils a marine teleost fish rich in (n-3) polyunsaturated fatty acids. *Comparative Biochemistry and Physiology* **87B** 733–739.
- Uhing RJ, Cowlen MS & Adams DO 1990 Mechanisms regulating the production of arachidonate metabolites in mononuclear phagocytes. *Current Topics in Membranes Transport* **35** 349–374.
- Vakharia K & Hinson JP 2005 Lipopolysaccharide directly stimulates cortisol secretion by human adrenal cells by a cyclooxygenase-dependent mechanism. *Endocrinology* **146** 1398–1402.
- Van Anholt RD, Spanings FAT, Koven WM & Wendelaar Bonga SE 2004 Dietary supplementation with arachidonic acid in tilapia (*Oreochromis mossambicus*) reveals physiological effects not mediated by prostaglandins. *General and Comparative Endocrinology* **139** 215–226.
- Wang H, Walker SW, Mason JI, Morley SD & Williams BC 2000 Role of arachidonic acid metabolism in ACTH-stimulated cortisol secretion by bovine adrenocortical cells. *Endocrine Research* **26** 705–709.
- Watanabe T 1982 Lipid nutrition in fish. *Comparative Biochemistry and Physiology* **73B** 3–15.
- Watanabe T, Izquierdo MS, Takeuchi T, Satoh S & Kitajima C 1989 Comparison between eicosapentaenoic and docosahexaenoic acids in terms of essential fatty acid efficacy in larval red seabream (*Nippon Suisan Gakkaishi*). *Bulletin Japan Social Science Fish* **55** 1635–1640.
- Wendelaar Bonga SE 1997 The stress response in fish. *Physiology Review* **77** 591–625.
- Welker TL & Congleton JL 2003 Relationship between dietary lipid source, oxidative stress, and physiological response to stress in sub-yearling Chinook salmon (*Oncorhynchus tshawytscha*). *Fish Physiology and Biochemistry* **29** 225–235.
- Yamazaki T, Higuchi K, Kominami S & Takemori S 2001 15-Lipoxygenase metabolite(s) of arachidonic acid mediates adrenocorticotropin action in bovine adrenal steroidogenesis. *Endocrinology* **137** 2670–2675.
- Yin Z, Lam TJ & Sin YM 1995 The effects of crowding stress on the non-specific immune response in fancy carp (*Cyprinus carpio* L.). *Fish and Shellfish Immunology* **5** 519–529.
- Zager RA, Iwata M, Conrad DS, Burkhart KM & Igarashi Y 1997 Altered ceramide and sphingosine expression during the induction phase ischemic acute renal failure. *Kidney International* **52** 60–70.

Received in final form 23 March 2006

Accepted 29 March 2006

Made available online as an Accepted Preprint
21 April 2006