Peptide YY (PYY)_{3–36} modulates thyrotropin secretion in rats

K J Oliveira, G S M Paula, R H Costa-e-Sousa, L L Souza, D C Moraes, F H Curty and C C Pazos-Moura

Laboratório de Endocrinologia Molecular, Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro 21949-900, Brazil
(Requests for offprints should be addressed to C C Pazos-Moura; Email: cpazosm@biof.ufrj.br)

Abstract

Peptide YY (PYY)_{3–36} is a gut-derived hormone, with a proposed role in central mediation of postprandial satiety signals, as well as in long-term energy balance. In addition, recently, the ability of the hormone to regulate gonadotropin secretion, acting at pituitary and at hypothalamus has been reported. Here, we examined PYY_{3–36} effects on thyrotropin (TSH) secretion, both *in vitro* and *in vivo*. PYY_{3–36}-incubated rat pituitary glands showed a dose-dependent decrease in TSH release, with 44 and 62% reduction at 10^{-8} and 10^{-6} M ($P < 0.05$ and $P < 0.001$ respectively), and no alteration in TSH response to thyrotropin-releasing hormone. *In vivo*, PYY_{3–36} i.p. single injection in the doses of 3 or 30 µg/kg body weight, administered to rats fed *ad libitum*, was not able to change serum TSH after 15 or 30 min.

However, in fasted rats, PYY_{3–36} at both doses elicited a significant rise (approximately twofold increase, $P < 0.05$) in serum TSH observed 15 min after the hormone injection. PYY_{3–36} treatment did not modify significantly serum T₄, T₃, or leptin. Therefore, in the present paper, we have demonstrated that the gut hormone PYY_{3–36} acts directly on the pituitary gland to inhibit TSH release, and in the fasting situation, *in vivo*, when serum PYY_{3–36} is reduced, the activity of thyroid axis is reduced as well. In such a situation, systemically injected PYY_{3–36} was able to acutely activate the thyrotrope axis, suggesting a new role for PYY_{3–36} as a regulator of the hypothalamic–pituitary–thyroid axis.

Journal of Endocrinology (2006) **191**, 459–463

Introduction

Peptide YY (PYY)_{3–36} is a gut-derived hormone, produced by endocrine L cells lining the distal small bowel and colon, released postprandially in proportion to the calories ingested (Adrian *et al.* 1985). Peripheral administration of PYY_{3–36} in rodents or humans has been reported to induce a marked inhibition of food intake (Batterham *et al.* 2002, 2003). PYY_{3–36} inhibits food intake at plasma concentrations that are within the normal physiological range seen in man after meals (Batterham *et al.* 2002, 2003) and therefore, a role for the hormone in central mediation of postprandial satiety signals has been proposed. Moreover, long-term administration has been demonstrated to induce a reduction in body weight (Batterham *et al.* 2002, 2003) although this is still a matter of controversy (Tschöp *et al.* 2004).

In addition, recent studies suggested that PYY_{3–36} may act as a neuroendocrine regulator. It has been demonstrated in prepubertal rats that PYY_{3–36} stimulated prolactin, luteinizing hormone (LH), and FSH secretion acting directly at pituitary gland, although it seems to have an inhibitory action at the hypothalamic level (Aguilar *et al.* 2004, Fernandez-Fernandez *et al.* 2005). Therefore, similar to other hormones involved in energy homeostasis, PYY_{3–36} is also able to influence the reproductive axis. However, there is no information concerning the thyrotrope axis.

PYY_{3–36} is a member of the neuropeptide Y (NPY) family and it is an agonist of receptor subtypes NPY-Y2 and NPY-Y5

(Keire *et al.* 2000). Experimental evidence suggests that circulating PYY_{3–36} inhibits appetite by acting directly on the arcuate nucleus via the Y2 receptor, a presynaptic inhibitory autoreceptor (Batterham *et al.* 2002, Talsania *et al.* 2005). NPY is a potent stimulator of food intake and PYY_{3–36} was able to inhibit the electrical activity of arcuate NPY neurons as well as to reduce the expression of the NPY mRNA (Batterham *et al.* 2002, Challisa *et al.* 2003). In addition, NPY originating in the hypothalamic arcuate nucleus exerts a profound inhibitory effect on the thyroid axis via effects on hypophysiotropic thyrotropin-releasing hormone (TRH) neurons. Chronic intracerebroventricular administration of NPY to normally fed rats resulted in reduction of circulating levels of thyroid hormones with inappropriately normal or low thyrotropin (TSH), and suppression of proTRH mRNA in the hypothalamic paraventricular nucleus (PVN; Fekete *et al.* 2001). NPY effects were reproduced by NPY-Y1 and NPY-Y5 analogs injected into the third cerebral ventricle, suggesting that both the receptors mediate NPY suppression of the hypothalamic–pituitary–thyroid axis (Fekete *et al.* 2002).

During fasting, as a means of conserving energy, the hypothalamic–pituitary–thyroid axis is suppressed and the activation of arcuate NPY neurons is a major component of the regulatory mechanism that causes a decline in proTRH mRNA in PVN, reducing circulating levels of TSH and thyroid hormones. Fasting is associated with reduced serum concentrations of PYY_{3–36} (Tovar *et al.* 2004, Chan *et al.* 2005), and peripherally injected PYY_{3–36} partly reversed the

fasting-induced c-Fos expression in arcuate nucleus neurons of mice (Riediger *et al.* 2004), suggesting a role for the peptide in fasting adaptation.

Therefore, considering NPY effects on thyroid axis, and the presence of Y5 receptors, and the mRNA encoding Y2 and Y5 receptors in pituitary gland (Parker *et al.* 2000, Fernandez-Fernandez *et al.* 2005), we tested the hypothesis that PYY₃₋₃₆ may act directly at the pituitary to modulate TSH secretion. In addition, we also investigated whether PYY₃₋₃₆, injected systemically, may modify the thyrotrope axis in fed and fasting rats.

Materials and Methods

Animals

Adult male Wistar rats, weighing 250–300 g, were kept under controlled lighting (12 h light:12 h darkness cycle, lights on at 0700 h) and controlled temperature (23 ± 1 °C). All experimental protocols were approved by our institutional animal care committee.

In vitro experiments

Ad libitum fed rats were killed by decapitation, and their anterior pituitaries were quickly dissected out for *in vitro* testing as described before (Rettori *et al.* 1989, Ortiga-Carvalho *et al.* 2002). Each hemi-pituitary was immediately transferred to a flask containing 1 ml Krebs–Ringer bicarbonate medium (pH 7.4) at 37 °C in an atmosphere of 95% O₂/5% CO₂ in a Dubnoff metabolic shaker. After a 30-min preincubation period, medium was changed to 1 ml medium alone (control) or medium containing PYY₃₋₃₆ (Bachem California, Inc., Torrance, CA, USA) to a final concentration of 10^{-10} , 10^{-8} , or 10^{-6} M. At the end of a 2 h incubation period, an aliquot was removed for TSH measurement. In another set of experiments, after 2 h incubation, a small aliquot was removed for TSH measurement and TRH (Bachem California, Inc.) was added to a final concentration of 50 nM in all tubes. The incubation was continued for 30 min to determine the TRH-stimulated TSH release in the absence or presence of the different concentrations of PYY₃₋₃₆.

In vivo experiments

Ad libitum fed rats were divided into three groups that received a single i.p. injection of 3 or 30 µg/kg body weight (BW) PYY₃₋₃₆ or 0.2 ml saline vehicle (control group). Another set of rats was fasted for 3 days before they received a single i.p. injection of 3 or 30 µg/kg BW PYY₃₋₃₆ or 0.2 ml saline vehicle (control group). In both experiments, rats were killed by decapitation, 15 or 30 min after the injection. Experiments were performed in the morning between 1000 and 1100 h. Serum was obtained from trunk blood for hormone measurements.

Quantification of TSH

TSH concentration in the serum and in the incubation medium was measured by specific RIA, employing reagents supplied by the National Institute of Diabetes, Digestive and Kidney Diseases, National Hormone & Peptide Program (NIDDK–NHPP) (Torrance, CA, USA), as previously described (Chard 1987, Ortiga-Carvalho *et al.* 1996), and was expressed in terms of the reference preparation 3 (RP3). Within-assay variation was 5.7%. Samples of the same experiment were measured within the same assay. Minimum assay detection was 0.52 ng/ml.

Quantification of serum concentrations of T₄, T₃, and leptin

Serum T₄ and T₃ were detected by RIA (MP Biomedicals Inc, Irvine, CA, USA). Detection limits were: 1 µg/dl for T₄ and 25 ng/dl for T₃. Serum leptin was measured using a specific rat RIA by LINCO Research (St Charles, MO, USA). Minimum detectable level was 0.5 ng/ml. Intraassay variation was less than 7% for all hormone measurements and all the samples were run within the same assay.

Statistical analysis

Data are reported as means $\pm$ S.E.M. One-way ANOVA followed by a Student–Newman–Keuls multiple comparisons test was employed for the assessment of significance of data. Serum TSH was analyzed after logarithmic transformation (Zar 1996). Differences were considered to be significant at $P < 0.05$.

Results

In vitro experiments

PYY₃₋₃₆-incubated hemi-pituitary glands showed, after 2 h incubation, a dose-dependent decrease in TSH release, statistically significant at 10^{-8} M ($P < 0.05$) and 10^{-6} M ($P < 0.001$), with a reduction of 44 and 62% respectively (Fig. 1). However, the TSH response to TRH was not significantly different among groups, presenting an increment around four times above the basal levels (TSH before TRH) in all groups.

In vivo experiments

The i.p. administration of 3 or 30 µg/kg BW of PYY₃₋₃₆ to *ad libitum* fed rats induced no changes in serum TSH either after 15 or 30 min (Fig. 2). Also, serum leptin and serum T₄ (Table 1) and T₃ concentrations (data not shown) were not significantly modified by the treatment with PYY₃₋₃₆. However, in fasted rats (Fig. 3), a significant rise in serum TSH was detected 15 min after PYY₃₋₃₆ administration at both doses (approximately twofold increase, $P < 0.05$). This

Figure 1 TSH release from isolated rat hemi-anterior pituitary glands incubated in absence (C, control) or presence of PYY₃₋₃₆ at different concentrations (10^{-10} , 10^{-8} , 10^{-6} M) for 2 h. $n=10$ hemi-pituitaries per group. * $P<0.05$, † $P<0.001$ vs control group. Data represent means $\pm$ S.E.M. Figure representative of two independent experiments.

effect was transitory, since no statistically significant effect was observed 30 min after PYY₃₋₃₆ injection of both doses in fasted rats. Serum concentrations of T₄ and leptin were not changed by PYY₃₋₃₆ treatment at any time point or doses (Table 1). Fasted rats presented approximately 11% reduction in body weight and in most of them serum T₃ was below 25 ng/ml.

Discussion

Here, we first demonstrated that PYY₃₋₃₆ acts directly at the pituitary to inhibit TSH secretion. Y2 and Y5 mRNA receptors had been detected at low levels in pituitary gland (Parker *et al.* 2000, Fernandez-Fernandez *et al.* 2005), however, binding studies indicated the presence of Y5 (Parker *et al.* 2000) and the absence of Y2 receptors (Sheikh *et al.* 1998). Therefore, it is probable that PYY₃₋₃₆ elicited the inhibitory effect on TSH secretion acting through activation of pituitary Y5 receptors. This would be consistent with the fact that an Y5 analog, injected centrally in rats, was able to suppress the hypothalamus–pituitary–thyroid axis. In addition, the present data are in agreement with a previous study showing that NPY, also a Y5 ligand, acts directly at rat pituitary to reduce β -TSH mRNA levels, as demonstrated by the incubation of pituitary cell cultures with NPY

Figure 2 Serum TSH concentration in fed rats after intraperitoneal administration of PYY₃₋₃₆ 3 (A) and 30 (B) µg/kg BW or saline (C). Determinations were performed at 15 and 30 min after injection of the peptide. $n=9-10$ animals per group. Data are presented as means $\pm$ S.E.M.

(Chowdhury *et al.* 2004). However, TRH action at the thyrotropes seems to be preserved, since the fold increment on TSH release induced by TRH was not significantly modified by the presence of PYY₃₋₃₆ in the incubation medium.

Contrary to what could be expected in view of the inhibitory pituitary action of PYY₃₋₃₆ on TSH release, systemic i.p. administration of the hormone in normally fed rats (3 and 30 µg/kg) failed to significantly modify serum TSH levels. Previous work of Fernandez-Fernandez *et al.* (2005), using a similar PYY₃₋₃₆ administration protocol, was also unable to demonstrate *in vivo* effects of PYY₃₋₃₆ on LH secretion, although the peptide had stimulated LH release from incubated rat pituitaries. Similar findings were reported concerning PYY₃₋₃₆ effects on PRL secretion (Aguilar *et al.* 2004). The doses of PYY₃₋₃₆ that we employed can be considered low and moderate, and it is possible that a higher dose would be necessary to modify TSH secretion. However, as demonstrated in previous studies (Batterham *et al.* 2002, Challisa *et al.* 2003, Talsania *et al.* 2005), similar PYY₃₋₃₆ doses were sufficient to inhibit food intake in rats, suggesting that these doses were enough to activate arcuate nucleus Y2 receptors. An alternative explanation could be that the final effect of injected PYY₃₋₃₆ may be the result of its action at different targets. In this sense, it must be considered that systemic PYY₃₋₃₆ rapidly reaches the arcuate nucleus, an area partially outside the blood–brain barrier, where it binds with high affinity to autoinhibitory Y2 receptors in NPY neurons,

Table 1 Serum concentration of T₄ and leptin in fed and fasting rats after intraperitoneal administration of peptide YY (PYY)₃₋₃₆. Data are presented as means $\pm$ S.E.M. of two separated experiments (fed and fasting rats). $n=7-10$ per group.

		Control	PYY 3 µg 15 min	PYY 3 µg 30 min	Control	PYY 30 µg 15 min	PYY 30 µg 30 min
Group							
T ₄ µg/dl	Fed	3.71 $\pm$ 0.28	3.88 $\pm$ 0.15	4.02 $\pm$ 0.19	3.71 $\pm$ 0.31	4.02 $\pm$ 0.22	4.36 $\pm$ 0.40
T ₄ µg/dl	Fasted	2.86 $\pm$ 0.28	2.62 $\pm$ 0.10	2.18 $\pm$ 0.05	2.76 $\pm$ 0.18	2.67 $\pm$ 0.30	2.37 $\pm$ 0.19
Leptin ng/ml	Fed	2.64 $\pm$ 0.25	2.40 $\pm$ 0.16	3.03 $\pm$ 0.41	3.91 $\pm$ 0.30	3.56 $\pm$ 0.44	3.80 $\pm$ 0.32
Leptin ng/ml	Fasted	1.27 $\pm$ 0.20	1.17 $\pm$ 0.20	0.84 $\pm$ 0.17	0.97 $\pm$ 0.14	0.76 $\pm$ 0.07	0.78 $\pm$ 0.09

Figure 3 Serum TSH concentrations in fasted rats after intraperitoneal administration of PYY₃₋₃₆ 3 (A) and 30 (B) µg/kg BW or saline (C). Determinations were performed at 15 and 30 min after injection of the peptide. $n=8-10$ animals per group. * $P<0.05$, † $P<0.001$ vs C (control vehicle-injected). Data are presented as means $\pm$ S.E.M.

which is the main postulated mechanism for its anorexigenic effect (Batterham *et al.* 2002). Therefore, by reducing NPY neuron activity, PYY₃₋₃₆ potentially may decrease NPY inhibitory input into TRH neurons. Thereby, it is possible that PYY₃₋₃₆ systemic administration had induced a higher TRH secretion that was counteracted by the direct pituitary inhibitory action, resulting in no alteration in serum TSH.

However, in fasted rats, it is likely that inhibition of NPY neurons via arcuate Y2 receptors is the predominant mechanism leading to the increase in serum TSH. In fasting conditions, there is a higher NPY inhibitory tonus on TRH neurons (Fekete *et al.* 2001) and, in addition, fasting was associated with reduced PYY₃₋₃₆ serum levels, as reported by others (Tovar *et al.* 2004, Chan *et al.* 2005), which may render the tissues more sensitive to PYY₃₋₃₆ action. Other reports support the concept that fasting enhances PYY₃₋₃₆ effects, such as those concerning the peptide action on gonadotropin secretion (Pinilla *et al.* 2006) and on food intake (Challisa *et al.* 2003, Riediger *et al.* 2004).

PYY₃₋₃₆ effect on TSH release was observed 15 min after the hormone administration, coincidentally with previous reported time for peak PYY₃₋₃₆ serum levels after a single i.p. injection in the dose of 3 µg/kg BW (Batterham *et al.* 2002). This fast action is consistent with a central action of PYY₃₋₃₆ through activation of Y2 receptors in arcuate nucleus, since these receptors have high affinity for PYY₃₋₃₆. However, the effect was transitory, which may reflect the short-life of the peptide, and the fact that a single dose would not be enough to normalize the reduced levels of PYY₃₋₃₆ in fasting rats. However, it cannot be excluded that the activation of inhibitory Y5 receptors pathways in the pituitary may counteract the arcuate nucleus Y2-mediated stimulus in thyrotrope axis.

As expected, fasted rats exhibited lower serum thyroid hormone levels and leptin; however, the effect of PYY₃₋₃₆ on TSH release could not be attributed to changes in these hormone levels, since they remained similar among groups.

Reduction in serum leptin during fasting is believed to have a major role as a peripheral signal of energy insufficiency, resulting

in an integrated response at the central nervous system which includes the activation of NPY-producing neurons in the hypothalamic arcuate nucleus. Considering that peripherally injected PYY₃₋₃₆ partly reversed the fasting-induced c-Fos expression in arcuate nucleus neurons of mice (Riediger *et al.* 2004), it is possible that reduction in serum PYY₃₋₃₆ also plays a role in neuroendocrine adaptation to fasting, which is an important component in the suppression of thyrotrope axis, a hypothesis that remains to be fully explored.

In conclusion, in the present paper we have demonstrated that the gut hormone PYY₃₋₃₆ acts directly at the pituitary gland to inhibit TSH release and that *in vivo*, systemically injected PYY₃₋₃₆ is able to acutely activate the thyrotrope axis during fasting. Therefore, we propose a new role for PYY₃₋₃₆ as a regulator of the hypothalamic–pituitary–thyroid axis.

Funding

This work was supported by Conselho Nacional de Pesquisa e Desenvolvimento and Fundação Carlos Chagas Filho de Amparo a Pesquisa do Estado do Rio de Janeiro. There is no conflict of interest that would prejudice its impartiality.

References

- Adrian TE, Ferri GL, Bacarese-Hamilton AJ, Fuessl HS, Polak JM & Bloom SR 1985 Human distribution and release of a putative new gut hormone, peptide YY. *Gastroenterology* **89** 1070–1077.
- Aguilar E, Fernandez-Fernandez R, Tena-Sempere M & Panilla L 2004 Effects of Peptide YY(3-36) on PRL secretion: pituitary and extra-pituitary actions in the rat. *Peptides* **25** 1147–1152.
- Batterham RL, Cowley MA, Small CJ, Herzog H, Cohen MA, Dakin CL, Wren AM, Brynes AE, Low MJ, Ghatge MA *et al.* 2002 Gut hormone PYY(3-36) physiologically inhibits food intake. *Nature* **418** 650–654.
- Batterham RL, Cohen MA, Ellis SM, Le Roux CW, Withers DJ, Frost GS, Ghatge MA & Bloom SR 2003 Inhibition of food intake in obese subjects by peptide YY3-36. *New England Journal of Medicine* **349** 941–948.
- Challisa BG, Pinnock SB, Colla AP, Carter RN, Dickson SL & O'Rahilly S 2003 Acute effects of PYY₃₋₃₆ on food intake and hypothalamic neuropeptide expression in the mouse. *Biochemical and Biophysical Research Communications* **311** 915–919.
- Chan JL, Stoyneva V, Kelesidis T, Raciti P & Mantzoros CS 2005 Peptide YY levels are decreased by fasting and elevated following caloric intake but are not regulated by leptin. *Diabetologia* **49** 169–173.
- Chard T 1987 An introduction to radioimmunoassay and related techniques. In *Laboratory Techniques in Biochemistry and Molecular Biology*, pp 1–74. Eds RH Burdon & PH van Kinsioenberg. Amsterdam: Elsevier Science.
- Chowdhury I, Chien JT, Chatterjee A & Yu JYL 2004 Effects of leptin and neuropeptide-Y on transcript levels of thyrotropin beta and common alpha subunits of rat pituitary cells *in vitro*. *Life Science* **75** 2897–2909.
- Fekete C, Kelly J, Mihály E, Sarkar S, Rand WM, Légrádi G, Emerson CH & Lechan RM 2001 Neuropeptide Y has a central inhibitory action on the hypothalamic–pituitary–thyroid axis. *Endocrinology* **142** 2606–2613.
- Fekete C, Sarkar S, Rand WM, Harney JW, Emerson CH, Bianco AC, Beck-Sickingner A & Lechan RM 2002 Neuropeptide Y1 and Y5 receptors mediate the effects of neuropeptide Y on the hypothalamic–pituitary–thyroid axis. *Endocrinology* **143** 4513–4519.

- Fernandez-Fernandez R, Aguilar E, Tena-Sempere M & Pinilla L 2005 Effects of polypeptide YY₃₋₃₆ upon luteinizing hormone-releasing hormone and gonadotropin secretion in prepubertal rats: *in vivo* and *in vitro* studies. *Endocrinology* **146** 1403–1410.
- Keire DA, Mannon P, Kobayashi M, Walsh JH, Solomon TE & Reeve JR 2000 Primary structures of PYY, [Pro³⁴]PYY, and PYY-(3–36) confer different conformations and receptor selectivity. *American Journal of Physiology. Gastrointestinal and Liver Physiology* **279** G126–G131.
- Ortiga-Carvalho TM, Polak J, McCann S & Pazos-Moura CC 1996 Effect of thyroid hormones on pituitary neuromedin B and possible interaction between thyroid hormones and neuromedin B on thyrotropin secretion. *Regulatory Peptides* **67** 47–53.
- Ortiga-Carvalho TM, Oliveira KJ, Soares BA & Pazos-Moura CC 2002 The role of leptin in the regulation of TSH secretion in the fed state: *in vivo* and *in vitro* studies. *Journal of Endocrinology* **174** 121–125.
- Parker MS, Wang JJ, Fournier A & Parker SL 2000 Upregulation of Pancreatic polypeptide – sensitive neuropeptide Y (NPY) receptors in estrogen-induced hypertrophy of the anterior pituitary gland in the Fischer-344 rat. *Molecular and Cellular Endocrinology* **164** 239–249.
- Pinilla L, Fernandez-Fernandez R, Vigo E, Navarro VM, Roa J, Castellano JM, Pineda R, Tena-Sempere M & Aguilar E 2006 The stimulatory effect of PYY₃₋₃₆ on gonadotropin secretion is potentiated in fasted rats. *American Journal of Physiology. Endocrinology and Metabolism* **290** 1162–1171.
- Rettori V, Milenkovic L, Fahim AM, Polak J, Bloom SR & McCann SM 1989 Role of neuromedin B in the control of the release of thyrotropin in the rat. *PNAS* **86** 4789–4792.
- Riediger T, Bothe C, Bescsei C & Lutz TA 2004 Peptide YY directly inhibits ghrelin-activated neurons of the arcuate nucleus and reverses fasting-induced c-Fos expression. *Neuroendocrinology* **79** 317–326.
- Sheikh SP, Feldthus N, Orkild H, Goke R, McGregor GP, Turner D, Moller M & Stuenkel EL 1998 Neuropeptide Y2 receptors on nerve endings from the rat neurohypophysis regulate vasopressin and oxytocin release. *Neuroscience* **82** 107–115.
- Talsania T, Anini Y, Siu S, Drucker DJ & Brubaker PL 2005 Peripheral exendin-4 and peptide YY₃₋₃₆ synergistically reduce food intake through different mechanisms in mice. *Endocrinology* **146** 3748–3756.
- Tovar SA, Seoane LM, Caminos JE, Nogueiras R, Casanueva FF & Diéguez C 2004 Regulation of peptide YY levels by age, hormonal, and nutritional status. *Obesity Research* **12** 1944–1950.
- Tschöp M, Castañeda TR, Joost HG, Thöne-Reineke C, Ortman S, Klaus S, Hagan MM, Chandler PC, Oswald KD, Benoit SC *et al.* 2004 Physiology: does gut hormone PYY₃₋₃₆ decrease food intake in rodents? *Nature* **430** 165.
- Zar JH 1996 *Data Transformations – Biostatistical Analysis*, edn 3 pp 277–279. London: Prentice Hall International.

Received in final form 17 August 2006

Accepted 22 August 2006

Made available online as an Accepted Preprint

15 September 2006