Cortisol regulation of ion transporter mRNA in Atlantic salmon gill and the effect of salinity on the signaling pathway

Pia Kiilerich, Karsten Kristiansen¹ and Steffen S Madsen

Institute of Biology and ¹Department of Biochemistry and Molecular Biology, University of Southern Denmark, Odense, Campusvej 55, DK-5230 Odense M, Denmark

(Correspondence should be addressed to S S Madsen; Email: steffen@biology.sdu.dk)

Abstract

Based on real-time RT-PCR, analysis of transcripts of selected ion-regulatory proteins (Na^+ , K^+ -ATPase $\alpha 1a$ and $\alpha 1b$ subunit, Na^+ , K^+ , 2Cl^- cotransporter, cystic fibrosis transmembrane conductance regulator (CFTR), and H^+ -ATPase B-subunit), the regulatory role of cortisol and the associated receptor signaling pathway (glucocorticoid (GR) versus mineralocorticoid (MR)) of cortisol was investigated in the salmon gill. Using a gill organ culture technique, the effect of cortisol with and without addition of specific hormone receptor antagonists (RU486 and spironolactone) was analyzed in gills from freshwater (FW) and seawater (SW) acclimated fish. The effect of cortisol was highly dependent on acclimation to salinity. In FW, cortisol stimulated the transcript levels of CFTR-I and Na^+ , K^+ -ATPase $\alpha 1a$ and $\alpha 1b$. Addition of RU486 totally abolished the cortisol effects on

CFTR-I and Na^+ , K^+ -ATPase $\alpha 1b$, suggesting that signaling was mediated via GR. Interestingly, both spironolactone and RU486 were able to inhibit the cortisol effect on Na^+ , K^+ -ATPase $\alpha 1a$ indicating a role for both MR and GR in regulation of this target gene. In SW, cortisol increased the transcript levels of CFTR-I, CFTR-II, Na^+ , K^+ -ATPase $\alpha 1a$ and $\alpha 1b$, and NKCC. Interestingly, the effect of cortisol on CFTR-I and Na^+ , K^+ -ATPase $\alpha 1a$ was mediated through GR and MR respectively, while both GR and MR signaling were required in the regulation of CFTR-II and Na^+ , K^+ -ATPase $\alpha 1b$. In FW gills, GR1 and MR transcript levels were not significantly affected by cortisol. In SW gills, GR1 and MR transcripts were downregulated by cortisol; GR1 was regulated via the MR and MR regulation was mediated via GR.

Journal of Endocrinology (2007) **194**, 417–427

Introduction

Cortisol, the major corticosteroid in teleost fish, is considered a dual regulator of chloride cell differentiation and function. During seawater (SW) acclimation, cortisol stimulates chloride cell numbers and Na^+ , K^+ -ATPase enzymatic activity in the gill and increases overall salinity tolerance in synergy with insulin-like growth factor-I (IGF-I) and growth hormone (GH; reviewed by McCormick 2001). In freshwater (FW) environment, cortisol presumably cooperates with prolactin (PRL) to ensure ion-osmotic homeostasis (reviewed by McCormick 2001). Cortisol alone stimulates overall Na^+ , K^+ -ATPase enzyme activity and mRNA levels of certain ion transport proteins in the gill *in vitro*: Na^+ , K^+ , 2Cl^- cotransporter (NKCC), $\alpha 1a$ and $\alpha 1b$ subunits of the Na^+ , K^+ -ATPase (McCormick & Bern 1989 (FW coho salmon, *Oncorhynchus kisutch*); Tipsmark *et al.* 2002 (FW brown trout, *Salmo trutta*, and Atlantic salmon, *Salmo salar*); Shrimpton & McCormick 1999 (FW rainbow trout, *Oncorhynchus mykiss*); and Deane & Woo 2005 (SW silver sea bream, *Sparus sarba*)).

The mineralocorticoid-like role of cortisol is unique to fish, since in vertebrates ranging from amphibians to

mammals aldosterone serves this function by acting through the mineralocorticoid receptor (MR; Agarwal & Mirshahi 1999). The absence of aldosterone in fish (see Prunet *et al.* 2006) may be compensated for by the ability of cortisol to bind both glucocorticoid receptors (GR) and MR in fish (Bury *et al.* 2003, Sturm *et al.* 2005), and that these receptors bind identical response elements in target gene promoters (Farman & Rafestin-Oblin 2001). The possibility that differential signaling by cortisol may occur through GR and MR in fish has yet to be elucidated. Recently, 11-deoxycorticosterone (DOC), an intermediate in aldosterone synthesis in mammals, was identified in fish as a potential mineralocorticoid hormone. DOC has high affinity for MR (Sturm *et al.* 2005) and seems to be the ancestral mineralocorticoid (Bridgham *et al.* 2006). The information concerning the function and significance of DOC is very limited, but the emergence of this potential hormone further emphasizes the requirement to investigate the signaling pathways of GR and MR in fish.

The knowledge of osmoregulatory gene regulation by cortisol in fish gill is scarce and little is known about exact cortisol signaling mechanisms through GR and MR. This is the first study that specifically addresses these questions. In

order to investigate the specific cortisol effect on gene expression in Atlantic salmon gill, an *in vitro* gill organ culture system was used. Using this technique, isolated effects of cortisol on gene expression were investigated in intact gill tissue, where cells remain in their natural cellular and junctional environment. QPCR analysis was applied to examine differential regulation of ion transport genes representing SW (Na^+ , K^+ -ATPase $\alpha 1b$, NKCC, the cystic fibrosis transmembrane conductance regulator isoform I and II (CFTR-I and CFTR-II)) and FW gills (Na^+ , K^+ -ATPase $\alpha 1a$ and the V-type H^+ -ATPase B-subunit). Specific antagonists for GR and MR (RU486 and spironolactone respectively) were applied to further examine the signaling pathway of cortisol.

Materials and Methods

Fish and sampling

In September, 1·5-year-old post-smolt *S. salar* (Burishoole stock) from Danmarks Center for Vildlaks (Randers, Denmark) were acclimated at the Odense Campus (University of Southern Denmark) at 14° C, 12 h light:12 h darkness cycle in 500 l fiberglass tanks with either flow through tap water or 30 ppt artificial SW for at least 30 days prior to sampling. The fish were fed *ad libitum* once daily with pelleted trout feed. When sampled, fish were stunned with a blow to the head and killed by cutting the spinal cord and pithing the brain.

Gill block incubation

To examine the effect of cortisol on gene expression, a gill block incubation experiment modified from McCormick & Bern (1989) was conducted using a repeated measures design. Excised gill arches from each fish were placed in separate glass Petri dishes placed on ice with 25 ml cold preincubation medium (Salmon Ringer's containing 400 000 IU/l streptomycin/penicillin and 2 mg/l amphotericin B. Salmon Ringer: 140 mM NaCl, 15 mM NaHCO_3 , 2·5 mM KCl, 1·5 mM CaCl_2 , 1·0 mM KH_2PO_4 , 0·8 mM MgSO_4 , 10 mM D-glucose, and 5·0 mM EPPS (pH 7·8); equilibrated with 99% O_2 /1% CO_2 for 10 min) for minimum half-an-hour. The gill filaments were dissected away from the cartilage arch and blocks of three to four filament pairs were cut out. Upon dissection, five to six gill blocks were frozen in liquid N_2 for RNA isolation to serve as preincubation controls. Groups of 10–12 gill blocks were carefully transferred to 6-well plates containing 2 ml incubation medium (Salmon Ringer's with 100 000 IU/l streptomycin/penicillin, and 2 mg/l amphotericin B) with or without the following agonist/antagonist combination: control, DMSO only (0·003 v/v); cortisol hemisuccinate + DMSO (F), 10 μg /ml (28 μM); RU486, 20 μg /ml (47 μM); Spironolactone (Spi), 20 μg /ml (48 μM); F + RU486, F + Spi. Cortisol hemisuccinate sodium salt (Sigma–Aldrich) was dissolved in Salmon Ringer, whereas RU486 and spironolactone (Sigma–Aldrich) were dissolved in

DMSO and mixed with Ringer to obtain the above concentrations. The 6-well plates were incubated at 14° C in a 99% O_2 /1% CO_2 atmosphere with shaking for 3 days. The gas in the incubation chamber was changed every day; the incubation medium was changed on day 2. Mucus was carefully removed by wiping the gill blocks on a paper towel and the gill blocks were stored at –80° C until analysis.

RNA purification, cDNA synthesis, and real-time semi-quantitative PCR

Total RNA was purified by the TRIzol procedure (Invitrogen) using 1 ml TRIzol/100 mg gill tissue according to manufacturer's recommendation. RNA concentration and purity was determined by measuring A_{260}/A_{280} . One microgram RNA was treated with 1 U RQ1 DNase (Promega) for 30 min at 37° C in a total volume of 20 μl followed by 5 min at 75° C to inactivate RQ1 DNase. Reverse transcription was carried out on 1 μg DNase-treated RNA with 2 μg random hexamers (GE Healthcare Bio-Sciences, Little Chalfont, UK) and 200 U. Moloney murine leukemia virus (MMLV) reverse transcriptase (Invitrogen) for 1 h at 37° C in the presence of 40 U RNAGuard (GE Healthcare) in a total volume of 25 μl . At the end the cDNA was diluted with 50 μl milliQ H_2O . Semi-quantitative real-time PCR analysis using SYBR Green detection was carried out on a Mx3000P instrument (Stratagene, La Jolla, CA, USA) using standard software settings including adaptive baseline for background detection, moving average, and amplification-based threshold settings with the built-in FAM/SYBR filter (excitation wavelength: 492 nm and emission wavelength: 516 nm). Reactions were carried out with 1 μl cDNA, 7·5 pmol glucocorticoid receptor (GR1) and mineralocorticoid receptor (MR) or 5 pmol (Na^+ , K^+ -ATPase $\alpha 1a$, Na^+ , K^+ -ATPase $\alpha 1b$, CFTR-I, CFTR-II, H^+ -ATPase, and NKCC) forward and reverse primer (DNA technology A/S, Denmark), 12·5 μl 2 $\times$ Brilliant SYBR green master mix (Stratagene) in a total volume of 25 μl . Cycling conditions: 95° C for 30 s, 60° C for 60 s, and 72° C for 5 s in 50 cycles (GR1 and MR) or 40 cycles (Na^+ , K^+ -ATPase $\alpha 1a$, Na^+ , K^+ -ATPase $\alpha 1b$, CFTR-I, CFTR-II, H^+ -ATPase, and NKCC). Melting curve analysis was carried out routinely with 30 s for each 1° C interval from 55 to 95° C.

Amplification efficiency, normalization, and calculations

For each primer set, cDNA was diluted 4, 16, 64, 256, and 1024 times in duplicates and analyzed by QPCR to determine amplification efficiency calculated as the slope of the resultant linear graph of threshold cycle (C_t) versus log cDNA concentration. The amplification efficiency (E_a) for each primer set was used for calculation of relative copy numbers of the respective target gene. EF1 α was used as normalization gene according to Olsvik *et al.* (2005). To verify the

normalization, 18S rRNA was used as a second normalization gene. No appreciable differences were observed between the results obtained with the two normalization genes (not shown). Relative copy number of the target gene cDNAs was calculated as $2^{(-Ct/Ea)}$, where Ct is the threshold cycle number and E_a is the amplification efficiency. Normalized units were obtained by dividing the relative copy number of the target gene cDNA with the relative copy number of the normalization gene cDNA.

Primers

Primer sequences are listed in Table 1. CFTR-I, CFTR-II, Na^+ , K^+ -ATPase $\alpha 1a$, NKCC, H^+ -ATPase B-subunit, GR1, MR, elongation factor (EF)1 α , and 18S rRNA primers were designed using the NetPrimer software (Premier Biosoft International, Palo Alto, CA, USA) with standard settings and double checked using the Primer3 software (Rozen & Skaletsky 2000) and BLASTed against the GenBank no. and EST sequences. Na^+ , K^+ -ATPase $\alpha 1b$ primers are identical to the Na^+ , K^+ -ATPase subunit specific primers based on *O. mykiss* mRNA sequences described by Richards *et al.* (2003). These primers have previously been successfully used in gill tissue from *S. salar* (Bystriansky *et al.* 2006). All primers were tested for nonspecific product amplification and primer-dimer formation using both melting curve analysis and agarose gel verification. In case of lack of *S. salar* mRNA sequence information, PCR primers based on conserved regions of *O. mykiss*, mRNA sequences were used when necessary (Na^+ , K^+ -ATPase $\alpha 1a$ and $\alpha 1b$, GR1, and MR). The high degree of conservation of nucleotide sequence between salmon and trout permit the use of primers between these two species. Moreover, the functionality and significance of the NKCC and H^+ -ATPase genes have yet to be validated since only partial sequences of these target genes is available so far.

All primers were synthesized by DNA Technology A/S (Århus, Denmark).

Statistical analysis

Data were analyzed using a repeated measures one-way ANOVA followed by a Bonferroni adjusted Fisher's least significant difference (LSD) test where appropriate, taking into account the total number of pairwise comparisons. When necessary, data were transformed to obtain normality and homogeneity of variances (Zar 1999). In all cases a significance level of $\alpha=0.05$ was used. All tests were performed using SAS (Version 9.1 for Windows, by SAS Institute Inc., Cary, NC, USA).

Results

Cortisol increased CFTR-I mRNA levels in FW- and SW-adapted salmon gill (Fig. 1A and B). The GR antagonist, RU486, decreased CFTR-I mRNA levels and totally abolished the effect of cortisol. The MR antagonist spironolactone had no effect on CFTR-I transcript levels and did not interfere with the effect of cortisol. While there was no effect of cortisol on CFTR-II mRNA levels in FW gills, an increase was observed in SW gills (Fig. 2A and B). RU486 reduced CFTR-II mRNA levels when compared with control and abolished the stimulatory effect of cortisol observed in SW gills. Spironolactone had no effect in FW gills (Fig. 2A) but reduced CFTR-II levels and abolished the effect of cortisol in SW gills (Fig. 2B). In FW and SW gills, Na^+ , K^+ -ATPase $\alpha 1a$ transcript level increased after cortisol treatment (Fig. 3A and B). Addition of both RU486 and spironolactone abolished the cortisol effect in FW gill, whereas only spironolactone antagonized the cortisol effect in SW gill. Cortisol increased Na^+ , K^+ -ATPase $\alpha 1b$ subunit transcript levels in both FW and SW gills (Fig. 4A and B), an effect which was fully antagonized by RU486 at both salinities. Spironolactone had no effect in FW gills but abolished the cortisol effect in SW gills (Fig. 4B).

In FW gills no effect of cortisol on NKCC mRNA level was observed (Fig. 5A). GR and MR antagonists both increased NKCC transcript levels in FW gills. Cortisol

Table 1 Sequences and target genes of primers used to investigate mRNA levels after *in vitro* incubation of gills from freshwater (FW) and seawater (SW) adapted Atlantic salmon

Target sequence	Forward primer 5'-3'	Reverse primer 5'-3'	Gene bank acc. number
<i>Salmo salar</i> CFTR-I	CCTTCTCCAATATGTTGAAGAGGCAAG	GCACTTGGATGAGTCAGCAG	AF155237
<i>Salmo salar</i> CFTR-II	GCCTTATTTCTCTATTGTATGCACTT	GCCACCATGAAAACTAAA GAGTACCTCAG	AF161070
<i>O. mykiss</i> Na^+ , K^+ -ATPase $\alpha 1a$	CCCAGGATCACTCAATGTCAC	CCAAAGGCAAATGGGTGTTAAT	AY319391
<i>O. mykiss</i> Na^+ , K^+ -ATPase $\alpha 1b$	CTGCTACATCTCAACCAACAACATT	CACCATCACAGTGTTCACTTGGAT	AY319390
<i>Salmo salar</i> NKCC	ACGTGTCCCACATCTCAG	GAGGGCTTGGATGAGTCT	AJ417890
<i>Salmo salar</i> H^+ -ATPase B-subunit	GAACATGGAACTGCTCG	TGGTAGGCCAGGTACTGA	AJ250811
<i>O. mykiss</i> GR1	ACGACGATGGAGCCGAAC	ATGGCTTTGAGCAGGGATAG	Z54210
<i>O. mykiss</i> MR	AGACTCGACCCCCACCAAG	CGTTAGTGGGACTGGTGCTC	AF209873
<i>Salmo salar</i> EF1 α	GAGAACCAATTGAGAATTCCGAGAAG	GCACCCAGGCATACCTGAAA	AF321836
<i>Salmo salar</i> 18S rRNA	TATTGTGCCGCTAGAGGTGA	CCTCCGACTTTCGTTCTTGA	AJ427629

Figure 1 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on CFTR-I mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. CFTR-I mRNA levels were normalized to mRNA levels of EF1α. Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. An overall effect of treatment was detected in the FW and the SW experiments ($P<0.0001$). #Indicates significant difference from the control group and * indicates a significant difference from the cortisol-treated group ($P<0.05$).

stimulated NKCC mRNA levels in SW gills (Fig. 5B). This was not antagonized by RU486 but tended to be diminished by the addition of spironolactone ($P=0.2082$; Fig. 5B).

No significant effect of cortisol or antagonists on H^+ -ATPase B-subunit transcript levels were found either in FW or in SW gills (Fig. 6A and B).

In FW gills there was no effect of cortisol on GR1 mRNA levels (Fig. 7A). In SW gills, cortisol decreased GR1 levels when compared with the control level (Fig. 7B). RU486 had no effect but spironolactone antagonized this effect completely. No significant effect of cortisol and/or antagonists was found on the MR mRNA level in FW gills (Fig. 8A). However, cortisol tended to lower the MR transcript level ($P=0.1374$). Also in SW gills, cortisol tended to lower the MR transcript level ($P=0.1356$), a tendency which was abolished by RU486

but not spironolactone (Fig. 8B). Yet, only an overall effect of treatment and no single significant differences was detected.

Stability of the investigated mRNA species during culture was investigated by comparing normalized mRNA levels in preincubation control gills and after 3-day incubation (Table 2). The level of CFTR-I, CFTR-II, Na^+ , K^+ -ATPase $\alpha 1b$, and NKCC transcripts decreased during incubation of FW and SW gills, whereas that of H^+ -ATPase mRNA increased. Na^+ , K^+ -ATPase $\alpha 1a$ transcript level decreased significantly during incubation of FW gills but increased in SW gills. The GR1 transcript level decreased during incubation of FW gills and was unchanged in SW gills. MR transcript levels were unchanged during incubation of FW gills but increased during incubation of SW gills.

A summary of all effects of cortisol is shown in Table 3.

Figure 2 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on CFTR-II mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. CFTR-II mRNA levels were normalized to mRNA levels of EF1α. Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD. An overall effect of treatment was detected in the FW and the SW experiment ($P<0.0001$). #Indicates significant difference from the control group and * indicates a significant difference from the cortisol-treated group ($P<0.05$).

Figure 3 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on Na^+ , K^+ -ATPase $\alpha 1a$ mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. Na^+ , K^+ -ATPase $\alpha 1a$ mRNA levels were normalized to mRNA levels of EF1 α . Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. An overall effect of treatment was detected in the FW and SW experiments ($P<0.0001$). [#]Indicates significant difference from the control group and ^{*} indicates a significant difference from the cortisol-treated group ($P<0.05$).

Discussion

Cortisol is considered as a fundamental regulator of chloride cell differentiation and proliferation, and numerous experiments confirm the pivotal role of cortisol in salinity acclimation of fish (reviewed by McCormick 2001). However, in terms of the basic signaling pathways and molecular mechanisms involved, cortisol regulation of gene expression in fish is still poorly understood. GR elements (GREs) have so far only been identified in very few piscine promoters (lactate hydrogenase-B in killifish, *Fundulus heteroclitus*: Schulte *et al.* 2000; α -amylase in seabass, *Lates calcarifer*: Ma *et al.* 2004; growth hormone 2 in rainbow trout: Yang *et al.* 1997 and PRL in rainbow trout: Argenton *et al.* 1996). No functional GRE has been identified in the

osmoregulatory target genes investigated here although GREs have been identified in the human GR and Na^+ , K^+ -ATPase $\alpha 1$ and $\beta 1$ promoters (Derfoul *et al.* 1998, Kolla *et al.* 1999, Yudit & Cidlowski 2002).

To determine the cortisol signaling pathway, the specific antagonists RU486 (GR) and spironolactone (MR) were used. These antagonists have previously been employed in several fish studies (Sloman *et al.* 2001, Bury *et al.* 2003, Scott *et al.* 2005, Shaw *et al.* 2007) without appreciably unspecific effects. However, the limited use and the insufficient validation of specificity of the antagonists in fish systems might impute some uncertainty to the conclusions. In fact, agonistic effects of spironolactone (Sturm *et al.* 2005) and RU486 (Mazon *et al.* 2004) have been reported, but there was no consistent evidence for such effects in this study when

Figure 4 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on Na^+ , K^+ -ATPase $\alpha 1b$ mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. Na^+ , K^+ -ATPase $\alpha 1b$ mRNA levels were normalized to mRNA levels of EF1 α . Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. An overall effect of treatment was detected in the FW and the SW experiment ($P<0.0001$). [#]Indicates significant difference from the control group and ^{*} indicates a significant difference from the cortisol-treated group ($P<0.05$).

Figure 5 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on NKCC mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. NKCC mRNA levels were normalized to mRNA levels of EF1α. Values (means ± S.E.M., n=6) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. An overall effect of treatment was detected in the FW and the SW experiments ($P<0.005$). #Indicates significant difference from the control group and * indicates a significant difference from the cortisol-treated group ($P<0.05$).

using similar antagonist concentrations. Despite lowering of the basal transcript level of some of the targets by antagonists, no overall pharmacological effect was observed.

The gill tissue organ culture

The main advantage of using the present organ culture approach is that isolated effects of cortisol may be studied in gill tissue, in which cells remain in their natural structural and junctional organization. Despite the fact that isolation of gill tissue from the organism removes all endogenous input (neural and hormonal), which may be vital for the integrity and functionality of the tissue, several studies have validated the application of this technique, and demonstrated good viability of the tissue even after 4-day incubation (McCormick & Bern 1989, Bury *et al.* 1998, Shrimpton & McCormick 1999, Mazon *et al.* 2004, Deane & Woo 2005). Our study confirms

that the tissue is indeed responsive to experimental manipulations. Three-day incubation *per se* had an effect on the majority of the investigated transcripts, but apparently there is no conformity in the direction of the changes in mRNA level. Among the transcripts investigated in the present paper, several transcripts (CFTR-I, CFTR-II, Na⁺, K⁺-ATPase α1b, and NKCC) showed decreased levels during 3-day control incubation. However, others were unchanged (MR) or even increased (V-type H⁺-ATPase) during incubation. This demonstrates the differential influence of the endogenous input to the gill.

Regulation of CFTR-I and CFTR-II

Two isoforms of the chloride transporter CFTR are present in Atlantic salmon (Chen *et al.* 2001), but their expression regulation and separate functions are largely unknown. Here

Figure 6 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on H⁺-ATPase B subunit mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. H⁺-ATPase mRNA levels were normalized to mRNA levels of EF1α. Values (means ± S.E.M., n=6) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. No overall effect of treatment was detected in the FW or the SW experiments ($P<0.05$).

Figure 7 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on GR1 mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. GR1 mRNA levels were normalized to mRNA levels of EF1 α . Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. No overall effect of treatment was detected in the FW experiment ($P>0.05$). An overall effect of treatment was detected in the SW experiment ($P<0.001$). #Indicates significant difference from the control group and * indicates a significant difference from the cortisol-treated group ($P<0.05$).

we show that the two isoforms are differentially regulated by cortisol. Cortisol increased the CFTR-I transcript level in both FW and SW, whereas CFTR-II levels were only affected in the SW gill. This agrees with a report by Singer *et al.* (2003) where cortisol treatment of FW Atlantic salmon stimulates CFTR-I but not CFTR-II mRNA levels. The CFTR-I isoform is also upregulated during smoltification and after SW-transfer in salmon, while CFTR-II is only transiently affected under these conditions (reviewed by Marshall & Singer 2002; T. Nilsen, Univ. of Bergen, pers. comm.). Altogether, the two CFTR isoforms seem to be differentially regulated and expressed in the gill. The effect of cortisol on CFTR-I transcript levels in salmon is opposite to that seen in FW-adapted striped bass gill (*Morone saxatilis*), where cortisol caused a downregulation of the CFTR mRNA level *in vitro* (Madsen *et al.* 2007). The effect of cortisol on CFTR-I and CFTR-II levels in FW and SW gills was consistently abolished

by RU486, indicating that cortisol signals through the GR in both hyper- and hypo-osmoregulating gill tissue. An inhibitory effect of RU486 was also shown for CFTR in the gill of SW-acclimating (Shaw *et al.* 2007), but not FW-acclimating (Scott *et al.* 2005) killifish. The present reduction of CFTR levels by RU486 alone when compared with control levels may be due to blocking of endogenous cortisol left in the tissue or due to a nonspecific effect of the antagonist. Putative GR-binding sites have been identified in the promoter region of the *F. heteroclitus* CFTR gene (Schulte & Keir 2003); however, the authors reported that these binding sites were nonfunctional when tested in cell culture and transgenic fish.

There was no evidence for cortisol signaling through MR in CFTR-I and CFTR-II regulation in FW-acclimated salmon in this study. However, in the SW gill cortisol stimulation of CFTR-II transcript levels was blocked by

Figure 8 Effect of cortisol (F) and GR (RU486) and MR (Spi) antagonists on MR mRNA levels in gill blocks from (A) FW- and (B) SW-acclimated Atlantic salmon. MR mRNA levels were normalized to mRNA levels of EF1 α . Values (means $\pm$ S.E.M., $n=6$) were analyzed with a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test. No overall effect of treatment was detected in the FW experiment ($P>0.05$). An overall effect of treatment was detected in the SW experiment ($P<0.05$) but no single significant effects was detected ($P>0.05$).

Table 2 Comparison of mRNA levels before (preincubation controls) and after (3-day controls) incubation of freshwater (FW)- and seawater (SW)-acclimated Atlantic salmon gill

	FW		SW	
	Preincubation	3-Day control	Preincubation	3-Day control
Target gene				
CFTR-I	2.8±0.6	0.9±0.1 ^a	5.3±0.5	1.0±0.1 ^a
CFTR-II	5.2±0.6	1.5±0.1 ^a	4.0±0.2	1.4±0.3 ^a
Na ⁺ , K ⁺ -ATPase α1a	19.7±8.0	6.6±1.2 ^a	76.5±13.9	170.9±25.9 ^a
Na ⁺ , K ⁺ -ATPase α1b	34.4±2.3	8.1±0.5 ^a	129.3±18.9	9.5±1.6 ^a
NKCC	36.0±8.7	19.6±3.4 ^a	233.0±25.5	46.9±8.0 ^a
H ⁺ -ATPase B	0.7±0.3	8.6±2.3 ^a	13.2±3.3	104.4±38.5 ^a
GR	46.5±3.0	1.9±1.2 ^a	46.1±2.6	40.2±2.1
MR	5.6±0.2	4.7±0.2	3.6±0.3	5.1±0.2 ^a

^aDenotes significant difference between preincubation and control values as determined by a repeated measures one-way ANOVA followed by Bonferroni adjusted Fisher's LSD test conducted on the datasets from Figs 1 to 8. Values are means±S.E.M. of six fish. mRNA levels are depicted as relative copy number of the target gene normalized to the relative copy number of EF1α.

spironolactone suggesting additional regulation through MR. Thus far the role of MR in CFTR regulation has only been investigated in killifish where no effect of spironolactone was reported after BW to FW transfer (Scott *et al.* 2005).

Regulation of Na⁺, K⁺-ATPase α1 isoforms

Recently, two Na⁺-ATPase α1 isoforms were identified in the rainbow trout gill displaying reverse responses to salinity changes (Richards *et al.* 2003): the α1a isoform, characteristic of the FW-gill and the α1b isoform which increases during SW acclimation. This pattern has been characterized in other salmonids as well (*S. salar* and Arctic char, *Salvelinus alpinus*; Bystriansky *et al.* 2006). Cortisol stimulated both transcripts in gills from both salinities thus agreeing with its dual role in osmoregulation (reviewed by McCormick 2001). Cortisol has previously been shown to increase overall Na⁺, K⁺-ATPase

protein (Seidelin *et al.* 1999) and α-subunit mRNA (Tipsmark *et al.* 2002) levels in the salmonid gill, but this is the first study to distinguish cortisol effects on the distinct Na⁺, K⁺-ATPase α1 isoforms and to give evidence for the involvement of both GR and MR. Curiously, the signaling pathway is opposite at the two salinities. The α1a isoform is consistently regulated by the MR (in FW and SW gills) and by additional GR signaling in the FW gill. The α1b isoform is consistently regulated via GR signaling (in FW and SW) and by additional MR signaling in the SW gill. This pattern of regulation complies with the hypothesis that GR signaling has a primary role in developing the gill for ion excretion, while MR signaling may be more important in stimulating ion retention (Sloman *et al.* 2001, Scott *et al.* 2005, Kiilerich *et al.* 2007). Future experiments should investigate to what degree the present cortisol effects may be modulated *in vivo* by interaction with other hormones in salinity acclimation, such as PRL, growth hormone, and IGF-I.

Table 3 Summary of cortisol action and the deduced signaling pathway. Arrows indicate upregulation (↑), downregulation (↓) and no regulation (→) of the target mRNA.

	FW		SW	
	Cortisol	Receptor	Cortisol	Receptor
Target cDNA				
CFTR-I	↑	GR	↑	GR
CFTR-II	→	–	↑	GR and MR
Na ⁺ , K ⁺ -ATPase α1a	↑	GR and MR	↑	MR
Na ⁺ , K ⁺ -ATPase α1b	↑	GR	↑	GR and MR
NKCC	→	–	↑	(MR)
H ⁺ -ATPase	→	–	→	–
GR	→	–	↓	MR
MR	(↓)	(GR and MR)	(↓)	(GR)

Brackets denote that the effect was statistically nonsignificant. See text for details

Regulation of Na^+ , K^+ , 2Cl^- cotransporter

Consistent with the SW-adaptive role of NKCC (reviewed by McCormick 2001, Tipsmark *et al.* 2002), cortisol increased NKCC mRNA levels but only in gills from SW-acclimated salmon. Previously, cortisol stimulated NKCC mRNA in FW gill *in vitro* (Tipsmark *et al.* 2002) and NKCC protein level *in vivo* (Pelis & McCormick 2001). A reason for the disparity in response to cortisol in the FW gill may be found in seasonal variation of sensitivity of the gill, since post-smolt salmon were used in this study whereas Tipsmark *et al.* (2002) used smolting salmon. Thus, stimulation of this particular SW-type ion transporter may depend on the developmental stage of chloride cells. No antagonist was able to efficiently inhibit the effect of cortisol in SW gill, but blocking of MR slightly lowered the NKCC transcript level when compared with cortisol alone. On the other hand, both RU486 and spironolactone increased the NKCC transcript level significantly in the FW gill. The reason for this is unknown, and further conclusions concerning the signaling pathway should await further research. Nonetheless, the lack of antagonist effect on NKCC mRNA expression in SW gill is in good agreement with Scott *et al.* (2005), where neither RU486 nor spironolactone implants affected the suppression of NKCC mRNA levels in BW- to FW-transferred killifish (Scott *et al.* 2005).

Regulation of H^+ -ATPase B-subunit

In addition to participating in acid–base regulation, vacuolar-type H^+ -ATPase is suspected to facilitate Na^+ uptake across the gill from hypo-osmotic environments (Marshall 2002, Hirose *et al.* 2003). Accordingly, protein levels have been shown to increase upon FW acclimation (Wilson *et al.* 2002). There is only limited information on the endocrine regulation on H^+ -ATPase expression, and only at the protein level. Based on *N*-ethylmaleimide-sensitive inhibition of V-type H^+ -ATPase activity, cortisol injection was shown to stimulate enzyme activity in the FW rainbow trout gill (Lin & Randall 1993). Our data, however, did not confirm this effect of cortisol at the mRNA level, when applied *in vitro*.

Regulation of GR and MR

Downregulation of SW gill GR1 transcript level by cortisol reported here is in good agreement with the downregulation of GR protein levels after corticosteroid treatment of gill, brain and liver of coho salmon, and rainbow trout (Pottinger 1990, Maule & Schreck 1991, Lee *et al.* 1992, Shrimpton & Randall 1994). However, an increase in GR mRNA levels upon cortisol treatment of rainbow trout hepatocytes (Sathiyaa & Vijayan 2003) was also reported. Interestingly, a dual mode of GR feedback regulation was recently reported by Takahashi *et al.* (2006) where chronic cortisol treatment *in vivo* increased GR levels, whereas a single cortisol injection

caused a decrease in GR transcript levels (Takahashi *et al.* 2006). Curiously, the cortisol regulation of GR1 transcript levels seen in SW gill is signaled through the MR. The GR has been linked to SW acclimation since the GR1 mRNA levels and the corticosteroid-binding ability increase in the gill during smoltification and after SW transfer in salmonids (reviewed by Prunet *et al.* 2006, Kiilerich *et al.* 2007). Consistent with this is the fact that cortisol only affects the gill GR1 mRNA levels in SW-acclimated salmon.

The MR was recently discovered in fish (Colombe *et al.* 2000, Sloman *et al.* 2001, Greenwood *et al.* 2003, Sturm *et al.* 2005) and its function is still puzzling. There is evidence that MR plays a role in chloride cell proliferation and differentiation in FW rainbow trout and killifish (Sloman *et al.* 2001, Scott *et al.* 2004, 2005) and a FW-adaptive role for MR has been proposed in salmon (Kiilerich *et al.* 2007). The present data show no convincing effects of cortisol on the transcript level of MR, but only a trend of a downregulation by cortisol is to occur via the GR, at least in the SW gill. Future experiments should include the possible role of DOC in modulating MR level.

Conclusion and perspectives

The present study demonstrated the important role of cortisol in regulating osmoregulatory gene expression in the salmon gill. Most transcripts investigated were stimulated by cortisol in both FW and SW (Na^+ , K^+ -ATPase $\alpha 1\text{b}$, CFTR-I) or in SW only (CFTR-II, NKCC). The difference in responsiveness may be due to developmental and functional differences of the target cells involved. This may come down to a matter of functional equipment with corticosteroid receptors. The involvement of GR and MR varies with respect to gene of interest and acclimation salinity, thus strongly supporting a role for both signaling pathways in salinity acclimation as well as interaction between them. All effects of cortisol and antagonists in FW and SW are listed in Table 3.

The differences in target gene regulation may be due to several molecular differences along the hormone signaling pathway, e.g., differences in GRE composition of individual promoters as has been described in killifish (Schulte *et al.* 2000). Colocalization of the 11β hydroxysteroid dehydrogenase 2 enzyme, which converts cortisol to the receptor inactive cortisone (Funder *et al.* 1988) with the MR, may add another level of control of cortisol signaling through the MR. This is particularly interesting because of the recent identification of DOC – an intermediate in cortisol synthesis in fish – as a potent and specific ligand for MR in fish (Sturm *et al.* 2005, Prunet *et al.* 2006). Future studies should emphasize structural analyses of promoter and GRE sequences in osmoregulatory genes in fish in conjunction with functional studies of GR/MR effects of cortisol and DOC.

Acknowledgements

The authors would like to thank Leyla Güler for excellent technical assistance. P K was supported by a grant from the School of Aquatic Sciences (SOAS, Denmark). S S M was supported by a grant from The Danish National Research Council (272-06-0526). The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Agarwal MK & Mirshahi M 1999 General overview of mineralocorticoid hormone action. *Pharmacology and Therapeutics* **84** 273–326 (Review).
- Argenton F, Ramoz N, Charlet N, Bernardini S, Colombo L & Bortolussi M 1996 Mechanisms of transcriptional activation of the promoter of the rainbow trout prolactin gene by GHF1/Pit1 and glucocorticoid. *Biochemical and Biophysical Research Communications* **224** 57–66.
- Bridgham JT, Carroll SM & Thornton JW 2006 Evolution of hormone-receptor complexity by molecular exploitation. *Science* **312** 97–101.
- Bury NR, Jie L, Flik G, Lock RAC & Wendelaar Bonga SE 1998 Cortisol protects against copper induced necrosis and promotes apoptosis in fish gill chloride cells *in vitro*. *Aquatic Toxicology* **40** 193–202.
- Bury NR, Sturm A, Le RP, Lethimonier C, Ducouret B, Guiguen Y, Robinson-Rechavi M, Laudet V, Rafestin-Oblin ME & Prunet P 2003 Evidence for two distinct functional glucocorticoid receptors in teleost fish. *Journal of Molecular Endocrinology* **31** 141–156.
- Bystriansky JS, Richards JG, Schulte PM & Ballantyne JS 2006 Reciprocal expression of gill Na⁺/K⁺-ATPase alpha-subunit isoforms alpha1a and alpha1b during seawater acclimation of three salmonid fishes that vary in their salinity tolerance. *Journal of Experimental Biology* **209** 1848–1858.
- Chen JM, Cutler C, Jacques C, Boeuf G, Denamur E, Lecointre G, Mercier B, Cramb G & Ferec C 2001 A combined analysis of the cystic fibrosis transmembrane conductance regulator: implications for structure and disease models. *Molecular Biology and Evolution* **18** 1771–1788.
- Colombe L, Fostier A, Bury N, Pakdel F & Guiguen Y 2000 A mineralocorticoid-like receptor in the rainbow trout, *Oncorhynchus mykiss*: cloning and characterization of its steroid binding domain. *Steroids* **65** 319–328.
- Deane EE & Woo NY 2005 Cloning and characterization of sea bream Na⁺-K⁺-ATPase α and β subunit genes: *in vitro* effects of hormones on transcriptional and translational expression. *Biochemical and Biophysical Research Communications* **17** 1229–1238.
- Derfoul A, Robertson NM, Lingrel JB, Hall DJ & Litwack G 1998 Regulation of the human Na/K-ATPase β 1 gene promoter by mineralocorticoid and glucocorticoid receptors. *Journal of Biological Chemistry* **273** 20702–20711.
- Farman N & Rafestin-Oblin ME 2001 Multiple aspects of mineralocorticoid selectivity. *American Journal of Physiology. Renal Physiology* **280** F181–F192 (Review).
- Funder JW, Pearce PT, Smith R & Smith AI 1988 Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science* **242** 583–585.
- Greenwood AK, Butler PC, White RB, DeMarco U, Pearce D & Fernald RD 2003 Multiple corticosteroid receptors in a teleost fish: distinct sequences, expression patterns, and transcriptional activities. *Endocrinology* **144** 4226–4236.
- Hirose S, Kaneko T, Naito N & Takei Y 2003 Molecular biology of major components of chloride cells. *Comparative Biochemistry and Physiology. Part B, Biochemistry and Molecular Biology* **136** 593–620 (Review).
- Kiilerich P, Kristiansen K & Madsen SS 2007 Hormone receptors in gills of smolting Atlantic salmon *Salmo salar*: expression of growth hormone, prolactin, mineralocorticoid and glucocorticoid receptors and 11 β -hydroxysteroid dehydrogenase type 2. *General and Comparative Endocrinology* **152** 295–303.
- Kolla V, Robertson NM & Litwack G 1999 Identification of a mineralocorticoid/glucocorticoid response element in the human Na/K-ATPase α 1 gene promoter. *Biochemical and Biophysical Research Communications* **266** 5–14.
- Lee PC, Goodrich M, Struve M, Yoon HI & Weber D 1992 Liver and brain glucocorticoid receptor in rainbow trout, *Oncorhynchus mykiss*: down-regulation by dexamethasone. *General and Comparative Endocrinology* **87** 222–231.
- Lin H & Randall DJ 1993 H⁺-ATPase activity in crude homogenates of fish gill tissue: inhibitor sensitivity and environmental and hormonal regulation. *Journal of Experimental Biology* **180** 163–174.
- Ma P, Liu Y, Reddy KP, Chan WK & Lam TJ 2004 Characterization of the seabass pancreatic α -amylase gene and promoter. *General and Comparative Endocrinology* **137** 78–88.
- Madsen SS, Jensen LN, Tipsmark CK, Kiilerich P & Borski RJ 2007 Differential regulation of cystic fibrosis transmembrane conductance regulator and Na⁺, K⁺-ATPase in gills of striped bass, *Morone saxatilis*: effect of salinity and hormones. *Journal of Endocrinology* **192** 249–260.
- Marshall WS 2002 Na(+), Cl(-), Ca(2+) and Zn(2+) transport by fish gills: retrospective review and prospective synthesis. *Journal of Experimental Zoology* **293** 264–283 (Review).
- Marshall WS & Singer TD 2002 Cystic fibrosis transmembrane conductance regulator in teleost fish. *Biochimica et Biophysica Acta* **1566** 16–27 (Review).
- Maule AG & Schreck CB 1991 Stress and cortisol treatment changed affinity and number of glucocorticoid receptors in leukocytes and gill of coho salmon. *General and Comparative Endocrinology* **84** 83–93.
- Mazon AF, Nolan DT, Lock RAC, Fernandes MN & Wendelaar Bonga SE 2004 A short-term *in vitro* gill culture system to study the effects of toxic (copper) and non-toxic (cortisol) stressors on the rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Toxicology In Vitro* **18** 691–701.
- McCormick SD 2001 Endocrine control of osmoregulation in teleost fish. *American Zoologist* **41** 781–794 (Review).
- McCormick SD & Bern HA 1989 *In vitro* stimulation of Na⁺-K⁺-ATPase activity and ouabain binding by cortisol in coho salmon gill. *American Journal of Physiology* **256** R707–R715.
- Olsvik PA, Lie KK, Jordal AE, Nilsen TO & Hordvik I 2005 Evaluation of potential reference genes in real-time RT-PCR studies of Atlantic salmon. *BMC Molecular Biology* **6** 21.
- Pelis RM & McCormick SD 2001 Effects of growth hormone and cortisol on Na(+)-K(+)-2Cl(-) cotransporter localization and abundance in the gills of Atlantic salmon. *General and Comparative Endocrinology* **124** 134–143.
- Pottinger TG 1990 The effect of stress and exogenous cortisol on receptor-like binding of cortisol in the liver of rainbow trout, *Oncorhynchus mykiss*. *General and Comparative Endocrinology* **78** 194–203.
- Prunet P, Sturm A & Milla S 2006 Multiple corticosteroid receptors in fish: from old ideas to new concepts. *General and Comparative Endocrinology* **147** 17–23 (Review).
- Richards JG, Semple JW, Bystriansky JS & Schulte PM 2003 Na⁺/K⁺-ATPase α -isoform switching in gills of rainbow trout (*Oncorhynchus mykiss*) during salinity transfer. *Journal of Experimental Biology* **206** 4475–4486.
- Rozen S & Skaletsky H 2000 Primer3 on the WWW for general users and for biologist programmers. *Methods in Molecular Biology* **132** 365–386.
- Sathiyaa R & Vijayan MM 2003 Autoregulation of glucocorticoid receptor by cortisol in rainbow trout hepatocytes. *American Journal of Physiology. Cell Physiology* **284** C1508–C1515.
- Schulte PM & Keir K 2003 The cystic fibrosis transmembrane conductance regulator and the evolution of euryhalinity in fish. (Abstract). *Integrative and Comparative Biology* **43** 978.
- Schulte PM, Glemet HC, Fiebig AA & Powers DA 2000 Adaptive variation in lactate dehydrogenase-B gene expression: role of a stress-responsive regulatory element. *PNAS* **97** 6597–6602.
- Scott GR, Richards JG, Forbush B, Isenring P & Schulte PM 2004 Changes in gene expression in gills of the euryhaline killifish *Fundulus heteroclitus* after abrupt salinity transfer. *American Journal of Physiology. Cell Physiology* **287** C300–C309.

- Scott GR, Keir KR & Schulte PM 2005 Effects of spironolactone and RU486 on gene expression and cell proliferation after freshwater transfer in the euryhaline killifish. *Journal of Comparative Physiology. B* **175** 499–510.
- Seidelin M, Madsen SS, Byrjalsen A & Kristiansen K 1999 Effects of insulin-like growth factor-I and cortisol on Na^+ , K^+ -ATPase expression in osmoregulatory tissues of brown trout (*Salmo trutta*). *General and Comparative Endocrinology* **113** 331–342.
- Shaw JR, Gabor K, Hand E, Lankowski A, Durant L, Thibodeau R, Stanton CR, Barnaby R, Coutermarsh B, Karlson KH *et al.* 2007 Role of glucocorticoid receptor in acclimation of killifish (*Fundulus heteroclitus*) to seawater and effects of arsenic. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **292** R1052–R1060.
- Shrimpton JM & McCormick SD 1999 Responsiveness of gill Na^+/K^+ -ATPase to cortisol is related to gill corticosteroid receptor concentration in juvenile rainbow trout. *Journal of Experimental Biology* **202** 987–995.
- Shrimpton JM & Randall DJ 1994 Downregulation of corticosteroid receptors in gills of coho salmon due to stress and cortisol treatment. *American Journal of Physiology* **267** R432–R438.
- Singer TD, Finstad B, McCormick SD, Wiseman SD, Schulte PM & McKinley RS 2003 Interactive effects of cortisol treatment and ambient seawater challenge on gill Na^+ , K^+ -ATPase and CFTR expression in two strains of Atlantic salmon smolts. *Aquaculture* **222** 15–28.
- Sloman KA, Desforges PR & Gilmour KM 2001 Evidence for a mineralocorticoid-like receptor linked to branchial chloride cell proliferation in freshwater rainbow trout. *Journal of Experimental Biology* **204** 3953–3961.
- Sturm A, Bury N, Dengreville L, Fagart J, Flouriot G, Rafestin-Oblin ME & Prunet P 2005 11-Deoxycorticosterone is a potent agonist of the rainbow trout (*Oncorhynchus mykiss*) mineralocorticoid receptor. *Endocrinology* **146** 47–55.
- Takahashi H, Sakamoto T, Hyodo S, Shepherd BS, Kaneko T & Grau EG 2006 Expression of glucocorticoid receptor in the intestine of a euryhaline teleost, the Mozambique tilapia (*Oreochromis mossambicus*): effect of seawater exposure and cortisol treatment. *Life Sciences* **78** 2329–2335.
- Tipsmark CK, Madsen SS, Seidelin M, Christensen AS, Cutler CP & Cramb G 2002 Dynamics of $\text{Na}(+)$, $\text{K}(+)$, $2\text{Cl}(-)$ cotransporter and $\text{Na}(+)$, $\text{K}(+)$ -ATPase expression in the branchial epithelium of brown trout (*Salmo trutta*) and Atlantic salmon (*Salmo salar*). *Journal of Experimental Zoology* **293** 106–118.
- Wilson JM, Whiteley NM & Randall DJ 2002 Ionoregulatory changes in the gill epithelia of coho salmon during seawater acclimation. *Physiological and Biochemical Zoology* **75** 237–249.
- Yang BY, Chan KM, Lin CM & Chen TT 1997 Characterization of rainbow trout (*Oncorhynchus mykiss*) growth hormone 1 gene and the promoter region of growth hormone 2 gene. *Archives of Biochemistry and Biophysics* **340** 359–368.
- Yudt MR & Cidlowski JA 2002 The glucocorticoid receptor: coding a diversity of proteins and responses through a single gene. *Molecular Endocrinology* **16** 1719–1726 (Review).
- Zar JH 1999 *Biostatistical Analysis*. edn 4, Upper Saddle River, NJ: Prentice-Hall.

Received in final form 25 May 2007

Accepted 4 June 2007

Made available online as an Accepted Preprint
6 June 2007