Human adrenal corticocarcinoma NCI-H295R cells produce more androgens than NCI-H295A cells and differ in 3 β -hydroxysteroid dehydrogenase type 2 and 17,20 lyase activities

Elika Samandari, Petra Kempná, Jean-Marc Nuoffer¹, Gaby Hofer, Primus E Mullis and Christa E Flück

Pediatric Endocrinology and Diabetology, University Children's Hospital Bern, Freiburgstrasse 15, G3 812, 3010 Bern, Switzerland

¹Department of Clinical Chemistry, University Hospital Inselspital, 3010 Bern, Switzerland

(Correspondence should be addressed to C E Flück; Email: christa.flueck@dkf.unibe.ch)

Abstract

The human adrenal cortex produces mineralocorticoids, glucocorticoids, and androgens in a species-specific, hormonally regulated, zone-specific, and developmentally characteristic fashion. Most molecular studies of adrenal steroidogenesis use human adrenocortical NCI-H295A and NCI-H295R cells as a model because appropriate animal models do not exist. NCI-H295A and NCI-H295R cells originate from the same adrenocortical carcinoma which produced predominantly androgens but also smaller amounts of mineralocorticoids and glucocorticoids. Research data obtained from either NCI-H295A or NCI-H295R cells are generally compared, although for the same experiments no direct comparison between the two cell lines has been performed. Therefore, we compared the steroid profile and the expression pattern of important genes involved in steroidogenesis in both cell lines. We found that steroidogenesis differs profoundly. NCI-H295A cells produce more mineralocorticoids, whereas NCI-H295R cells produce more androgens. Expression of the 3 β -hydroxysteroid

dehydrogenase (HSD3B2), cytochrome b5, and sulfonyltransferase genes is higher in NCI-H295A cells, whereas expression of the cytochrome P450c17 (CYP17), 21-hydroxylase (CYP21), and P450 oxidoreductase genes does not differ between the cell lines. We found lower 3 β -hydroxysteroid dehydrogenase type 2 but higher 17,20-lyase activity in NCI-H295R cells explaining the 'androgenic' steroid profile for these cells and resembling the zona reticularis of the human adrenal cortex. Both cell lines were found to express the ACTH receptor at low levels consistent with low stimulation by ACTH. By contrast, both cell lines were readily stimulated by 8Br-cAMP. The angiotensin type 1 receptor was highly expressed in NCI-H295R than NCI-H295A cells and angiotensin II stimulated steroidogenesis in NCI-H295R but not NCI-H295A cells. Our data suggest that comparative studies between NCI-H295A and NCI-H295R cells may help find important regulators of mineralocorticoid or androgen biosynthesis.

Journal of Endocrinology (2007) **195**, 459–472

Introduction

Human androgens are predominantly produced in the gonads (ovaries and testes) and in the adrenal cortex. The adult adrenal cortex consists of three morphologically and biochemically distinct zones (Miller 1988, Rainey *et al.* 2004): the outer zona glomerulosa (ZG), which produces mineralocorticoids for the regulation of water and salt homeostasis; the middle zona fasciculata (ZF), which produces glucocorticoids controlling body fuel metabolism and stress response; and the inner zona reticularis (ZR), which produces adrenal androgens for which the specific role and regulation are unknown. Generally, androgens are essential for the normal sexual development in males during fetal life and puberty. In adulthood, they are essential for sexual behavior, fertility, and thus maintenance of the human species.

All steroid hormones are produced from cholesterol (Fig. 1). The transfer of cholesterol to the mitochondria for the first

and rate limiting step of steroidogenesis, the conversion of cholesterol to pregnenolone (Preg), is supported by the steroidogenic acute regulatory protein (StAR). Enzymes involved in adrenal steroidogenesis are P450 side-chain cleavage (CYP11A1), 3 β -hydroxysteroid dehydrogenase type 2 (3 β HSDII; HSD3B2), 21-hydroxylase (P450c21; CYP21A2), 11 β -hydroxylase (P450c11 β ; CYP11B1), and aldosterone synthase (P450c11AS; CYP11B2) for the production of mineralocorticoids in the ZG. Additionally, the 17 α -hydroxylase activity of cytochrome P450c17 (CYP17) is needed for the production of glucocorticoids in the ZF. Finally, both the 17 α -hydroxylase and 17,20-lyase activities of P450c17 and the activity of the sulfonyltransferase (SULT2A1) are required to produce the C19 steroids dehydroepiandrosterone (DHEA) and its sulfate (DHEAS). P450c17 plays a key role in the zone-specific production of steroids. Presence of the 17 α -hydroxylase activity of P450c17 alone in the ZF results in the production of cortisol, while presence of both 17 α -hydroxylase and 17,20-lyase

Figure 1 Adrenal steroidogenesis. The initial and rate-limiting step of steroidogenesis is the conversion of cholesterol to Preg by the P450_{scc} enzyme. Preg may undergo 17α-hydroxylation to 17OH-Preg by the 17α-hydroxylase activity of the P450_{c17} enzyme or may be converted to Prog by the 3βHSDII enzyme. Prog may be converted to deoxycorticosterone (DOC) and aldosterone by sequential conversion by the P450_{c21}, P450_{c11β}, and P450_{c11AS} enzymes, which are all present in the human adrenal ZG. Alternatively, 17OH-Preg may be converted stepwise to 17OH-Prog, 11DOC and cortisol by 3βHSDII, P450_{c21}, and P450_{c11β} in the human adrenal ZF. Moreover, 17OH-Preg may be converted to DHEA by the 17,20-lyase activity of P450_{c17} followed by conversion to DHEAS by SULT2A1, or to Δ4A by 3βHSDII; all present in the human adrenal ZR. POR is the obligate electron transfer cofactor for P450_{c21} and P450_{c17}, while b5 fosters 17,20-lyase activity through allosteric mechanisms. In human adrenal H295 cells (as in the human testes), Δ4A may be converted to testosterone by 17βHSDIII/V. Dashed box shows reactions which are not described for the normal human adrenal cortex.

activities in the ZR results in the production of androgens (Miller *et al.* 1997). The 17,20 lyase activity of P450_{c17} in the ZR is fostered by the expression of the flavoprotein NADPH cytochrome P450 oxidoreductase (POR) and cytochrome b5 (b5) in the ZR, as well as by phosphorylation of P450_{c17} (Lin *et al.* 1993, Zhang *et al.* 1995, Auchus *et al.* 1998, Lam *et al.* 2001, Rainey *et al.* 2002, Pandey *et al.* 2003).

Steroid hormone biosynthesis is regulated in a species-specific, developmentally characteristic, tissue-specific, and a cAMP-dependent fashion (Waterman & Keeney 1996, Sewer & Waterman 2001, Fluck & Miller 2004), but not all regulation is understood in great details. The adrenocorticotrophic hormone (ACTH)/cAMP pathway plays a major role in the regulation of steroid hormone biosynthesis (John *et al.* 1986b, Sewer & Waterman 2003, Stocco *et al.* 2005). Upon release from the pituitary gland, ACTH stimulates the G-protein-coupled melanocortin 2 receptor (MC2R) which is located almost exclusively in the adrenal cortex (Forti *et al.* 2006). Activation of MC2R triggers the production of cAMP, which in turn activates protein kinase A and possibly crosstalk with other signaling pathways (Sewer & Waterman 2002). For adrenal steroidogenesis, it is known that ACTH/cAMP stimulation increases gene transcription of most steroidogenic enzymes (John *et al.* 1986a, Sewer & Waterman 2003, Stocco *et al.* 2005). Yet the exact downstream signalling cascade remains elusive as most genes encoding steroidogenic enzymes do not harbor typical cAMP response elements (CRE) in their promoters.

Aldosterone synthesis may be influenced by ACTH, but it is mainly regulated by the renin–angiotensin system as well as by the concentration of potassium (Goodman 2003). Angiotensin II (AngII) stimulates type 1 membrane receptors (AT1) of adrenal cells which act through inositol trisphosphate, diacylglycerol, as well as calcium to promote the formation of Preg from cholesterol and the synthesis of aldosterone through stimulation of P450_{c11AS}.

The fetal and adult human adrenals differ principally. The fetal adrenal consists of a definitive zone (the later adult adrenal) and a larger fetal zone, principally producing DHEA and DHEAS. After birth, the fetal zone disappears by the age of 6 months and the adrenal cortex first consists of the ZF and ZG. During ‘adrenarche’ by 3–4 years of age, the ZR appears histologically and starts to produce androgens by 6–8 years of age (Dhom 1973a,b, Cutler *et al.* 1978, Havelock *et al.* 2004, Campbell 2006). The role and the regulation of human adrenal androgen biosynthesis in general and adrenarche in particular are still poorly understood. Animal models may not be suitable for studies, because first of all, adrenal steroidogenesis varies fundamentally from species to species and secondly, adrenarche occurs only in higher-order primates (Havelock *et al.* 2004). Thus, most investigators use human cell models for their studies, i.e. NCI-H295A (A) and NCI-H295R (R) cells.

Human adrenocortical carcinoma A and R cells are two human adrenocortical cell lines that derive from the same adrenocortical tumor cell line NCI-H295 (H295; Gazdar

et al. 1990, Staels *et al.* 1993, Rainey *et al.* 2004). H295 cells were originally cultured from an adrenocortical carcinoma in a 48-year-old black female (Gazdar *et al.* 1990). H295 cells express all genes for the steroidogenic enzymes found in all three layers of an adult adrenal cortex (Fig. 1; Staels *et al.* 1993). Using different growth conditions, the two substrains A and R cells were produced in two different laboratories (Gazdar *et al.* 1990, Rodriguez *et al.* 1997). In contrast to the parental H295 cells, both A and R cells grow as a tight monolayer. Similar to H295 cells, A and R cells express all genes involved in adrenal steroidogenesis (Staels *et al.* 1993, Rodriguez *et al.* 1997, Dardis & Miller 2003). However, all investigators who study adrenal steroidogenesis use either A or R cells and compare their research data directly assuming that the two cell lines are identical. So far, no data are available comparing the two cell systems in the same experiments. Hence, we asked the question whether the two commonly used adrenal cell models are identical; and if they were different, what differences could be observed especially with respect to androgen production. To compare the two cell lines we studied their growth pattern, the steroid profiles and the expression pattern of genes involved in steroidogenesis. As we found major differences in androgen production between the two cell lines, we suggest that comparative studies of the two cell systems will give new insight into the regulation of adrenal androgen biosynthesis.

Materials and Methods

Materials

We purchased 8-bromo-cAMP (8Br-cAMP) and AngII from Fluka (Sigma-Aldrich). Synacthen^R (ACTH; beta¹⁻²⁴-corticotropin) was obtained from Novartis (Basel, Switzerland). Radio-labeled [³H]-Preg (14 Ci/mmol) and [³H]-DHEA (63 Ci/mmol) were bought from Perkin-Elmer (Boston, MA, USA). Radio-labeled [³H]-17 α -hydroxypregnenolone (17OHPreg; 50 Ci/mmol) was procured from American Radiolabel Chemicals Inc. (St. Louis, MO, USA). Trilostane was extracted in absolute ethanol (EtOH) from tablets commercially available as Modrenal[®] (Bioenvision, NY, USA).

Cell cultures

Human adrenal A cells were kindly provided by Prof. W L Miller and R cells were purchased from the American Type Culture Collection (number CRL-2128). Standard medium for A cells is RPMI-1640 (Gibco BRL Invitrogen) supplemented with 2% fetal calf serum, 0.1% selenium/insulin/transferrin (SIT), and penicillin/streptomycin (antibiotics; Rodriguez *et al.* 1997). Standard medium for R cells is Dulbecco's modified Eagle's/Ham's F12 (DMEM/F12; Gibco BRL Invitrogen) supplemented with 5% NuI serum, 0.1% SIT, and antibiotics (all from Gibco BRL).

Cell proliferation

A and R cells were cultured in six-well plates (750×10³ cells/well). After 72 h, cells were washed with PBS, trypsinized, and collected in PBS for counting with a hemocytometer.

Steroid profiling

A and R cells were grown in six-well plates (750×10³ cells/well). Steroid metabolism was labeled by adding [³H]-Preg, [³H]-17OHPreg, or [³H]-DHEA (~200 000 c.p.m./well), all dissolved in EtOH. To block the 3 β HSD activity, cells were treated with 1 μ M trilostane for 4 h prior to labeling steroidogenesis (Mahesh & Brann 1992). Steroids were extracted from the media as previously described (Auchus *et al.* 1998) and separated on thin layer chromatography (TLC) plates (Whatman, Bottingen, Switzerland) using chloroform:ethylacetate in the ratio of 3:1 as the running system. TLC plates were then exposed on screens and visualized on a Storm PhosphorImager (GE Healthcare, Wankesha, WI, USA). Radioactive spots were densitometrically quantified using either the ImageQuant (Molecular Dynamics, Sunnyvale, CA, USA) or the PDQuest Software (Bio-Rad). Steroid conversion was assessed by calculating percentage of radioactivity measured from a specific steroid hot spot in relation to the total radioactivity added to the reaction. To compare enzyme activities unstimulated versus stimulated, the conversion ratio of substrate to product was estimated.

RNA extraction and semi-quantitative RT-PCR

Total RNA from A and R cells was extracted using the Nucleospin RNA kit from Macherey Nagel (Oensingen, Switzerland). Normal human adrenal tissues were obtained from patients undergoing nephrectomy/adrenalectomy for kidney disorders with the permission of the local ethical committee. RNA from the human adrenal tissues was extracted using the Trizol method as described by the manufacturer (Gibco BRL). Extracted RNA was reverse transcribed to cDNA with the Improm RNA Transcriptase kit (Promega) using oligo dT primers or random hexamers (0.5 μ g/1 μ g RNA). Synthesized cDNA was amplified by Go-Taq polymerase (Promega) and specific primers (Table 1) in a final volume of 25–50 μ l. PCR conditions were 45 s at 95 °C, 45 s at 59 °C, 45 s at 72 °C for 28–35 cycles. Aliquots of RT-PCR products were electrophoresed on a 1.5% agarose gel, visualized with ethidium bromide, then scanned and quantified using the Alpha Imager 3400 (Alpha Innotech, San Leandro, CA, USA). The housekeeping genes glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18s ribosomal RNA (18SrRNA) served as internal controls. Primers were purchased from Microsynth (Balgach, Switzerland).

Table 1 List of primers used for conventional PCR and SYBR green real-time PCR assays

	Oligonucleotide sequences	NCBI accession number (5' start site for S, 3' for AS)
Name		
HSD17B3	S: 5'-GAGATCGAGCGGACTACAGG-3' AS: 5'-TACTCAGCTTCCAAGGCGTT-3'	BC034281 (303) BC034281 (643)
HSD17B5	S: 5'-CCAGTTGACTGCAGAGGACA-3' AS: 5'-TGTCTACCAGAAGCCCTGTG-3'	NM_003739 (927) NM_003739 (1080)
POR	S: 5'-TTCTGTTTTCGCTCATCGTG-3' AS: 5'-GGACTTCTACGACTGGCTGC-3'	BC034277 (110) BC034277 (493)
SULT2A1	S: 5'-TTCGTGATAAGGGATGAAGATGTA-3' AS: 5'-GAGATTCTCTGCCTGATGCA-3'	NM_003167 (167) NM_003167 (252)
b5	S: 5'-TGCACCACAAGGTGTACGAT-3' AS: 5'-AGGATGTCGGGCACTCTACA-3'	NM_148923 (200) NM_148923 (321)
CYP21A2	S: 5'-AGAGGGATCACATCGTGGAG-3' AS: 5'-CTCTGGACAGCTCCTGGAAG-3'	BC125182 (791) BC125182 (931)
18SrRNA	S: 5'-CTCAACACGGGAAACCTCAC-3' AS: 5'-AGACAAATCGTCCACCAAC-3'	NR_003286 (1247) NR_003286 (1364)
GAPDH	S: 5'-GAGTCAACGGATTGTCGT-3' AS: 5'-CTGAGAACGGGAAGCTTGTC-3'	BC083511 (72) BC083511 (256)
MC2R	S: 5'-GACCGCTACATCACCATCTT-3' AS: 5'-CCCTGGTTTTAGAATCCA-3'	NM_000529 (556) NM_000529 (1099)
AT1	S: 5'-TCAATGCTCAAGCCAGTGAC-3' AS: 5'-AGTGGCTGCTCAGTCTGTT-3'	NT_005612 (1283129) NT_005612 (1283321)

Real-time PCR

The mRNA levels of several genes involved in steroid production were quantitatively analyzed by real-time PCR using an ABI 7000 Sequence Detection System (Applied Biosystems, Foster City, CA, USA). Briefly, PCRs were performed in 96-well plates (Abgene, Epsom, UK) using cDNA prepared as described above (50 ng/20 µl), TaqMan Universal PCR Mix and 1 µl specific primers and probes obtained as assay-on-demand gene expression products (Applied Biosystems, www.appliedbiosystems.com). Alternatively, ABsolute QPCR SYBR Green Mix (ABgene) and specific primers were used (Table 1). GAPDH or 18SrRNA served as internal controls (reference genes). Correction for internal control, ΔC_t , was calculated as $C_{t\text{gene of interest}} - C_{t\text{reference gene}}$. Further, expression of each gene was calculated by the formula $2^{-\Delta C_t}$ (Livak & Schmittgen 2001).

Aldosterone, cortisol, and free testosterone measurements

Aldosterone measurement was done by competitive RIA on a gamma counter (Diagnostic Products Corporation, Los Angeles, CA, USA). Cortisol was measured by the competitive immunoassay Modular Analytics E170 (Roche Diagnostics). Measurement of the free testosterone was carried out by the Coat-A-Count solid phase ¹²⁵I RIA as described by the manufacturer (Diagnostic Products Corporation).

Plasmid constructs and luciferase reporter assays

Transient transfection of A and R cells with promoter constructs fused to a luciferase reporter gene was used to characterize promoter activities of CYP17, HSD3B2, b5, and SULT2A1. Constructs of the long (−3.7 kb) and short basal

promoters (−227 bp) of CYP17 in pMG3 vector, and of the HSD3B2 and the b5 promoters were described previously (Lin *et al.* 2001, Huang *et al.* 2005, Kempna *et al.* 2006). The reporter construct pCREluc, which is a luciferase expression vector under the control of 16 CRE, has been described earlier (Vaisse *et al.* 2000). For the construction of the −1400/+50 SULT2A1 promoter, the following primers were used: sense 5'-CTAGCTAGCTAGCTTTTGC-TCTTGCCCTGCTG-3' and antisense 5'-CCGCTC-GAGCTGCGTGGTGTGAGGGTTTC-3', the fragment was cloned into the pGL3 basic vector using *XhoI* and *NheI* restriction sites. For transient transfection, A and R cells were sub-cultured into 24-well plates at a density of about 250 000 cells/well. Prior to lipofectamin transfections, growth medium was replaced by the transfection medium DMEM/F12 containing 10% NuI serum supplemented with 0.1% SIT. Transfection was carried out for 6 h at 37 °C using 1.6 µl Lipofectamin2000 (Invitrogen), 10 ng *Renilla* luciferase in pRL-TK vector, and 0.5 µg specific *Firefly* luciferase reporter plasmids per well. After transfection, medium was replaced by growth medium. Some cells were stimulated with 100 nM ACTH or 0.5 mM 8Br-cAMP for 24 h. Finally, cells were lysed and assayed for Dual luciferase activities as described by the manufacturer (Promega). Co-transfection with *Renilla* luciferase was used to control for the transfection efficiency. As an additional control, cells were transfected with the empty vectors.

Statistical analysis

Data represent the mean of at least three independent experiments. Experimental variation is given as S.E.M. For statistics, to test whether data were parametric, *F*-test was

performed. As we found that the variance was not significantly different between the datasets, we used parametric statistical testing, namely the unpaired two-tailed Student's *t*-test (Prism 4.00, GraphPad software Inc., San Diego, CA, USA). Statistical significance: * $P \leq 0.05$ and ** $P \leq 0.01$.

Results

R cells produce more androgens than A cells

Human adrenal A and R cells are originally cultured in RPMI-1640 and DMEM/F12 media respectively. To compare the effect of different growth media on cell proliferation, both cell lines were grown in either of the two growth media. A and R cells showed higher proliferation in DMEM/F12 growth medium compared with RPMI-1640 medium. However, no difference between the two cell lines was observed (Fig. 2A). Hence, we decided to use DMEM/F12 medium as growth medium for both A and R cell lines for all further comparative experiments.

In order to characterize A and R cells, the steroid profile was assessed using [3 H]-Preg as a substrate for labeling the steroidogenesis. Interestingly, we found profound differences in the steroid profile of A and R cells (Fig. 2B and C). A cells mainly produced 17 α -hydroxyprogesterone (17OHProg) and 11-deoxycortisol (11DOC), which are precursors of glucocorticoids. In contrast, R cells produced not only 17OHProg, 17OHProg, and 11DOC, but also DHEA and androstenedione (Δ 4A), which are the adrenal androgens (Fig. 1). In addition, little progesterone (Prog) was detected in both cell lines, more in A than R cells. These results suggested that both adrenal cell lines produce similar amounts of mineralocorticoids and glucocorticoids, but that R cells produce markedly more adrenal androgens.

To confirm the observed differences in steroidogenesis of A and R cells, we measured the amount of mineralocorticoids, glucocorticoids, and androgens in the culture medium of both cell lines. A cells produced more aldosterone than R cells, while cortisol production was similar between the two cell lines

Figure 2 A and R cells have different steroid profiles even under the same growth conditions. (A) Proliferation of A and R cells. Equal numbers of cells were grown in either DMEM/F12 or RPMI-1640 medium for 72 h. Thereafter, cells were counted with a hemocytometer. Each counting was performed in duplicates. No difference was observed ($P=0.61$ for A cells grown in DMEM versus RPMI; $P=0.16$ for R cells in DMEM versus RPMI). (B and C) Steroid profiles of A and R cells. Cells were grown in DMEM/F12 medium. Steroidogenesis was labeled with [3 H]-Preg ($\sim 220\,000$ c.p.m./well) as substrate for 90 min. Steroids were extracted from the growth medium, separated on TLC plates, and then visualized. (B) Representative TLC. (C) Quantification of specific steroids produced by A and R cells assessed as percentage of total labeled steroids. (D) Production of mineralocorticoids, glucocorticoids, and androgens. Equal numbers of A and R cells were grown in DMEM/F12 medium supplemented with serum. After 50 h, supernatants of the cells were collected and concentrations of aldosterone, cortisol, and testosterone were determined. All data represent the mean $\pm$ s.e.m. of three independent experiments. * $P < 0.05$, ** $P < 0.01$.

(Fig. 2D). Moreover, testosterone was only measurable in the culture medium of R cells (Fig. 2D). Thus, it seems that A cells produce more mineralocorticoids, whereas R cells produce more androgens. Hence, high mineralocorticoid production in A cells may be due to high 3β HSDII activity and low P450c17 activities, as this combination of enzyme activities favors the conversion of Preg to Prog towards the mineralocorticoid pathway (Fig. 1). In contrast, high activities of 17α -hydroxylase and $17,20$ -lyase are essential for the production of androgens (Fig. 1). Therefore, higher androgen production in R cells may be due to higher activity of P450c17 and lower activity of 3β HSDII in these cells.

The expression of genes encoding steroidogenic enzymes may differ among human adrenocortical cell lines

Being predominantly interested in differences in the androgen production between the two adrenal cell lines, we investigated

the expression of the HSD3B2, CYP21A2, CYP17, SULT2A1, microsomal b5, and POR genes at the mRNA level performing qualitative RT-PCR. We found that all these genes were expressed in both A and R cells (data not shown). In order to compare the expression of the genes involved in the synthesis of androgens quantitatively, we performed real-time PCR experiments. A cells expressed higher levels of HSD3B2, b5, and SULT2A1 when compared with R cells (Fig. 3A and B). However, no differences in the expression of the CYP21A2, CYP17, and POR genes were observed (Fig. 3A and B). Higher expression of the HSD3B2 gene in A cells together with similar CYP21A2 gene expression in A and R cells support our hypothesis that 3β HSDII activity may be higher in A cells than in R cells. However, the observed gene expression data do not suggest higher P450c17 activities in R cells for which one would expect not only lower expression of HSD3B2, but also higher expression of CYP17, b5, and POR (Fig. 1).

Figure 3 Expression and promoter activity of genes involved in steroidogenesis. (A and B) Quantitative assessment of gene expression in A and R cells using Taqman (A) or SYBR green (B) real-time PCR assays. For Taqman 18SrRNA and for SYBR green GAPDH served as internal control. Data represent the mean $\pm$ S.E.M. of 4–6 independent experiments. * $P < 0.05$, ** $P < 0.01$. (C) Semiquantitative assessment of the expression of the HSD17B3 and HSD17B5 genes in two different human adrenal tissues (hAd1 and hAd2), and in A and R cells (internal control 18SrRNA). Agarose gel depicting one representative experiment, confirmed three times independently. (D) Activities of different promoter luciferase reporter constructs of steroidogenic genes in A and R Cells. Cells were transfected with promoter constructs as indicated. PRL-TK served as an internal control for transfection efficiency, and transfection of empty vectors served as an additional control. Data represent the mean $\pm$ S.E.M. of 4–6 independent experiments. * $P < 0.05$, ** $P < 0.01$.

Human adrenal tumors often produce androgens (Geissler *et al.* 1994, Miller 2002, Allolio *et al.* 2004). We were able to detect testosterone in the cell culture medium of R cells but not A cells. Activities of the 17 α -hydroxysteroid dehydrogenase type III (17 β HSDIII; *HSD17B3*) and/or 17 α -hydroxysteroid dehydrogenase type V (17 β HSDV; *HSD17B5*) enzymes are required for the production of testosterone from Δ 4A (Geissler *et al.* 1994, Dufort *et al.* 1999). Therefore, we studied the expression of the *HSD17B3* and *HSD17B5* genes in A and R cells and in human adrenal tissues (Fig. 3C). We confirmed the absence of the *HSD17B3* gene expression in two independent tissue samples of normal human adrenals. However, we found that both human adrenal cell lines express the *HSD17B3* gene. Slightly higher expression of the *HSD17B3* gene in R cells may partially explain the higher levels of testosterone measured in cell culture medium of R cells. Expression of the *HSD17B5* gene was detected in two independent human adrenal tissues as well as in both A and R cells (Fig. 3C). As the expression of *HSD17B5* was similar between the cell lines, the activity of 17 β HSDV may not explain the observed difference in testosterone production.

To compare transcriptional activation of different steroid genes in both cell lines, A and R cells were transiently transfected with specific promoter-luciferase reporter constructs. We found significantly higher activities of the -1050 bp *HSD3B2*, -227 bp *CYP17*, -3.7 kb *CYP17* and -1300 bp *b5* promoter constructs in R cells (Fig. 3D). The activity of the -1400 bp *SULT2A1* promoter reporter was similar in both cell lines (Fig. 3D). From our previous observations from the gene expression patterns (Figs 2 and 3A–C), we would have suggested similar activity of the *CYP17* promoter in A and R cells, but higher activities of *HSD3B2*, *b5*, and *SULT2A1* promoters in A cells. Yet data obtained from transient transfection of reporter constructs in cells should be interpreted with caution for many reasons such as specific choice of the promoter region or the vectors used (Kempna *et al.* 2006).

Activities of 3 β HSDII and P450c17 in A and R cells

High expression of the *HSD3B2* gene in A cells at the mRNA level suggests high activity of 3 β HSDII. To confirm this hypothesis, we performed functional studies. Cells were incubated with [3 H]-DHEA, a substrate for 3 β HSDII that is converted to Δ 4A (Fig. 1). Our studies revealed twofold higher activity of the 3 β HSDII enzyme in A cells than in R cells (Fig. 4A). This finding is consistent with the gene expression studies (Fig. 3A). Thus, A cells express more *HSD3B2* than R cells resulting in a higher 3 β HSDII activity, which may lead steroid production towards the mineralocorticoid pathway and result in lower androgen production.

Further, functional studies were performed to compare the 17 α -hydroxylase and 17,20-lyase activities of P450c17 in A and R cells. To assess P450c17 enzyme activity specifically, 3 β HSD was blocked by trilostane. Cells were incubated with [3 H]-Preg to study both the 17 α -hydroxylase and 17,20-lyase activities, or with [3 H]-17OHPreg to study the 17,20-lyase activity alone (Fig. 1). Using [3 H]-Preg as a substrate, we did

not observe differences in 17 α -hydroxylase or 17,20-lyase activities between A and R cells (Fig. 4B). In contrast, using [3 H]-17OHPreg as a direct substrate for the 17,20-lyase, we found significantly higher 17,20-lyase activity in R cells when compared with A cells (Fig. 4C). This difference in 17,20-lyase activity was not observed using [3 H]-Preg as a substrate because of the two steps in the reaction and the shorter incubation time (90 vs 480 min). Similar *CYP17* gene expression and 17 α -hydroxylase activity for A and R cells is in line with equal production of glucocorticoids in both cell lines. Furthermore, higher 17,20-lyase activity in R cells explains higher androgen production in R cells.

Expression of AT1 and MC2R, and the ACTH/cAMP signaling

Expression of the AT1 receptor is described in R cells but not investigated in A cells (Rainey *et al.* 2004). We found that the AT1 receptor is expressed in both cell lines but at lower levels in A cells (Fig. 5A). R cells are known to express low levels of the ACTH receptor (MC2R; Mountjoy *et al.* 1994), while no data are available for A cells. Thus, we studied MC2R expression in A and R cells at the mRNA level. Low expression of the MC2R gene was observed in both cell lines compared with the abundantly expressed *POR* gene or the housekeeping gene *18S rRNA* (Fig. 5A). Further, we investigated the effect of ACTH and 8Br-cAMP stimulation on A and R cells. The classic ACTH/cAMP signaling pathway exerts its effect through PKA activation resulting in phosphorylation of CRE-binding proteins, which will then bind to CRE in targeted genes. Therefore, cells were transiently transfected with a pCREluc construct to assess transcriptional activation by ACTH or 8Br-cAMP stimulation in both cell lines. Eight Br-cAMP stimulated pCREluc activity in both cell lines about 9- to 15-fold (Fig. 5B). Generally, the activation of the pCREluc reporter tended to be higher in R cells but this difference did not reach statistical significance. By contrast, only a two- to threefold stimulation was observed by ACTH stimulation in both cell lines (Fig. 5B). This result may be explained by low expression of the MC2R in A and R cells. Alternatively, studies by others suggested that poor stimulation by ACTH may be due to impaired trafficking of the receptor from the endoplasmic reticulum to the cell surface (Noon *et al.* 2002).

Steroid production in A and R cells after stimulation with ACTH, 8Br-cAMP, and AngII

Previous studies have shown that H295 cells are unable to respond to ACTH or AngII stimulation (Staels *et al.* 1993). By contrast, steroidogenesis of R cells is reported to be regulated by AngII (Bird *et al.* 1993, 1994), potassium (Rainey *et al.* 1994, Pezzi *et al.* 1996, Hilbers *et al.* 1999), and ACTH (Rainey *et al.* 1993, Staels *et al.* 1993, Mountjoy *et al.* 1994). These investigators observed that the responsiveness of R cells is highly dependent on growth conditions (Rainey *et al.* 2004). However, no data are available comparing A and

Figure 4 P450c17 and 3βHSDII activities. (A) Activity of 3βHSDII in A versus R cells. Cells were incubated with [3 H]-DHEA for 60 min. Steroids were then extracted from the growth medium and separated on TLC plates. Representative TLC plate and quantitative analysis of three independent experiments. (B and C) Activities of 17 α -hydroxylase and 17,20-lyase. Cells were incubated with [3 H]-Preg (B) or [3 H]-17OHPreg (C) (~220 000 c.p.m./well) for 90 or 480 min to assess both the 17 α -hydroxylase and 17,20-lyase activities (B) or the 17,20-lyase activity alone (C). Trilostane was used to block 3βHSD activity. Representative TLCs and quantitative analysis. Data represent the mean $\pm$ s.e.m. of three independent experiments. * P <0.05, ** P <0.01.

Figure 5 Expression of MC2R and AT1, and ACTH/cAMP stimulation. (A) Agarose gel showing the expression of the MC2R and the AT1 gene in A and R cells after 35 cycles of PCR. For comparison, expression of POR is depicted. Expression of 18S rRNA served as internal control. Shown is one representative experiment, confirmed three times independently. (B) Activity of the pCRELuc reporter in A and R cells after stimulation with either 0.5 mM 8Br-cAMP or 100 nM ACTH for 24 h. Cells were transiently transfected with the pCRELuc reporter, and luciferase activity was assessed by the Dual luciferase assay (Promega). No difference was observed ($P=0.39$ for cAMP stimulation in A versus R cells; $P=0.25$ for ACTH stimulation in A versus R cells). Data represent the mean $\pm$ S.E.M. of 3–4 independent experiments. * $P<0.05$, ** $P<0.01$.

R cells under the exactly same growth and stimulation conditions. Therefore, we stimulated both cell lines under standardized growth conditions with either ACTH, 8Br-cAMP, or AngII, and assessed aldosterone, cortisol, and testosterone production, as well as steroid profiles. We found that ACTH stimulation did not change the production of aldosterone, cortisol, and testosterone in A and R cells (Fig. 6A). By contrast, stimulation with 8Br-cAMP stimulated the production of all measured steroids significantly (Fig. 6B). While this stimulatory effect of cAMP was moderate for R cells, it was more than tenfold for A cells.

AngII stimulation did not change steroid production in A cells, but increased aldosterone and cortisol production in R cells (Fig. 6C). Testosterone production seemed to decrease after AngII stimulation, but this change did not reach statistical significance ($P=0.08$). All these effects of ACTH, cAMP, and AngII on steroidogenesis in A and R cells were confirmed by steroid profiling experiments (Fig. 7A). While ACTH stimulation did not change the steroid profile of A and R cells (data not shown), 8Br-cAMP stimulated steroidogenesis in both (Fig. 7B). Substrate to product calculations from these data suggested a stimulatory effect of cAMP on the enzyme activities of P450c21 (A cells, 250%; R cells, 130% of unstimulated) and P450c17 (A cells, 190%; R cells, 190% of unstimulated). AngII stimulation did not change the steroid profile of A cells, but increased 11DOC and decreased 17OHPreg as well as DHEA production in R cells (Fig. 7C). Accordingly, substrate to product calculations suggested a stimulatory effect of AngII on the activities of P450c21 (130% of unstimulated) and 3 β HSDII (150% of unstimulated) but an inhibitory effect on P450c17 activity (70% of unstimulated).

Discussion

Most molecular studies of human adrenal steroidogenesis use human adrenocortical A and R cells as cellular models for the adrenal cortex. As both cell lines originate from the adrenocortical H295 tumor cell line (Gazdar *et al.* 1990, Staels *et al.* 1993), it is generally believed that those three cell lines have the same characteristics concerning steroidogenesis. Several investigators reported expression of key steroidogenic genes for all three cell lines and described the steroid profile of these cell lines (Fig. 1). However, no data of a direct comparison between these cell lines were available. We show in this study that human adrenocortical A and R cells differ significantly in their steroidogenesis. Overall, A cells produce more mineralocorticoids, while R cells produce more adrenal androgens (DHEA, $\Delta 4A$) and testosterone. Interestingly, we found equal cortisol production for both A and R cells. The cell line-specific 'signatures' are mainly explained by higher expression and activity of 3 β HSDII in A cells and lower expression and activity of 3 β HSDII in combination with higher activity of 17,20 lyase in R cells. These findings underline the importance of the balance between 3 β HSDII and P450c17 activities for steroidogenesis. First, the 17 α -hydroxylase and 17,20 lyase activities of P450c17 are the qualitative regulators of steroidogenesis, which are essential for glucocorticoid and androgen production (Miller 1988). To favor androgen production over glucocorticoid production, the 17,20 lyase activity of P450c17 is enhanced by cytochrome b5, POR, and phosphorylation (Lin *et al.* 1993, Zhang *et al.* 1995, Pandey *et al.* 2003, Pandey & Miller 2005). Second, 3 β HSDII and P450c17 have common substrates and are therefore competitors (Fig. 1). In previous competition experiments using recombinantly produced proteins expressed in yeast, we showed that decreased 3 β HSDII activity in relation to P450c17 activities will shift steroidogenesis towards enhanced

Figure 6 Production of mineralocorticoids, glucocorticoids, and androgens after stimulation with ACTH, 8Br-cAMP, and AngII. Equal numbers of A and R cells were grown in DMEM/F12 medium, initially with serum for 24 h, then serum-free for 24 h before stimulation. After stimulation supernatants of the cells were collected and concentrations of aldosterone, cortisol, and testosterone were measured. (A) Summary of the results for A cells and R cells 24 h after stimulation with 100 nM ACTH. (B) Summary of the results 24 h after stimulation with 0.5 mM 8Br-cAMP. (C) Summary of the results 72 h after stimulation with 100 nM AngII. Note that aldosterone was only detected after 72 h incubation in serum-free medium. All data represent mean $\pm$ s.e.m. of three independent experiments. * $P < 0.05$.

Figure 7 Steroid profiles after stimulation with ACTH, 8Br-cAMP, and AngII. Equal numbers of A and R cells were grown in DMEM/F12 medium, initially with serum for 24 h, then serum-free. Cells were stimulated and steroidogenesis was labeled with [3 H]-Prog ($\sim 220\,000$ c.p.m./well). Steroids were extracted from the supernatants, separated on TLC plates, and then visualized. (A) Representative TLC. (B and C) Quantification of specific steroids produced by A and R cells assessed as percentage of total labeled steroids after stimulation with 0.5 mM 8Br-cAMP for 24 h (B) or 100 nM AngII for 72 h (C). Data for ACTH after 24-h stimulation are not shown as no change was observed upon stimulation for both cell lines. Data represent mean $\pm$ S.E.M. of 2–3 independent experiments, each performed in duplicates. * $P < 0.05$.

$\Delta 4$ A production from Preg, thus favoring the sex steroid over the mineralocorticoid pathway (Fluck *et al.* 2005). Thus, higher androgen production in R cells despite equal expression of P450c17 for A and R cells may be explained—at least in part—by lower expression of HSD3B2 in R cells. This concept of higher P450c17/3 β HSDII ratio stressing androgen production was suggested previously by Conley & Bird (1997) and supported by the observation that adrenarche is characterized by a drop in HSD3B2 expression (Gell *et al.* 1996, 1998).

Physiologically, human adrenal mineralocorticoid production is acutely regulated by AngII and glucocorticoid production by ACTH. Both A and R cells do not express high levels of the ACTH receptor. Accordingly, we found that steroidogenesis of both cell lines may not be stimulated by ACTH. By contrast, stimulation with cAMP directly increased steroid production in both cell lines indicating that the cAMP signaling is intact. We found higher expression of the AT1 receptor in R cells and AngII was able to stimulate steroidogenesis in R cells but not A cells, although basally A cells produced more aldosterone than R cells. This suggests that the regulation of AngII on aldosterone production is disrupted in A cells. However, from our data we cannot say at what level this signaling is disturbed.

Our finding that A and R cells are different may have two major implications. First, research data obtained from studies using one cell line may not apply to the other cell line. Second, studying the difference between A and R cells in further detail may help us gain important information on how androgen production is regulated.

To understand the exact regulation of androgen biosynthesis including the concerted activities of 3 β HSDII and P450c17 is important, but little is known. Recent studies have suggested that the expression and activity of 3 β HSDII from weeks 7 to 12 of fetal life is essential for safeguarding the sexual development of a female fetus (Goto *et al.* 2006). Outside this fetal time window, the human fetal adrenals lack 3 β HSDII activity until about 23 weeks, gestation (Narasaka *et al.* 2001), and therefore produce high amounts of DHEAS. After birth, adrenal androgen production is shut down with the change from the fetal to the postnatal adrenal cortex, which consists of the ZG and ZF producing relatively constant amounts of mineralocorticoids and glucocorticoids throughout life. In contrast, adrenal androgen production starts after the development of the ZR (adrenarche), reaches a sex-specific plateau around age 30 years, and decreases constantly thereafter (adrenopause; Rainey *et al.* 2002). Immunohistochemical studies of the human adrenal cortex revealed that during adrenarche, the ZR shows specific changes in the expression of b5, SULT2A1, and HSD3B2 (Suzuki *et al.* 2000). SULT2A1 and b5 are markedly enhanced and HSD3B2 is grossly diminished after adrenarche. In addition, epidemiologic data suggest that girls with premature exaggerated adrenarche have a significantly higher risk to develop the polycystic ovary syndrome (PCOS), the most common hyperandrogenic disorder affecting 5–10% of women (Escobar-Morreale *et al.* 2005). Elevated serum

androgens in PCOS have been shown to originate from both the ovaries and the adrenals (Lachelin *et al.* 1982, Barnes *et al.* 1989, Ehrmann *et al.* 1992). Studies of the steroid metabolism of ovarian theca cells isolated from PCOS women demonstrated enhanced 3 β HSDII and P450c17 activities in PCOS compared with normal cells (Nelson *et al.* 1999, 2001). Additionally, northern blot and quantitative real-time PCR analyses revealed an increase in the abundance of the mRNAs for CYP17, HSD3B2, and CYP11A1 in PCOS compared with normal cells, but no difference for StAR and 17 β HSDV was observed (Nelson *et al.* 1999, 2001). In addition, there is experimental evidence that enhanced expression of b5 together with abundance in POR as well as phosphorylation of P450c17 lead to an increase in 17,20 lyase activity (Lin *et al.* 1993, Zhang *et al.* 1995, Pandey *et al.* 2003, Pandey & Miller 2005). Considering all these research data describing characteristics of androgen production, our comparison between A and R cells suggests that R cells are more 'androgenic' than A cells. The profile of R cells may be compared with the steroid profile of adrenarche for low expression and activity of 3 β HSDII and high activity of 17,20 lyase. However, R cells do not have higher expression of b5 or SULT2A1 as found in the ZR after adrenarche. Similarly, R cells share the steroid profile of PCOS with higher activity of 17,20 lyase; but in contrast to PCOS expression of HSD3B2 and P450c17 is not enhanced when compared with A cells.

A and R cells are tumor cells. The original H295 cell line was isolated from an adrenocortical carcinoma of a middle-aged woman suffering from elevated serum aldosterone, cortisol, and androgens (Gazdar *et al.* 1990). The normal human adrenal cortex seems not to produce relevant amounts of testosterone (Miller 2002). However, 50% of human adrenocortical tumors are hormonally active and about half of them produce testosterone (Allolio *et al.* 2004). Original H295 cells are reported to secrete large quantities of adrenal androgens and smaller amounts of testosterone (Gazdar *et al.* 1990). Therefore, it has been suggested that the major steroid pathway in H295 cells is androgen formation, although mineralocorticoids and glucocorticoids were also detected (Gazdar *et al.* 1990). However, the steroid profile of H295 was reported to be influenced by serum conditions (Gazdar *et al.* 1990). Thus, the observed differences between A and R cells may have been caused by different growth conditions and handling in several laboratories over time. Both cell lines were subcultured and manipulated to grow adherently in different laboratories under different conditions (Gazdar *et al.* 1990, Rodriguez *et al.* 1997). From our studies using the same growth medium for all experiments, we can exclude that factors present in the growth medium were responsible for the observed differences. In our experience, passage number may play a role for both cell lines beyond 30–35 passages. R cells seem to resemble their parental H295 cells more than A cells. For both H295 and R cells, testosterone production has been reported (Gazdar *et al.* 1990, Danesi *et al.* 1996, Hecker *et al.* 2006). We were able to detect testosterone in the growth medium of R cells but not A cells. Two enzymes 17 β HSDIII and 17 β HSDV are involved in the

synthesis of testosterone from $\Delta 4A$ or DHEA (Fig. 1; Miller 2002). Normally, the HSD17B3 gene is only expressed in the human testis (Geissler *et al.* 1994, Miller 2002), whereas the HSD17B5 gene is expressed in several human tissues such as the testes and the adrenals (Dufort *et al.* 1999). To date no data was available regarding the expression of these genes in A and R cells. We show that both HSD17B3 and HSD17B5 genes are expressed in A as well as R cells. Therefore, the difference in testosterone production between A and R cells may be mainly explained by lower 3β HSDII and higher 17,20 lyase activity in R cells leading to an increase in substrate availability for 17β HSDIII/V.

In conclusion, human adrenocortical R cells produce more androgens than A cells even though they derive from the same tumor. Thus, studying the differences between these two cell lines may serve as an excellent model to find key regulators of androgen production.

Acknowledgements

We thank Prof. Walter L Miller for kindly providing the NCI-H295A cells for our study, and Dr Chantal Cripe-Mamie for proofreading the manuscript. This work was supported by grants of the Swiss National Science Foundation (3232B0103178/9 and 320000-116299) and by the Foundation Valeria and Ettore Rossi, Bern, Switzerland. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Allolio B, Hahner S, Weismann D & Fassnacht M 2004 Management of adrenocortical carcinoma. *Clinical Endocrinology* **60** 273–287.
- Auchus RJ, Lee TC & Miller WL 1998 Cytochrome b5 augments the 17,20-lyase activity of human P450c17 without direct electron transfer. *Journal of Biological Chemistry* **273** 3158–3165.
- Barnes RB, Rosenfield RL, Burstein S & Ehrmann DA 1989 Pituitary–ovarian responses to nafarelin testing in the polycystic ovary syndrome. *New England Journal of Medicine* **320** 559–565.
- Bird IM, Hanley NA, Word RA, Mathis JM, McCarthy JL, Mason JI & Rainey WE 1993 Human NCI-H295 adrenocortical carcinoma cells: a model for angiotensin-II-responsive aldosterone secretion. *Endocrinology* **133** 1555–1561.
- Bird IM, Mason JI & Rainey WE 1994 Regulation of type 1 angiotensin II receptor messenger ribonucleic acid expression in human adrenocortical carcinoma H295 cells. *Endocrinology* **134** 2468–2474.
- Campbell B 2006 Adrenarche and the evolution of human life history. *American Journal of Human Biology* **18** 569–589.
- Conley AJ & Bird IM 1997 The role of cytochrome P450 17α -hydroxylase and 3β -hydroxysteroid dehydrogenase in the integration of gonadal and adrenal steroidogenesis via the $\Delta 5$ and $\Delta 4$ pathways of steroidogenesis in mammals. *Biology of Reproduction* **56** 789–799.
- Cutler GB, Glenn M, Bush M, Hodgen GD, Graham CE & Loriaux DL 1978 Adrenarche: a survey of rodents, domestic animals, and primates. *Endocrinology* **103** 2112–2118.
- Danesi R, La Rocca RV, Cooper MR, Ricciardi MP, Pellegrini A, Soldani P, Kragel PJ, Paparelli A, Del Tacca M & Myers CE 1996 Clinical and experimental evidence of inhibition of testosterone production by suramin. *Journal of Clinical Endocrinology and Metabolism* **81** 2238–2246.
- Dardis A & Miller WL 2003 Dexamethasone does not exert direct intracellular feedback on steroidogenesis in human adrenal NCI-H295A cells. *Journal of Endocrinology* **179** 131–142.
- Dhom G 1973a Morphology of the human adrenarche. *Acta Endocrinologica. Supplementum* **173** 27.
- Dhom G 1973b The prepuberal and puberal growth of the adrenal (adrenarche). *Beiträge zur Pathologie* **150** 357–377.
- Dufort I, Rheault P, Huang XF, Soucy P & Luu-The V 1999 Characteristics of a highly labile human type 5 17β -hydroxysteroid dehydrogenase. *Endocrinology* **140** 568–574.
- Ehrmann DA, Rosenfield RL, Barnes RB, Brigell DF & Sheikh Z 1992 Detection of functional ovarian hyperandrogenism in women with androgen excess. *New England Journal of Medicine* **327** 157–162.
- Escobar-Morreale HF, Luque-Ramirez M & San Millan J 2005 The molecular-genetic basis of functional hyperandrogenism and the polycystic ovary syndrome. *Endocrine Reviews* **26** 251–282.
- Fluck CE & Miller WL 2004 GATA-4 and GATA-6 modulate tissue-specific transcription of the human gene for P450c17 by direct interaction with Sp1. *Molecular Endocrinology* **18** 1144–1157.
- Fluck CE, Yaworsky DC & Miller WL 2005 Effects of anticonvulsants on human p450c17 (17α -hydroxylase/ $17,20$ lyase) and 3β -hydroxysteroid dehydrogenase type 2. *Epilepsia* **46** 444–448.
- Forti FL, Dias MH & Armelin HA 2006 ACTH receptor: ectopic expression, activity and signaling. *Molecular and Cellular Biochemistry* **293** 147–160.
- Gazdar AF, Oie HK, Shackleton CH, Chen TR, Triche TJ, Myers CE, Chrousos GP, Brennan MF, Stein CA & La Rocca RV 1990 Establishment and characterization of a human adrenocortical carcinoma cell line that expresses multiple pathways of steroid biosynthesis. *Cancer Research* **50** 5488–5496.
- Geissler WM, Davis DL, Wu L, Bradshaw KD, Patel S, Mendonca BB, Elliston KO, Wilson JD, Russell DW & Andersson S 1994 Male pseudohermaphroditism caused by mutations of testicular 17β -hydroxysteroid dehydrogenase 3. *Nature Genetics* **7** 34–39.
- Gell JS, Atkins B, Margraf L, Mason JI, Sasano H, Rainey WE & Carr BR 1996 Adrenarche is associated with decreased 3β -hydroxysteroid dehydrogenase expression in the adrenal reticularis. *Endocrine Research* **22** 723–728.
- Gell JS, Carr BR, Sasano H, Atkins B, Margraf L, Mason JI & Rainey WE 1998 Adrenarche results from development of a 3β -hydroxysteroid dehydrogenase-deficient adrenal reticularis. *Journal of Clinical Endocrinology and Metabolism* **83** 3695–3701.
- Goodman HM 2003. Adrenal glands. In: *Basic Medical Endocrinology*, pp 111–163. Academic Press, Elsevier.
- Goto M, Piper Hanley K, Marcos J, Wood PJ, Wright S, Postle AD, Cameron IT, Mason JI, Wilson DI & Hanley NA 2006 In humans, early cortisol biosynthesis provides a mechanism to safeguard female sexual development. *Journal of Clinical Investigation* **116** 953–960.
- Havelock JC, Auchus RJ & Rainey WE 2004 The rise in adrenal androgen biosynthesis: adrenarche. *Seminars in Reproductive Medicine* **22** 337–347.
- Hecker M, Newsted JL, Murphy MB, Higley EB, Jones PD, Wu R & Giesy JP 2006 Human adrenocarcinoma (H295R) cells for rapid *in vitro* determination of effects on steroidogenesis: hormone production. *Toxicology and Applied Pharmacology* **217** 114–124.
- Hilbers U, Peters J, Bornstein SR, Correa FM, Johren O, Saavedra JM & Ehrhart-Bornstein M 1999 Local renin–angiotensin system is involved in K^+ -induced aldosterone secretion from human adrenocortical NCI-H295 cells. *Hypertension* **33** 1025–1030.
- Huang N, Pandey AV, Agrawal V, Reardon W, Lapunzina PD, Mowat D, Jabs EW, Van Vliet G, Sack J, Fluck CE *et al.* 2005 Diversity and function of mutations in P450 oxidoreductase in patients with Antley–Bixler syndrome and disordered steroidogenesis. *American Journal of Human Genetics* **76** 729–749.
- John ME, John MC, Boggaram V, Simpson ER & Waterman MR 1986a Transcriptional regulation of steroid hydroxylase genes by corticotropin. *PNAS* **83** 4715–4719.
- John ME, Simpson ER, Waterman MR & Mason JI 1986b Regulation of cholesterol side-chain cleavage cytochrome P-450 gene expression in adrenal cells in monolayer culture. *Molecular and Cellular Endocrinology* **45** 197–204.

- Kempna P, Hofer G, Mullis PE & Fluck CE 2007 Pioglitazone inhibits androgen production in NCI-H295R cells by regulating gene expression of CYP17 and HSD3B2. *Molecular Pharmacology* **71** 787–798.
- Lachelin GC, Judd HL, Swanson SC, Hauck ME, Parker DC & Yen SS 1982 Long term effects of nightly dexamethasone administration in patients with polycystic ovarian disease. *Journal of Clinical Endocrinology and Metabolism* **55** 768–773.
- Lam CW, Arlt W, Chan CK, Honour JW, Lin CJ, Tong SF, Choy KW & Miller WL 2001 Mutation of proline 409 to arginine in the meander region of cytochrome P450c17 causes severe 17 α -hydroxylase deficiency. *Molecular Genetics and Metabolism* **72** 254–259.
- Lin D, Black SM, Nagahama Y & Miller WL 1993 Steroid 17 α -hydroxylase and 17,20-lyase activities of P450c17: contributions of serine106 and P450 reductase. *Endocrinology* **132** 2498–2506.
- Lin CJ, Martens JW & Miller WL 2001 NF-1C, Sp1, and Sp3 are essential for transcription of the human gene for P450c17 (steroid 17 α -hydroxylase/17,20 lyase) in human adrenal NCI-H295A cells. *Molecular Endocrinology* **15** 1277–1293.
- Livak KJ & Schmittgen TD 2001 Analysis of relative gene expression data using real-time quantitative PCR and the $2(-\Delta\Delta C(T))$ method. *Methods* **25** 402–408.
- Mahesh VB & Brann DW 1992 Interaction between ovarian and adrenal steroids in the regulation of gonadotropin secretion. *Journal of Steroid Biochemistry and Molecular Biology* **41** 495–513.
- Miller WL 1988 Molecular biology of steroid hormone synthesis. *Endocrine Reviews* **9** 295–318.
- Miller WL 2002 Disorders of androgen biosynthesis. *Seminars in Reproductive Medicine* **20** 205–216.
- Miller WL, Auchus RJ & Geller DH 1997 The regulation of 17,20 lyase activity. *Steroids* **62** 133–142.
- Mountjoy KG, Bird IM, Rainey WE & Cone RD 1994 ACTH induces up-regulation of ACTH receptor mRNA in mouse and human adrenocortical cell lines. *Molecular and Cellular Endocrinology* **99** R17–R20.
- Narasaka T, Suzuki T, Moriya T & Sasano H 2001 Temporal and spatial distribution of corticosteroidogenic enzymes immunoreactivity in developing human adrenal. *Molecular and Cellular Endocrinology* **174** 111–120.
- Nelson VL, Legro RS, Strauss JF & McAllister JM 1999 Augmented androgen production is a stable steroidogenic phenotype of propagated theca cells from polycystic ovaries. *Molecular Endocrinology* **13** 946–957.
- Nelson VL, Qin KN, Rosenfield RL, Wood JR, Penning TM, Legro RS, Strauss JF, III & McAllister JM 2001 The biochemical basis for increased testosterone production in theca cells propagated from patients with polycystic ovary syndrome. *Journal of Clinical Endocrinology and Metabolism* **86** 5925–5933.
- Noon LA, Franklin JM, King PJ, Goulding NJ, Hunyady L & Clark AJ 2002 Failed export of the adrenocorticotrophin receptor from the endoplasmic reticulum in non-adrenal cells: evidence in support of a requirement for a specific adrenal accessory factor. *Journal of Endocrinology* **174** 17–25.
- Pandey AV & Miller WL 2005 Regulation of 17,20 lyase activity by cytochrome b5 and by serine phosphorylation of P450c17. *Journal of Biological Chemistry* **280** 13265–13271.
- Pandey AV, Mellon SH & Miller WL 2003 Protein phosphatase 2A and phosphoprotein SET regulate androgen production by P450c17. *Journal of Biological Chemistry* **278** 2837–2844.
- Pezzi V, Clark BJ, Ando S, Stocco DM & Rainey WE 1996 Role of calmodulin-dependent protein kinase II in the acute stimulation of aldosterone production. *Journal of Steroid Biochemistry and Molecular Biology* **58** 417–424.
- Rainey WE, Bird IM, Sawetawan C, Hanley NA, McCarthy JL, McGee EA, Wester R & Mason JI 1993 Regulation of human adrenal carcinoma cell (NCI-H295) production of C19 steroids. *Journal of Clinical Endocrinology and Metabolism* **77** 731–737.
- Rainey WE, Bird IM & Mason JI 1994 The NCI-H295 cell line: a pluripotent model for human adrenocortical studies. *Molecular and Cellular Endocrinology* **100** 45–50.
- Rainey WE, Carr BR, Sasano H, Suzuki T & Mason JI 2002 Dissecting human adrenal androgen production. *Trends in Endocrinology and Metabolism* **13** 234–239.
- Rainey WE, Saner K & Schimmer BP 2004 Adrenocortical cell lines. *Molecular and Cellular Endocrinology* **228** 23–38.
- Rodriguez H, Hum DW, Staels B & Miller WL 1997 Transcription of the human genes for cytochrome P450csc and P450c17 is regulated differently in human adrenal NCI-H295 cells than in mouse adrenal Y1 cells. *Journal of Clinical Endocrinology and Metabolism* **82** 365–371.
- Sewer MB & Waterman MR 2001 Insights into the transcriptional regulation of steroidogenic enzymes and StAR. *Reviews in Endocrine and Metabolic Disorders* **2** 269–274.
- Sewer MB & Waterman MR 2002 cAMP-dependent transcription of steroidogenic genes in the human adrenal cortex requires a dual-specificity phosphatase in addition to protein kinase A. *Journal of Molecular Endocrinology* **29** 163–174.
- Sewer MB & Waterman MR 2003 cAMP-dependent protein kinase enhances CYP17 transcription via MKP-1 activation in H295R human adrenocortical cells. *Journal of Biological Chemistry* **278** 8106–8111.
- Staels B, Hum DW & Miller WL 1993 Regulation of steroidogenesis in NCI-H295 cells: a cellular model of the human fetal adrenal. *Molecular Endocrinology* **7** 423–433.
- Stocco DM, Wang X, Jo Y & Manna PR 2005 Multiple signaling pathways regulating steroidogenesis and steroidogenic acute regulatory protein expression: more complicated than we thought. *Molecular Endocrinology* **19** 2647–2659.
- Suzuki T, Sasano H, Takeyama J, Kaneko C, Freije WA, Carr BR & Rainey WE 2000 Developmental changes in steroidogenic enzymes in human postnatal adrenal cortex: immunohistochemical studies. *Clinical Endocrinology* **53** 739–747.
- Vaisse C, Clement K, Durand E, Hercberg S, Guy-Grand B & Froguel P 2000 Melanocortin-4 receptor mutations are a frequent and heterogeneous cause of morbid obesity. *Journal of Clinical Investigation* **106** 253–262.
- Waterman MR & Keeney DS 1996 Signal transduction pathways combining peptide hormones and steroidogenesis. *Vitamins and Hormones* **52** 129–148.
- Zhang LH, Rodriguez H, Ohno S & Miller WL 1995 Serine phosphorylation of human P450c17 increases 17,20-lyase activity: implications for adrenarche and the polycystic ovary syndrome. *PNAS* **92** 10619–10623.

Received in final form 17 September 2007

Accepted 27 September 2007

Made available online as an Accepted Preprint
27 September 2007