Expression of vasopressin receptors in ACTH-independent macronodular bilateral adrenal hyperplasia causing Cushing's syndrome: molecular, immunohistochemical and pharmacological correlates

E Louiset¹, V Contesse¹, L Groussin², D Cartier¹, C Duparc¹, V Perraudin¹, J Bertherat² and H Lefebvre^{1,3}

¹INSERM U413, Laboratory of Cellular and Molecular Neuroendocrinology, European Institute for Peptide Research (IFRMP 23), University of Rouen, 76821 Mont-Saint-Aignan, France

²INSERM U567, CNRS UMR8104, Department of Endocrinology-Metabolism-Cancer, Institut Cochin, Université Paris V-René Descartes, 75014 Paris, France

³Department of Endocrinology, Institute for Biomedical Research, Rouen University Hospital, 76031 Rouen Cedex, France

(Correspondence should be addressed to H Lefebvre; Email: herve.lefebvre@chu-rouen.fr)

Abstract

Cortisol secretion in ACTH-independent macronodular adrenal hyperplasia (AIMAH) causing Cushing's syndrome can be controlled by illegitimate receptors. The aim of the present study was to characterize the molecular, immunohistochemical, and pharmacological profiles of vasopressin receptors in cells derived from three patients with AIMAH (H1–H3), in order to evaluate the role of ectopic vasopressin receptors in the physiopathology of hypercortisolism. Expression of mRNAs encoding the vasopressin receptor types (V_{1a} , V_{1b} , and V_2) were analyzed by RT-PCR in adrenal tissues. The presence of V_{1a} and V_2 receptors was studied by immunohistochemistry on adrenal sections. The pharmacological profiles of vasopressin receptors involved in the control of cortisol secretion were investigated using the V_{1a} receptor antagonist SR49059 and the V_2 receptor agonist [deamino-Cys¹, Val⁴, D-Arg⁸]-vasopressin on cultured

cells. The V_{1a} receptor protein was present and functional in H1 and H3 tissues, whereas the V_{1b} receptor was not expressed in any of the tissues. RT-PCR experiments revealed that V_2 receptor mRNAs were detected in the three tissues. In contrast, immunohistochemical and cell incubation studies showed that the V_2 receptor was involved in the stimulatory effect of AVP on cortisol secretion in H1 and H2, but not in H3 cells. Taken together, these data show that expression of functional ectopic V_2 receptors and repression of eutopic V_{1a} receptor can coexist in some hyperplastic corticosteroidogenic tissues. They also reveal that immunohistochemical and incubation studies are essential for the characterization of ectopic receptors actually involved in the control of cortisol secretion by AIMAHs.

Journal of Endocrinology (2008) **196**, 1–9

Introduction

Expression of ectopic and overexpression of eutopic G-protein-coupled receptors for peptide hormones have been reported in different lesions, including hyperplasia, tumors, and cancers of various tissues (Qu *et al.* 2004, Havt *et al.* 2005, Weng *et al.* 2006). Activation of aberrant receptors initiates intracellular signals responsible for excessive cellular activity and cell proliferation. Characterization of aberrant receptors may be used to help shed light on the pathophysiological mechanism of gain-of-function disorders and may allow the development of therapeutic approaches to reduce the activity of hyperfunctioning cells. In Cushing's syndrome, due to unilateral adrenal adenoma or adrenocorticotropin (ACTH)-independent bilateral macronodular adrenal hyperplasia (AIMAH), overproduction of cortisol, which leads to suppression of ACTH release, is under the control of illegitimate G-protein-coupled receptors (Lacroix *et al.* 2004). These receptors include ectopic receptors for gastric inhibitory

polypeptide (GIP), luteinizing hormone (LH) or serotonin₇ (5-HT₇) receptors, and abnormally active eutopic receptors like vasopressin V_1 and 5-HT₄ receptors (Lacroix *et al.* 2004). Illegitimate receptors were initially detected by clinical studies aimed at evaluating the plasma cortisol responses to various physiological and pharmacological stimuli (Lacroix *et al.* 1999). In particular, aberrant expression of vasopressin receptors has been described in some cases of AIMAH responding to physiological stimuli of endogenous vasopressin. In these reports, elevation of plasma AVP provoked by upright posture, insulin-induced hypoglycemia, or hypertonic saline infusion test was followed by an increase in plasma cortisol (Lacroix *et al.* 1997, Daidoh *et al.* 1998, Miyamura *et al.* 2003). An abnormal response of cortisol to injection of AVP or lysine vasopressin (LVP) was also observed (Lacroix *et al.* 1997, Daidoh *et al.* 1998, Miyamura *et al.* 2003, Bertherat *et al.* 2005). It was conceivable that hypersensitivity of cortisol secretion to AVP in these patients might result from activation of illegitimate receptors,

i.e., overexpressed eutopic V_{1a} receptors and/or ectopic V_{1b} and V_2 receptors. In agreement with this hypothesis, molecular studies have shown overexpression of V_{1a} receptor and abnormal expression of V_{1b} and V_2 receptor mRNAs in whole tissue explants removed from AIMAHs (Miyamura *et al.* 2002, Mune *et al.* 2002, Lee *et al.* 2005). However, RT-PCR data are ambiguous since it has been reported that adrenal chromaffin cells and blood vessels express the V_{1b} and V_2 receptors respectively (Aldasoro *et al.* 1997, Grazzini *et al.* 1999, Medina *et al.* 1999). In addition, ectopic vasopressin receptor mRNAs may not be translated into functional proteins in adrenal hyperplastic tissues. In this respect, administration of the V_2 receptor agonist desmopressin failed to affect plasma cortisol in a case of vasopressin-sensitive AIMAHs expressing the V_2 receptor mRNA (Miyamura *et al.* 2003, Vezzosi *et al.* 2007). It can also be noticed that the effect of V_2 and V_{1b} receptor ligands on cortisol production has never been studied *in vitro* in vasopressin-sensitive AIMAHs expressing V_{1b} and/or V_2 receptor mRNAs. Thus, there is currently no clear evidence for the involvement of ectopic AVP receptors in the abnormal cortisol response to AVP in AIMAH tissues.

We have previously investigated *in vivo* the responsiveness of cortisol secretion to upright and LVP stimulation tests in three patients with AIMAHs and Cushing's syndrome (Bertherat *et al.* 2005). The aim of the present study was to determine *in vitro* the type of receptors mediating the corticotropic action of AVP in the tissues removed from the three patients. For this purpose, expression of AVP receptors has been investigated using RT-PCR and immunohistochemical approaches. Moreover, pharmacological characterization of vasopressin receptors was performed in cultured hyperplasia cells.

Materials and Methods

Patients

Three previously described patients with AIMAH causing overt Cushing's syndrome were studied (patients 1, 2, and 4 from Bertherat *et al.* 2005). The diagnosis of ACTH-independent Cushing's syndrome was based on the results of hormonal investigations. Briefly, increase in 24-h urinary cortisol excretion, alteration of plasma cortisol circadian rhythm, lack of cortisol suppression under a low-dose dexamethasone test (2 mg/day for 2 days), and suppression of basal plasma ACTH levels (below 1 pmol/l (5 pg/ml)) were observed in the three patients. The patients underwent a series of clinical tests aimed at detecting the expression of illegitimate receptors followed by bilateral adrenalectomy. Pathological examination of the adrenal tissues confirmed the diagnosis in all cases. The effect of AVP on cortisol secretion by hyperplasia cells has been investigated *in vitro*. The clinical presentation of the patients as well as the results of clinical testing and cell incubation studies are summarized in Table 1.

In the present study, normal adrenal tissue explants (control tissues) were obtained from patients undergoing expanded

Table 1 Clinical characteristics, endocrine evaluation, and radiological findings in the three patients with adrenocorticotropin (ACTH)-independent macronodular adrenal hyperplasia (AIMAH; H1–H3)

	H1	H2	H3
Sex	Female	Female	Female
Age (year)	34	45	54
Symptoms of hypercortisolism	Hypokalemia, hypertension, diabetes mellitus, faciotruncal adiposity	Hypokalemia, hypertension, Asthenia, depressive mood, facial plethora, truncal adiposity, facial hirsutism	Hypertension, ecchymotic lesions, central obesity, facial hirsutism, proximal myopathy
Maximum diameter of adrenal nodules at CT scan (cm)	Left 6, right 4	Left 4, right 3.5	Left 5.4, right 5
Urinary free cortisol (nmol/d; 55 < N < 250)	1172	1448	1951
Plasma cortisol (nmol/l)			
0800 h (275 < N < 550)	510	166	439
2000 h	466	663	933
Plasma ACTH 0800 h (pmol/l; 4 < N < 13)	< 1	1	1
Plasma cortisol response to upright test ^a	+62%	+29%	–15%
Plasma cortisol response to terlipressin (0.5 mg i.v.) ^a	+52%	–10%	+15%
In vitro AVP efficiency on cortisol secretion ^a	+373.3 ± 17.8%	+126.8 ± 16.2%	+47.9 ± 2.7%

N, normal value.

^aPercentage of basal level.

nephrectomy for kidney cancer. Pituitary corticotrophic adenoma tissue was obtained from a patient undergoing hypophysectomy for Cushing's disease. Fragments of tissue were either transported to the laboratory in culture medium for primary culture, immediately stored at -80°C until RNA extraction or fixed in formalin and embedded in paraffin for immunohistochemical study. The protocol of collection of the tissues and the experimental procedures were approved by the regional ethics committee and written informed consent was obtained from all subjects.

Reagents

Collagenase (type IA), DNase I, insulin, apotransferrin, L-ascorbic acid, [Arg⁸]-vasopressin (AVP), [deamino-Cys¹, Val⁴, D-Arg⁸]-vasopressin (dDAVP), and Tri Reagent were purchased from Sigma. The rabbit V_{1a} and V₂ receptor antibodies were obtained from Santa Cruz Biotechnology (Le Perray en Yvelines, France). The nutrient medium Ham's F-12 (HAM) and Dulbecco's modified Eagle medium (DMEM) were obtained from Life Technologies Inc. (Paisley, Scotland, UK). The antibiotic-antimycotic solution and fetal bovine serum (FBS) were from Bio-Whittaker (Walkersville, MD, USA). (2S)-1-[(2R,3S)-5-chloro-3-(2-chlorophenyl)-1-(3,4-dimethoxybenzene-sulfonyl)-3-hydroxy-2,3-dihydro-1H-indole-2-carbonyl]-pyrrolidine-2-carboxamide (SR49059) was supplied by Sanofi Recherche. SuperScript II and DNA Taq Polymerase were from Life Technologies.

RNA extraction and conventional RT-PCR

Expression of genes encoding vasopressin (V_{1a}, V_{1b}, and V₂) receptors in adrenal hyperplasia tissues, normal adrenal gland, and pituitary corticotrophic adenoma was analyzed by RT-PCR. Amplification of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA was used as a control of quality of reverse-transcribed mRNAs. The RT-PCRs were performed according to the method and experimental conditions previously described (Vezzosi *et al.* 2007). Briefly,

total RNAs were extracted by using Tri Reagent. Total RNA (1 µg) from each preparation was converted to single stranded cDNA by SuperScript II with oligo(dT)₁₂₋₁₈ primer. PCRs were carried out using gene-specific primers for each sequence (Table 2) and DNA Taq Polymerase. The PCRs were performed for 40 cycles (94 °C, 40 s; 50 °C, 60 s; 72 °C, 90 s). The PCR products were separated on agarose gels, blotted on nylon membranes, and hybridized with [³²P]ATP-labeled internal gene-specific oligonucleotides (Table 2).

Immunohistochemistry

Deparaffinized sections from the three hyperplasias and three normal adrenal glands were incubated overnight at 4 °C in a humidified atmosphere with rabbit polyclonal antibodies directed against V_{1a} and V₂ receptors (1:200). The sections were then incubated with a streptavidin-biotin-peroxidase complex (Dako Corporation, Carpinteria, CA, USA) and the enzymatic activity was revealed with diaminobenzidine. The specificities of the immunoreactions were controlled by substituting the primary antisera with non-immune serum. The tissue sections were counterstained with hematoxylin, mounted in Eukitt (Kindler GmbH & Co., Freiburg, Germany), coverslipped, and examined on an Eclipse E-600 microscope equipped with a CCD DXC950 camera (Nikon, Les Ulis, France).

Cell culture

Cell culture experiments were conducted as previously described (Bertherat *et al.* 2005). Briefly, hyperplasia and normal adrenal gland fragments free of fat were immediately immersed in DMEM supplemented with 0.5% antibiotic-antimycotic solution. After dissection of medullary tissues with scissors, the adrenocortical fragments were enzymatically dissociated in DMEM containing collagenase (2 mg/ml) and desoxyribonuclease I (70 µg/ml) for 45 min at 37 °C. Adrenocortical cells were cultured at 37 °C in 100% relative humidity in a 5% CO₂-95% (v/v) air atmosphere. Incubation

Table 2 RT-PCR analysis of vasopressin (V_{1a}, V_{1b}, and V₂) receptor mRNAs

	Accession number in Genbank	Primer sequence (5'-3')	Primer position
Receptor			
V _{1a}	AF 101725	s: CGGCTTCATCTGCTACAACATC as: CGAGTCCTTCCACATACCCGT p: ATGTGGTCTGCTGGGATCCC	731-752 1215-1235 957-977
V _{1b}	AF 101726	s: CAGCAGCATCAACACCATCT as: CCATGTAGATCCAGGGGTTG p: GATTCCACCAATGTGGCTTTC	921-940 1122-1141 1063-1083
V ₂	NM 000054	s: GTGGCCAAGACTGTGAGGAT as: CATAGATCCAGGGGTTGGTG p: GTGCTACTCATGTTGCTGGCC	1031-1050 1192-1211 1157-1177
GAPDH	M17701	s: TGCTGAGTAYGTCGTGGAGTC as: TTGGTGGTGCAAGKGCATTGC	297-317 467-488

Oligonucleotide sequences for sense (s), antisense (as), and hybridization probes (p).

experiments of cells were conducted after 2 days in culture with fresh DMEM (control experiments) or DMEM with either AVP or dDAVP. AVP was incubated in the absence or presence of SR49059. Cells were incubated with each secretagogue for 24 h at 37 °C. Following each incubation period, aliquots of the culture medium were taken and immediately frozen at -20°C until cortisol RIA (Lefebvre *et al.* 1992). Results are expressed as mean $\pm$ S.E.M. from four experiments and statistical significance was assessed by Bonferroni test after one-way ANOVA. The dose-response curves were generated using the Prism software (GraphPad Software, San Diego, CA, USA).

Results

Molecular characterization of AVP receptors in AIMAH tissues

Expression of genes encoding V_{1a} , V_{1b} , and V_2 receptors in adrenal hyperplasia, normal adrenal gland, and pituitary corticotrophic adenoma tissues were determined by RT-PCR. V_{1a} receptor mRNAs were detected in normal adrenal gland (NA), and in both adrenal hyperplastic tissues from patients 1 and 3 (H1 and H3), but not in H2 tissue (Fig. 1). V_{1b} receptor mRNA was not present in any of the adrenal tissues, but its occurrence was observed in the pituitary corticotrophic adenoma (control experiment). PCR products corresponding to V_2 receptor mRNA were visualized in hyperplastic tissues from the three patients.

Immunohistochemical localization of V_{1a} and V_2 receptors in AIMAH tissues

Presence of V_{1a} and V_2 receptors in normal adrenal gland and hyperplasia tissues was examined by immunohistochemistry. Incubation of normal adrenal tissue slices with anti- V_{1a} receptor antibodies produced intense labeling of the zona fasciculata (Fig. 2A and B). V_{1a} receptor-immunoreactive cells were also observed in hyperplasia H1 and H3 (Fig. 2C and G).

Figure 1 Expression of AVP receptor mRNAs in AIMAH causing Cushing's syndrome. RT-PCR analysis of mRNA expression encoding the AVP receptor subtypes V_{1a} , V_{1b} , and V_2 , and GAPDH in normal adrenal gland (NA) and hyperplasia H1, H2, and H3. Pituitary corticotrophic adenoma (CA) was used as a positive control for V_{1b} mRNA.

V_{1a} receptor-positive cells had the morphological characteristics of spongiocytic cells, i.e., cells with abundant cytoplasm and numerous lipid droplets. Immunolabeling was present in the cytoplasm and at the periphery of the spongiocytic cells (Fig. 2D and H). In contrast, the H2 tissue did not display V_{1a} receptor immunoreactivity (Fig. 2E and F).

Incubation of normal adrenal tissue sections with V_2 receptor antibodies did not give rise to immunolabeling (Fig. 3A and B). At variance, clusters of V_2 receptor-immunoreactive small compact and/or large spongiocytic cells were detected in nodules of hyperplasia H1 and H2 (Fig. 3C and E). Immunolabeling was present in the cytoplasm and at the periphery of the spongiocytic cells (Fig. 3D and F). Incubation of H3 tissue slices with the V_2 receptor antiserum did not reveal immunoreactive material in small compact and large spongiocytic cells (Fig. 3G and H). No labeling was observed in all tissues when primary antibodies were substituted with non-immune serum (data not shown).

Pharmacological properties of AVP receptors in AIMAH tissues

AVP receptors expressed in AIMAH tissues and normal adrenal glands were characterized by a pharmacological approach. As previously reported (Perraudin *et al.* 1993), AVP stimulated cortisol production in cultured cells derived from normal adrenal gland (Fig. 4A). The stimulatory effect of AVP was blocked by the V_{1a} receptor antagonist SR49059 (10^{-6} M). Application of graded concentrations of AVP (10^{-12} – 10^{-7} M) to cultured H1 cells induced a dose-related increase in cortisol secretion with high potency (mean $\text{pEC}_{50} = 10.1 \pm 0.1$; $n=4$) and efficacy ($E_{\text{max}} = +373.3 \pm 17.8\%$; $n=4$; Fig. 4B). SR49059 (10^{-7} M) shifted the AVP response curve to the right yielding a mean pEC_{50} of 8.9 ± 0.3 ($n=4$; $P<0.001$; Fig. 4B). AVP (10^{-12} – 10^{-7} M) also stimulated cortisol production in a dose-dependent manner ($\text{pEC}_{50} = 10.2 \pm 0.4$; $E_{\text{max}} = 126.8 \pm 16.2\%$; $n=4$) in cultured H2 cells (Fig. 4C). However, SR49059 (10^{-7} M) did not significantly modify the stimulatory effect of AVP in H2 cells ($\text{pEC}_{50} = 9.9 \pm 0.4$; $n=4$; $P>0.05$; Fig. 4C). In cultured H3 cells, AVP (10^{-12} – 10^{-7} M) provoked an increase in cortisol release ($\text{pEC}_{50} = 10.9 \pm 0.2$; $E_{\text{max}} = 47.9 \pm 2.7\%$; $n=4$; Fig. 4D). The V_2 receptor agonist dDAVP (10^{-12} – 10^{-7} M) had no effect on cortisol production (Fig. 4D). The stimulatory effect of AVP on H3 cells was inhibited by SR49059 ($\text{pEC}_{50} = 7.5 \pm 0.3$; $n=4$; $P<0.0001$; Fig. 4D).

Discussion

Both clinical and molecular studies have demonstrated the presence of multiple illegitimate receptors including AVP receptors in AIMAHs causing subclinical and/or overt Cushing's syndrome (Lacroix *et al.* 2004). In the present study, we have investigated the expression of AVP receptor subtypes in normal adrenal gland and three hyperplasia tissues. We have also examined the pharmacological profile of AVP receptors in cultured cells derived from the three AIMAHs.

Figure 2 Immunohistochemical localization of V_{1a} receptor in AIMAH causing Cushing's syndrome. V_{1a} receptor-immunoreactive cells in the zona fasciculata of two normal adrenal glands (NA1, A; $\times 260$ and NA2, B; $\times 280$). V_{1a} receptor-immunopositive cells in H1 and H3 tissues (H1, C; $\times 220$ and H3, G; $\times 220$). Higher magnification view of hyperplasia sections showing the presence of V_{1a} receptor immunoreactivity in the cytoplasm and at the periphery of spongiocytic cells (H1, D; $\times 440$ and H3, H; $\times 360$). Absence of V_{1a} receptor immunoreactivity in H2 tissue (E; $\times 220$) and (F; $\times 220$). Scale bars: 50 μ m.

Figure 3 Immunohistochemical localization of V_2 receptor in AIMAH causing Cushing's syndrome. Absence of V_2 receptor-immunoreactive cell in zona fasciculata of two normal adrenal glands (NA1, A; $\times 220$ and NA2, B; $\times 220$). Clusters of V_2 receptor-immunopositive cells in H1 and H2 tissues (H1, C; $\times 320$ and H2, E; $\times 220$). Higher magnification view of hyperplasia sections showing the presence of V_2 receptor immunoreactivity in the cytoplasm and at the periphery of spongiocytic cells (H1, D; $\times 380$ and H2, F; $\times 440$). Absence of V_2 receptor immunoreactivity in H3 tissue at low (G; $\times 100$) and high (H; $\times 340$) magnifications. Scale bars: 50 μ m.

Figure 4 Effect of AVP and vasopressinergic ligands on cortisol secretion from cultured AIMAH cells. Effect of the V_{1a} receptor antagonist SR49059 (10^{-7} M) on cortisol secretion induced by 10^{-8} M AVP from cultured normal adrenocortical cells (NA, A). Effect of graded concentrations of AVP (from 10^{-12} to 10^{-7} M) on cortisol secretion from H1 (B) and H2 (C) cells in the absence (●) or the presence of SR49059 (10^{-6} M) (○). Effect of graded concentrations (from 10^{-12} to 10^{-7} M) of AVP and the V_2 receptor agonist dDAVP (Δ) on cortisol production from H3 cells (D). AVP was administered alone (●) or in combination with SR49059 (10^{-6} M) (○). Data values represent mean $\pm$ S.E.M. from four experiments.

It is well documented that AVP stimulates cortisol secretion, through activation of V_{1a} receptors, in normal adrenal gland (Perraudin *et al.* 1993, Gallo-Payet & Guillon 1998) and some AIMAH tissues (Lacroix *et al.* 1997, Daidoh *et al.* 1998). In agreement with these data, our RT-PCR experiments revealed the presence of V_{1a} receptor mRNA in both normal tissue and hyperplasias H1 and H3. In addition, V_{1a} receptor-like immunoreactivity was visualized in the normal cortex and a subpopulation of steroidogenic cells in H1 and H3 tissues. The stimulatory effect of AVP on cortisol production from cultured cells, derived from either normal adrenal gland or hyperplasia H3, was blocked by the V_1 receptor antagonist SR49059, indicating that the corticotrophic action of the nonapeptide was mediated by eutopic V_{1a} receptors. SR49059 exerted a weaker inhibitory effect on H1 cells, suggesting that eutopic V_{1a} receptors partially relayed the AVP response in this lesion. By contrast, the presence of V_{1a} receptor mRNA or protein was not detected in the H2 tissue. Consistent with the lack of V_{1a} receptor, our *in vitro* experiments revealed that SR49059 failed to inhibit AVP-induced cortisol secretion in H2 cultured cells. These data demonstrate the absence of eutopic V_{1a} receptor in an adrenal hyperplastic lesion and indicate that the stimulatory effect of AVP is actually mediated by ectopic receptors in H2

tissue. Such loss of eutopic receptor and gain of function due to abnormal expression of another receptor subtype in AIMAH have already been reported for 5-HT receptors but have never been described for AVP receptors (Louiset *et al.* 2006). This observation is also consistent with the previously shown underexpression of the melanocortin type 2 receptor in AIMAH tissues overexpressing the GIP receptor (Lampron *et al.* 2006). It has been proposed that these combined molecular defects may be related to abnormal activity or mutations of transcription factors involved in the regulation of the expression of multiple membrane receptors (Lacroix *et al.* 2001).

Expression of the V_{1b} receptor gene has been shown in some AIMAH tissues (Mune *et al.* 2002, Miyamura *et al.* 2003, Lee *et al.* 2005). The fact that V_{1b} receptor mRNA cannot be detected in present RT-PCR experiments excludes the involvement of ectopic V_{1b} receptors in AVP-induced cortisol production in the three AIMAH tissues.

As previously reported in some cases of AVP-sensitive AIMAH (Mune *et al.* 2002, Miyamura *et al.* 2003, Lee *et al.* 2005, Vezzosi *et al.* 2007), we found that V_2 receptor mRNA was produced in the three hyperplastic tissues. Immunohistochemical experiments revealed the presence of V_2 receptor protein in steroidogenic cells

of H1 and H2 tissues. The AVP-induced cortisol secretion observed in cultured H2 cells, which did not possess V_{1a} and V_{1b} receptors, provides the first evidence for the occurrence of functional illegitimate V_2 receptors in an adrenocortical hyperplasia causing Cushing's syndrome. The V_2 receptor gene is also expressed in hyperplasia H3. However, several data indicate that V_2 receptors are not involved in the vasopressinergic control of steroidogenesis in this tissue: i) immunohistochemical studies failed to detect any V_2 receptor-like immunoreactivity in steroidogenic cells; ii) the stimulatory effect of AVP on corticosteroidogenesis was suppressed by SR49059; and iii) the V_2 receptor agonist dDAVP had no influence on cortisol secretion. These results suggest that V_2 receptor mRNAs present may not be translated into functional proteins in AIMAH, as previously observed for the 5-HT₄ receptor (Louiset *et al.* 2006). The lack of V_2 receptor protein in H3 tissue may result from decreased mRNA stability. Various molecular partners, such as RNA-stabilizing proteins and small RNAs, may potentially be involved in this process (Chu & Rana 2007, Eberhardt *et al.* 2007).

We have previously demonstrated the presence of numerous cells producing AVP in the hyperplastic cortex of patients with AIMAH (Bertherat *et al.* 2005). Thus, it can be speculated that eutopic V_{1a} and ectopic V_2 receptors mediate a direct intra-adrenal vasopressinergic tone involving paracrine and autocrine mechanisms. Our observations may be pathophysiologically relevant since it has been reported that AVP increases proliferation of different cell types, including rat adrenocortical cells, through activation of V_{1a} receptors (Chiu *et al.* 2002, Lagumdzija *et al.* 2004, Trejter *et al.* 2005). These data suggest that the nonapeptide, locally produced in AIMAH tissues, may stimulate proliferation of steroidogenic cells via activation of V_{1a} receptors. Moreover, expression of ectopic V_2 receptors, which are classically coupled to adenylyl cyclase (Birnbaumer 2000), indicates that V_2 receptors may also have contributed to the physiopathology of AIMAH by stimulating cell steroidogenesis and mitogenesis. In support of this hypothesis, it has been recently described that transfer of adenylyl cyclase-coupled GIP and LH receptors in bovine adrenocortical cells causes cortisol hypersecretion and adenomatous hyperproliferation in a mouse xenotransplantation model (Mazzuco *et al.* 2006a,b).

In conclusion, the present study provides the first immunohistochemical characterization of AVP receptors in AIMAH tissues. Combined with molecular and pharmacological approaches, immunohistochemistry revealed that the steroidogenic response to AVP can be mediated by ectopic V_2 receptor alone or in association with eutopic V_{1a} receptor. Our data also clearly indicate that the sole RT-PCR data are not sufficient for the determination of the receptor subtype(s) actually involved in the steroidogenic effect of a regulatory factor in hyperplastic adrenocortical tissues.

Acknowledgements

This work was supported by the Conseil Régional de Haute-Normandie, IFRMP 23, INSERM U413, the CHU

de Rouen, the Réseau COMETE (PHRC AOM 02068), and the Assistance Publique-Hôpitaux de Paris (Grant CRC 02005). This work is dedicated to the memory of Vincent Contesse, a brilliant researcher and a beloved friend. We are indebted to M Gras and H Lemonnier for technical assistance. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Aldasoro M, Medina P, Vila JM, Otero E, Martinez-Leon JB & Lluch S 1997 Endothelium-dependent relaxation of human saphenous veins in response to vasopressin and desmopressin. *Journal of Vascular Surgery* **25** 696–703.
- Bertherat J, Contesse V, Louiset E, Barrande G, Duparc C, Groussin L, Emy P, Bertagna X, Kuhn JM, Vaudry H *et al.* 2005 *In vivo* and *in vitro* screening for illegitimate receptors in adrenocorticotropin-independent macronodular adrenal hyperplasia causing Cushing's syndrome: identification of two cases of gonadotropin/gastric inhibitory polypeptide-dependent hypercortisolism. *Journal of Clinical Endocrinology and Metabolism* **90** 1302–1310.
- Birnbaumer M 2000 Vasopressin receptors. *Trends in Endocrinology and Metabolism* **11** 406–410.
- Chiu T, Wu SS, Santiskulvong C, Tangkijvanich P, Yee HF Jr & Rozengurt E 2002 Vasopressin-mediated mitogenic signaling in intestinal epithelial cells. *American Journal of Physiology. Cell Physiology* **282** C434–C450.
- Chu CY & Rana TM 2007 Small RNAs: regulators and guardians of the genome. *Journal of Cell Physiology* **213** 412–419.
- Daidoh H, Morita H, Hanafusa J, Mune T, Murase H, Sato M, Shibata T, Suwa T, Ishizuka T & Yasuda K 1998 *In vivo* and *in vitro* effects of AVP and V_{1a} receptor antagonist on Cushing's syndrome due to ACTH-independent bilateral macronodular adrenocortical hyperplasia. *Clinical Endocrinology* **49** 403–409.
- Eberhardt W, Doller A, Akool EL-S & Pfeilschifter J 2007 Modulation of mRNA stability as a novel therapeutic approach. *Pharmacology and Therapeutics* **114** 56–73.
- Gallo-Payet N & Guillon G 1998 Regulation of adrenocortical function by vasopressin. *Hormone and Metabolic Research* **30** 360–367.
- Grazzini E, Breton C, Derick S, Andres M, Raufaste D, Rickwaert F, Boccard G, Colson P, Guérineau NC, Serradeil-le Gal C *et al.* 1999 Vasopressin receptors in human adrenal medulla and pheochromocytoma. *Journal of Clinical Endocrinology and Metabolism* **84** 2195–2203.
- Havt A, Schally AV, Halmos G, Varga JL, Toller GL, Horvath JE, Szepeshazi K, Köster F, Kovitz K, Groot K *et al.* 2005 The expression of the pituitary growth hormone-releasing hormone receptor and its splice variants in normal and neoplastic human tissues. *PNAS* **102** 17424–17429.
- Lacroix A, Tremblay J, Touyz RM, Deng LY, Larivière R, Cusson JR, Schiffrin EL & Hamet PJ 1997 Abnormal adrenal and vascular responses to vasopressin mediated by a V_1 -vasopressin receptor in a patient with adrenocorticotropin-independent macronodular adrenal hyperplasia, Cushing's syndrome, and orthostatic hypotension. *Journal of Clinical Endocrinology and Metabolism* **82** 2414–2422.
- Lacroix A, Mircescu H & Hamet P 1999 Clinical evaluation of the presence of abnormal hormone receptors in adrenal Cushing's syndrome. *Endocrinologist* **9** 9–15.
- Lacroix A, Ndiaye N, Tremblay J & Hamet P 2001 Ectopic and abnormal hormone receptors in adrenal Cushing's syndrome. *Endocrine Reviews* **22** 75–110.
- Lacroix A, Baldacchino H, Bourdeau I, Hamet P & Tremblay J 2004 Cushing's syndrome variants secondary to aberrant hormone receptors. *Trends in Endocrinology and Metabolism* **15** 375–382.
- Lagumdzija A, Bucht E, Stark A, Hulting AL & Petersson M 2004 Arg-vasopressin increases proliferation of human osteoblast-like cells and decreases production of interleukin-6 and macrophage colony-stimulating factor. *Regulatory Peptides* **121** 41–48.
- Lampron A, Bourdeau I, Hamet P, Tremblay J & Lacroix A 2006 Whole genome expression profiling of glucose-dependent insulinotropic peptide

- (GIP)- and adrenocorticotropin-dependent adrenal hyperplasias reveals novel targets for the study of GIP-dependent Cushing's syndrome. *Journal of Clinical Endocrinology and Metabolism* **91** 3611–3618.
- Lee S, Hwang R, Lee J, Rhee Y, Kim DJ, Chung UI & Lim SK 2005 Ectopic expression of vasopressin V1b and V2 receptors in the adrenal glands of familial ACTH-independent macronodular adrenal hyperplasia. *Clinical Endocrinology* **63** 625–630.
- Lefebvre H, Contesse V, Delarue C, Feuilloley M, Héry F, Grise P, Raynaud G, Verhofstad AAJ, Wolf LM & Vaudry H 1992 Serotonin-induced stimulation of cortisol secretion from human adrenocortical tissue is mediated through activation of a serotonin₄ receptor subtype. *Neuroscience* **47** 999–1007.
- Louiset E, Contesse V, Groussin L, Cartier D, Duparc C, Barrande G, Bertherat J, Vaudry H & Lefebvre H 2006 Expression of serotonin₇ receptor and coupling of ectopic receptors to protein kinase A and ionic currents in adrenocorticotropin-independent macronodular adrenal hyperplasia causing Cushing's syndrome. *Journal of Clinical Endocrinology and Metabolism* **91** 4578–4586.
- Mazzuco TL, Chabre O, Feige JJ & Thomas M 2006a Aberrant expression of human luteinizing hormone receptor by adrenocortical cells is sufficient to provoke both hyperplasia and Cushing's syndrome features. *Journal of Clinical Endocrinology and Metabolism* **91** 196–203.
- Mazzuco TL, Chabre O, Sturm N, Feige JJ & Thomas M 2006b Ectopic expression of the gastric inhibitory polypeptide receptor gene is a sufficient genetic event to induce benign adrenocortical tumor in a xenotransplantation model. *Endocrinology* **147** 782–790.
- Medina P, Segarra G, Vila JM, Chuan P, Domenech C & Lluch S 1999 V2-receptor-mediated relaxation of human renal arteries in response to desmopressin. *American Journal of Hypertension* **12** 188–193.
- Miyamura N, Taguchi T, Murata Y, Taketa K, Iwashita S, Matsumoto K, Nishikawa T, Toyonaga T, Sakakida M & Araki E 2002 Inherited adrenocorticotropin-independent macronodular adrenal hyperplasia with abnormal cortisol secretion by vasopressin and catecholamines: detection of the aberrant hormone receptors on adrenal gland. *Endocrine* **19** 319–326.
- Miyamura N, Tsutsumi A, Senokuchi H, Nakamaru K, Kawashima J, Sakai K, Taguchi T, Tokunaga H, Nishida K, Uehara M *et al.* 2003 A case of ACTH-independent macronodular adrenal hyperplasia: simultaneous expression of several aberrant hormone receptors in the adrenal gland. *Endocrine Journal* **50** 333–340.
- Mune T, Murase H, Yamakita N, Fukuda T, Murayama M, Miura A, Suwa T, Hanafusa J, Daido H, Morita H *et al.* 2002 Ectopic overexpression of vasopressin V1a receptor in adrenocorticotropin-independent macronodular adrenal hyperplasia. *Journal of Clinical Endocrinology and Metabolism* **87** 5706–5713.
- Perraudin V, Delarue C, Lefebvre H, Contesse V, Kuhn JM & Vaudry H 1993 Vasopressin stimulates cortisol secretion from human adrenocortical tissue through activation of V1 receptors. *Journal of Clinical Endocrinology and Metabolism* **76** 1522–1528.
- Qu X, Xiao D & Weber HC 2004 Transcriptional activation of the human gastrin-releasing peptide receptor gene in gastrointestinal and prostatic epithelial cancer cells. *Journal of Molecular Neuroscience* **22** 75–82.
- Trejter M, Carraro G, Rucinski M, Hochol A, Rebuffat P, Nussdorfer GG & Malendowicz LK 2005 Arginin-vasopressin regulates proliferative activity of the regenerating rat adrenal cortex. *International Journal of Molecular Medicine* **15** 993–997.
- Vezzosi D, Cartier D, Régnier C, Otal P, Bennet A, Parmentier F, Plantavid M, Lacroix A, Lefebvre H & Caron P 2007 Familial adrenocorticotropin-independent macronodular adrenal hyperplasia with aberrant serotonin and vasopressin adrenal receptors. *European Journal of Endocrinology* **156** 21–31.
- Weng J, Wang J, Hu X, Wang F, Ittmann M & Liu M 2006 PSGR2, a novel G-protein coupled receptor, is overexpressed in human prostate cancer. *International Journal of Cancer* **118** 1471–1480.

Received in final form 2 October 2007

Accepted 10 October 2007

Made available online as an Accepted Preprint
10 October 2007