Stem cell factor/c-Kit signalling in normal and androgenetic alopecia hair follicles

Valerie A Randall, Tracey J Jenner, Nigel A Hibberts, Isabel O De Oliveira and Tayyebah Vafaei

Division of Biomedical Sciences, School of Life Sciences, University of Bradford, Bradford BD7 1DP, UK

(Correspondence should be addressed to V A Randall; Email: v.a.randall@bradford.ac.uk)

Abstract

Androgens stimulate many hair follicles to alter hair colour and size via the hair growth cycle; in androgenetic alopecia tiny, pale hairs gradually replace large, pigmented ones. Since stem cell factor (SCF) is important in embryonic melanocyte migration and maintaining adult rodent pigmentation, we investigated SCF/c-Kit signalling in human hair follicles to determine whether this was altered in androgenetic alopecia. Quantitative immunohistochemistry detected three melanocyte-lineage markers and c-Kit in four focus areas: the epidermis, infundibulum, hair bulb (where pigment is formed) and mid-follicle outer root sheath (ORS). Colocalisation confirmed melanocyte c-Kit expression; cultured follicular melanocytes also exhibited c-Kit. Few ORS cells expressed differentiated melanocyte markers or c-Kit, but NK1/beteb antibody, which also recognises early melanocyte-lineage antigens, identified fourfold more cells, confirmed by colocalisation. Occasional similar bulbar cells were seen. Melanocyte distribution,

concentration and c-Kit expression were unaltered in balding follicles. Androgenetic alopecia cultured dermal papilla cells secreted less SCF, measured by ELISA, than normal cells. This identifies three types of melanocyte-lineage cells in human follicles. The c-Kit expression by dendritic, pigmented, bulbar melanocytes and rounded, differentiated, non-pigmenting ORS melanocytes implicate SCF in maintaining pigmentation and migration into regenerating hair bulbs. Less differentiated, c-Kit-independent cells in the mid-follicle ORS stem cell niche and occasionally in the bulb, presumably a local reserve for long scalp hair growth, implicate other factors in activating stem cells. Androgens appear to reduce alopecia hair colour by inhibiting dermal papilla SCF production, impeding bulbar melanocyte pigmentation. These results may facilitate new treatments for hair colour changes in hirsutism, alopecia or greying.

Journal of Endocrinology (2008) **197**, 11–23

Introduction

Hair colour is often very important and may need to alter to allow appropriate changes, e.g. signalling of sexual maturity (Jansen & van Baalen 2006) or seasonal camouflage (Flux 1970). In men, highly pigmented beard hair replaces the almost colourless, tiny vellus hairs on the child's face after puberty (Marshall & Tanner 1970), while the older, mature male is frequently distinguished by the reverse process on the scalp causing androgenetic alopecia or male pattern baldness. Androgens are the main regulator of human hair follicles (reviewed in Randall (2007)) causing the changes signalling adulthood and, paradoxically, also androgenetic alopecia (Hamilton 1951, 1958). Darker lion manes also correlate with higher testosterone levels (West & Packer 2002) and testosterone increases pigmentation in Mallard feathers (Haase *et al.* 1995) and deer hair (Bubenik & Bubenik 1985).

Hair follicles can change the size and colour of the hairs they produce via their unique ability to regenerate themselves during the regular cycles of growth (anagen), regression (catagen) and rest (telogen) during which the hair is shed and

replaced (Kligman 1959). The new hair may be identical in colour to the previous one, as frequently occurs on normal human scalp, or very different like the Scottish hare's brown summer coat (Flux 1970). Hair colour depends on the amount and type of pigment present in the hair shaft, including the balance of brown/black eumelanin and red/yellow pheomelanin. Melanocytes located in the epithelial cell matrix of the anagen hair bulb, around the top of the dermal papilla, synthesise melanins in specialised organelles, melanosomes, exporting them to adjacent keratinocytes in the bulb (Chase 1958; reviewed in Castanet & Ortonne (2000), Slominski *et al.* 2004, Lin & Fisher 2007). Unlike the epidermis, the follicular melanin synthesis is cyclical, closely coupled to the anagen hair manufacture (Burnett *et al.* 1969, Commo & Bernard 2000, Slominski *et al.* 2004, Sharov *et al.* 2005).

Although hormones are known to coordinate with seasonal and age/sex-related changes in the hair follicles (Randall 2007), our understanding of how follicle pigmentation is controlled is limited. Androgens are believed to act on follicles predominantly via the mesenchyme-derived dermal papilla,

situated within the hair bulb at the follicle base (Randall 2007); this regulates many aspects of follicular activity and determines the hair type produced (Reynolds & Jahoda 2004, Waters *et al.* 2007). In the current mechanism, androgens alter dermal papilla production of regulatory paracrine factors, e.g. growth factors and/or extracellular matrix components, which influence the follicular target cells (Randall *et al.* 1991, Randall 1994, 2007). Follicular melanocytes would be the targets in hair colour alterations.

Local factors can influence not only epidermal melanocytes, particularly the melanocortins and adrenocorticotrophic hormone (ACTH), but also β -endorphin, endothelin, prostaglandins and catecholamines etc (reviewed in Slominski *et al.* (2004), Lin & Fisher 2007). Such factors regulating follicular melanocytes are not so well established, although pro-opiomelanocortin, ACTH and α -melanocyte-stimulating hormone (α -MSH) seem to have local roles, particularly in balancing eumelanin and pheomelanin proportions (reviewed in Slominski *et al.* (2004), Tobin & Kausar 2005, Costin & Hearing 2007, Rousseau *et al.* 2007). In addition, the cytokine, stem cell factor (also called SCF, mast cell growth factor, steel factor or c-Kit ligand) and its cell surface receptor, c-Kit, play important roles in both epidermal and hair pigmentation (Williams *et al.* 1992). Both are essential to enable melanocyte-lineage cell migration from the neural crest into developing murine hair follicles (Zsebo *et al.* 1990, Geissler *et al.* 1988, Jordan & Jackson 2000) and people with the white forelock of piebaldism exhibit mutations in the *c-Kit* gene (Fleischman *et al.* 1991, Spritz *et al.* 1992). The SCF/c-Kit system is also involved in post-natal rodent hair cycles; c-Kit antibodies inhibit murine hair pigmentation (Nishikawa *et al.* 1991), promote apoptosis of follicular melanocytes (Okura *et al.* 1995) and prevent regeneration of the follicle bulb pigment units in the next anagen (Botchkareva *et al.* 2001). Similarly, overexpression of epidermal SCF causes epidermal hyperpigmentation in mice and atypical melanocyte distribution (Kunisada *et al.* 1998a,b). Since melanocytes synthesising follicle pigment are physically close to the dermal papilla in the bulb, we hypothesised that it was a local source of SCF; this is supported by secretion of SCF by cultured dermal papilla cells (Hibberts *et al.* 1996, Lu *et al.* 2006).

Most research focuses on laboratory rodent follicles, but these have significant differences to human follicles: no colour changes after puberty, no major responses to androgens, much smaller size, anagen involving weeks not the years of scalp growth and no epidermal melanocytes. We aimed to further investigate SCF/c-Kit signalling in normal human anagen hair follicles and to compare it in androgenetic alopecia to see if the androgen-promoted reduced pigmentation in balding involved changes in the SCF/c-Kit system. Two approaches were employed: follicular melanocyte-lineage cell distribution and expression of the SCF receptor, c-Kit were compared in normal and androgenetic alopecia hair follicles using immunohistochemistry, and the ability of cultured dermal papilla cells derived from normal and balding scalp to produce SCF measured. Initially, the expression of antigens

associated with stages of melanocyte-lineage cell development and c-Kit were compared in frozen sections to determine whether this varied with follicular location or with the follicle's response to androgens *in vivo*, focusing particularly on the site of pigment synthesis, the hair bulb, the melanocyte stem cell niche in the mid-follicle outer root sheath (ORS), i.e. the lower part of the permanent follicle, ORS, below the sebaceous duct (Staricco 1959, Sugiyama & Kukita 1976, Horikawa *et al.* 1996, Narisawa *et al.* 1997, Commo & Bernard 2000, Botchkareva *et al.* 2001, Nishimura *et al.* 2002, 2005, Steingrimsdottir *et al.* 2005), the infundibulum and epidermis. Three antibodies to melanocyte-lineage cells were used to discriminate between differentiation stages; NK1/beteb recognises all melanocyte-lineage cells including very immature ones, as it binds 7 and 100 kDa antigens in premelanosome-like structures, premelanosomes and melanosomes (Hayashibe *et al.* 1986), while HMB-45 and Mel-2 only recognise mature melanocytes binding to 10 kDa melanosome-associated and cell surface protein antigens respectively (Kapur *et al.* 1992); Mel-2 was selected for direct comparison with c-Kit, a membrane receptor. Antibodies to tyrosinase-related protein (TRP)-2 used to locate immature melanocytes in rodents (Botchkareva *et al.* 2001) were not employed, as these did not stain human follicular melanocytes here or elsewhere (Commo *et al.* 2004a). The number of cells stained with each antibody in every region was quantified in each individual studied; using two antibodies to different mature melanocyte antigens on parallel sections of each sample provided an internal control to check quantitation reliability. To confirm, the human follicle melanocyte expression of c-Kit colocalisation studies of c-Kit and melanocyte-lineage antigens and immunocytochemistry of primary lines of cultured epidermal and follicular melanocytes were also carried out. Finally, primary lines of dermal papilla cells were derived from normal scalp follicles and intermediate androgenetic alopecia follicles and their SCF production measured by ELISA after culture in the presence, and absence, of testosterone. This enabled the comparison of any *in vivo* or *in vitro* effects of androgens. Since hair colour normally remains unchanged within a cycle even over the years required for long hair, *in vitro* effects were not anticipated.

Materials and Methods

Skin samples

Scalp skin by-products of routine surgical procedures were collected from apparently healthy Caucasian adults. This included: five women and three men, aged 38–52 for the main immunohistochemical study and one man and four women aged 34–48 for melanocyte culture. Non-balding (occipital) scalp of 12 men, some with male pattern baldness, aged 38–59, were used for the colocalisation immunohistological studies and for non-balding dermal papilla cell

culture. Balding vertex scalp samples were obtained from 13 men with androgenetic alopecia aged 24–46 years for immunohistochemical analysis and dermal papilla cell culture. Foreskin samples were obtained similarly from four males aged 2–12 years. Declaration of Helsinki protocols were followed and appropriate patient consent and local Ethical Committee approval obtained. There was no evidence of hair thinning or loss of pigmentation in the non-balding samples, but balding skin contained much smaller hair follicles producing finer, less pigmented hairs. Skin samples were collected in sterile culture medium or PBS and transported to the laboratory at 4 °C taking up to 6 h for normal skin, and up to 24 h for androgenetic alopecia samples, before embedding in OCT and freezing at –80 °C or processing for cell culture.

Immunohistochemistry

Vertical frozen sections (5–8 µm thick) were cut using a Leica cryostat (CM 1800) and collected on poly-L-lysine-coated slides to increase adherence (Marzia *et al.* 1975). Special care was needed during sectioning to orientate the specimen to obtain longitudinal sections of hair follicles. Poly-L-lysine (BDH Chemicals Ltd, Poole, Dorset, UK) diluted 1:10 with double distilled water was applied to very clean, dry slides previously soaked in methanol. After 5 min, the excess was removed and the slides were left to air dry. Sections were fixed in acetone at 4 °C for 10 min and air dried for 3–24 h. All staining took place in a moistened chamber at room temperature.

Endogenous peroxidase activity was blocked by adding five drops of 3% hydrogen peroxide:methanol (1:4) solution for 20 min. After rinsing with PBS, followed by immersion in further PBS for 10 min, non-specific binding sites were blocked by incubation with mouse serum (1:100) (Dako, Buckinghamshire, UK) for 20 min, followed by 1-h incubation with various primary mouse anti-human monoclonal antibodies. After investigating various dilutions and incubation times, antibodies were used at the following dilutions: NKI/beteb 1:10 (Monosan, Uden, The Netherlands), HMB-45 1:10 (Dako), Mel-2 1:20 (Serotec, Kidlington, Oxford, UK) and c-Kit 1:10 (Serotec). Positive controls of parallel incubations of foreskin sections and two negative controls of scalp skin with either PBS or an antibody (1:20) to *Aspergillus niger* glucose oxidase, a protein absent from mammalian tissues (Dako), were carried out with each series. After washing with PBS for 10 min (also repeated after each of the next stages), sections were incubated with a secondary antibody, biotinylated anti-rabbit and anti-mouse immunoglobulins, (Dako) for 20 min. Excess liquid was removed after washing by blotting carefully with tissue and four to six drops of horseradish peroxidase-streptavidin complex (Dako) per section were incubated for 20 min. Similar amounts of chromagen 3-amino-5-ethylcarbazole (AEC; Dako) followed the next wash for about 20 min; red colour appearance was monitored microscopically. Final rinsing with PBS was followed by mounting using an aqueous mountant, Glycergel (Dako).

Analysis of quantitative results All sections were examined on a Leitz Ortolux light microscope using wide field Periplan NG/GF eyepieces and the number of stained cells per field were counted in a 0.45 mm field under 400× magnification; care was taken to check that actual cells were counted by altering the focal depth. The numbers of positive cells for each region were quantified from at least 5, and up to 15, sections for each antibody for each of the eight non-balding and eight balding individuals. These were averaged for each individual and then the mean ± S.E.M. obtained for each antibody in each region. Initially, the normal and balding results were analysed separately and data were checked using the SPSS statistical package to determine if a Gaussian normal distribution would enable the use of parametric statistical analysis. Since there were three different factors, i.e. the donor, the site and the antibody, which could affect the numbers of stained cells, three-way ANOVA was carried out by testing donor against site and antibody. Since there was no effect of donor, the results from each site and each antibody were analysed by two-way ANOVA. Further, two-way ANOVAs were carried out to determine whether there were any differences for any antibodies at any site between normal and balding skin.

Colocalisation studies

Dual staining experiments were carried out using two primary antibodies and blue and red chromagens. Sections were incubated with 3% BSA for 30 min followed by an avidin–biotin blocking kit (Vector Laboratories, Peterborough, UK) involving 15 min avidin, 1 min PBS and 15 min biotin to block non-specific staining. The same antibodies were used, as previously described, except that Mel-2 was replaced by Mel-5 that recognises a 75 kDa glycoprotein (Bhawan 1997; Signet Laboratories, Dedham, MA, USA; 02026). The first primary antibody was incubated for 60 min, followed by the biotinylated secondary antibody, horse anti-mouse IgG for 30 min and the 0.2 ml ABC–AP complex (Vector Laboratories) for 30 min with four drops of substrate Vector blue (Vector Laboratories); sections were washed carefully with PBS between steps. The second primary antibody was applied and all steps were repeated, but using vector red (Vector laboratories) as the second substrate.

Antigen expression by cultured melanocytes

Epidermal and hair follicular melanocytes were isolated from scalp skin using the method established by Tobin *et al.* (1995). Briefly, after 1 h in Eagle's minimum essential medium (EMEM; Gibco) containing penicillin (100 IU/ml) and streptomycin (100 µg/ml; both Sigma Chemical Co.) to reduce infection risk, the epidermis and ~1 mm of the upper dermis were removed, cut into 1 cm³ pieces and incubated in medium supplemented with 0.5% collagenase type V (Sigma) and 5% fetal bovine serum (FBS; Globepharm, Surrey, UK)

for 1–2 h at 37 °C to separate the epidermis. Hair follicles were washed in PBS until microscopic examination confirmed removal of contaminating dermal tissue and then tissues were incubated with 0.05% trypsin with 0.53 mM EDTA (Sigma) for 5–10 min at 37 °C to produce single cell suspensions. The cells were cultured in 35 mm tissue culture dishes (Scientific Laboratory Supplies, Nottingham, UK) in the EMEM as described above, supplemented with 10% FBS, 0.2 µg/ml cholera toxin (Sigma), 50 nM 12-*o*-tetradecanoyl phorbol-13-acetate (TCS Biologicals, Buckinghamshire, UK) and keratinocyte serum-free medium (KSFM; TCS Biologicals; EMEM: KSFM, 2:1). Any contaminating fibroblasts were removed with geneticin treatment (TCS Biologicals; Halaban & Alfano 1984) and keratinocytes by differential trypsinisation (Eisinger & Marko 1982). Melanocyte cultures at passage 2 were maintained in medium without phorbol esters or cholera toxin for 48 h prior to immunocytochemistry. The cells (1×10^3) were seeded onto sterile 12-well tissue-culture treated 5 mm slides with removable lids for immunocytochemistry and incubated at 37 °C overnight. The cells were fixed and stained as described for frozen sections above.

Comparison of SCF production by normal and balding dermal papilla cells

Culture of dermal papilla cells Skin samples were transported in E199 medium (Sigma) and transferred to fresh medium on arrival. Primary cultures of dermal papilla cells were established from small pieces of full depth scalp skin by the methods established for non-balding and balding dermal papilla cells described previously (Messenger 1984, Randall *et al.* 1996). Balding dermal papillae were isolated from the small intermediate follicles from large samples of balding scalp, rather than the tiny vellus follicles, because these are actually undergoing changes in response to androgens and also because of the extreme technical difficulties of isolating dermal papillae from miniature vellus follicles. Briefly, skin samples were microdissected in 100 mm sterile Petri dishes containing E199 medium, supplemented with glutamine (2 mmol/ml), penicillin (100 IU/ml), streptomycin (100 µg/ml) and amphotericin B (2.5 µg/ml); all supplements supplied by Gibco. Under a Wild dissecting microscope (Leitz), each follicle was removed separately and individually microdissected to isolate the dermal papilla; each individual papilla was then transferred to a 35 mm tissue culture-treated Petri dish (Bibby Sterilin, Stone, Staffordshire, UK) in the same medium supplemented with 20% serum. Depending on the sample size, three to eight dermal papillae were placed in one dish.

Primary cultures were left untouched in a humidified incubator at 37 °C in 95% air: 5% CO₂ for 2 weeks to establish. They were then examined under phase contrast microscopy and the medium was changed weekly until sufficient cells had grown from the papillae (about 4–5

weeks), for subculture into a 25 cm² tissue culture-treated flask (Bibby Sterilin) in the complete medium to establish primary cell lines; this was termed passage 1. Thereafter, the cells were further subcultured into 25 cm² flasks with a split ratio of 1:3 when almost confluent, generally after about 2 weeks. The cells were stored frozen in liquid nitrogen if not required immediately for experimental purposes.

Preparation of conditioned medium Primary cell lines of non-balding scalp ($n=6$) and balding scalp ($n=5$) dermal papilla cells at passage 4 were grown to confluence in 25 cm² flasks in E199 medium supplemented with 10% FCS. Each flask was washed with PBS (5 ml; $\times 4$) to remove any remnants of serum and parallel flasks of each cell line were supplied with 5 ml serum-free medium containing 10 nM testosterone in 0.001% ethanol or the ethanol vehicle alone. After 15 min, this 'conditioned' medium was removed from each cell line to determine whether there was any SCF in the system initially. This was replaced with further serum-free medium (5 ml) with, or without, 10 nM testosterone for 24 h. The medium was centrifuged at 1000 *g* to pellet cell debris, sterile filtered (0.2 µm) and stored at –20 °C.

Measurement of SCF The concentration of SCF was measured in serum-free medium and the various dermal papilla cell conditioned media using a sandwich ELISA kit, designed for cell culture medium and based on a monoclonal antibody capable of detecting natural human and recombinant SCF (R&D Systems Europe, Abingdon, UK). A 10-point standard curve ranging from 0 to 100 pg/ml SCF in serum-free medium was measured in every assay. All points were assayed in duplicate according to the manufacturer's instructions. The absorbance of each well was measured at 450 nM with wavelength correction set at 540 nM on an ELISA plate reader zeroed against the assay buffer. The minimum detectable level of SCF was 3 pg/ml with a sensitivity of 3 pg/ml; no significant cross-reactivity or interference between other cytokines has been detected. The SCF content was expressed as pg per ml/µg of the medium protein content measured by the Bradford assay (Bradford 1976). Statistical analyses of the differences between SCF concentrations in the media were performed using the non-paired, one-tailed Student's *t*-test after confirmation of normal distribution.

Results

Immunohistochemical localisation of melanocyte antigens in scalp skin samples

Large amounts of readily visible brown pigment were seen in the hair cortex and in the epithelial matrix of the hair bulb above, and around, the upper portions of the dermal papilla of normal scalp follicles. In contrast, pigment was not obvious in the scalp epidermis. In balding samples less pigment was

generally visible in the hair bulb and hair, but some was sometimes seen in the epidermis. Sectioning androgenetic alopecia follicles was even more difficult than normal ones due to their small size. Using a red chromagen enabled visualisation of staining amongst the melanin pigment in the hair bulb. All sections stained with the negative control antibody showed no positive staining, but appropriate positive intermittent, red, cellular staining was seen in the melanocytes of the basal layer of the epidermis in the foreskin sections with each antibody, confirming the specificity of the staining. All staining in scalp sections using single antibodies also showed red, intermittent, cellular staining. All results were normally distributed and the ANOVA tests showed that there was no discontinuity, i.e. variation within the donors at any site or with any antibody.

Non-balding scalp skin

NKI/beteb, an antibody that recognises melanocyte-lineage cells including very early development stages, stained cells in the basal layer of the epidermis (epidermis 10.80 ± 0.30 ; mean $\pm$ S.E.M.; Fig. 1), but not in the upper layers. This pattern continued into the outer layer of the ORS of the follicle infundibulum (Fig. 2a) where the mean number of cells present per field was slightly, but insignificantly, higher (14.00 ± 0.50 , Fig. 1). Although there was some similar staining in the outer layer of the ORS of the mid-follicle, i.e. below the junction point of the sebaceous gland (Fig. 2b),

the number of cells detected was much lower (4.00 ± 0.10) than in the epidermis ($P < 0.05$), infundibulum ($P < 0.05$) and hair bulb ($P < 0.001$; Fig. 1). In the hair bulb, staining was predominantly seen in the matrix above, and around, the upper portion of the dermal papilla amongst the pigment (Fig. 2c). Staining was also observed on the outer edges of the matrix, mainly below the base of the dermal papilla, but no visible pigment was associated with these cells. More cells were detected in the hair bulb (11.00 ± 1.10) than in the epidermis ($P < 0.001$), but similar numbers were observed in the infundibulum (Fig. 1). No staining was observed in other parts of the skin.

Antibodies to the more differentiated melanocyte markers, HMB-45 and Mel-2, produced a similar distribution to NKI/beteb throughout the skin being confined to the basal layers of the epidermis, the ORS and the hair bulb (Fig. 2d-i). Both stained very similar numbers of cells in all sites (Fig. 1), detecting significantly fewer cells than NKI/beteb in all regions (HMB-45, Mel-2 epidermis $P < 0.01$, 0.005 ; infundibulum $P < 0.01$, 0.01 ; mid-follicle $P < 0.001$, 0.001 ; hair bulb $P < 0.01$, 0.01). The proportions of HMB-45 or Mel-2 to NKI/beteb were $1:1.4$ – $2:2$, except in the mid-follicle ORS where this increased dramatically to $1:4$ ($P < 0.001$). Staining with the antibody to c-Kit was similarly distributed to the melanocyte-lineage antibodies (Fig. 2j-l), but additional staining of individual cells, probably mast cells, was also observed in the dermis. Stained cell numbers were very similar to those detected by HMB-45 and Mel-2 and

Figure 1 Comparison of the numbers of cells stained by antibodies to three melanocyte markers and c-Kit in the epidermis and hair follicle of normal and androgenetic alopecia scalp. Results are the mean $\pm$ S.E.M. values from eight non-balding (black bars) and eight balding (grey bars) individuals of the number of cells stained with each antibody per field at high magnification. Although there were marked differences between the concentrations of cells in each area of the follicle, there were no differences in staining between normal and androgenetic alopecia individuals in any area, including the hair bulb, with any antibody.

Figure 2 Antibodies to three melanocyte-lineage antigens plus c-Kit (the receptor for SCF) stained variable numbers of cells in the basal layer of scalp epidermis, the outer layer of the outer root sheath of the infundibulum and mid-follicle and the matrix of the hair bulb. (a–c) NK1/beteb (Mag.); (d–f) HMB-45; (g–i) Mel-2 (Mag.); (j–l) c-Kit. Positive staining is red and melanin pigment is brown. (a, d, g, j) show staining in the epidermis (E) and infundibulum (I) but not in the dermis (D); (b, e, h and k) in the outer root sheath of the permanent middle part of the follicle below the sebaceous gland, hair cortex (C); (c, f, i and l) show red cellular staining plus brown melanin in the matrix of the hair bulb (HM) above and around the dermal papilla (DP). Occasional staining can be seen at the bottom of the bulb away from the pigmented regions (e.g. f). Some staining in each figure is highlighted by arrow. CTS indicates connective tissue, or dermal, sheath. Scale bar represents 100 μ m in c, e, f, h, i, k, l and 62.5 μ m in a, b, d, g, j.

significantly lower than those by NK1/beteb (epidermis, 6.80 ± 0.25 , $P < 0.01$; infundibulum, 9.50 ± 0.50 , $P < 0.01$; mid-follicle, 1.00 ± 0.7 , $P < 0.001$; hair bulb, 9.00 ± 0.42 , $P < 0.01$; Fig. 1).

Colocalisation analysis of melanocyte-lineage cells in scalp skin

Dual labelling studies showed that cells in all areas exhibiting single antibody staining coexpressed HMB-45, Mel-5 and c-Kit with NK1/beteb. The cells staining only with NK1/beteb were also seen in limited numbers in the epidermis (Fig. 2a and b), infundibulum (Fig. 2c) and hair bulb (Fig. 2e), but were the majority amongst the fewer cells in the mid-follicle ORS (Fig. 2d); there was no evidence of

staining with c-Kit antibody in many of these cells (Fig. 3A). The antibody to c-Kit also marked cells in the dermis outside the follicle, which were not recognised by NK1/beteb, presumably mast cells (see 2c, d & e; Costa *et al.* 1996).

Cultured melanocytes

Epidermal melanocyte cultures showed an increase in the number of differentiated dendritic cells that formed more pigment with time in culture. Parallel hair follicle melanocyte cultures contained two distinct populations: actively dividing bipolar, non-pigmenting melanocytes termed hair follicle melanocyte I or HFM-I by Tobin *et al.* (1995) and highly

Figure 3 (A) Colocalisation of c-Kit with melanocyte antigen NKI/beteb. Colocalisation studies confirmed that cells expressing NKI/beteb also simultaneously exhibited c-Kit in the epidermis (E), infundibulum (I), ORS and hair bulb. Some cells in all areas, and the majority in the mid-follicle ORS (d), also exhibited only NKI/beteb staining confirming at least two types of melanocyte-lineage cells in the follicle. Bar represents: (a) epidermis (62.5 μ m); (b) epidermis at greater magnification (25 μ m); (c) infundibulum (62.5 μ m); (d) mid-follicle ORS (62.5 μ m); (e) hair bulb (100 μ m); and (f) hair bulb at higher magnification (25 μ m). Red or blue cells show single staining, dual stained cells are purple. Red stain indicates NKI/beteb and blue indicates c-Kit. Brown/black in hair bulb is melanin pigment. Black arrows highlight NKI/beteb-only stained cells. Blue stained cells outside the hair follicle like those marked by white arrowheads (e.g. c, d and e) are not of melanocyte-lineage and are probably mast cells (D, dermis; DP, dermal papilla; E, epidermis; HM, Hair matrix (site of keratinocyte division and initial differentiation) in hair bulb; I, infundibulum). (B) Cultured hair follicle and epidermal melanocytes exhibit three melanocyte-lineage antigens and c-Kit. Melanocytes (passage 2) derived from scalp epidermis and hair follicles of the same individual exhibit staining for melanocyte antigens NKI/beteb, HMB-45 and Mel-2 and c-Kit by immunocytochemistry. Views were selected to show pigmenting (PD) and less pigmented, amelanotic (AM) epidermal melanocytes and both actively dividing bipolar melanocytes (HPM-I) and highly pigmented, differentiated cells (HPM-II). Epidermal melanocytes, upper row, hair follicular melanocytes, lower row. Scale bar represents 150 μ m.

pigmented, differentiated cells that were lost by passage 3, HFM-II. Cultured epidermal and hair follicle melanocytes derived from the same individuals exhibited staining with antibodies for all three melanocyte-lineage antigens, plus c-Kit (Fig. 3B); no staining occurred with the negative

control antibody with either cell type. Although staining in follicular melanocytes varied with some cells staining heavily and others to a lesser extent, there was no consistent difference between the two populations and bipolar melanocytes exhibited both patterns.

Androgenetic alopecia scalp skin

The distribution and pattern of the staining of each antibody in the epidermis, infundibulum, hair follicle and hair bulb were the same in balding scalp skin samples as in non-balding scalp. Quantitation and statistical analysis confirmed the same significant differences in distribution between NKI/beteb and the other two melanocyte markers, HMB-45 and Mel-2 plus c-Kit as seen in normal scalp, in all sites examined (Fig. 1). There were no significant differences between the numbers of cells per unit area in balding or non-balding scalp (Fig. 1), when the results were compared for each antibody in each region.

Reduction in SCF production by balding dermal papilla cells

Androgenetic alopecia dermal papillae were more difficult to isolate, due to the smaller size of the follicles, and to establish the cell cultures from normal scalp, as reported previously (Randall *et al.* 1996). The ELISA for SCF proved very reproducible with very little variation between duplicates or between different plates; serial dilution of pure SCF produced linear graphs when plotted against absorbance. Neither control serum-free medium nor media conditioned for 15 min by dermal papilla cells from either non-balding or balding scalp hair follicles contained detectable SCF. However, media conditioned for 24 h by all individual primary cell lines of scalp dermal papilla cells contained SCF (Fig. 4). Dermal papilla cells derived from androgenetic alopecia scalp follicles secreted significantly less SCF (2.25 ± 0.26 µg/mg protein) than those from non-balding scalp (3.09 ± 0.38 ; $P < 0.05$). Adding 10 nM testosterone *in vitro* had no effect on the amount produced by balding scalp cell lines at either time point (Fig. 4).

Figure 4 Dermal papilla cells from androgenetic alopecia follicles secrete less SCF in culture than those from non-balding scalp cells. The levels of SCF in serum-free medium conditioned for 24 h by non-balding (vertical shading) and balding scalp (horizontal shading) dermal papilla cells in the presence (darker shading), or absence (light shading), of 10 nM testosterone were measured by ELISA. An initial collection of 15-minute conditioned media contained no detectable SCF (data not shown). Results are the means of seven non-balding and five balding primary cell lines expressed as a percentage of the values for non-balding cells cultured in the absence of testosterone.

Discussion

Expression of c-Kit by melanocyte-lineage cells in human hair follicles

The cells exhibiting three melanocyte-lineage antigens, plus c-Kit, the SCF receptor, were located by immunohistochemistry in four main areas of normal and androgenetic alopecia scalp: the epidermal basal layer; the follicle infundibulum; the mid-follicle ORS; and the hair bulb. These antigens were also expressed by cultured human epidermal and hair follicle melanocytes. This concurs with earlier studies of human epidermis (Grichnik *et al.* 1996, 1998, Norris *et al.* 1996) and rodent hair follicles (Okura *et al.* 1995, Botchkareva *et al.* 2001), and with NKI/beteb distribution in human follicles (Horikawa *et al.* 1996, Commo & Bernard 2000) and cultured melanocytes (Tobin & Bystry 1996, Zhu *et al.* 2004, Ma *et al.* 2006).

There were slightly more stained cells in the infundibulum, than the epidermis (concurring with Grichnik *et al.* (1996)), and the hair bulb (Fig. 1). In the bulb, all antibodies stained cells amongst the matrix pigment around the upper dermal papilla (Fig. 2), the classical follicle pigment unit (Slominski *et al.* 2004); occasional, more rounded cells at the matrix edges, mostly slightly lower than the dermal papilla base, also expressed all antigens (e.g. Figure 2f and i). In the mid-follicle ORS small, oval cells were detected in much lower concentrations than elsewhere ($P < 0.001$; Figs 1–3A). The strong correlation in cell numbers detected by HMB-45 and Mel-2 (Fig. 1) and their staining of cultured melanocytes (Fig. 4) suggests that these antibodies recognised the same well-developed melanocytes (Adema *et al.* 1993, Horikawa *et al.* 1996), supporting the quantitation's validity. Interestingly, similar cell numbers also exhibited c-Kit in each area (Fig. 1); scalp epidermal melanocyte expression concurs with forearm (Norris *et al.* 1996) and foreskin studies (Grichnik *et al.* 1996). Melanocyte c-Kit expression was confirmed by staining in cultured epidermal and hair follicle melanocytes (Fig. 3B) and colocalisation with NKI/beteb (Fig. 3A). Thus, differentiated melanocytes in all sites can respond to SCF.

In marked contrast, NKI/beteb detected significantly more cells in every region (Fig. 1) as previously reported in the human epidermis (Grichnik *et al.* 1996) and the ORS (Horikawa *et al.* 1996, Commo *et al.* 2004b). Since NKI/beteb stains antigens in premelanosome-like structures as well as premelanosomes and melanosomes (Hayashibe *et al.* 1986), it is presumably recognising very early melanocyte-lineage cells in addition to the more differentiated melanocytes detected by the other antibodies (Adema *et al.* 1993, Horikawa *et al.* 1996). The greater number of cells expressing NKI/beteb than c-Kit, (Fig. 1), confirmed by colocalisation studies (Fig. 3A), indicates small numbers of much less differentiated SCF/c-Kit independent melanocyte-lineage cells in each area. A small percentage of less differentiated stem cells would be predicted within any renewing tissue like the follicle or epidermis. Since epidermal repigmentation

spreads out from hair follicles in treated vitiligo (Ortonne *et al.* 1979, Cui *et al.* 1991, Na *et al.* 1998) or after dermabrasion (Staricco 1960, 1961), undifferentiated infundibular cells could be moving from the mid-follicle or a local reserve of stem cells. In the hair bulb, the difference between NKI/beteb and other antigens' expression was greater (2.2:1) than in the infundibulum or epidermis (1.6:1) and some cells only expressed NKI/beteb during dual staining (Fig. 3e and f). These could be a stem cell reservoir for the bulb, to enable melanin production to continue throughout the very long anagen of human scalp hair; less differentiated melanocytes were detected in the lower bulb rim after dermabrasion (Staricco 1960). Since melanin synthesis produces damaging oxidative stress (Arck *et al.* 2006) replacement melanocytes may be necessary to maintain the pigment. Such less differentiated cells could also be the few melanocytes that persist after anagen, moving upwards in catagen to the telogen capsule and possibly populating the next anagen bulb (Commo & Bernard 2000, Sharov *et al.* 2005). The local signals that stimulate their maturation to pigment-producing melanocytes or migration would not involve SCF as they do not express c-Kit.

In human follicles, the mid-follicle ORS is believed to mirror the rodent follicle bulge as the stem cell niche for melanocyte-lineage cells (Staricco 1959, Sugiyama & Kukita 1976, Horikawa *et al.* 1996, Narisawa *et al.* 1997, Commo & Bernard 2000, Botchkareva *et al.* 2001, Nishimura *et al.* 2002, 2005, Steingrimsdottir *et al.* 2005). Follicular epithelial stem cells also reside in the bulge/permanent ORS (Costarelis *et al.* 1990, Lyle *et al.* 1998, Waters *et al.* 2007). Our mid-follicle ORS results confirm this stem cell niche as four times more cells were detected with NKI/beteb than the other markers, unlike the lower ratios elsewhere (Fig. 1), confirmed by colocalisation studies. These results support earlier observations of amelanocytic melanocytes (Staricco 1959, 1960, 1961, Staricco & Miller-Milinska 1962) and NKI/beteb expression (Horikawa *et al.* 1996, Narisawa *et al.* 1997, Commo & Bernard 2000) in human ORS; they indicate two types of melanocyte-lineage cells: immature cells expressing only NKI/beteb, often seen in groups, presumably stem cells; and small numbers of more differentiated cells expressing melanocyte and c-Kit antigens, probably the next stage of differentiation. This agrees with observations of melanin (Narisawa *et al.* 1997) and both rounded and dendritic cells (Commo & Bernard 2000) in the ORS. No c-Kit expression by most mid-follicle ORS melanocytes concurs with its absence from mice bulge melanocytes (Botchkareva *et al.* 2001) and the c-Kit-independence of mouse melanocyte stem cells (Nishimura *et al.* 2002). The undifferentiated human ORS melanocytes seen here presumably give rise to transient amplifying cells that express c-Kit and migrate under SCF control as in the mouse (Nishimura *et al.* 2002).

Melanin was normally only visible in the hairs and hair bulbs indicating that only matrix melanocytes were producing melanin. Occasional balding epidermis pigment was presumably stimulated by greater exposure to sunlight with

reduced hair cover. These dramatic differences in pigment, despite similar concentrations of differentiated melanocytes in the bulb, epidermis and infundibulum, indicate major variations in melanin-producing activity, presumably due to regional differences in the local regulators. Epidermal pigmentation increases locally in response to u.v. light with greater local production of paracrine regulators, including SCF (Hachiya *et al.* 2001, Slominski *et al.* 2004). The SCF/c-Kit pathway is definitely functional in adult epidermal pigmentation; SCF activates melanocytes *in vivo* (Grichnik *et al.* 1998, Costa *et al.* 1996) and in xenografts (Grichnik *et al.* 1998, Sriwiriyanont *et al.* 2006), while c-Kit antibodies inhibit xenograft melanocytes. Since c-Kit antibodies prevent hair pigmentation in adult mice follicles (Botchkareva *et al.* 2001), c-Kit expression by human follicular melanocytes strongly indicates a role for SCF/c-Kit signalling in human follicular pigmentation. Further support comes from reversible hair greying in cancer patients whose tyrosine kinase interfering chemotherapy (Moss *et al.* 2003, Routhouska *et al.* 2006) presumably blocks post-receptor c-Kit signalling, as c-Kit is a tyrosine kinase receptor.

Androgenetic alopecia

There were no differences between normal and balding scalp in staining patterns or cell concentrations with any antibody in the hair bulb or elsewhere (Fig. 1). Therefore, neither lower bulb melanocyte concentration nor c-Kit expression is responsible for reduced pigmentation during androgenetic alopecia miniaturisation. Although post-receptor signalling defects are possible, this would reduce melanocytes entering the bulb, like mouse anti-c-Kit antibody treatment (Botchkareva *et al.* 2001). A decrease in the local regulators seems more likely. The dermal papilla is a strong candidate to supply local pigment regulators as it determines what hair type is produced (Reynolds & Jahoda 2004), is the focus for the follicle pigment unit (Slominski *et al.* 2004) and the centre of androgen action (Randall 2007). This would enable a mechanism for circulating hormones to alter hair colour in tandem with environmental or developmental changes by altering dermal papilla production of SCF and/or other regulators.

Cultured dermal papilla cells from intermediate androgenetic alopecia follicles secreted about two-thirds less SCF than non-balding cells (Fig. 4). There is substantial evidence that cultured dermal papilla cell behaviour reflects that of the *in vivo* dermal papilla. Cultured human and rat dermal papilla cells induce new follicle formation when re-implanted *in vivo* (Reynolds & Jahoda 2004) and human cells derived from follicles with different androgen responses *in vivo* retain appropriate different characteristics *in vitro* (Randall *et al.* 2003) indicating androgen exposure *in vivo* established their gene expression. These include biologically relevant androgen receptor and 5 α -reductase gene expression in androgen-responsive dermal papilla cells (Ando *et al.* 1999). Cultured beard and androgenetic alopecia cells also exhibit

specific, high-affinity, low-capacity hormone binding, i.e. androgen receptors (Randall *et al.* 1992, Hibberts *et al.* 1998). Particularly importantly, cultured dermal papilla cell capacity to metabolise testosterone to the more potent 5 α -dihydro-testosterone reflects hair growth in 5 α -reductase deficiency with beard, but not pubic, axillary nor non-balding scalp, cells forming 5 α -dihydrotestosterone (Itami *et al.* 1990, Thornton *et al.* 1993, Hamada *et al.* 1996). Additionally, only androgen-responsive cells alter their protein production in response to androgen with effects reflecting their androgen responses *in vivo* (Itami *et al.* 1995, Hibberts & Randall 1996, Obana *et al.* 1997, Thornton *et al.* 1998, Sonada *et al.* 1999, Inui *et al.* 2002, Hibino & Nishiyama 2004). The reduced SCF production by balding dermal papilla cells (Fig. 4) parallels other differences between balding and non-balding scalp dermal papilla cells. Androgenetic alopecia cells contain higher levels of androgen receptors than non-balding cells (Hibberts *et al.* 1998), secrete less growth-promoting factors, produce inhibitory factors that delay rodent hair growth *in vivo* (Hamada & Randall 2006) and respond appropriately to testosterone *in vitro* (Hibberts & Randall 1996, Sonada *et al.* 1999, Inui *et al.* 2002, Hibino & Nishiyama 2004).

In vitro testosterone had no effect on the SCF amounts secreted by balding cells. Although drawing inferences from absent *in vitro* responses are difficult as conditions may not fully recapitulate *in vivo*, androgenetic alopecia cells contain androgen receptors (Hibberts *et al.* 1998) and respond to androgens *in vitro* by decreasing their growth-stimulatory capacity (Hibberts & Randall 1996) and altering their production of TGF- β (Hibino & Nishiyama 2004) and protease nexin-1 (Sonada *et al.* 1999) indicating functional receptor-mediated responses. There was also no alteration in beard dermal papilla cell SCF production with testosterone (Hibberts *et al.* 1996), although testosterone *in vitro* increased beard cell production of mitogenic factors (Thornton *et al.* 1998) and IGF-1 (Itami *et al.* 1995). Beard cells secreted much more SCF, about 250% of normal scalp, and more than 3.5 $\times$ these balding cell levels; beard hairs are normally the darkest hairs. This suggests that androgens alter dermal papilla cell *scf* gene expression either prior to, or during, the early stages of anagen thereby setting that cycle's pigment level. This seems likely as hair colour is normally relatively constant along

a hair, even during several years of growth. These changes in *scf* gene expression are alterable during future hair cycles; treatment with 5 α -reductase inhibitors stimulates regrowth of large, pigmented terminal hairs (Kaufman *et al.* 1998, Olsen *et al.* 2006) and antiandrogens reduce hair colour in hirsutism (Sawers *et al.* 1982). However, antiandrogens do not alter the colour or size of hair in hirsutism quickly, unlike their effect on the sebaceous glands; follicles with long cycles such as the beard also take longer to respond than those with shorter cycles on the thigh (Sawers *et al.* 1982). This supports the concept of androgens acting prior to, or at the beginning of the next anagen rather than affecting follicles during mid-anagen.

In summary, these experiments highlight three types of melanocyte in human scalp anagen hair follicles (Table 1). First, the classic hair bulb melanocyte expressing all melanocyte markers and c-Kit and actively producing melanin, presumably in response to the dermal papilla's SCF. Secondly, melanocytes expressing these antigens, but not making melanin, apparently differentiated cells ready for signals to produce pigment or migrate; these were found in similar numbers in the infundibulum and epidermis, but far fewer in the bulb and mid-follicle ORS. Thirdly, less differentiated cells, only expressing NKI/beteb, probably forms of melanocyte stem cell, were present in small numbers in all areas and were the majority in mid-follicle ORS. During follicular melanocyte culture two types were observed here and elsewhere (Hu *et al.* 1957, Tobin & Bystry, 1996, Ma *et al.* 2006, Na *et al.* 2006), including less differentiated, bipolar forms with great dividing capacity. These SCF/c-Kit-independent cells in the mid-follicle ORS, the follicle stem cell niche, correspond to c-Kit-negative stem cells in mice (Nishimura *et al.* 2002, Steingrimsso, 2005). Similar undifferentiated cells in the bulb may be a reservoir to maintain pigmentation through long scalp anagens and/or melanocytes, which survive catagen and repopulate the next bulb (Commo & Bernard 2000, Sharov *et al.* 2005).

What signals activate the c-Kit-independent, immature cells in the bulb or ORS melanocyte stem cell niche is unknown; other melanocyte modulators such as endothelin 1 and 3, hepatocyte growth factor and basic fibroblast growth factor are possibilities (reviewed Lin & Fisher 2007). Identifying these could facilitate development of useful

Table 1 Summary of pigment formation and distribution of melanocyte-lineage cells in human scalp epidermis and hair follicles. Results of immunohistochemical analysis of human scalp skin with antibodies, NKI/beteb, which recognise all melanocyte-lineage cells, HMB-45 and Mel-2, which recognise differentiated melanocytes, and to c-Kit, the receptor for stem cell factor

	Pigment	NKI/beteb	NKI/beteb, HMB-45, Mel-2, c-Kit
Epidermis	+	+++	+++++++
Infundibulum		++++	+++++++
Mid-follicle ORS		++++	+
Hair bulb	+++++	+++++	+++++++

Presence is indicated by +, with more + indicating greater numbers of cells or pigment, +? indicates epidermal pigment in normal caucasian scalp is usually very limited.

clinical treatments; increasing hair pigmentation in androgenetic alopecia would increase hair visibility even without hair enlargement, and inhibition would reduce it in hirsutism. Such therapies may also reduce canities, age-related greying.

There were no differences in distribution, expression pattern or cell concentration between normal and androgenetic alopecia follicles, but cultured dermal papilla cells from androgenetic alopecia follicles secreted less SCF. This suggests that the mechanism for reduced hair pigment in androgenetic alopecia involves androgens binding to androgen receptors in the dermal papilla, probably at the beginning of anagen, inhibiting SCF gene expression and synthesis by dermal papilla cells throughout anagen causing bulb melanocytes to produce less pigment. Although normal and androgenetic alopecia follicles contained the same concentration of melanocytes, the follicles were smaller and the total number of bulb melanocytes reduced. Since SCF is implicated in the movement of melanocyte-lineage cells into developing follicles and adult mouse bulbs (Jordan & Jackson 2000, Botchkareva *et al.* 2001), this would be expected if less SCF was produced by balding dermal papillae during hair bulb reformation in early anagen. Overall, these observations suggest that modifiers of SCF/c-Kit signalling, or the unidentified regulators of the c-Kit-negative melanocyte stem cells, could offer future routes to alter hair pigmentation for clinical effect in androgen-modulated hair disorders including hirsutism and androgenetic alopecia or cosmetically for greying.

Acknowledgements

The authors are grateful for helpful guidance with melanocyte culture from Prof Desmond Tobin and for beneficial discussions with Prof John Wood, both at the School of Life Sciences, University of Bradford. They also gratefully acknowledge a fellowship from CAPES, Brazil to support I O de O and a research studentship from the University of Bradford for T J J. The authors have no conflict of interest that would prejudice the impartiality of this paper.

References

Adema GJ, de Boer AJ, Van't Hullenaar R, Denijn M, Ruiter DJ, Vogel AM & Figdor CG 1993 Melanocyte-lineage specific antigens recognized by monoclonal antibodies NK1/beteb, HMB-50 and HMB-45 are encoded by a single cDNA. *American Journal of Pathology* **143** 1579–1585.

Ando Y, Yamaguchi Y, Hamada K, Yoshikawa K & Itami S 1999 Expression of mRNA for androgen receptor, 5 α -reductase and 17 β -hydroxysteroid dehydrogenase in human dermal papilla cells. *British Journal of Dermatology* **141** 840–845.

Arck PC, Overall R, Spatz K, Liezman C, Handjiski B, Klapp BF, Birch-Machin MA & Peters EMJ 2006 Towards a 'free radical theory of graying': melanocyte apoptosis in the aging human hair follicle is an indicator of oxidative stress induced tissue damage. *FASEB Journal* **20** E908–E920.

Bhawan J 1997 Mel-5: a novel antibody for differential diagnosis of epidermal pigmented lesions of the skin in paraffin-embedded sections. *Melanoma Research* **7** 43–46.

Botchkareva NV, Khlgatian M, Longley BJ, Botchkarev VA & Gilchrist BA 2001 SCF/c-Kit signaling is required for cyclic regeneration of the hair pigmentation unit. *FASEB Journal* **15** 645–658.

Bradford MM 1976 A rapid and sensitive method for the quantification of microgram quantities of protein utilising the principle of protein dye binding. *Analytical Biochemistry* **72** 248–254.

Bubenik GA & Bubenik AB 1985 Seasonal variations in hair pigmentation of white-tailed deer and their relationship to sexual activity and plasma testosterone. *Journal of Experimental Zoology* **235** 387–395.

Burnett JB, Holstein TJ & Quevedo WC 1969 Electrophoretic variations of tyrosinase in follicular melanocytes during the hair growth cycle in mice. *Journal of Experimental Zoology* **171** 369–376.

Castanet J & Ortonne J-P 2000 Hair pigmentation. In *Hair and its Disorders: Biology, Pathology and Management*, pp 49–65. Eds FM Camacho, VA Randall & VH Price. London: Martin Dunitz.

Chase HB 1958 The behaviour of pigment cells and epithelial cells in the hair follicle. In *The Biology of Hair Growth*, pp 29–237. Eds W Montagna & RA Ellis. New York: Academic Press.

Commo S & Bernard BA 2000 Melanocyte subpopulation turnover during human hair cycle: an immunohistochemical study. *Pigment Cell Research* **13** 253–259.

Commo S, Gaillard O & Bernard BA 2004a Absence of TRP-2 in melanogenic melanocytes of human hair. *Pigment Cell Research* **17** 488–497.

Commo S, Gaillard O & Bernard BA 2004b Human hair greying is linked to a specific depletion of hair follicle melanocytes affecting both the bulb and the outer root sheath. *British Journal of Dermatology* **150** 435–443.

Costa JJ, Demetri GD & Harrist TJ 1996 Recombinant human stem cell factor (kit ligand) promotes human mast cell and melanocyte hyperplasia and functional activation *in vivo*. *Journal of Experimental Medicine* **183** 2681–2686.

Costarelis G, Sun T-T & Lavker RM 1990 Label-retaining cells reside in the bulge area of pilosebaceous unit: implications for follicular stem cells, hair cycle and skin carcinogenesis. *Cell* **61** 1329–1337.

Costin GE & Hearing VJ 2007 Human skin pigmentation: melanocytes modulate skin colour in response to stress. *FASEB Journal* **21** 976–994.

Cui J, Shen LY & Wang GC 1991 Role of hair follicles in the repigmentation of vitiligo. *Journal of Investigative Dermatology* **97** 410–416.

Eisinger M & Marko O 1982 Selective proliferation of normal human melanocytes *in vitro* in the presence of phorbol ester and cholera toxin. *PNAS* **79** 2018–2022.

Fleischman RA, Saltman DL, Stastny V & Zneimer S 1991 Deletion of the c-Kit proto-oncogene in the human developmental defect piebald trait. *PNAS* **88** 10885–10889.

Flux JEC 1970 Colour change of mountain hares (*Lepus timidus scotius*) in north-east Scotland. *Journal of Zoology* **162** 345–358.

Geissler EN, Cheng SY, Gusella JF & Housmann D 1988 The dominant-white spotting (W) locus of the mouse encodes the c-kit proto-oncogene. *Cell* **55** 185–192.

Grichnik JM, Wazir NA, Burch JA, Byers JD, Garcia CA, Clark E & Shea CR 1996 KIT expression reveals a population of precursor melanocytes in human skin. *Journal of Investigative Dermatology* **106** 967–971.

Grichnik JM, Burch JA, Burchette J & Shea CR 1998 The SCF/KIT pathway plays a critical role in the control of normal human melanocyte homeostasis. *Journal of Investigative Dermatology* **111** 233–238.

Haase E, Ito S & Wakanatsu K 1995 Influences of sex, castration and androgens on the eumelanin and pheomelanin contents of different feathers in wild mallards. *Pigment Cell Research* **8** 164–170.

Hachiya A, Kobayashi A, Ohuchi A, Takema Y & Imokawa G 2001 The paracrine role of stem cell factor/c-kit signaling in the activation of human melanocytes in ultraviolet-b-induced pigmentation. *Journal of Investigative Dermatology* **116** 578–586.

Halaban R & Alfano FD 1984 Selective elimination of fibroblasts from cultures of normal human melanocytes. *In Vitro* **20** 447–450.

Hamada K & Randall VA 2006 Inhibitory autocrine factors produced by the mesenchyme-derived hair follicle dermal papilla may be a key to male pattern baldness. *British Journal of Dermatology* **54** 609–618.

- Hamada K, Thornton MJ, Laing I, Messenger AG & Randall VA 1996 The metabolism of testosterone by dermal papilla cells cultured from human pubic and axillary hair follicles concurs with hair growth in 5 α -reductase deficiency. *Journal of Investigative Dermatology* **106** 1017–1022.
- Hamilton JB 1951 Patterned loss of hair in man; types and incidence. *Annals of the New York Academy of Sciences* **53** 708–728.
- Hamilton JB 1958 Age, sex and genetic factors in the regulation of hair growth in man: a comparison of Caucasian and Japanese populations. In *The Biology of Hair Growth*, pp 399–433. Eds W Montagna & RA Ellis. New York: Academic Press.
- Hayashibe K, Mishima Y, Ichihashi M & Kawai M 1986 Melanosomal antigenic expression on the cell surface and intracellular subunits within melanogenic compartments of pigment cells: analysis by antimelanosome-associated monoclonal antibody. *Journal of Investigative Dermatology* **87** 89–94.
- Hibberts NA & Randall VA 1996 Testosterone inhibits the capacity of cultured cells from human balding scalp dermal papilla cells to produce keratinocyte mitogenic factors. In Degroot LJ, Jameson JL, editors, *Endocrinology* 5th Ed Section XIV Burger H G, editor. Male Reproduction, Philadelphia: W B Saunders Co. 303–306.
- Hibberts NA, Messenger AG & Randall VA 1996 Dermal papilla cells derived from beard hair follicles secrete more stem cell factor (SCF) in culture than scalp cells or dermal fibroblasts. *Biochemical and Biophysical Research Communications* **222** 401–405.
- Hibberts NA, Howell AE & Randall VA 1998 Dermal papilla cells from human balding scalp hair follicles contain higher levels of androgen receptors than those from non-balding scalp. *Journal of Endocrinology* **156** 59–65.
- Hibino T & Nishiyama T 2004 Role of TGF- β 2 in the human hair cycle. *Journal of Dermatology Science* **35** 9–18.
- Horikawa T, Norris DA, Johnson TW, Zekman T, Dunscomb N, Bennion SD, Jackson RL & Morelli JG 1996 Dopa-negative melanocytes in the outer root sheath of human hair follicles express premelanosomal antigens but not a melanosomal antigen or the melanosome-associated glycoproteins, tyrosinase, TRP-1 and TRP-2. *Journal of Investigative Dermatology* **106** 28–35.
- Hu F, Staricco RG, Pinkus H & Forsnaugh RP 1957 Human melanocytes in tissue culture. *Journal of Investigative Dermatology* **28** 15–32.
- Inui S, Fukuzato Y, Nakajima F, Yoshikawa K & Itami S 2002 Androgen-inducible TGF β 1 from balding dermal papilla cells inhibits epithelial cell growth: a clue to understand paradoxical effects of androgen on human hair growth. *FASEB Journal* **16** 1967–1969.
- Itami S, Kurata S & Takayasu S 1990 5- α -reductase activity in cultured human dermal papilla cells from beard compared with reticular dermal fibroblasts. *Journal of Investigative Dermatology* **94** 150–152.
- Itami S, Kurata S, Sonada T & Takayasu S 1995 Interactions between dermal papilla cells and follicular epithelial cells *in vitro*: effect of androgen. *British Journal of Dermatology* **132** 527–532.
- Jansen VAA & van Baalen M 2006 Altruism through beard chromodynamics. *Nature* **440** 663–666.
- Jordan SA & Jackson IJ 2000 MGF(KIT ligand) is a chemokinetic factor for melanoblast migration into hair follicles. *Developmental Biology* **225** 424–436.
- Kapur RP, Bigler SA, Skelly M & Gown AM 1992 Anti-melanoma monoclonal antibody HMB45 identifies an oncofetal glycoconjugate associated with immature melanosomes. *Journal of Histochemistry and Cytochemistry* **40** 207–212.
- Kaufman KD, Olsen EA, Whiting D, Savin R, Devillez R, Bergfeld W, Pricew VH, Van Neste D, Roberts JL, Hordinsky M *et al.* 1998 Finasteride in the treatment of men with androgenetic alopecia. *Journal of the American Academy of Dermatology* **39** 578–589.
- Kligman AG 1959 The human hair cycle. *Journal of Investigative Dermatology* **33** 307–316.
- Kunisada T, Yoshida H, Yamazaki H, Miyamoto A, Hemmi H, Nishimura E, Shultz LD, Nishikawa S & Hayashi S 1998a Transgene expression of steel factor in the basal layer of epidermis promotes survival, proliferation, differentiation and migration of melanocyte precursors. *Development* **125** 2915–2923.
- Kunisada T, Lu SZ, Yoshida H, Nishikawa S, Mizoguchi M, Hayashi S, Tyrrell L, Williams DA, Wang X & Longley BJ 1998b Murine cutaneous mastocytosis and epidermal melanocytosis induced by keratinocyte expression of transgenic stem cell factor. *Journal of Experimental Medicine* **187** 1565–1573.
- Lin JY & Fisher DE 2007 Melanocyte biology and skin pigmentation. *Nature* **445** 843–850.
- Lü ZF, Cai SQ, Wu JJ & Zheng M 2006 Biological characterisation of cultured dermal papilla cells and hair follicle regeneration *in vitro* and *in vivo*. *Chinese Medical Journal* **119** 275–281.
- Lyle S, Christofidou-Solomidou M, Lin Y, Elder DE, Albelda S & Costarelis G 1998 The C8/144B monoclonal antibody recognizes cytokeratin 15 and defines the location of human hair follicle stem cells. *Journal of Cell Science* **111** 3179–3188.
- Ma H-J, Yue X-Z, Wang D-G, Li C-R & Zhu W-Y 2006 A modified method for purifying amelanotic melanocytes from human hair follicles. *Journal of Dermatology* **33** 239–248.
- Marshall WA & Tanner JM 1970 Variations in the pattern of pubertal changes in boys. *Archives of Disease in Childhood* **45** 13–23.
- Marzia D, Schatten G & Sale W 1975 Adhesion of cells to surfaces coated with poly-L-lysine. *Journal of Cell Biology* **66** 198–203.
- Messenger AG 1984 The culture of dermal papilla cells from human hair follicles. *British Journal of Dermatology* **110** 685–689.
- Moss KG, Toner GC, Cherrington JM, Mendel DB & Laird AD 2003 Hair depigmentation is a biological readout for pharmacological inhibition of kit in mice and humans. *Journal of Pharmacology and Experimental Therapeutics* **307** 476–480.
- Na GY, Seo SK & Choi SK 1998 Single hair grafting for the treatment of vitiligo. *Journal of the American Academy of Dermatology* **38** 580–584.
- Na GY, Paek SH, Park BC, Kim DW, Lee WJ, Lee SJ, Kim MK & Kim JC 2006 Isolation and characterization of outer root sheath melanocytes of human hair follicles. *Journal of Dermatology* **155** 902–909.
- Narisawa Y, Kohda H & Tanaka T 1997 Three-dimensional demonstration of melanocyte distribution of human hair follicles: special reference to the bulge area. *Acta Dermato-Venerologica* **77** 97–101.
- Nishikawa S, Kusakabe M, Yoshinaga K, Ogawa M, Hayashi S, Kunisada T, Era T, Sakakura T & Nishikawa S 1991 *In utero* manipulation of coat colour formation by a monoclonal anti-c-Kit antibody: two distinct waves of c-Kit-dependency during melanocyte development. *EMBO Journal* **10** 2111–2118.
- Nishimura EK, Jordan S, Oshima H, Yoshida H, Osawa M, Moriyama M, Jackson IJ, Barrandon Y, Miyachi Y & Nishikawa S 2002 Dominant role of the niche in melanocyte stem-cell fate determination. *Nature* **416** 854–860.
- Nishimura EK, Scott RG & Fisher DE 2005 Mechanisms of hair graying: incomplete melanocyte stem cell maintenance in the niche. *Science* **307** 720–724.
- Norris A, Todd C, Graham A, Quinn AG & Thody AJ 1996 The expression of the c-Kit receptor by epidermal melanocytes may be reduced in vitiligo. *British Journal of Dermatology* **134** 299–306.
- Obana N, Chang C & Uno H 1997 Inhibition of hair growth by testosterone in the presence of dermal papilla cells from the frontal bald scalp of the post-pubertal stump-tailed macaque. *Endocrinology* **138** 356–361.
- Okura M, Maeda H, Nishikawa S & Mizoguchi M 1995 Effects of monoclonal anti-c-Kit antibody (ACK2) on melanocytes in newborn mice. *Journal of Investigative Dermatology* **105** 322–328.
- Olsen EA, Hodinsky M, Whiting D, Stough D, Hobbs S, Ellis ML, Wilson T & Rittmaster RS, Dutasteride Alopecia Research Team 2006 The importance of dual 5 α -reductase inhibition in the treatment of male pattern hair loss: results of a randomized placebo-controlled study of dutasteride versus finasteride. *Journal of the American Academy of Dermatology* **55** 1014–1023.
- Ortonne JP, MacDonald DM, Micoud A & Thivolet J 1979 PUVA-induced repigmentation of vitiligo: a histochemical (split-DOPA) and ultrastructural study. *British Journal of Dermatology* **101** 1–12.
- Randall VA 1994 Androgens and human hair growth. *Clinical Endocrinology* **40** 439–457.
- Randall VA 2007 Hormonal regulation of hair follicles exhibits a biological paradox. *Seminars in Cell and Developmental Biology* **18** 274–285.

- Randall VA, Thornton MJ, Hamada K, Redfern CP, Nutbrown M, Ebling FJ & Messenger AG 1991 Androgens and the hair follicle: cultured human dermal papilla cells as a model system. *Annals of the New York Academy of Sciences* **642** 355–375.
- Randall VA, Thornton MJ & Messenger AG 1992 Cultured dermal papilla cells from androgen-dependant human hair follicles (e.g. beard) contain more androgen receptors than those from non-balding areas of scalp. *Journal of Endocrinology* **133** 141–147.
- Randall VA, Hibberts NA & Hamada K 1996 A comparison of the culture and growth of dermal papilla cells derived from normal and balding (androgenetic alopecia) scalp. *British Journal of Dermatology* **134** 437–444.
- Randall VA, Sundberg JP & Philpott MP 2003 Animal and *in vitro* models for the study of hair follicles. *Journal of Investigative Dermatology. Symposium Proceedings* **8** 39–45.
- Reynolds AJ & Jahoda CAB 2004 Cultured human and rat tooth papilla cells induce hair follicle regeneration and fiber growth. *Differentiation* **72** 566–575.
- Rousseau K, Kausar S, Pritchard LE, Warhurst A, Oliver RL, Slominski A, Wei ET, Thody AJ, Tobin DJ & White A 2007 Proopiomelanocortin (POMC), the ACTH/melanocortin precursor, is secreted by human epidermal keratinocytes and melanocytes and stimulates melanogenesis. *FASEB Journal* **21** 1844–1856.
- Routhousa S, Gilliam AC & Mirirani P 2006 Hair depigmentation during chemotherapy with a class III/V receptor tyrosine kinase inhibitor. *Archives of Dermatology* **142** 1477–1479.
- Sawers RS, Randall VA & Iqbal MJ 1982 Studies on the clinical and endocrine aspects of antiandrogens. In *Hirsutism*, pp 145–168. Ed. SL Jeffocate. London: Wiley.
- Sharov AA, Tobin DJ, Sharova TY, Atoyan R & Botchkarev VA 2005 Changes in different melanocyte population during hair follicle involution (catagen). *Journal of Investigative Dermatology* **125** 1259–1267.
- Slominski A, Tobin DJ, Shibahara S & Wortsman J 2004 Melanin pigmentation in mammalian skin and its hormonal regulation. *Physiological Reviews* **84** 1155–1228.
- Sonada T, Asada Y, Kurata S & Takayasu S 1999 The mRNA for protease nexin-1 is expressed in human dermal papilla cells and its level is affected by androgen. *Journal of Investigative Dermatology* **113** 308–313.
- Spritz RA, Holmes SA, Itin P & Kuster W 1992 Novel mutations of the KIT (mast/stem cell growth factor receptor) proto-oncogene in human piebaldism. *Journal of Investigative Dermatology* **101** 22–25.
- Sriwiriyanont P, Ohuchi A, Hachiya A, Visscher MO & Boissy RE 2006 Interaction between stem cell factor and endothelin-1: effects on melanogenesis in human skin xenografts. *Laboratory Investigation* **86** 1115–1125.
- Staricco RG 1959 Amelanotic melanocytes in the outer sheath of the human hair follicle—preliminary and short report. *Journal of Investigative Dermatology* **33** 295–297.
- Staricco RG 1960 The melanocytes and the hair follicle. *Journal of Investigative Dermatology* **35** 185–194.
- Staricco RG 1961 Mechanism of migration of the melanocytes from the hair follicle into the epidermis following dermabrasion. *Journal of Investigative Dermatology* **36** 99–104.
- Staricco RG & Miller-Milinska A 1962 Activation of the amelanotic melanocytes in the outer root sheath of the hair follicle following ultra violet rays exposure. *Journal of Investigative Dermatology* **39** 163–164.
- Steingrimsdottir E, Copeland GC & Jenkins N 2005 Melanocyte stem cell maintenance and hair graying. *Cell* **121** 9–12.
- Sugiyama S & Kukita A 1976 Melanocyte reservoir in the hair follicles during the hair growth cycle: an electron microscopic study. In *Biology and Disease of the Hair*, pp 181–200. Ed. K Toda. Baltimore: University Park Press.
- Thornton MJ, Laing I, Hamada K, Messenger AG & Randall VA 1993 Differences in testosterone metabolism by beard and scalp hair follicle dermal papilla cells. *Clinical Endocrinology* **39** 633–639.
- Thornton MJ, Hamada K, Messenger AG & Randall VA 1998 Beard, but not scalp, dermal papilla cells secrete autocrine growth factors in response to testosterone *in vitro*. *Journal of Investigative Dermatology* **111** 727–732.
- Tobin DJ & Bystryń JC 1996 Different populations of melanocytes are present in hair follicles and epidermis. *Pigment Cell Research* **9** 304–310.
- Tobin DJ & Kausar S 2005 Hair melanocytes as neuro-endocrine sensors—pigments for our imagination. *Molecular and Cellular Endocrinology* **243** 1–11.
- Tobin D, Colen SR & Bystryń JC 1995 The isolation and long term culture of human hair follicle melanocytes. *Journal of Investigative Dermatology* **104** 86–89.
- Waters JM, Richardson GD & Jahoda CA 2007 Hair follicle stem cells. *Seminars in Cell and Developmental Biology* **18** 245–254.
- West PM & Packer C 2002 Sexual selection, temperature and the lion's mane. *Science* **297** 1339–1343.
- Williams DE, de Vries P, Namen AE, Widmer MB & Lyman SD 1992 The steel factor. *Developmental Biology* **151** 368–376.
- Zhu W, Zhang R, Ma H & Wang D 2004 Isolation and culture of amelanotic melanocytes from human hair follicles. *Pigment Cell Research* **17** 668–673.
- Zsebo KM, Williams DA, Geissler EN, Broudy VC, Martin FH, Atkins HL, Hsu RY, Birkett NC, Okino NH, Murdock DC *et al.* 1990 Stem cell factor is encoded at the Sl locus of the mouse and is the ligand for the c-Kit tyrosine kinase receptor. *Cell* **63** 213–224.

Received in final form 20 November 2007

Accepted 24 January 2008

Made available online as an Accepted Preprint
24 January 2008