Cardiac corticosteroid receptors mediate the enlargement of the ovine fetal heart induced by chronic increases in maternal cortisol

Seth A Reini¹, Garima Dutta, Charles E Wood¹ and Maureen Keller-Wood

Department of Pharmacodynamics, College of Pharmacy, University of Florida, Box 100487, Gainesville, Florida 32610-0487, USA

¹Department of Physiology and Functional Genomics, College of Medicine, University of Florida, Gainesville, Florida 32610-0274, USA

(Correspondence should be addressed to M Keller-Wood; Email: kellerwd@cop.ufl.edu)

Abstract

Previous studies have demonstrated that modest, physiologically relevant increases in maternal cortisol in late gestation result in enlargement of the fetal heart. In this study, we investigated the role of mineralocorticoid receptor (MR) or glucocorticoid receptor (GR) in this enlargement. Ewes with single fetuses were randomly assigned at ~120 days of gestation to one of four groups: maternal cortisol infusion (1 mg/kg per day, cortisol); maternal cortisol infusion with fetal intrapericardial infusion of the MR antagonist (MRa) potassium canrenoate (600 µg/day; cortisol+MRa); maternal cortisol infusion with fetal intrapericardial infusion of the GR antagonist (GRa) mifepristone (50 µg/day, cortisol+GRa); and maternal saline infusion (control). At ~130 days of gestation, fetal heart to body weight ratio and right ventricular (RV) and left ventricular (LV) free wall thicknesses were increased in the cortisol group

when compared with control group. Fetal hearts from the cortisol+MRa group weighed significantly less, with thinner LV, RV, and interventricular septum walls, when compared with the cortisol group. Fetal hearts from the cortisol+GRa group had significantly thinner RV walls than the cortisol group. Fetal arterial pressure and heart rate were not different among groups at 130 days. Picrosirius red staining of fetal hearts indicated that the increased size was not accompanied by cardiac fibrosis. These results suggest that physiologic increases in maternal cortisol in late gestation induce fetal cardiac enlargement via MR and, to a lesser extent, by GR, and indicate that the enlargement is not secondary to an increase in fetal blood pressure or an increase in fibrosis within the fetal heart.

Journal of Endocrinology (2008) **198**, 419–427

Introduction

In late gestation, normal fetal growth and fetal cardiovascular homeostasis is dependent on the proper regulation of maternal cortisol levels. Although reductions in maternal cortisol prevent the normal increases in maternal plasma volume and uteroplacental blood flow and reduce fetal growth (Jensen *et al.* 2002a, 2005), increases in maternal cortisol also alter fetal growth. Chronically increased maternal cortisol levels, within the range that occurs with maternal stress, reduce fetal growth rates while increasing heart growth in fetal sheep (Jensen *et al.* 2002b, 2005).

The mechanisms by which chronically elevated maternal cortisol levels increase the size of the fetal heart are not known. Giraud *et al.* (2006) have shown that cortisol chronically infused directly into the coronary artery increased cell cycle activity in myocytes of late gestation sheep fetuses, suggesting a direct induction by cortisol of hyperplastic growth rather than hypertrophic growth. Conversely, it has been demonstrated that large doses of cortisol infused directly into the fetus in late gestation causes left ventricular (LV) hypertrophy along with an increase in fetal arterial pressure

and cardiac expression of angiotensinogen mRNA (Lumbers *et al.* 2005). We have shown that maternal cortisol infusion in sheep during late gestation caused an increase in fetal heart size and wall thickness without increasing fetal arterial pressure or cardiac angiotensinogen; we found an increase in the ratio of angiotensin type 2 receptor (AT2 receptor) to AT1 receptor mRNA in the fetal heart, suggesting that the renin-angiotensin system (RAS) may play a key role in the enlargement process. Furthermore, in the same study it was observed that LV expression of 11β-hydroxysteroid dehydrogenase (11β-HSD2) mRNA, the enzyme that converts cortisol into cortisone, decreased in the fetal hearts in response to the elevated cortisol, suggesting that cortisol can act directly on mineralocorticoid receptors (MRs) or glucocorticoid receptors (GRs) to induce the cardiac enlargement (Reini *et al.* 2006).

In adult hearts, both the MR and the RAS have been implicated in cardiac fibrosis and hypertrophy after injury (Fraccarollo *et al.* 2003, 2005, Xiao *et al.* 2004). We propose that corticosteroid receptors also play a role in cardiac enlargement in the fetal heart, although by mechanisms independent of cardiac injury and fibrosis. The purpose of this

study was to test the hypothesis that increase in fetal heart weight and wall thickness in response to increased maternal cortisol is mediated by cardiac corticosteroid receptors, MR and/or GR, and to determine if cardiac fibrosis accompanies the cardiac enlargement in response to cortisol. We hypothesized that cortisol acts within the myocardium on MRs, and to a lesser degree on GRs, to induce enlargement of the fetal heart. We also hypothesized that cardiac fibrosis is not involved in the enlargement of the heart observed in the fetuses of cortisol-infused ewes.

Materials and Methods

Experimental design

Ewes (*Ovis aries*) pregnant with single fetuses were studied. All animal use was approved by the University of Florida Institutional Animal Care and Use Committee and conformed to the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Ewes and their fetuses were operated on between 118 and 123 days of gestation (term ~148 days). Animals were randomly assigned to one of four groups at the time of surgery. The first group consisted of six control ewes infused with saline; the second group consisted of five ewes to which cortisol (hydrocortisone hemisuccinate; Sigma) was administered by continuous i.v. infusion (1 mg/kg per day; cortisol); the third group consisted of six ewes to which cortisol was infused, with infusion of the MR antagonist potassium canrenoate (Sigma; 600 µg/day; cortisol+MRa) directly into the pericardial space of the fetus; and the fourth group consisted of four ewes to which cortisol was infused, with infusion of the GR antagonist mifepristone (Sigma; 50 µg/day; cortisol+GRa) directly into the pericardial space of the fetus. For the control and cortisol groups, there were no infusions into the pericardial space. The intrapericardial infusions were performed by use of Alzet minipumps (model 2ML2; 5 µl/h; Durect Corporation, Cupertino, CA, USA) in order to achieve continuous infusion of the antagonists into the pericardial space without any appreciable increase in pericardial fluid volume (0.12 ml/day). The doses of MR and GR antagonists were calculated based on their effective systemic doses, and scaled to reflect the smaller distribution volume of the fetal heart (20 g). Because these drugs are steroid (mifepristone) or lactone (canrenoate) derivatives, they are able to distribute throughout the tissue over the 10 days of study after mixing in the pericardial fluid. Effects of the MR and GR antagonists were confirmed using immunohistochemistry to confirm the expected cellular redistribution of receptors with antagonist administration (see below).

The cortisol dose and the duration of cortisol infusion (10 days) were determined based on a previous study in this laboratory (Jensen *et al.* 2005) showing that infusion at this rate and duration produces levels similar to mild maternal stressors and results in enlargement of the fetal heart.

Surgical procedures

Halothane (1.5–2.5%) in oxygen was used to anesthetize ewes during surgery. Fetal femoral tibial artery catheters and an amniotic fluid catheter were placed as described previously (Wood & Rudolph 1983, Jensen *et al.* 2002a). Catheters were also placed in the fetal pericardial space for the delivery of the drug as described previously (Wood 2002). In each case, an incision was made in the uterus over the left side of the fetal chest and an incision was made between the third and fourth fetal ribs. The fetal skin was marsupialized to the uterus to prevent leakage of amniotic fluid. The fetal heart was exposed and a small incision was made in the pericardium, through which a silastic catheter (0.76 mm i.d., 1.65 mm o.d.; Dow Corning, Midland, MI, USA) was placed and held in place with the use of a purse-string suture (4-0 Tevdek; Teleflex Medical, Mansfield, MA, USA). For infusion of potassium canrenoate, the silastic catheter was connected to a Tygon tubing connector (1.27 mm o.d.; St Gobain Performance plastics; Akron, OH, USA) which was connected at its other end to the Alzet pump containing the drug (50 mg/ml in 0.9% saline). Because mifepristone is not soluble in aqueous solution and therefore cannot be directly loaded into the pump reservoir, mifepristone was dissolved in 47.5% ethanol in saline (0.42 mg/ml ethanol–saline); this solution was placed in a polyethylene tubing (1.40 mm i.d., 1.90 mm o.d.) which was then connected to the silastic pericardial catheter on one end and to the Alzet minipump on the other end using smaller gauge polyethylene tubing. The Alzet pump, filled with saline, provided the flow to pump the mifepristone solution from the tubing into the pericardium. The pump was placed under the skin of the fetus near the scapula. In the control group, five of six fetuses also had pericardial catheters placed, but no infusion was delivered; in the cortisol group, three of five fetuses had pericardial catheters placed, but no infusion was delivered.

After closure of the uterus, catheters were placed in the maternal femoral artery and vein and routed to the maternal flank. All ewes were treated with flunixinam (1 mg/kg i.m.; Fort Dodge Animal Health, Fort Dodge, IA, USA) at the end of the surgical procedure, before recovery from anesthesia.

Ewes were returned to their pen after recovery from anesthesia. At this time, the i.v. infusion of cortisol (1 mg/kg per day cortisol as cortisol hemisuccinate in normal saline; Sigma) or infusion of saline to the ewe was initiated. Maternal infusions were delivered through a 0.22 µm filter (Millipore Corp. Medford, MA, USA) via a syringe pump at the rate of 1.17 ml/h. Animals were housed in individual pens with access to water, food, and salt blocks *ad libitum*. Ampicillin (500 mg i.m. bid; Webster Veterinary) was administered for 3 days postoperatively. Flunixinamine was administered on the morning after surgery.

Experimental protocol

Fetuses were studied from the day of surgery until death on 129–132 days of gestation. All cortisol-infused ewes and their

fetuses were killed on day 10 of infusion. Fetal and maternal blood samples were withdrawn on day 5 (124–126 days of gestation) and day 10 (129–132 days of gestation) after the start of the infusion for determination of blood gases, plasma cortisol, and plasma adrenocorticotropin (ACTH) concentrations. All blood samples were taken immediately after entering the room in which the ewes were housed in order to minimize the effect of handling on plasma ACTH and cortisol. On day 10 of infusion, maternal and fetal blood pressure and heart rate were recorded over a 40-min interval using LabView software (National Instruments, Austin, TX, USA) and disposable pressure transducers (Transpac; Hospira, Lake Forest, IL, USA). Amniotic fluid pressures were subtracted from fetal intra-arterial pressures in order to calculate fetal arterial pressure. In two animals, one in the cortisol group and one in the cortisol + MRa group, we were unable to reliably measure fetal heart rate; data from those two fetuses are excluded from analysis.

The ewe was euthanized on day 10 using an overdose of euthanasia solution containing pentobarbital, and the fetus was removed and weighed. The fetal heart was also immediately dissected, blotted to remove blood from the chambers, and weighed. Ventricular and septal wall thicknesses were measured using a micrometer at a standardized site on the heart, taking care to exclude measurement at the level of the papillary muscles or valves.

Analysis

Blood gases and pH were measured with a blood gas/electrolyte analyzer (ABL77; Radiometer America, Westlake, OH, USA). Electrolytes (sodium and potassium) were measured using an electrolyte analyzer (Roche 9180). For measurement of packed cell volume (PCV), blood was spun in microcapillary tubes for 3 min at 5 000 g. (Damon Division, International Equipment, Needham Heights, MA, USA). Plasma protein was determined using a refractometer.

Plasma ACTH was measured by RIA, using an antibody to ACTH (1–39) (Bell *et al.* 1991), and plasma cortisol concentration was measured using a commercially available enzyme immunoassay kit (EA 65, Oxford Biomedical, Oxford, MI, USA) that has minimal cross-reactivity with cortisone (2.08%).

Immunohistochemical localization of MR and GR

At the time of killing, a section of the fetal heart was fixed in 4% buffered paraformaldehyde overnight. Hearts were dehydrated with increasing concentrations of reagent alcohol followed by xylene, embedded in paraffin wax, cut into 10 µm thick sections on a Zeiss rotary microtome, and placed on poly-L-lysine coated slides. The sections were stained with anti-GR (Santa Cruz Biotechnology, Santa Cruz, CA, USA M-20) or anti-MR (M1-18, 6G1, gift of Gomez-Sanchez *et al.* (2006)) as described previously (Reini *et al.* 2006). This analysis was performed to assess the ability of the drugs to act

in the heart and cause the expected changes in cytonuclear localization of the receptors. The MR antagonist potassium canrenoate acts in a similar manner to spironolactone and would therefore be expected to prevent nuclear localization of MR (Lombes *et al.* 1994, Fejes-Toth *et al.* 1998); conversely the GR antagonist mifepristone (also known as RU486) causes nuclear localization even in the absence of agonist (Jewell *et al.* 1995, Scheuer *et al.* 2004). The localization observed (Fig. 1) is consistent with these effects. In the control fetuses GRs were primarily located in the cytosol, whereas MR was apparent in the cytosol and nucleus. A dramatic increase in MR localization to the nucleus was apparent in the cortisol-treated fetuses, indicating cortisol activation of MR. We did not find as dramatic an increase in nuclear GR with cortisol, indicating that fewer GR are activated. In the case of MR antagonist administration, fewer MR were apparent in the nucleus than with cortisol alone, whereas with GR antagonist, equivalent MR localization to the nucleus occurred as with cortisol alone. Consistent with the known effect of mifepristone, in GR antagonist-treated ewes, there was more nuclear GR than in the case of cortisol alone or cortisol + MRa.

Figure 1 Immunohistochemical localization of MR (left column) and GR (right column) in representative hearts from fetuses of control, cortisol, cortisol + MRa, and cortisol + GRa groups. All photos at 40× power.

Collagen staining

Sections from each group ($n=4-6$) were stained with picosirius red (Sigma) in order to determine collagen content. Sections were hydrated and immersed in Sirius red (0.1% in saturated picric acid). The sections were then washed in acidified water (0.5% glacial acetic acid), dehydrated, and mounted in permount. All images were visualized using an Olympus DP71 microscope and Olympus software. Ten pictures of LV, five of right ventricle (RV) and five of septum, were taken from each heart in areas without large blood vessels so that primarily interstitial, rather than perivascular, collagen deposition could be quantified. Picosirius red staining was quantified using Image J software (NIH, Bethesda, MD, USA) by three different people who were blinded as to the experimental group. The average value of the percentage of the image that stained red from these three observations was calculated.

Data analysis

Fetal heart weight was normalized to body weight. The heart weight to body weight ratio, LV, RV, and septal thickness, fetal and maternal blood pressure and heart rate, as well as fetal and maternal plasma ACTH and cortisol, sodium and potassium, and PCVs were analyzed by one-way ANOVA with multiple comparisons using Duncan's method (Zar 1984). Plasma hormone (cortisol and ACTH) and protein concentrations were also analyzed by one-tailed *t*-test, comparing the data from all three groups of cortisol-treated ewes to the data from the control group (Zar 1984). Average cortisol values were calculated from the day 5 and day 10 values and were log transformed before analysis. The Mann-Whitney rank sum test was used for maternal plasma protein analysis at day 10 (Zar 1984).

Values for the picosirius red staining were analyzed by two-way ANOVA in order to determine significance across the cortisol treatment groups and areas of the heart (LV, RV, and septum); the percent stained area data were transformed using arcsine prior to analysis to correct for heteroscedascity (Winer 1971).

For all analyses, $P<0.05$ was used as the criterion for statistical significance.

Results

Maternal physiology

Maternal cortisol concentrations were significantly increased in the ewes treated with cortisol when compared with the non-treated ewes (average of days 5 and 10, 9.0 ± 0.9 vs 5.9 ± 1.4 ng/ml). When the four groups were compared individually, there was a trend for each cortisol-treated group to have increased cortisol concentrations when compared with the control ewes (Table 1), but there were no differences among

groups. ACTH levels were not significantly altered in response to cortisol treatment, although there was a trend for cortisol-treated ewes to have lower ACTH concentrations than the control group (Table 1).

Maternal sodium, potassium, and PCV values were not different between the groups at day 5 or day 10 (data not shown). Maternal plasma protein concentrations were significantly elevated in the cortisol-treated ewes on days 5 (8.2 ± 0.2 vs 7.6 ± 0.2 g 100/ml) and 10 (7.9 ± 0.1 vs 7.4 ± 0.1 g 100/ml) when compared with the control ewes.

Maternal arterial pressures and heart rates were not different between the four groups (data not shown).

Fetal physiology

The average plasma cortisol concentrations (days 5 and 10) were significantly elevated in the fetuses whose mothers were infused with cortisol compared with control (3.4 ± 0.6 vs 1.5 ± 0.6 ng/ml). There was a trend for each cortisol-treated group to have increased cortisol concentrations when compared with the control fetuses when the four groups were compared individually (Table 1). ACTH levels were not significantly altered in response to cortisol treatment (Table 1).

There were no significant differences among the groups in the blood gas values or PCV (Table 2), nor were there effects on fetal electrolytes (data not shown). There were also no effects of treatment on fetal heart rate and blood pressures (Table 3).

Fetal heart measurements

Heart weight was significantly greater in the cortisol group when compared with the control group and cortisol+MRa group, but not the cortisol+GRa group (Fig. 2). LV and RV free wall thicknesses were significantly greater in the fetal hearts of the cortisol-treated group when compared with the control group. LV and RV free wall, as well as septum thicknesses were greater in the fetal hearts of the cortisol group when compared with the cortisol+MRa group (Fig. 2). LV free wall and septum thicknesses were not different in the cortisol group when compared with the cortisol+GRa group (Fig. 2). However, RV free wall thickness was greater in the cortisol group when compared with the cortisol+GRa group.

Collagen staining

Fetal heart sections were stained with picosirius red in order to measure the amount of interstitial collagen deposition (Fig. 3). The percentage of collagen staining in the LV, RV, septum, and whole heart was not significantly altered among the groups (Table 4, Fig. 3).

Table 1 Fetal and maternal cortisol concentrations (average of days 5 and 10) and ACTH concentration on day 10. Data are expressed as mean $\pm$ S.E.M

	Maternal cortisol (ng/ml)	Fetal cortisol (ng/ml)	Maternal ACTH (pg/ml)	Fetal ACTH (pg/ml)
Control	5.9 $\pm$ 1.4	1.5 $\pm$ 0.6	37 $\pm$ 8	36 $\pm$ 8
Cortisol	9.6 $\pm$ 2.3	2.7 $\pm$ 0.5	20 $\pm$ 1	27 $\pm$ 5
Cortisol+MRa	8.7 $\pm$ 0.4	3.6 $\pm$ 1.0	31 $\pm$ 11	38 $\pm$ 6
Cortisol+GRa	8.3 $\pm$ 1.5	3.9 $\pm$ 1.9	21 $\pm$ 1	57 $\pm$ 33

Discussion

We conclude that blockade of corticosteroid receptors in the fetal heart prevents the enlargement of the heart observed when maternal cortisol concentrations are chronically increased. We found that blockade of the MRs blocked the increase in heart weight, as well as wall thickness. Blockade of GRs significantly reduced RV enlargement, and produced smaller insignificant effects on thickness of the LV free wall and septum and on heart weight. Administration of neither MR nor GR blocker into the pericardium resulted in increases in fetal ACTH or fetal blood pressure, suggesting that the infusions of antagonist did not produce systemic effects. The results indicate that small increases in cortisol increase fetal heart size via an intracardiac action at the MRs and, to a lesser extent, GRs within the fetal heart. We also conclude that the increase in fetal heart weight in response to elevated cortisol occurs without an increase in collagen deposition (Fig. 3).

Role of MR and GR in the heart

Our laboratory has previously shown that both MR and GR are abundantly expressed in the heart in late gestation ovine fetus (Reini *et al.* 2006), suggesting a role for these receptors in fetal heart development *in vivo*. Other investigators have found that aldosterone directly stimulates myocyte surface area (Okoshi *et al.* 2004) and remodeling of myocyte membrane (Kliche *et al.* 2006) in cultures of neonatal myocytes, an effect presumed to be mediated by MR in the myocytes. Cortisol also increases expression of atrial natriuretic peptide in cultured neonatal myocytes, and both cortisol and aldosterone potentiate the effect of phenylephrine on hypertrophy in these cultures (Lister *et al.* 2006), also indicating an intracardiac action at MR in these cultures.

One of the major factors influencing the ability of cortisol to activate MR and/or GR is the local activity of the 11 β -HSD enzymes, 11 β -HSD1 and 11 β -HSD2. 11 β -HSD1 primarily converts cortisone into cortisol, while 11 β -HSD2 converts cortisol into cortisone, which is inactive at MR and GR (Krozowski *et al.* 1999). We had previously shown that mRNA expression of 11 β -HSD2 mRNA is relatively low compared with 11 β -HSD1 within the fetal heart (Reini *et al.* 2006). Using immunohistochemistry, we also found that although MR, GR, and 11 β -HSD1 are abundantly expressed in both myocytes and blood vessels within the fetal heart, 11 β -HSD2 seemed to be localized in blood vessels more abundantly than in myocytes. This suggested that cortisol has access to both MR and GR within the fetal heart, and that when plasma cortisol levels are increased, as in the present study, action of cortisol at MR and GR in the heart would also increase. Our present study demonstrates that the effect of cortisol is blocked by antagonists of the MR and/or GR, suggesting a role of intracardiac corticosteroid receptors. This is consistent with the ability of cortisol to alter myocyte growth in cultured myocytes. We also hypothesized that blockade of MR would have a greater effect in inhibiting the effect of cortisol than would blockade of GR, because MR has been shown to have greater affinity for cortisol than GR (Reul & DeKloet 1985, Richards *et al.* 2003). Indeed, this is what we observed in the present study: in the cortisol group, there was a 14% increase in heart weight relative to body weight when compared with the control group; this enlargement was completely blocked when MR antagonist was administered to the heart, whereas there was only 44% blockade of the increase in weight after administration of the GR antagonist. Similarly, in the cortisol group, LV, RV, and septum thicknesses were $\sim$ 20% thicker than control fetuses and the MR antagonist produced a 95, 149, and 114% reduction of this increase in thickness of the LV, RV, and

Table 2 Fetal blood gas and packed cell volume. Data are expressed as mean $\pm$ S.E.M

	Fetal PO ₂ (mmHg)	Fetal PCO ₂ (mmHg)	Fetal pH	Fetal packed cell volume (fractional volume of red blood cells)
Control	21.7 $\pm$ 1.0	56 $\pm$ 1	7.34 $\pm$ 0.01	0.313 $\pm$ 0.007
Cortisol	21.5 $\pm$ 1.0	53 $\pm$ 2	7.35 $\pm$ 0.01	0.326 $\pm$ 0.013
Cortisol+MRa	20.7 $\pm$ 1.1	54 $\pm$ 2	7.30 $\pm$ 0.03	0.348 $\pm$ 0.007
Cortisol+GRa	21.9 $\pm$ 0.4	55 $\pm$ 1	7.32 $\pm$ 0.02	0.325 $\pm$ 0.009

Table 3 Fetal arterial pressure and fetal heart rate on day 10. Data are expressed as mean ± S.E.M

	Fetal arterial pressure (mmHg)	Fetal heart rate (beats per min)
Control	47.5 ± 2.7	170 ± 6
Cortisol	46.9 ± 2.7	168 ± 11
Cortisol + MRa	43.0 ± 0.9	172 ± 4
Cortisol + GRa	46.0 ± 1.0	164 ± 7

septum respectively, whereas the GR antagonist group produced 63, 110, and 65% reductions of thickness. Overall, GR blockade was approximately half as effective as MR blockade in inhibiting the increase in heart weight or wall thickness.

The relative differences in effectiveness of MR and GR blockade are consistent with the expected relative binding of fetal cortisol at these receptors. The MRs are higher affinity receptors with greater occupancy at low cortisol concentrations (Reul & DeKloet 1985), and therefore a greater effect would be expected after blockade of these receptors. Based on the expected free fraction of cortisol in the fetuses, we calculate that the free cortisol concentrations would be ~0.8 nM in the control fetuses and 1.9 nM in the fetuses of the cortisol-infused ewes. Based on previous studies of cortisol binding at ovine MR and GR (Richards *et al.* 2003), we would predict that these free concentrations would result in ~65% occupancy of MR and 35% occupancy of GR in the control fetuses, and 85% occupancy of MR and 60%

Table 4 Collagen content determined by picrosirius red staining (fraction of total area) in left ventricle (LV), right ventricle (RV), and septum. Data are expressed as mean ± S.E.M

	LV	RV	Septum
Control	0.037 ± 0.005	0.039 ± 0.004	0.037 ± 0.006
Cortisol	0.050 ± 0.009	0.057 ± 0.009	0.052 ± 0.010
Cortisol + MRa	0.049 ± 0.011	0.037 ± 0.007	0.043 ± 0.008
Cortisol + GRa	0.060 ± 0.012	0.058 ± 0.013	0.059 ± 0.013

occupancy of GR in the cortisol-infused fetuses. Thus, these levels would be expected to exert more effects via MR than via GR activation if both act at GRE to induce genes responsible for cardiac growth.

Role of MR in hypertrophy in the adult heart

In adult rats, the MR is thought to induce cardiac hypertrophy and fibrosis occurring in response to ischemia; systemic administration of MR blockers have been shown to reduce markers of inflammation and fibrosis in hearts of adult rats (Brilla *et al.* 1993, Sun *et al.* 2002, Fraccarollo *et al.* 2005). It has been established that in adult humans with severe heart failure, there is a reduction in the severity of cardiac hypertrophy and an increase in survival rate after treatment with the MR antagonists, eplerenone or spironolactone (Pitt *et al.* 1999, 2001). The effect of MR blockers on survival rate appears to be the result of a decrease in cardiac fibrosis (Fraccarollo *et al.* 2004); increases in interstitial collagen content are a feature of adult cardiac hypertrophy (Pearlman *et al.* 1981), particularly in the case of hypertension or myocardial infarction (Young *et al.* 2007). The mechanism for the *in vivo* effect of MR in contributing to inflammation and subsequent fibrosis is not clear. It has been suggested that the effect is through a non-genomic action, and that the effect in ischemic tissue is predominately on vascular cells expressing MR, rather than on fibroblasts or on myocytes (Mihailidou & Funder 2005, Young *et al.* 2007). It is generally assumed that the protective effect of the MR antagonist results from blocking the action of aldosterone at MR. It has been suggested, however, that many heart failure patients without elevated plasma aldosterone levels still benefit from MR blockade, indicating that aldosterone may not be the only relevant MR ligand (Young *et al.* 2007). Since plasma cortisol concentrations are much higher than aldosterone, and since there is not a significant amount of 11β-HSD2 expressed within the heart, it is reasonable to propose that cortisol may be playing a role in the fibrosis that is observed in heart failure patients.

In our studies the effects of cortisol do not appear to involve increase in fibrosis, as there was no increase in collagen content with maternal cortisol infusion, nor were there any effects of either MR or GR blockade. This suggests that the mechanism of the enlargement of the fetal heart in the present study may be fundamentally different from what is observed in

Figure 2 Mean fetal heart measurements from control (open bars), cortisol (solid bars), cortisol + MRa (diagonal striped bars), and cortisol + GRa (cross-hatched bars) groups taken at the time of killing: heart to body mass ratio (upper left), left ventricular (LV) free wall thickness (lower right), septal wall thickness (upper right), and right ventricular (RV) free wall thickness (lower left). Data are expressed as mean ± S.E.M. Horizontal lines between groups indicate that differences are statistically significant, *P* < 0.05.

Figure 3 Representative pictures showing picrosirius red staining of collagen in left ventricular wall of fetal hearts from (A) control, (B) cortisol, (C) cortisol + MRa, and (D) cortisol + GRa groups; all photos at 40 $\times$ power. Bar indicates 200 μ m. Arrows indicate the dark staining corresponding to positive Sirius red staining.

adult rat models or human pathology, in which ischemia is a contributing component.

Mechanisms of enlargement of the fetal heart

Due to the unique ability of the fetal heart to grow through both hyperplasia and hypertrophy, either mechanism could account for the cortisol-induced increases in fetal heart weight and wall thickness in our model. In early gestation, cardiac growth is mostly a result of the production of new myocytes originating through cell division and proliferation (Smolich 1995). After $\sim$ 115 days of gestation in sheep, however, cardiac growth results primarily from increases in myocyte size (Jonker *et al.* 2007). Myocytes lose their ability to divide and proliferate shortly after birth in an event in which there is nuclear division without subsequent cell division (Oparil *et al.* 1984). In fetal sheep, the number of terminally differentiated or binucleate myocytes increases from $\sim$ 115 days of gestation through term, and heart growth during this period is due to increases in both myocyte size and myocyte proliferation (Jonker *et al.* 2007). Theoretically, cortisol could be stimulating growth through either hypertrophy or hyperplasia, or possibly even both.

Rudolph *et al.* (1999) showed that cortisol (1.2 μ g/min) infusion for 72–80 h directly into the left coronary artery of the ovine fetus (124–131 days) decreased LV DNA content. This was interpreted as cortisol-induced inhibition of myocyte proliferation in preparation for life after birth. The fetal blood pressures from that study were not reported. In a study by Lumbers *et al.* (2005) high dose infusion of cortisol (72.1 mg/d for $\sim$ 60 h) increased LV myocyte size and increase cardiac angiotensinogen mRNA, suggesting an induction of hypertrophy. However, there was also a significant increase in blood pressure in these fetuses, suggesting that the cardiac hypertrophy may have resulted from elevated blood pressure.

Conversely, maternal dexamethasone administration (48 μ g/d from E17) increased relative heart weight and

increased myocyte proliferation in the fetal and newborn rat heart (Torres *et al.* 1997). In agreement with this, Giraud *et al.* (2006) showed that subpressor doses of cortisol (0.5 μ g/kg per min for 7 days) infused directly into the circumflex coronary artery of the fetus led to an increase in Ki-67 stained myocytes in both the left and RVs; as Ki-67 is expressed only in cells in the proliferative phase, this suggested that cortisol stimulated proliferation in these hearts. Hearts infused with cortisol weighed more than control hearts in this study, but there were no changes in myocyte size or percent binucleation. Interestingly, there were also no differences in aortic, right atrial, systolic, and diastolic pressures between the groups. These studies suggest that elevated fetal cortisol concentrations directly stimulate cardiomyocyte proliferation in the late-term fetus.

The present study does not provide direct evidence for cardiomyocyte proliferation as a means of cardiac enlargement in response to cortisol. It is important to note that in this study a subpressor dose of cortisol was used, as in the study by Giraud *et al.* (2006). We did not observe an increase in arterial pressure in response to the moderately elevated cortisol levels indicating that the fetal hearts in this study were not subjected to chronically increased systolic load, a possible trigger to myocyte hypertrophy seen in some other studies. Although in the present study blood pressure was only measured at 10 days of cortisol infusion, in our previous study (Jensen *et al.* 2005) fetal arterial blood pressure was not elevated at either 5 or 10 days of maternal cortisol infusion. The doses of cortisol administered in our study resulted in relatively small increases in fetal cortisol, well below those that have been shown to increase fetal blood pressure in other studies (Wood *et al.* 1987, Tangalakakis *et al.* 1992, Unno *et al.* 1999). Furthermore, in this study we observed no evidence within the fetal heart in support of interstitial collagen deposition, a symptom of cardiac hypertrophy in response to hypertension within the adult human heart (Diez 2007).

Conclusions

The data suggest that the enlargement of the fetal heart in response to a modest and chronic rise in maternal cortisol levels is mediated by MRs and, to a lesser extent, GRs within the fetal heart. Intrapericardial infusion of an MR antagonist completely prevented the increase in wall thickness and heart weight. The GR blockade was less effective, although GR blockade prevented the increase in RV free wall thickness, and tended to attenuate the increase in LV free wall thickness and whole heart weight. The cortisol-induced enlargement is not accompanied by an increase in interstitial collagen deposition within the fetal heart. This indicates the possibility of a different mechanism for the enlargement observed in the fetal heart than that observed in adult cardiac hypertrophy and fibrosis.

Declaration of Interest

The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

Funding

This work was supported by NIH grant DK62080 to M K-W and American Heart Association Florida–Puerto Rico Affiliate pre-doctoral fellowship grant 0615236B to S A R.

Acknowledgements

We acknowledge the technical assistance of Jarret McCartney for assistance in the postoperative care of the sheep and for assistance with necropsies. We also acknowledge the assistance of Kavi Patel and Sasha Gidwani for analysis of the collagen content.

References

- Bell ME, Wood CE & Keller-Wood M 1991 Influence of reproductive state on pituitary–adrenal activity in the ewe. *Domestic Animal Endocrinology* **8** 245–254.
- Brilla CG, Matsubara LS & Weber KT 1993 Antifibrotic effects of spironolactone in preventing myocardial fibrosis in systemic arterial hypertension. *American Journal of Cardiology* **71** 12A–16A.
- Diez J 2007 Mechanisms of cardiac fibrosis in hypertension. *Journal of Clinical Hypertension* **9** 546–550.
- Fejes-Toth G, Pearce D & Naray-Fejes-Toth A 1998 Subcellular localization of mineralocorticoid receptors in living cells: effects of receptor agonists and antagonists. *PNAS* **95** 2973–2978.
- Fraccarollo D, Galuppo P, Hildemann S, Christ M, Ertl G & Bauersachs J 2003 Additive improvement of left ventricular remodeling and neurohormonal activation by aldosterone receptor blockade with eplerenone and ACE inhibition in rats with myocardial infarction. *Journal of the American College of Cardiology* **42** 1666–1673.
- Fraccarollo D, Galuppo P & Bauersachs J 2004 Mineralocorticoid receptor antagonism and cardiac remodeling in ischemic heart failure. *Current Medicinal Chemistry – Cardiovascular and Hematological Agents* **2** 287–294.
- Fraccarollo D, Galuppo P, Schmidt I, Ertl G & Bauersachs J 2005 Additive amelioration of left ventricular remodeling and molecular alterations by combined aldosterone and angiotensin receptor blockade after myocardial infarction. *Cardiovascular Research* **67** 97–105.
- Giraud GD, Louey S, Jonker S, Schultz J & Thornburg KL 2006 Cortisol stimulates cell cycle activity in the cardiomyocyte of the sheep fetus. *Endocrinology* **147** 3643–3649.
- Gomez-Sanchez CE, de Rodriguez AF, Romero DG, Estess J, Warden MP, Gomez-Sanchez MT & Gomez-Sanchez EP 2006 Development of a panel of monoclonal antibodies against the mineralocorticoid receptor. *Endocrinology* **147** 1343–1348.
- Jensen E, Wood C & Keller-Wood M 2002a The normal increase in adrenal secretion during pregnancy contributes to maternal volume expansion and fetal homeostasis. *Journal of the Society for Gynecologic Investigation* **9** 362–371.
- Jensen EC, Gallaher BW, Breier BH & Harding JE 2002b The effect of a chronic maternal cortisol infusion on the late-gestation fetal sheep. *Journal of Endocrinology* **174** 27–36.
- Jensen E, Wood CE & Keller-Wood M 2005 Chronic alterations in ovine maternal corticosteroid levels influence uterine blood flow and placental and fetal growth. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **288** R54–R61.
- Jewell CM, Webster JC, Burnstein KL, Sar M, Bodwell JE & Cidlowski JA 1995 Immunocytochemical analysis of hormone mediated nuclear translocation of wild type and mutant glucocorticoid receptors. *Journal of Steroid Biochemistry and Molecular Biology* **55** 135–146.
- Jonker SS, Zhang L, Louey S, Giraud GD, Thornburg KL & Faber JJ 2007 Myocyte enlargement, differentiation, and proliferation kinetics in the fetal sheep heart. *Journal of Applied Physiology* **102** 1130–1142.
- Kliche K, Kuhn M, Hillebrand U, Ludwig Y, Stock C & Oberleithner H 2006 Direct aldosterone action on mouse cardiomyocytes detected with atomic force microscopy. *Cellular Physiology and Biochemistry* **18** 265–274.
- Krozowski Z, Li KX, Koyama K, Smith RE, Obeyesekere VR, Stein-Oakley A, Sasano H, Coulter C, Cole T & Sheppard KE 1999 The type I and type II 11 β -hydroxysteroid dehydrogenase enzymes. *Journal of Steroid Biochemistry and Molecular Biology* **69** 391–401.
- Lister K, Autelitano DJ, Jenkins A, Hannan RD & Sheppard KE 2006 Cross talk between corticosteroids and alpha-adrenergic signalling augments cardiomyocyte hypertrophy: a possible role for SGK1. *Cardiovascular Research* **70** 555–565.
- Lombes M, Binart N, Delahaye F, Baulieu EE & Rafestin-Oblin ME 1994 Differential intracellular localization of human mineralocorticosteroid receptor on binding of agonists and antagonists. *Biochemical Journal* **302** 191–197.
- Lumbers ER, Boyce AC, Joulianos G, Kumarasamy V, Barner E, Segar JL & Burrell JH 2005 Effects of cortisol on cardiac myocytes and on expression of cardiac genes in fetal sheep. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **288** R567–R574.
- Mihailidou AS & Funder JW 2005 Nongenomic effects of mineralocorticoid receptor activation in the cardiovascular system. *Steroids* **70** 347–351.
- Okoshi MP, Yan X, Okoshi K, Nakayama M, Schuldt AJ, O'Connell TD, Simpson PC & Lorell BH 2004 Aldosterone directly stimulates cardiac myocyte hypertrophy. *Journal of Cardiac Failure* **10** 511–518.
- Oparil S, Bishop SP & Clubb FJ Jr 1984 Myocardial cell hypertrophy or hyperplasia. *Hypertension* **6** III38–III43.
- Pearlman ES, Weber KT & Janicki JS 1981 Quantitative histology of the hypertrophied human heart. *Federation Proceedings* **40** 2042–2047.
- Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, Palensky J & Wittes J 1999 The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *New England Journal of Medicine* **341** 709–717.
- Pitt B, Williams G, Remme W, Martinez F, Lopez-Sendon J, Zannad F, Neaton J, Roniker B, Hurley S, Burns D *et al.* 2001 The EPHEUS trial: eplerenone in patients with heart failure due to systolic dysfunction complicating acute myocardial infarction. Eplerenone Post-AMI Heart Failure Efficacy and Survival Study. *Cardiovascular Drugs and Therapy* **15** 79–87.
- Reini SA, Wood CE, Jensen E & Keller-Wood M 2006 Increased maternal cortisol in late gestation ewes decreases fetal cardiac expression of 11 β -HSD2 mRNA and the ratio of AT1 to AT2 receptor mRNA. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **21** R1708–R1716.
- Reul JMHM & DeKloet ER 1985 Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* **117** 2505–2511.
- Richards EM, Hua Y & Keller-Wood M 2003 Pharmacology and physiology of ovine corticosteroid receptors. *Neuroendocrinology* **77** 2–14.
- Rudolph AM, Roman C & Gournay V 1999 Perinatal myocardial DNA and protein changes in the lamb: effect of cortisol in the fetus. *Pediatric Research* **46** 141–146.
- Scheuer DA, Bechtold AG, Shank SS & Akana SF 2004 Glucocorticoids act in the dorsal hindbrain to increase arterial pressure. *American Journal of Physiology. Heart and Circulatory Physiology* **286** H458–H467.
- Smolich JJ 1995 Ultrastructural and functional features of the developing mammalian heart: a brief overview. *Reproduction, Fertility, and Development* **7** 451–461.
- Sun Y, Zhang J, Lu L, Chen SS, Quinn MT & Weber KT 2002 Aldosterone-induced inflammation in the rat heart: role of oxidative stress. *American Journal of Pathology* **161** 1773–1781.
- Tangalakis K, Lumbers ER, Moritz KM, Towstoles MK & Wintour EM 1992 Effect of cortisol on blood pressure and vascular reactivity in the ovine fetus. *Experimental Physiology* **77** 709–717.

- Torres A, Belser WW III, Umeda PK & Tucker D 1997 Indicators of delayed maturation of rat heart treated prenatally with dexamethasone. *Pediatric Research* **42** 139–144.
- Unno N, Wong CH, Jenkins SL, Wentworth RA, Ding XY, Li C, Robertson SS, Smotherman WP & Nathanielsz PW 1999 Blood pressure and heart rate in the ovine fetus: ontogenic changes and effects of fetal adrenalectomy. *American Journal of Physiology* **276** H248–H256.
- Winer BJ 1971 *Statistical Principles in Experimental Design*., New York: McGraw-Hill.
- Wood CE 2002 The ovine fetal endocrine reflex responses to haemorrhage are not mediated by cardiac nerves. *Journal of Physiology* **541** 613–622.
- Wood CE & Rudolph AM 1983 Negative feedback regulation of adrenocorticotropin secretion by cortisol. *Endocrinology* **112** 1930–1936.
- Wood CE, Cheung CY & Brace RA 1987 Fetal heart rate, arterial pressure, and blood volume responses to cortisol infusion. *American Journal of Physiology* **253** R904–R909.
- Xiao HD, Fuchs S, Campbell DJ, Lewis W, Dudley SC Jr, Kasi VS, Hoit BD, Keshelava G, Zhao H, Capecchi MR *et al.* 2004 Mice with cardiac-restricted angiotensin-converting enzyme (ACE) have atrial enlargement, cardiac arrhythmia, and sudden death. *American Journal of Pathology* **165** 1019–1032.
- Young MJ, Lam EY & Rickard AJ 2007 Mineralocorticoid receptor activation and cardiac fibrosis. *Clinical Science* **112** 467–475.
- Zar JH 1984 *Biostatistical Analysis*., Englewood Cliffs, NJ: Prentice-Hall.

Received in final form 22 April 2008

Accepted 20 May 2008

Made available online as an Accepted Preprint

21 May 2008