The role of calcium influx pathways in phospholipase D activation in bovine adrenal glomerulosa cells

Haixia Qin², Patricia Kent², Carlos M Isales^{2,4}, Peter M Parker², Mariya V Wilson²
and Wendy B Bollag^{1,2,3,4}

¹Charlie Norwood VA Medical Center, One Freedom Way, Augusta, Georgia 30904, USA

²Institute of Molecular Medicine and Genetics, ³Departments of Medicine and Cell Biology and Anatomy and ⁴Department of Orthopaedic Surgery, Medical College of Georgia, 1120 15th Street, Augusta, Georgia 30912, USA

(Correspondence should be addressed to W B Bollag; Email: wbollag@mcg.edu)

Abstract

The steroid hormone aldosterone maintains sodium homeostasis and is therefore important in the control of blood volume and pressure. Angiotensin II (AngII) and elevated extracellular potassium concentrations ($[K^+]_e$), the prime physiologic regulators of aldosterone secretion from adrenal glomerulosa cells, activate phospholipase D (PLD) in these cells. The role of Ca^{2+} in the activation by these agents is unknown, although nitrendipine, a voltage-dependent Ca^{2+} channel antagonist, does not inhibit AngII-elicited PLD activation, despite the fact that this compound blocked elevated $[K^+]_e$ -stimulated PLD activity. PLD activation triggered by AngII was also unaffected by the T-type calcium channel inhibitor nickel. Nevertheless, Ca^{2+} influx was required for AngII-induced PLD activation in both primary cultures of bovine adrenal glomerulosa cells and a glomerulosa cell model, the NCI H295R adrenocortical

carcinoma cell line. The involvement of store-operated Ca^{2+} (SOC) influx and Ca^{2+} release-activated Ca^{2+} (CRAC) influx pathways in PLD activation was investigated using thapsigargin, an endoplasmic reticulum Ca^{2+} pump inhibitor that empties the store to induce SOC influx, and the SOC inhibitor YM-58483 (BTP2), as well as a CRAC inhibitor, tyrphostin A9. In bovine glomerulosa cells, tyrphostin A9 inhibited AngII-induced PLD activation without affecting elevated $[K^+]_e$ -stimulated enzyme activity. On the other hand, differences were observed between the bovine adrenal glomerulosa and H295R cells in the involvement of Ca^{2+} influx pathways in PLD activation, with the involvement of the SOC pathway suggested in the H295R cells. In summary, our results indicate that Ca^{2+} entry only through certain Ca^{2+} influx pathways is linked to PLD activation.

Journal of Endocrinology (2009) **202**, 77–86

Introduction

The steroid hormone aldosterone controls sodium homeostasis and thus blood volume and pressure. Aldosterone is secreted from glomerulosa cells in the adrenal cortex in response to two major regulators, angiotensin II (AngII) and elevated extracellular potassium concentration ($[K^+]_e$). AngII induces aldosterone secretion by binding to the AngII type 1 receptor, which is coupled to phosphatidylinositol 4,5-bisphosphate hydrolysis resulting in the production of inositol 1,4,5-trisphosphate and diacylglycerol (DAG; reviewed in Barrett *et al.* (1989), Foster (2004), Spät & Hunyady (2004) and Rainey *et al.* (2008)). These second messengers, in turn, increase cytosolic calcium (Ca^{2+}) levels (to activate Ca^{2+} /calmodulin-dependent protein kinases) and stimulate protein kinase C (PKC) respectively (reviewed in Rainey *et al.* (2008)). Elevated $[K^+]_e$, on the other hand, triggers glomerulosa cell depolarization, which opens voltage-dependent Ca^{2+} channels and also increases cytosolic

Ca^{2+} levels (reviewed in Rainey *et al.* (2008)). We previously showed that not only AngII but also elevated $[K^+]_e$ activates phospholipase D (PLD; Betancourt-Calle *et al.* 2001), the activity of which can produce lipid messengers including phosphatidic acid and DAG (reviewed in Bollag & Zheng (2005)). Furthermore, inhibition of PLD-mediated signal generation inhibits both AngII- and elevated $[K^+]_e$ -elicited aldosterone secretion (Betancourt-Calle *et al.* 2001, Bollag *et al.* 2002, Zheng & Bollag 2003), suggesting the importance of this signaling enzyme to steroid hormone production. Therefore, it is important to understand the mechanisms that regulate PLD activity in response to AngII and elevated $[K^+]_e$.

The ability of both AngII and elevated $[K^+]_e$ to activate PLD suggests a possible involvement of Ca^{2+} in the process, since an increase in the cytosolic Ca^{2+} concentration is common to these two agonists (reviewed in Rainey *et al.* (2008)). However, the role of Ca^{2+} in PLD activation in other systems is unclear, and, in fact, Exton (1999) indicated that 'direct control of the enzyme by physiological changes

in cytosolic Ca^{2+} seems unlikely'. Investigators have shown an ability of cytosolic Ca^{2+} concentration to modulate PLD activity, with chelation of intracellular Ca^{2+} inhibiting PLD activation in response to some agonists and Ca^{2+} ionophores increasing enzyme activity (reviewed in [Exton \(1999\)](#)). The mechanism by which Ca^{2+} regulates PLD activity is not known but may involve calmodulin and/or the Ca^{2+} -sensitive PKC isoenzymes (reviewed in [Exton \(1999\)](#)). On the other hand, in primary bovine glomerulosa cells, the inability of the Ca^{2+} ionophores ionomycin or A23187 to activate PLD ([Bollag et al. 2002](#)) suggests that changes in cytosolic Ca^{2+} levels alone are not sufficient to stimulate PLD activity. Furthermore, the fact that elevated $[\text{K}^+]_e$, which functions through voltage-dependent Ca^{2+} channels, activates PLD in bovine adrenal glomerulosa cells ([Betancourt-Calle et al. 2001](#)) suggests a possible involvement of Ca^{2+} influx in regulating PLD activity. However, the lack of inhibition of AngII-induced PLD activation by nitrendipine ([Bollag et al. 2002](#)), a voltage-dependent Ca^{2+} channel antagonist, at a dose that inhibits aldosterone secretion ([Kojima et al. 1985b](#)), supports the idea that only certain Ca^{2+} pathways may be important in stimulating PLD in response to particular agonists.

The objectives of this study were to determine the role of Ca^{2+} influx pathways in activating PLD in response to AngII as well as to elevated $[\text{K}^+]_e$ in primary cultures of bovine adrenal glomerulosa cells. In addition, we sought to compare the effects of modulating Ca^{2+} influx on PLD activation and acute aldosterone secretion in this bovine system with the responses observed in the NCI H295R human adrenocortical carcinoma cell line.

Materials and Methods

Materials

The following were obtained from Sigma: phorbol 12-myristate 13-acetate (PMA), AngII, and 22(R)-hydroxycholesterol. UltroSer G was acquired from BioSeptra (Cergy St Christophe, France) under a permit from the US Department of Agriculture. [^3H]Oleic acid was purchased from Dupont NEN (Boston, MA, USA). Silica gel 60 thin-layer chromatography plates with concentrating zones were obtained from EM Science through VWR (West Chester, PA, USA) and phosphatidylethanol (PEt) and phosphatidic acid standards from Avanti Polar Lipids (Alabaster, AL, USA). ITS+ premix (12.5 mg insulin, 12.5 mg transferrin, 12.5 µg selenous acid, 10.7 µg linoleic acid, and 2.5 mg BSA) was purchased from BD Biosciences (San Jose, CA, USA). Thapsigargin, YM-58483 (BTP2) and tyrphostin A9 were obtained from Calbiochem (San Diego, CA, USA).

Primary culture of bovine adrenal glomerulosa cells

Bovine adrenal glomerulosa cells were prepared and cultured as described in [Bollag et al. \(2007\)](#). Briefly, the glomerulosa layer was dissected from adrenal glands of near-term fetal calves

obtained from a local meat-packing plant. Glomerulosa cells were released from tissue slices by enzymatic and mechanical means and collected by centrifugation. Cells were plated in Primaria plates (BD Falcon, Franklin Lakes, NJ, USA) overnight in a medium consisting of DMEM/Ham's F12 (1:1 vol:vol) containing 10% horse serum, 2% fetal bovine serum, 100 µM ascorbate, 1.2 µM α -tocopherol, 0.05 µM Na_2SeO_3 , 50 µM butylated hydroxyanisole, 5 µM metyrapone, 100 U/ml penicillin, 100 µg/ml streptomycin, and 0.25 µg/ml fungizone. The cells were then incubated for 20–24 h in serum-free DMEM/Ham's F12 (containing 0.2% BSA, 100 µM ascorbate, 1.2 µM α -tocopherol, 0.05 µM Na_2SeO_3 , 50 µM butylated hydroxyanisole, 5 µM metyrapone, and antibiotics) prior to experimentation.

Culture of NCI H295R human adrenocortical carcinoma cells

NCI H295R cells were cultured as described in [Zheng & Bollag \(2003\)](#). Briefly, cells were grown in DMEM/Ham's F12 (1:1 vol:vol) containing 1% ITS+ (12.5 mg insulin, 12.5 mg transferrin, 12.5 µg selenous acid, 10.7 µg linoleic acid, and 2.5 mg BSA from BD Biosciences), 2% UltroSer G, 100 U/ml penicillin, 100 µg/ml streptomycin, and 0.25 µg/ml fungizone, to ~70–75% confluence. The cells were then incubated for 20–24 h in serum-free medium (consisting of DMEM/Ham's F12 containing 0.01% BSA and antibiotics) prior to experimentation.

Measurement of aldosterone secretion

Cells were incubated for the indicated times with the appropriate agents in bicarbonate-buffered Krebs' Ringer containing 2.5 mM sodium acetate (KRB^+). Aldosterone release into the medium was then assayed using a RIA kit from Diagnostic Products Corporation (Los Angeles, CA, USA) as described in [Bollag et al. \(2007\)](#). For stimulation with an elevated $[\text{K}^+]_e$, KCl was substituted iso-osmotically for NaCl, as in [Betancourt-Calle et al. \(2001\)](#).

PLD activity assay

Bovine glomerulosa or H295R cells were prelabeled with 2.5–5 µCi/ml [^3H]oleate in the appropriate serum-free medium for 20–24 h. The cells were then preincubated for 30 min in KRB^+ prior to stimulation in the presence of 0.5% ethanol with and without AngII in KRB^+ containing or lacking 1.2 mM CaCl_2 or in the presence and absence of 50 µM NiCl_2 2 µM thapsigargin or 10 µM tyrphostin A9 (or DMSO as the vehicle control) for 5 (H295R cells) or 30 min (bovine adrenal glomerulosa cells). Again, stimulation with elevated $[\text{K}^+]_e$ was performed by iso-osmotic substitution. Phospholipids were extracted with chloroform/methanol and separated by thin-layer chromatography, and radiolabeled PEt was quantified by liquid scintillation spectrometry as described in [Bollag \(1998\)](#).

Statistical analysis

Experiments were performed a minimum of three times in duplicate or triplicate. Data were analyzed by ANOVA with a Student–Newmann–Keuls *post hoc* test using the computer program Instat (Graphpad, San Diego, CA, USA), except where indicated in the figure legends. A *P* value ≤ 0.05 was considered significant.

Results

We have previously shown that elevated [K⁺]_e stimulates PLD activity (Betancourt-Calle *et al.* 2001). Because of the importance of PLD to sustained aldosterone secretion (Bollag *et al.* 2002, Zheng & Bollag 2003), we wanted to characterize the mechanism by which elevated [K⁺]_e and AngII activate PLD. We previously showed that the voltage-dependent Ca²⁺ channel antagonist, nitrendipine, does not alter AngII-induced PLD activation (Bollag *et al.* 2002), even at a concentration (1 μM) reported to inhibit both transient, or T-type, and long-lasting, or L-type, Ca²⁺ channels (Barrett *et al.* 1995). The role of voltage-dependent Ca²⁺ channels in elevated [K⁺]_e-induced PLD activation was determined by monitoring the effect of nitrendipine on the elevated [K⁺]_e-stimulated increase in the levels of [³H]PEt, a specific marker of PLD activity (Thompson *et al.* 1991), in [³H]oleate-prelabeled primary bovine adrenal glomerulosa cells (Bollag *et al.* 1990, 1992). Nitrendipine (1 μM) completely abolished elevated [K⁺]_e-induced changes in PLD activity (Fig. 1),

Figure 1 Inhibition of Ca²⁺ influx with nitrendipine completely inhibited elevated [K⁺]_e-elicited PLD activation. [³H]Oleate-prelabeled cells were incubated for 30 min with KRB⁺ containing 3.5 mM KCl (3.5 mM K⁺–KRB⁺, control), 1 μM nitrendipine (NTR) in 3.5 mM K⁺–KRB⁺, 15 mM K⁺–KRB⁺, or 1 μM NTR in 15 mM K⁺–KRB⁺ in the presence of 0.5% ethanol. (Note that NaCl in the medium was substituted for by KCl to maintain osmolarity in the 15 mM K⁺–KRB⁺ as in Betancourt-Calle *et al.* (2001)). Reactions were terminated by the addition of 0.2% SDS containing 5 mM EDTA, and [³H]phosphatidylethanol was extracted, separated by thin-layer chromatography and quantified as in Jung *et al.* (1998). Values are expressed as -fold over the control and represent the means (± s.e.m.) of five separate experiments; **P* < 0.05 versus the control value.

although it had no effect on either basal (Fig. 1; Bollag *et al.* 2002) or AngII-stimulated PLD activity (Bollag *et al.* 2002), indicating that Ca²⁺ influx through voltage-dependent Ca²⁺ channels is absolutely required for the effect of elevated [K⁺]_e on PLD activity.

To further define the involvement of T-type versus L-type channels in the PLD response to AngII and elevated [K⁺]_e, we used the T-type channel inhibitor, nickel, at a dose reported to inhibit T-type Ca²⁺ influx, specifically through α-1H T-type Ca²⁺ channels (Schrier *et al.* 2001). We observed no effect of 50 μM nickel on PLD activity either under basal conditions or upon stimulation with AngII (Fig. 2A). On the other hand, elevated [K⁺]_e-stimulated PLD activity, and nickel returned this increase to a level not significantly different from the control value, causing an approximate 43% inhibition (Fig. 2B). Together, these results are consistent with the idea that voltage-dependent Ca²⁺ channels are not involved in AngII-induced PLD activation but are critical for the activity stimulated by elevated [K⁺]_e.

To determine whether Ca²⁺ influx was involved in AngII's activation of PLD, AngII-stimulated PLD activity was then examined in the presence or absence of extracellular Ca²⁺. Interestingly, we observed a reduction in radiolabeled PEt production in the absence of Ca²⁺ in primary bovine adrenal glomerulosa cells (Fig. 3). This result suggests that whereas elevated [K⁺]_e requires Ca²⁺ influx through voltage-dependent Ca²⁺ channels, AngII-elicited PLD activation is independent of Ca²⁺ influx through voltage-dependent Ca²⁺ channels but nevertheless requires Ca²⁺ entry, presumably through store-operated Ca²⁺ (SOC) or Ca²⁺ release-activated Ca²⁺ (CRAC) channels.

To test the role of SOC influx in AngII- and elevated extracellular [K⁺]_e-induced PLD activation, we determined the effects on PLD activity of thapsigargin, a pharmacologic agent that inhibits the sarcoplasmic/endoplasmic reticulum Ca²⁺ pump to release Ca²⁺ from these stores and activate SOC influx (reviewed in Spät & Hunyady (2004)). We selected a dose of thapsigargin (2 μM) that has previously been shown to increase SOC Ca²⁺ influx in primary cultures of bovine adrenal glomerulosa cells (Burnay *et al.* 1998, Aptel *et al.* 1999). We found that thapsigargin had little effect on AngII-induced PLD activation (Fig. 4A) in primary bovine adrenal glomerulosa cells, whereas this compound stimulated elevated [K⁺]_e-elicited PLD activity (Fig. 4B). Interestingly, these results were mirrored by the effects of thapsigargin on aldosterone secretion in that the compound had little effect on AngII-induced steroidogenesis but augmented elevated [K⁺]_e-elicited aldosterone secretion (Fig. 5A), consistent with a previous report in bovine adrenal glomerulosa cells (Burnay *et al.* 1994). In addition, thapsigargin enhanced the aldosterone secretory response to the phorbol ester, PMA, which mimics DAG action in primary bovine adrenal glomerulosa cells (Fig. 5B). This latter result is consistent with the proposed requirement for two signals mediating sustained aldosterone secretion: a DAG second messenger and a Ca²⁺ influx signaling pathway (reviewed in Rasmussen *et al.* (1995)).

Figure 2 Inhibition of α -1H T-type Ca^{2+} channels with nickel inhibited elevated $[\text{K}^+]_e$, but not AngII-, elicited PLD activation. (A) $[\text{H}]$ Oleate-prelabeled cells were incubated for 30 min with KRB^+ containing 0.5% ethanol and with and without 10 nM AngII in the presence or absence of 50 μM nickel (Ni) for 30 min. Reactions were terminated by the addition of 0.2% SDS containing 5 mM EDTA, and $[\text{H}]$ phosphatidylethanol was extracted, separated by thin-layer chromatography and quantified as in Jung *et al.* (1998). Values are expressed as -fold over the control and represent the means ($\pm$ S.E.M.) of four separate experiments; * $P < 0.05$ versus the control value. (B) $[\text{H}]$ Oleate-prelabeled cells were incubated for 30 min with 3.5 mM K^+ - KRB^+ or 15 mM K^+ - KRB^+ containing 0.5% ethanol in the absence or presence of 50 μM nickel (Ni) for 30 min, and PLD activity monitored as above. Values are expressed as -fold over the control and represent the means ($\pm$ S.E.M.) of four separate experiments; ** $P < 0.01$ versus the control value.

To further examine the role of SOC in AngII- and elevated $[\text{K}^+]_e$ -induced PLD activation, we treated cells with these agonists in the presence and absence of the SOC inhibitor, YM-58483, also known as BTP2. We found that BTP2 had

Figure 3 AngII-induced PLD activation was modulated by Ca^{2+} in primary bovine adrenal glomerulosa cells. Bovine adrenal glomerulosa cells were prelabeled for 20–24 h with 2.5–5 $\mu\text{Ci}/\text{ml}$ $[\text{H}]$ oleate in serum-free medium. Cells were then stimulated with or without 10 nM AngII in the presence of 0.5% ethanol in medium containing 1.2 mM Ca^{2+} or no added Ca^{2+} for 30 min. Lipids were extracted and separated by thin-layer chromatography. PLD was measured as the radioactivity in phosphatidylethanol (PE) relative to the control, and values are expressed as the means ($\pm$ S.E.M.) of four separate experiments; *** $P < 0.001$ versus the control value; †† $P < 0.01$ versus the value in the presence of Ca^{2+} .

no effect on PLD activation elicited in response to AngII (Fig. 6A). However, this agent actually seemed to increase elevated $[\text{K}^+]_e$ -induced PLD activation, converting a non-significant response to elevated $[\text{K}^+]_e$ alone to a significant increase in the presence of BTP2 (Fig. 6B). This result suggests that whereas SOC can enhance the activation of PLD, i.e. in conjunction with elevated $[\text{K}^+]_e$, it is not necessary for PLD activation in response to either agonist.

The archetypical SOC pathway is CRAC, mediated by Stim and Orai proteins (reviewed in Potier & Trebak (2008)). To examine the possible role of CRAC channels in AngII-induced PLD activation, we determined the effect of the CRAC channel inhibitor, tyrphostin A9 (Denys *et al.* 2004) on this signaling event. As shown in Fig. 7A, we found that tyrphostin A9 inhibited AngII-elicited PLD activation in primary bovine adrenal glomerulosa cells, without affecting basal PLD activity. By contrast, tyrphostin A9 had no effect on elevated $[\text{K}^+]_e$ -induced PLD activation in the primary glomerulosa cells (Fig. 7B). In experiments to determine the ability of tyrphostin A9 to modulate the aldosterone secretory response, we found that tyrphostin A9 completely blocked AngII- and elevated $[\text{K}^+]_e$ -induced aldosterone secretion (data not shown). However, we also observed that tyrphostin A9 significantly inhibited the ability of 22(R)-hydroxycholesterol to trigger steroidogenesis (Fig. 7C). Because 22(R)-hydroxycholesterol can directly enter mitochondria to access the rate-limiting enzyme of aldosterone

Figure 4 Thapsigargin had little or no effect on AngII-induced PLD activation, but enhanced the effect of elevated $[K^+]_e$ on PLD activity in bovine adrenal glomerulosa cells. (A) Bovine adrenal glomerulosa cells were prelabeled for 20–24 h with 2.5–5 $\mu\text{Ci}/\text{ml}$ [^3H]oleate in serum-free medium. Cells were then stimulated with or without 10 nM AngII in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.1–0.2% DMSO) in the presence of 0.5% ethanol for 30 min. Lipids were extracted and separated by thin-layer chromatography. Values represent the radioactivity in phosphatidylethanol relative to the control and are expressed as the means $\pm$ s.e.m. of nine separate experiments; * $P < 0.05$ versus the control value. (B) [^3H]Oleate-prelabeled bovine adrenal glomerulosa cells were incubated with 3.5 mM K^+ –KRB⁺ or 15 mM K^+ –KRB⁺ in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.1–0.2% DMSO) in the presence of 0.5% ethanol for 30 min. Lipids were extracted and separated by thin-layer chromatography. Values represent the radioactivity in phosphatidylethanol relative to the control and are expressed as the means $\pm$ s.e.m. of five separate experiments; ** $P < 0.01$ versus the control value; † $P < 0.05$ versus K^+ alone.

synthesis, thereby bypassing signaling mechanisms, inhibitory effects on secretion induced by this compound indicate that tyrphostin A9 either inhibits aldosterone biosynthetic enzymes or affects cell health. However, the fact that tyrphostin A9 did not alter basal or elevated $[K^+]_e$ -elicited PLD activity indicates that the inhibitor is not simply cytotoxic and suggests instead that the compound inhibits an enzyme in the aldosterone synthetic pathway.

We have previously shown that AngII activates PLD in H295R cells (Zheng & Bollag 2003), as in primary cultures of bovine adrenal glomerulosa cells (Bollag *et al.* 1990), and that this activity mediates, at least in part, the ability of AngII to induce steroidogenesis (Zheng & Bollag 2003), as in the bovine cells (Bollag *et al.* 2002). However, in contrast to results in primary bovine adrenal glomerulosa cells, in which AngII-induced PLD activation is sustained (Jung *et al.* 1998), in H295R cells AngII elicits only transient PLD activation (Zheng & Bollag 2003). We wished to examine whether H295R cells exhibited other differences in the mechanism of PLD activation in response to aldosterone agonists. Initially, we examined the Ca^{2+} dependence of AngII-induced PLD activation in the H295R cells. Similar to the primary bovine adrenal glomerulosa cells (Fig. 2), a lack of extracellular Ca^{2+} also inhibited AngII-elicited PLD activation in H295R cells (Fig. 8). Importantly, however, we were unable to detect PLD activation in response to elevated $[K^+]_e$ in the H295R cells under a variety of conditions (data not shown), including at different time points and following pretreatment with aldosterone, reported to increase T-type channel expression (Lesouhaitier *et al.* 2001).

On the other hand and in contrast to the primary bovine adrenal glomerulosa cells (Fig. 4), thapsigargin enhanced AngII-induced PLD activation in the H295R cells (Fig. 9A). The effects on AngII-induced PLD activation in the H295R cells were not accompanied by comparable changes in aldosterone secretion, as thapsigargin had little or no effect on AngII-induced aldosterone production (Fig. 9B), although this result was similar to that observed in the primary bovine adrenal glomerulosa cells (Fig. 4). However, in contrast to the bovine cells, thapsigargin did not alter elevated $[K^+]_e$ -elicited steroidogenesis in the H295R cells (Fig. 9B). Taken together, these results suggest that while similarities exist between the mechanisms of AngII-induced PLD activation in primary bovine adrenal glomerulosa cells versus the human adrenocortical carcinoma H295R cells, there are also distinct differences. However, which AngII-elicited PLD activation scenario is most similar to the situation occurring *in vivo*, i.e. in human patients, is at present unknown.

Discussion

We have previously shown that PLD-mediated signal generation is required for maximal AngII-induced secretion in both primary cultures of bovine adrenal glomerulosa cells and the NCI H295R human adrenocortical carcinoma cell

Figure 5 Thapsigargin had little or no effect alone or on AngII-stimulated aldosterone secretion but enhanced steroidogenesis in response to elevated $[\text{K}^+]_e$ and PMA in primary cultures of bovine adrenal glomerulosa cells. (A) Bovine adrenal glomerulosa cells were incubated for 1 h with a bicarbonate-buffered Krebs Ringer solution with 2.5 mM sodium acetate (KRB^+) containing no additions (Con), 15 mM KCl (K^+ ; iso-osmotic substitution with Na^+) or 10 nM AngII in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.1–0.2% DMSO). Supernatants were collected and assayed for aldosterone secretion by RIA. Values represent the means $\pm$ S.E.M. of four to five separate experiments; * $P < 0.05$, ** $P < 0.01$ versus control by ANOVA and a Student–Newmann–Keuls *post hoc* test. (B) Bovine adrenal glomerulosa cells were incubated for 1 h with a bicarbonate-buffered Krebs Ringer solution with 2.5 mM sodium acetate (KRB^+) containing no additions (Con) or 100 nM PMA in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.1–0.2% DMSO). Supernatants were collected and assayed for aldosterone secretion by RIA. Values represent the means $\pm$ S.E.M. of four to five separate experiments, expressed as the percentage of the maximal aldosterone secretory response; * $P < 0.05$, ** $P < 0.01$ versus control by ANOVA and a Student–Newmann–Keuls *post hoc* test.

Figure 6 Inhibition of store-operated Ca^{2+} influx with BTP2 (YM-58483) had no effect on AngII-induced, and increased elevated $[\text{K}^+]_e$ -elicited PLD activation. (A) [^3H]Oleate-prelabeled cells were incubated for 30 min with KRB^+ containing 0.5% ethanol and vehicle (0.1% DMSO) or 10 nM AngII in the presence or absence of 1 μM BTP2 for 30 min. Reactions were terminated by the addition of 0.2% SDS containing 5 mM EDTA, and [^3H]phosphatidyl-ethanol was extracted, separated by thin-layer chromatography and quantified as in Jung *et al.* (1998). Values are expressed as -fold over the control and represent the means ($\pm$ S.E.M.) of four separate experiments; * $P < 0.05$ versus the control value. (B) [^3H]Oleate-prelabeled cells were incubated for 30 min with 3.5 mM K^+ – KRB^+ or 15 mM K^+ – KRB^+ containing 0.5% ethanol with vehicle (0.1% DMSO) or 1 μM BTP2 for 30 min, and PLD activity monitored as above. Values are expressed as -fold over the control and represent the means ($\pm$ S.E.M.) of four separate experiments; * $P < 0.05$ versus the control value.

line (Bollag *et al.* 2002, Zheng & Bollag 2003; as well as elevated $[\text{K}^+]_e$ -induced steroidogenesis in bovine cells (Betancourt-Calle *et al.* 2001)). Because of the importance of PLD signal generation to sustained aldosterone secretion, we sought to determine the mechanism by which AngII and elevated $[\text{K}^+]_e$ modulate PLD's activity. Our results indicate that Ca^{2+} influx through voltage-dependent Ca^{2+} channels is absolutely required for elevated $[\text{K}^+]_e$ -induced PLD activation. This finding is not particularly surprising considering that elevated $[\text{K}^+]_e$ is thought to exert its effects on adrenal glomerulosa cells by depolarizing the cells, thereby opening voltage-dependent calcium channels (reviewed in Rainey *et al.* (2008)). Perhaps a bit more unexpected is the observation that AngII-induced PLD activity also requires Ca^{2+} influx, although this Ca^{2+} influx is not mediated by voltage-dependent Ca^{2+} channels, as neither nifedipine nor nickel has an inhibitory effect on AngII-induced PLD activation (Bollag *et al.* 2002). However, removal of extracellular Ca^{2+} inhibits AngII-elicited PLD activation in both primary bovine adrenal glomerulosa cells (Fig. 3) and H295R cells (Fig. 8), suggesting that Ca^{2+} influx plays some

role in PLD activation. Nevertheless, the importance of Ca²⁺ in regulating PLD activity was somewhat unanticipated since we previously found no effect of Ca²⁺ ionophores on PLD activity in primary glomerulosa cells nor inhibition of

AngII-induced enzyme activation by a voltage-dependent Ca²⁺ channel antagonist (Bollag *et al.* 2002). On the other hand, we have previously shown that in both cell models, PKC appears to mediate, at least in part, PLD activation in response to AngII (Bollag *et al.* 2002, Zheng & Bollag 2003), as has been reported for other cell systems (reviewed in Frohman & Morris (1996) and Exton (1998)). Thus, exposure to PMA is sufficient to induce PLD activation in both bovine and H295R cells (Bollag *et al.* 2002, Zheng & Bollag 2003), and PKC inhibitors reduce AngII-stimulated PLD activity (Zheng & Bollag 2003). Thus, the requirement for Ca²⁺ influx may be related to activation of PKC, presumably a classical, Ca²⁺-dependent PKC isoenzyme, as suggested by the results of Kojima *et al.* (1994) who demonstrated that Ca²⁺ can, indeed, regulate PKC activity.

Although voltage-dependent Ca²⁺ channels do not appear to be involved in AngII-stimulated PLD activation (Bollag *et al.* 2002), AngII also increases Ca²⁺ influx through other pathways, such as SOC and CRAC (reviewed in Spät & Hunyady (2004)). Stimulation of SOC by thapsigargin has no effect alone or together with AngII on PLD activation in primary bovine adrenal glomerulosa cells (Fig. 3). We also observed no effect of thapsigargin on AngII-elicited aldosterone secretion, in contrast to a previous report in rat adrenal glomerulosa cells (Hajnoczky *et al.* 1991). The reason for this disparity is unclear but may be related to the species and/or the use of submaximal concentrations of AngII and thapsigargin in the rat cells (Hajnoczky *et al.* 1991). Nevertheless, the 2 µM thapsigargin dose used herein was sufficient to enhance the elevated [K⁺]_e-stimulated PLD

Figure 7 Inhibition of Ca²⁺ release-activated Ca²⁺ influx with tyrphostin A9 inhibited AngII-, but not elevated [K⁺]_e-, elicited PLD activation. (A) [³H]Oleate-prelabeled cells were incubated for 30 min with KRB⁺ containing 0.5% ethanol and vehicle (0.1% DMSO, control) or 10 nM AngII (+0.1% DMSO) in the presence or absence of 10 µM tyrphostin A9 (TyrA9) for 30 min. Reactions were terminated by the addition of 0.2% SDS containing 5 mM EDTA, and [³H]phosphatidylethanol was extracted, separated by thin-layer chromatography, and quantified as in Jung *et al.* (1998). Values are expressed as -fold over the control and represent the means (±S.E.M.) of five separate experiments; ***P*<0.01, ****P*<0.001 versus the control value, ††*P*<0.01 versus AngII alone. (B) [³H]Oleate-prelabeled cells were incubated for 30 min with 3.5 mM K⁺-KRB⁺ or 15 mM K⁺-KRB⁺ containing 0.5% ethanol and vehicle (0.1% DMSO, control) or 10 µM tyrphostin A9 (TyrA9) for 30 min, and PLD activity monitored as above. Values are expressed as -fold over the control and represent the means (±S.E.M.) of four separate experiments; **P*<0.05, ***P*<0.01 versus the control value. (C) Bovine adrenal glomerulosa cells were incubated for 1 h with KRB⁺ containing no additions (Con) or 10 µM 22(R)-hydroxycholesterol in the presence or absence of 10 µM tyrphostin A9 (TyrA9) or vehicle (0.1–0.2% DMSO). Supernatants were collected and assayed for aldosterone secretion by RIA. Values represent the means ±S.E.M., expressed as the ng aldosterone/ml per 60 min, of three separate experiments; **P*<0.03 versus 22(R)-hydroxycholesterol alone, as determined using an unpaired Student's *t*-test. Note that 22(R)-hydroxycholesterol induced a significant, ~11 000-fold increase over the control aldosterone secretory value.

Figure 8 AngII-induced PLD activation was modulated by Ca^{2+} in NCI H295R adrenocortical carcinoma cells. H295R cells were prelabeled for 20–24 h with 2.5–5 $\mu\text{Ci}/\text{ml}$ [^3H]oleate in serum-free medium. Cells were then stimulated with or without 100 nM AngII in KRB^+ containing 1.2 mM Ca^{2+} or no added Ca^{2+} in the presence of 0.5% ethanol for 5 min. Lipids were extracted and separated by thin-layer chromatography. Values represent the radioactivity in phosphatidylethanol (PEt) relative to the control and are expressed as the means $\pm$ S.E.M. of four separate experiments; ** $P < 0.01$, *** $P < 0.01$ versus the control value, $^{\dagger}P < 0.05$ versus the value in the presence of Ca^{2+} .

activity and aldosterone secretory rate (Fig. 4), indicating its efficacy in promoting emptying of the intracellular Ca^{2+} store, as has been previously demonstrated in these same cells (Burnay *et al.* 1998, Aptel *et al.* 1999). On the other hand, however, thapsigargin was able to enhance AngII's activation of PLD in H295R cells (Fig. 9). It is not clear why there is the disparity between the two cell types, although we have previously reported that AngII induces a transient PLD activation in H295R cells (Zheng & Bollag 2003) but a sustained activity in bovine adrenal glomerulosa cells (Jung *et al.* 1998). Similarly, Bird *et al.* (1993) have shown that the production of inositol bis- and trisphosphates in response to AngII in the H295R cells is also transient, whereas in bovine adrenal glomerulosa cells AngII elicits a sustained phosphoinositide hydrolysis response (Kojima *et al.* 1984). These data suggest that phosphoinositide-specific phospholipase C is only transiently activated in H295R cells by AngII, such that over time the endoplasmic reticulum Ca^{2+} pool might be expected to refill upon extinction of the transient signal, thereby terminating SOC. Thus, thapsigargin, by prolonging SOC influx could enhance PLD activation (Fig. 9A) in response to AngII in the H295R cells, without affecting these parameters in primary bovine adrenal glomerulosa cells (Fig. 5), which exhibit sustained AngII-elicited phospholipase C activation and therefore maintained (i.e. already maximal) SOC. However, despite its enhancement of AngII-induced PLD activation in H295R cells, thapsigargin did not alter

Figure 9 Thapsigargin enhanced the effect of AngII on PLD activation but had no effect on AngII- (or elevated $[\text{K}^+]_e$ -) stimulated aldosterone secretion in H295R cells. (A) H295R cells were prelabeled for 20–24 h with 2.5–5 $\mu\text{Ci}/\text{ml}$ [^3H]oleate in serum-free medium. Cells were then stimulated with or without 10 or 100 nM AngII in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.2% DMSO) in the presence of 0.5% ethanol. Lipids were extracted and separated by thin-layer chromatography. Values represent the radioactivity in phosphatidylethanol relative to the control and are expressed as the means $\pm$ S.E.M. of three separate experiments; * $P < 0.05$ versus the control value. (B) H295R cells were incubated for 5 h with medium containing no additions (Con), 15 mM KCl (K^+), or 100 nM AngII in the presence or absence of 2 μM thapsigargin (Thaps) or vehicle (0.1–0.2% DMSO) as indicated. Supernatants were collected and assayed for aldosterone secretion by RIA. Values represent the means $\pm$ S.E.M. of four to five separate experiments; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus the control value.

AngII- (or elevated [K⁺]_e-) stimulated aldosterone secretion in these cells (Fig. 9B). Furthermore, the SOC inhibitor BTP2 (YM-5848) had no effect on AngII-induced PLD activation and appeared to increase that elicited by elevated [K⁺]_e (Fig. 6), suggesting that Ca²⁺ influx mediated by SOC is not necessary for PLD activation in response to either agonist (Fig. 6).

SOC Ca²⁺ influx occurs through a pathway exhibiting heterogeneous characteristics, with BTP2 reported to inhibit transient receptor potential (TRP) channels, with a half-maximal inhibitory concentration of 0.1–0.3 μM (He *et al.* 2005). On the other hand, CRAC represents an archetypical SOC (reviewed in Potier & Trebak (2008)). Based on our results, CRAC may be a key AngII-induced Ca²⁺ influx pathway involved in regulating PLD activation in primary bovine adrenal glomerulosa cells. Thus, a CRAC inhibitor, tyrphostin A9 (Denys *et al.* 2004), inhibited AngII-stimulated PLD activity but had no effect on the basal or elevated [K⁺]_e-induced value (Fig. 7A and B). Tyrphostin A9 has also been reported to inhibit platelet-derived growth factor receptor tyrosine kinase activity (Marhaba *et al.* 1996); thus, the possibility remains that the effects of this compound on AngII-induced PLD activation may be related to inhibition of tyrosine kinase. However, in H295R cells, we were previously unable to detect an effect of the general tyrosine kinase inhibitor, genistein (or the Janus kinase inhibitor AG490 or Src kinase inhibitor PP2), on AngII-stimulated PLD activity (Zheng & Bollag 2003), suggesting the likelihood that tyrphostin A9's inhibition of CRAC mediates the inhibition of PLD activation in response to AngII. Unfortunately, it was not possible to determine whether CRAC was important in AngII-elicited steroidogenesis. Although tyrphostin A9 completely inhibited AngII-induced aldosterone secretion, as well as that elicited in response to elevated [K⁺]_e (data not shown), it also inhibited secretion in response to 22(R)-hydroxycholesterol (Fig. 7C), indicating either effects on cell health or on one or more aldosterone biosynthetic enzymes. However, the inability of tyrphostin A9 to alter basal or elevated [K⁺]_e-stimulated PLD activity (Fig. 6) suggests that cytotoxicity was not responsible for the inhibition. An ability of inhibitors to affect aldosterone biosynthetic enzymes is not without precedent, as Rainey *et al.* have demonstrated that genistein inhibits 3β-hydroxysteroid dehydrogenase (Sirianni *et al.* 2001).

As expected, in primary bovine adrenal glomerulosa cells, elevated [K⁺]_e-induced PLD activation was dependent on voltage-dependent Ca²⁺ channels (Fig. 1). Interestingly, SOC also appeared to stimulate elevated [K⁺]_e-elicited PLD activation and aldosterone secretion in bovine adrenal glomerulosa cells, as shown by the ability of thapsigargin to stimulate these processes (Figs 4 and 5). There is some controversy as to whether elevated [K⁺]_e triggers phosphoinositide hydrolysis (Hunyady *et al.* 1982, Kojima *et al.* 1985a, Underwood *et al.* 1988, Ganguly *et al.* 1990, Hajnoczky *et al.* 1992, Betancourt-Calle *et al.* 2001). Our data, however, would suggest that if it does so, elevated [K⁺]_e is not

particularly effective at inducing SOC, since thapsigargin can augment the effects of this ion on aldosterone secretion in primary bovine adrenal glomerulosa (Fig. 5). On the other hand, we were unable to detect elevated [K⁺]_e-induced PLD activation in H295R cells, under a variety of conditions (data not shown), suggesting differences in signal transduction, particularly with regards to the Ca²⁺ signal, in the bovine adrenal glomerulosa cells versus the H295R cells.

Together, our results suggest that: 1) PLD activation in response to elevated [K⁺]_e, but not AngII, is dependent on Ca²⁺ influx via voltage-dependent Ca²⁺ channels, but 2) AngII-induced PLD activation nevertheless requires Ca²⁺, likely entering the cell through CRAC channels. Furthermore, our results highlight the importance of the specific Ca²⁺ influx pathway in coupling of the aldosterone agonists, AngII and elevated [K⁺]_e to PLD activation in adrenal glomerulosa cells.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by a National Institutes of Health Award #HL70046 and American Heart Association Grant-in-Aid Award #0350166N.

References

- Aptel HB, Burnay MM, Rossier MF & Capponi AM 1999 The role of tyrosine kinases in capacitative calcium influx-mediated aldosterone production in bovine adrenal zona glomerulosa cells. *Journal of Endocrinology* **163** 131–138.
- Barrett PQ, Bollag WB, Isales CM, McCarthy RT & Rasmussen H 1989 Role of calcium in angiotensin II-mediated aldosterone secretion. *Endocrine Reviews* **10** 1–22.
- Barrett PQ, Ertel EA, Smith MM, Nee JJ & Cohen CJ 1995 Voltage-gated calcium currents have two opposing effects on the secretion of aldosterone. *American Journal of Physiology* **268** C985–C992.
- Betancourt-Calle S, Jung EM, White S, Calle RA, Rasmussen H & Bollag WB 2001 Elevated K⁺ induces MARCKS phosphorylation and phospholipase D activation in glomerulosa cells. *Molecular and Cellular Endocrinology* **184** 65–76.
- Bird IM, Hanley NA, Word RA, Mathis JM, McCarthy JL, Mason JI & Rainey WE 1993 Human NCI-H295 adrenocortical carcinoma cells: a model for angiotensin-II-responsive aldosterone secretion. *Endocrinology* **133** 1555–1561.
- Bollag WB 1998 Measurement of phospholipase D activity. *Methods in Molecular Biology* **105** 151–160.
- Bollag WB & Zheng X 2005 The role of phospholipase D in keratinocyte biology. In *Trends in Protein Research*. Hauppauge, New York: NovaScience Publishers.
- Bollag WB, Barrett PQ, Isales CM, Liscovitch M & Rasmussen H 1990 A potential role for phospholipase-D in the angiotensin-II-induced stimulation of aldosterone secretion from bovine adrenal glomerulosa cells. *Endocrinology* **127** 1436–1443.

- Bollag WB, Barrett PQ, Isales CM, Liscovitch M & Rasmussen H 1992 Signal transduction mechanisms involved in carbachol-induced aldosterone secretion from bovine adrenal glomerulosa cells. *Molecular and Cellular Endocrinology* **86** 93–101.
- Bollag WB, Jung EM & Calle RA 2002 Mechanism of angiotensin II-induced phospholipase D activation in adrenal glomerulosa cells. *Molecular and Cellular Endocrinology* **192** 7–16.
- Bollag WB, Kent P, White S, Malinova M, Isales CM & Calle RA 2007 Characterization and phospholipase D mediation of the angiotensin II priming response in adrenal glomerulosa cells. *Endocrinology* **148** 585–593.
- Burnay MM, Python CP, Vallotton MB, Capponi AM & Rossier MF 1994 Role of the capacitative calcium influx in the activation of steroidogenesis by angiotensin-II in adrenal glomerulosa cells. *Endocrinology* **135** 751–758.
- Burnay MM, Vallotton MB, Capponi AM & Rossier MF 1998 Angiotensin II potentiates adrenocorticotrophic hormone-induced cAMP formation in bovine adrenal glomerulosa cells through a capacitative calcium influx. *Biochemical Journal* **330** 21–27.
- Denys A, Aires V, Hichami A & Khan NA 2004 Thapsigargin-stimulated MAP kinase phosphorylation via CRAC channels and PLD activation: inhibitory action of docosahexaenoic acid. *FEBS Letters* **564** 177–182.
- Exton JH 1998 Phospholipase D. *Biochimica et Biophysica Acta* **1436** 105–115.
- Exton JH 1999 Regulation of phospholipase D. *Biochimica et Biophysica Acta* **1439** 121–133.
- Foster RH 2004 Reciprocal influences between the signalling pathways regulating proliferation and steroidogenesis in adrenal glomerulosa cells. *Journal of Molecular Endocrinology* **32** 893–902.
- Frohman MA & Morris AJ 1996 Phospholipid signalling: rho is only ARF the story. *Current Biology* **6** 945–947.
- Ganguly A, Chiou S & Davis JS 1990 Intracellular mediators of potassium-induced aldosterone secretion. *Life Sciences* **46** 173–180.
- Hajnoczky G, Varnai P, Hollo Z, Christensen SB, Balla T, Enyedi P & Spät A 1991 Thapsigargin-induced increase in cytoplasmic Ca^{2+} concentration and aldosterone production in rat adrenal glomerulosa cells: interaction with potassium and angiotensin-II. *Endocrinology* **128** 2639–2644.
- Hajnoczky G, Varnai P, Buday L, Farago A & Spät A 1992 The role of protein kinase-C in control of aldosterone production by rat adrenal glomerulosa cells: activation of protein kinase-C by stimulation with potassium. *Endocrinology* **130** 2230–2236.
- He LP, Hewavitharana T, Soboloff J, Spassova MA & Gill DL 2005 A functional link between store-operated and TRPC channels revealed by the 3,4-bis(trifluoromethyl)pyrazole derivative, BTP2. *Journal of Biological Chemistry* **280** 10997–11006.
- Hunyady L, Balla T, Nagy K & Spät A 1982 Control of phosphatidylinositol turnover in adrenal glomerulosa cells. *Biochimica et Biophysica Acta* **713** 352–357.
- Jung EM, Betancourt-Calle S, Mann-Blakeney R, Foushee T, Isales CM & Bollag WB 1998 Sustained phospholipase D activation in response to angiotensin II but not carbachol in bovine adrenal glomerulosa cells. *Biochemical Journal* **330** 445–451.
- Kojima I, Kojima K, Kreutter D & Rasmussen H 1984 The temporal integration of the aldosterone secretory response to angiotensin occurs via two intracellular pathways. *Journal of Biological Chemistry* **259** 14448–14457.
- Kojima I, Kojima K & Rasmussen H 1985a Intracellular calcium and adenosine 3',5'-cyclic monophosphate as mediators of potassium-induced aldosterone secretion. *Biochemical Journal* **228** 69–76.
- Kojima I, Kojima K & Rasmussen H 1985b Role of calcium fluxes in the sustained phase of angiotensin II-mediated aldosterone secretion from adrenal glomerulosa cells. *Journal of Biological Chemistry* **260** 9177–9184.
- Kojima I, Kawamura N & Shibata H 1994 Rate of calcium entry determines the rapid changes in protein kinase C activity in angiotensin II-stimulated adrenal glomerulosa cells. *Biochemical Journal* **297** 523–528.
- Lesouhaitier O, Chiappe A & Rossier MF 2001 Aldosterone increases T-type calcium currents in human adrenocarcinoma (H295R) cells by inducing channel expression. *Endocrinology* **142** 4320–4330.
- Marhaba R, Mary F, Pelassy C, Stancescu AT, Aussel C & Breitmayer JP 1996 Tyrphostin A9 inhibits calcium release-dependent phosphorylations and calcium entry via calcium release-activated channel in Jurkat T cells. *Journal of Immunology* **157** 1468–1473.
- Potier M & Trebak M 2008 New developments in the signaling mechanisms of the store-operated calcium entry pathway. *Pflügers Archiv* **457** 405–415.
- Rainey WE, Bollag WB & Isales CM 2008 Aldosterone regulation. In *Textbook of Nephro-Endocrinology*, Eds AJ Singh & GH Williams. Atlanta, GA: Elsevier, Inc.
- Rasmussen H, Isales CM, Calle R, Throckmorton D, Anderson M, Gasalla-Herraz J & McCarthy R 1995 Diacylglycerol production, Ca^{2+} influx, and protein kinase C activation in sustained cellular responses. *Endocrine Reviews* **16** 649–681.
- Schrier AD, Wang H, Talley EM, Perez-Reyes E & Barrett PQ 2001 Alpha1H T-type Ca^{2+} channel is the predominant subtype expressed in bovine and rat zona glomerulosa. *American Journal of Physiology. Cell Physiology* **280** C265–C272.
- Sirianni R, Sirianni R, Carr BR, Pezzi V & Rainey WE 2001 A role for src tyrosine kinase in regulating adrenal aldosterone secretion. *Journal of Molecular Endocrinology* **26** 207–215.
- Spät A & Hunyady L 2004 Control of aldosterone secretion: a model for convergence in cellular signaling pathways. *Physiological Reviews* **84** 489–539.
- Thompson NT, Bonser RW & Garland LG 1991 Receptor-coupled phospholipase D and its inhibition. *Trends in Pharmacological Sciences* **12** 404–408.
- Underwood RH, Greeley R, Glennon ET, Menachery AI, Braley LM & Williams GH 1988 Mass determination of polyphosphoinositides and inositol triphosphate in rat adrenal glomerulosa cells with a microspectrophotometric method. *Endocrinology* **123** 211–219.
- Zheng X & Bollag WB 2003 AngII induces transient phospholipase D activity in the H295R glomerulosa cell model. *Molecular and Cellular Endocrinology* **206** 113–122.

Received in final form 22 March 2009

Accepted 15 April 2009

Made available online as an Accepted Preprint
16 April 2009