Regulation and action of fibroblast growth factor 17 in bovine follicles

M F Machado, V M Portela¹, C A Price¹, I B Costa, P Ripamonte², R L Amorim³ and J Buratini Jr²

Departamento de Reprodução Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista, Botucatu, São Paulo 18618-000, Brazil

¹Centre de recherche en reproduction animale, Faculté de médecine vétérinaire, Université de Montréal, St-Hyacinthe, Quebec J2S 7C6, Canada

²Departamento de Fisiologia, Instituto de Biociências, Universidade Estadual Paulista, Rubião Júnior, Botucatu, São Paulo 18618-000, Brazil

³Departamento de Clínica Veterinária, Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista, Botucatu, São Paulo, Brazil

(Correspondence should be addressed to J Buratini Jr; Email: buratini@ibb.unesp.br)

Abstract

Fibroblast growth factor 17 (FGF17) is a member of the FGF8 subfamily that appears to be relevant to folliculogenesis and oogenesis, as the prototype member FGF8 is an oocyte-derived protein that signals to cumulus cells. FGF8 has structural and receptor-binding similarities to FGF17, whose expression in the ovary has not been reported. In this study, we demonstrate localization of FGF17 protein to the oocyte of preantral follicles, and to the oocyte and granulosa cells of antral follicles. Real-time PCR demonstrated the presence of mRNA in oocytes and, to a lesser extent, in

granulosa and theca cells. *FGF17* mRNA abundance was low in granulosa and theca cells from healthy follicles and increased significantly in atretic follicles. Addition of FSH or IGF-I to granulosa cells *in vitro* decreased *FGF17* mRNA abundance, and treatment with FGF17 inhibited estradiol and progesterone secretion from granulosa cells in relation to control cultures without these additives. We conclude that FGF17 is a potential mediator of granulosa cell differentiation.

Journal of Endocrinology (2009) **202**, 347–353

Introduction

Follicle development is under the control of gonadotropins, steroids, and a variety of locally produced peptides including fibroblast growth factors (FGFs; Webb *et al.* 2003). Several FGFs and their receptors (FGFR) have been detected in ovarian follicles, suggesting roles in the regulation of folliculogenesis (van Wezel *et al.* 1995, Berisha *et al.* 2004, Buratini *et al.* 2005a,b, 2007). The 22 known FGFs have been grouped into seven subfamilies with distinct receptor-binding properties. The FGFR proteins are encoded by five different genes, three of which, *FGFR1*, 2, and 3, undergo alternative splicing to produce two functional variants (B and C; Ornitz *et al.* 1996, Itoh & Ornitz 2004). Paracrine roles within the follicle have been explored for FGF2 and for the FGF7 subfamily, containing also FGF10. FGF2, which is predominantly expressed by theca cells (Berisha *et al.* 2000), stimulates proliferation and inhibits steroidogenesis in both theca and granulosa cells (Lavranos *et al.* 1994, Vernon & Spicer 1994, Spicer & Stewart 1996, Nilsson *et al.* 2001). *FGF7* and *FGF10* are expressed in the theca cell layer but not in granulosa cells, and the receptor FGFR2B is predominantly expressed in granulosa cells (Parrott & Skinner 1998, Berisha *et al.* 2004, Buratini *et al.* 2007). Both FGFs inhibit estradiol (E₂) secretion from cultured granulosa cells (Parrott & Skinner 1998, Buratini *et al.* 2007).

Another subfamily that may be of interest for potential paracrine signaling is the FGF8 subfamily. In adult rodents, *Fgf8* gene expression is largely confined to the oocyte (Valve *et al.* 1997), and was reported in oocytes as well as somatic follicle cells in cattle (Buratini *et al.* 2005b). FGF8 activates FGFR3C, which was found in theca and granulosa cells in cattle, and FGFR4, which was localized only to theca cells (Ornitz *et al.* 1996, Buratini *et al.* 2005b). Although the roles of the FGF8 subfamily in the control of the ovarian activity are still poorly understood, it has been recently shown that oocyte-derived FGF8 cooperates with bone morphogenetic protein 15 (BMP15) to promote glycolytic activity in cumulus cells in mice (Sugiura *et al.* 2007).

The FGF8 subfamily also contains FGF17 (Itoh & Ornitz 2004), which also preferentially activates FGFR3C and FGFR4 (Ford-Perriss *et al.* 2001, Zhang *et al.* 2006). *FGF17* gene expression was first detected in the embryonic brain, and is most associated with neurogenesis (O'Leary *et al.* 2007) and skeletal development (Krejci *et al.* 2007). Very little is known about the pattern of expression of *FGF17* in the reproductive system. mRNA encoding *FGF17* was detected in human prostatic epithelial cells (Polnaszek *et al.* 2004), human placenta, and in mouse oocytes and embryos (Zhong *et al.* 2006).

The objective of the present work was to test the hypothesis that FGF17 is a candidate for paracrine signaling within the follicle. Specifically, we sought to localize *FGF17* mRNA in follicular cell types, to determine whether mRNA

expression is under the control of the major regulators of follicle development, FSH and insulin-like growth factor-I (IGF-I), and to gain insight into the potential role of FGF17 in the regulation of follicle steroidogenesis.

Materials and Methods

Tissues

Follicles of diameter ≥ 5 mm were dissected from the ovaries of adult cows (predominantly Nellore, *Bos indicus*) obtained in an abattoir local to the Sao Paulo State University campus in Botucatu and transported to the laboratory in saline on ice. Follicular fluid was aspirated, centrifuged, and frozen for progesterone (P₄) and E₂ assays. The antral cavity was flushed repeatedly with cold saline, and granulosa cells were recovered by centrifugation at 1200 g for 1 min and pooled with the follicular fluid pellet. The remaining granulosa cells adhering to the follicle wall were removed by gently scraping with a blunt Pasteur pipette, and the theca layer was removed with forceps and washed in saline by repeated passages through a 1 ml syringe. Samples were collected in Trizol (Invitrogen Life Technologies), homogenized with a Polytron, and submitted immediately to total RNA extraction according to the manufacturer's protocol.

Follicles were classed according to E₂:P₄ ratios of >1, 1–0.01, and <0.01 (Ireland *et al.* 1994), which we defined as healthy, transitional, and highly atretic respectively (Grimes & Ireland 1986). Mean follicle fluid steroid concentrations and follicle diameters for each follicle class are given in Table 1. Cross-contamination of theca and granulosa cells was tested by the detection of mRNA encoding cytochromes P450 aromatase (*CYP19A1*) and 17α-hydroxylase (*CYP17A1*) mRNA in each sample by PCR as described (Buratini *et al.* 2005b). The detection of *CYP19A1* amplicons in theca samples or of *CYP17A1* amplicons in granulosa samples indicated cross-contamination, and such samples were discarded.

Cumulus–oocyte complexes (grades 1 and 2; Leibfried & First 1979) were aspirated from antral follicles (2–8 mm) collected at an abattoir, and oocytes were mechanically isolated by careful and repeated pipetting until no adhering

cumulus cells could be observed under a stereomicroscope. Total RNA was extracted from pools of 20 oocytes with the RNeasy kit (Qiagen).

Cell culture

Granulosa cell culturing was performed as described (Gutiérrez *et al.* 1997) with modifications (Manuel Silva & Price 2000). All materials were obtained from Invitrogen, except where otherwise stated. Follicles with a ≤ 5 mm diameter were dissected from ovaries of *Bos taurus* cows obtained at an abattoir local to the University of Montreal in St Hyacinthe, and transported to the laboratory in PBS at 35 °C containing penicillin (100 IU/ml) and streptomycin (100 µg/ml). Follicles with obvious signs of atresia (avascular theca, debris in antrum) were discarded. Cells were collected by repeatedly passing the follicle wall through a pipette, washed twice by centrifugation at 980 g for 20 min each, and suspended in DMEM/F12 containing Hepes (20 mM), sodium selenite (4 ng/ml), BSA (0.1%; Sigma–Aldrich), penicillin (100 IU/ml), streptomycin (100 µg/ml), transferrin (2.5 µg/ml), non-essential amino acid mix (1.1 mM), androstenedione (10^{−7} M at the start of culture and 10^{−6} M at each medium change), and insulin (10 ng/ml). Cell viability was estimated with 0.4% trypan blue stain. Cells were seeded into 24-well tissue culture plates (Sarstedt, Newton, NC, USA) at a density of 10⁶/well in 1 ml medium. Cultures were maintained at 37 °C in 5% CO₂ in air for 6 days, with 700 µl medium being replaced every 2 days.

To determine the regulation of *FGF17* mRNA expression, cells were stimulated with graded doses of FSH (AFP-5332B, NIDDK, Bethesda, MD, USA; 0, 0.1, 1, 10, and 100 ng/ml) or IGF-I analog (LR3; Sigma–Aldrich; 0, 5, 10, 50, and 100 ng/ml) starting on day 2 of culture. At the end of the culture period, cells were collected in Trizol and stored at 70 °C until RNA extraction. Data were derived from three independent cultures performed at different times.

To determine the potential role of FGF17, granulosa cells were treated with FSH (10 ng/ml) to stimulate *FGFR3c* mRNA levels (Buratini *et al.* 2005b) and with graded doses of FGF17 (PeproTech, Rocky Hill, NJ, USA) starting on day 2 of culture. To measure steroid secretion, the medium was removed for steroid assay on day 6 and stored at −20 °C. The cells were lysed with 200 µl of 1 M NaOH for 2 h followed by neutralization with 200 µl of 1 M HCl for total cell protein measurement with the Bradford protein assay (Bio–Rad). These cultures were performed on three independent cultures performed at different times, and were performed several months after the FSH/IGF dose–response cultures.

Reverse transcription (RT)-PCR

Theca and granulosa cell RNA (1 µg) samples were incubated with DNase I (1 U/µg RNA; Invitrogen), and then reverse transcribed with SuperScript III (200 U/µl; Invitrogen) and oligo-d(T) primer. The RNA yield from pools of twenty oocytes was too low to be accurately quantified by

Table 1 Mean (±S.E.M.) follicular fluid estradiol (E₂) and progesterone (P₄) concentrations and follicle diameters of follicles classed as healthy, transitional, and atretic based on E₂:P₄ ratio

	Healthy (n=16)	Transitional (n=17)	Atretic (n=10)
Estradiol (ng/ml)	540±249*	16±3 ⁺	0.3±0.1*
Progesterone (ng/ml)	19±3*	44±7 ⁺	290±59*
Diameter (mm)	9±1	9±1	8±1
Diameter range (mm)	5–13	6–11	5–11

Within rows, means with different superscripts are significantly different (P<0.05).

spectrometry, so 8 µl aliquots of RNA, corresponding approximately to five oocytes, were incubated with DNase I (1 U/sample; Invitrogen) to produce 10 µl of RNA solution, which is the maximum volume for the RNA sample in the reverse transcription (RT) protocol (SuperScript III, Invitrogen).

Primers for *FGF17* mRNA were designed based on the predicted bovine sequences, and amplicons were sequenced to confirm identity. Relative real-time RT-PCR analysis was performed with an ABI 7500 using TaqMan Assay by Design (Applied Biosystems, Sao Paulo, Brazil) for the target gene (*FGF17*), as it provided a higher amplification efficiency in comparison with SYBR Green. As previous studies have shown that these two detection systems produce similar results for high abundance messages (Jeong *et al.* 2005), we used Power SYBR Green PCR Master Mix (Applied Biosystems) for housekeeping genes. Amplification efficiencies for target and housekeeping genes were similar. The primer sequences, fragment size, and annealing temperature for each gene are shown in Table 2. Reactions were optimized to provide maximum amplification efficiency for each gene. PCR was performed on 0.5 µl cDNA in 25 µl reaction volumes in duplicate, and the specificity of each PCR product was determined by melting curve analysis (for housekeeping genes) and confirmation of the amplicon size using electrophoresis with 2% agarose gels (for *FGF17* and housekeeping genes). Negative controls (water replacing cDNA) were run in every plate. The relative expression of each target gene was calculated using the $\Delta\Delta C_t$ method with efficiency correction (Pfaffl 2001); the control was a cDNA sample from each cell type analyzed. An initial analysis of *FGF17* mRNA across follicle cell types was performed with glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) as housekeeping gene, as this gene is expressed at similar levels in theca and granulosa cells (Buratini *et al.* 2005b). To select the most stable housekeeping gene for detailed analyses of each cell type, peptidylprolyl isomerase A (*PPIA*), *GAPDH*, and histone H2AFZ (*H2AFZ*) amplification profiles were compared using the geNorm applet for Microsoft Excel (medgen.ugent.be/genorm; Ramakers *et al.* 2003); the most stable housekeeping genes were *PPIA* for theca cells and oocytes, *GAPDH* for granulosa, and *H2AFZ* for cultured granulosa cells.

Immunohistochemistry

Bovine ovaries were collected from an abattoir, bisected, and fixed in paraformaldehyde. Fixed tissues were embedded in paraffin, and 3 µm sections were placed on poly-L-lysine-coated slides. Sections were deparaffinized in xylene twice for 20 min, and hydrated in successive 3 min washes in 95 and 85% ethanol. Antigen retrieval was achieved by incubating in 0.5 mM Tris-EDTA pH 9.0 at 96 °C for 30 min. Endogenous peroxidase was quenched by incubation in methanol with 5% hydrogen peroxide for 10 min, and then rinsed ten times in distilled water and twice for 5 min in 0.5 M Tris pH 7.4. Slides were then incubated with polyclonal FGF17 antibody (1.25 µg/ml; 500-P152; PeproTech) for 2 h at room temperature in a humidified chamber. Slides were washed in 0.5 M Tris pH 7.4, then incubated with HRP-conjugated secondary antibody for 40 min (EnVision Dual Link System, Dako, Carpinteria, CA, USA). Immunostaining was revealed with liquid DAB (3,3' diamino benzidine; Dako), and sections were counterstained with Harris hematoxylin. Six to seven sections were examined from each of three ovaries. Negative controls were performed by preincubating FGF17 antibody with twice the concentration of recombinant FGF17 protein for 2 h at room temperature (2.5 µg/ml; 100-27; PeproTech). The immunogen used for antibody production has at most 67% homology with other human FGFs. No bovine FGF8 family proteins are available, so it is not possible to test antibody specificity experimentally.

Steroid assays

E₂ and P₄ were assayed in follicular fluids using iodinated tracers and the antibodies furnished in the Third Generation Estradiol RIA and Progesterone RIA kit (Diagnostic Systems Laboratories Inc., Webster, TX, USA) respectively with a revised protocol (Buratini *et al.* 2005b). The standard curves and samples were diluted in PBS-gelatin. The intra- and inter-assay coefficients of variation (CV) were 7.4 and 13.5% respectively for E₂, and 6.8 and 7.0% respectively for P₄. The sensitivities of the assays were 0.3 ng/ml for E₂ and 0.2 ng/ml for P₄. E₂ was measured in conditioned medium in

Table 2 Details of primers used for real-time PCR

Target	Sequence	Fragment size (bp)	Annealing temperature (°C)
<i>GAPDH</i>	F: 5' ggc gtg aac cac gag aag tat aa 3' R: 5' ccc tcc acg atg cca aag t 3'	119	62
<i>PPIA</i>	F: 5' gcc atg gag cgc ttg gg 3' R: 5' cca cag tca gca atg gtg atc t 3'	65	60
<i>H2AFZ</i>	F: 5' gag gag ctg aac aag ctg ttg 3' R: 5' ttg tgg tgg ctc tca gtc ttc 3'	74	60
<i>FGF17</i>	F: 5' ccg ggt gcg cat caa g 3' R: 5' gct tgc ccc tct tat tca tac aga t 3' Probe FAM 5' ctg aga gtg aga aat ac 3'	62	60

F, forward primer; R, reverse primer.

duplicate as described (Bélanger *et al.* 1990) without solvent extraction. Intra- and inter-assay CV were 6 and 9% respectively. P_4 was measured in conditioned medium in duplicate as described (Lafrance & Goff 1985) with mean intra- and inter-assay CV were 7.2 and 18% respectively. The sensitivity of these assays was 10 and 4 pg per tube for E_2 and P_4 , equivalent to 0.3 and 20 ng/ μ g protein respectively. Steroid concentrations in culture medium were corrected for cell number by expressing per unit mass of total cell protein.

Statistical analysis

The data were transformed to logarithms if not normally distributed. ANOVA was used to compare follicle size and intrafollicular steroid concentrations across follicle classes, to test the effect of follicle class on *FGF17* mRNA abundance in granulosa cells, to test the effects of FSH and IGF-I on *FGF17* mRNA levels in cultured granulosa cells, and to test the effect of FGF17 on steroid secretion from cultured granulosa cells. Means comparisons were performed with the Tukey–Kramer HSD test. Non-parametric ANOVA was used to test the effect of follicle class on *FGF17* mRNA abundance in theca cells, and to compare *FGF17* mRNA abundance across follicle cell types as data were not normally distributed even after transformation to logarithms. Means comparisons were performed with the Kruskal–Wallis test. Data are presented as means $\pm$ S.E.M. Analyses were performed with JMP software (SAS Institute, Cary, NC, USA).

Results

Immunohistochemistry revealed the presence of FGF17 predominantly in oocytes and granulosa cells of preantral and antral follicles (Fig. 1). Staining was predominant in the nucleus of oocytes in preantral follicles, and was more intense in the cytoplasm of oocytes from antral follicles (Fig. 1A–C). Staining was weak in the theca cell layer, detected at background levels in deep ovarian stroma (Fig. 1E), and also clearly observed in the surface epithelium. Atretic follicles with only a few layers of granulosa cells and many pycnotic nuclei also stained strongly for FGF17 (Fig. 1D). No staining was observed when FGF17 antibody was preincubated with excess FGF17 (Fig. 1F).

A survey of *FGF17* expression in follicle cells by real-time PCR showed the presence of mRNA in pooled oocytes and at comparatively low levels in both granulosa and theca cells. We initially compared *FGF17* mRNA across follicle cell types using *GAPDH* as housekeeping gene. Relative abundance was higher in oocytes (51 ± 14) compared with granulosa and theca cells (0.025 ± 0.008 and 0.005 ± 0.001 respectively) from healthy and transitional follicles. A detailed assay of *FGF17* in granulosa and theca cells collected from healthy, transitional, and atretic follicles showed that mRNA abundance was significantly higher in granulosa and theca cells from atretic follicles than from healthy or transitional follicles (Fig. 2). There was no effect of diameter on *FGF17* expression within any of the health status classes.

Figure 1 Immunohistological detection of FGF17 in bovine ovaries. Staining is observed in the oocyte of primordial (A, arrow), primary (B), and antral (C) follicles. The granulosa cell layer was well stained in healthy (C and E) and atretic (D) antral follicles, whereas the theca layer was weakly stained. No staining was observed in the presence of excess FGF17 protein (F). Bars=20 μ m (A, B, and D), 30 μ m (C), and 50 μ m (E and F). g, granulosa; t, theca; s, stroma.

As FGF17 protein and mRNA were detected in granulosa cells, we determined whether expression can be regulated by FSH and IGF-I. Cells were cultured in serum-free medium and treated for 6 days with graded doses of FSH or IGF-I. Both gonadotropic hormones significantly inhibited *FGF17* mRNA abundance while increasing E_2 secretion (Fig. 3). Given this apparent inverse relationship between E_2 and FGF17, we sought to determine whether FGF17 can regulate E_2 secretion from granulosa cells. FGF17 inhibited E_2 and P_4 secretion from cultured granulosa cells in a dose-dependent manner (Fig. 4).

Discussion

These data demonstrate for the first time the expression of FGF17 protein and mRNA in the ovary. The most significant findings are the higher abundance of *FGF17* mRNA in granulosa and theca cells of atretic follicles compared with healthy follicles, the inhibition of *FGF17* mRNA abundance in granulosa cells by FSH and IGF-I, and the inhibitory effect of FGF17 on granulosa cell steroid secretion. These data point to an inhibitory role for FGF17 in follicular steroidogenesis and its involvement in the mechanisms controlling follicle atresia.

Figure 2 Follicular *FGF17* mRNA abundance varies with follicle health. Follicles were classified as healthy, transitional, or atretic based on $E_2:P_4$ ratios in follicle fluid and RNA measured by real-time PCR in isolated granulosa and theca cells. Data are presented as mean ($\pm$ S.E.M.) values relative to the housekeeping gene and to a calibrator sample present in all PCR runs, which were calculated using the $\Delta\Delta C_t$ method with efficiency correction. Bars with different letters are significantly different ($P < 0.05$). Numbers in parentheses denote the number of samples analyzed.

In the female reproductive system, *FGF17* mRNA has been described only in isolated mouse oocytes and embryos (Zhong *et al.* 2006). In the present study, we describe *FGF17* mRNA predominantly in oocytes, compared with lower expression in granulosa and theca cells. This conclusion is drawn from an analysis of relative expression (calculated as $\Delta\Delta C_t$ value) in all three cell types using *GAPDH* as housekeeping gene. This is based on the assumption that this housekeeping gene is expressed at similar levels in all three cell types, and our previous data have shown this to be valid for granulosa and theca cells (Buratini *et al.* 2005b). Analysing the same data with *PPIA* as housekeeping gene gave the same results; mRNA abundance was at least 100-fold greater in oocytes compared with granulosa and theca cells. This conclusion is consistent with the oocyte localization of the prototype member of this FGF subfamily, FGF8 (Valve *et al.* 1997, Buratini *et al.* 2005b, Sugiura *et al.* 2007). The higher expression of *FGF17* in the oocyte does not imply that this cell is the predominant intrafollicular source of FGF17 as the number of somatic cells is greatly superior within the follicle, but rather suggests that oocyte-derived FGF17 may play a role in the mechanisms controlling cumulus cells differentiation. In fact FGF8 was shown to synergize with BMP15 to promote glycolysis in murine cumulus cells (Sugiura *et al.* 2007).

Other FGFs have been localized to the bovine oocyte, including FGF2 and FGF10 (van Wezel *et al.* 1995, Buratini *et al.* 2007), but this is not a FGF-wide phenomenon as *FGF7* mRNA was readily detected in theca cells but not in oocytes (Buratini *et al.* 2007).

Protein was predominantly localized to the nucleus of oocytes in primordial and primary follicles, whereas in follicles at later stages of development, FGF17 protein was also clearly detected in the ooplasm. This suggests translocation of protein to the nucleus in oocytes, which is more apparent in primordial and primary follicles. Nuclear translocation of endogenous FGF has been reported for FGF1, -2, and -3 (Zhan *et al.* 1992, Kiefer & Dickson 1995, Sheng *et al.* 2004). FGF17 protein was also present in the granulosa cell layer in healthy follicles, and to a lesser extent in theca cells, which is in agreement with greater mRNA abundance in granulosa cells. Strong staining in the cytoplasm of granulosa cells may also reflect binding of FGF17 to receptors (FGFR2C and FGFR3C; Berisha *et al.* 2004, Buratini *et al.* 2005b) and internalization. This is supported by evidence for receptor-mediated internalization of exogenous FGFs in several cell lines (Belleudi *et al.* 2002, Olsnes *et al.* 2003, Wesche *et al.* 2006). Therefore, the staining pattern combined with mRNA data suggests an autocrine action for FGF17 in granulosa cells within the follicle wall. Detection of FGF17 appears to be specific as preincubation of the antibody with FGF17 abolished staining, although we cannot rule out the possibility that the antibody reacts with other bovine FGFs.

Granulosa and theca cell abundance of *FGF17* mRNA was significantly increased in atretic follicles compared with

Figure 3 Effects of FSH and IGF-I on *FGF17* mRNA abundance and steroid secretion in granulosa cells. Cells from follicles 2 to 5 mm diameter were placed in serum-free culture, and the stated doses of FSH or IGF-I were added on day 2. Total RNA was collected on day 6, and *FGF17* mRNA abundance was measured by real-time PCR. Data are presented as mean ($\pm$ S.E.M.) values relative to a calibrator sample by the $\Delta\Delta C_t$ method with efficiency correction. Steroid concentrations were measured by RIA on day 6 and expressed relative to total protein at the end of culture. Bars with different letters are significantly different ($P < 0.05$). Data were derived from three independent cultures.

Figure 4 Effects of FGF17 on steroid secretion from granulosa cells. Cells from follicles 2 to 5 mm diameter were placed in serum-free culture and the stated doses of FGF17 were added on day 2. Media were collected on day 6 for measurement of steroid concentrations by RIA. Data are presented as mean ($\pm$ S.E.M.) values and expressed relative to total cell protein. Bars with different letters are significantly different ($P < 0.05$). Data were derived from three independent cultures.

healthy follicles. FGF17 protein staining was also strong in the granulosa layer in atretic follicles. It is not known whether this is part of an apoptotic pathway or dysregulation of gene expression during cell death. There was no effect of size on *FGF17* expression in atretic follicles ranging from 5 to 11 mm, suggesting that *FGF17* expression is enhanced as non-ovulatory dominant follicles and also non-selected recruited follicles enter atresia. This pattern of expression of *FGF17* differs from that of *FGF7* and *FGF10*: *FGF10* mRNA abundance was significantly lower in atretic follicles compared with healthy follicles, whereas that of *FGF7* did not change (Buratini *et al.* 2007). There is no information on the expression of other FGFs in atretic follicles.

As *FGF17* mRNA and protein were both detected in granulosa cells, we determined whether the major gonadotrophic hormones FSH and IGF-I regulate mRNA levels. Both hormones decreased *FGF17* mRNA abundance while increasing E_2 secretion, which is consistent with the low mRNA levels observed in healthy, estrogenic follicles. We are not aware of other reports describing the regulation of FGF expression by gonadotropins in granulosa cells, but microarray studies identified *FGF2* and *FGF5* as genes downregulated by FSH in the ovarian surface epithelium (Ji *et al.* 2004). It has been proposed that genes downregulated by FSH in granulosa cells are those primarily involved in cell cycle control and apoptosis (Sasson *et al.* 2003), again suggesting a role for FGF17 in atresia. Therefore, the ability of FSH and IGF-I to suppress FGF expression may significantly account for their well-established pro-survival action in antral follicles.

To explore the function of FGF17, we cultured bovine granulosa cells in serum-free medium with recombinant FGF17. In this estrogenic cell model, FGF2 and 10 have been demonstrated to inhibit E_2 secretion (Cao *et al.* 2006, Buratini *et al.* 2007), and we show here that FGF17 has a similar effect. Interestingly, FGF2 was more potent at inhibiting E_2 secretion than P_4 secretion (Vernon & Spicer 1994, Cao *et al.* 2006), whereas in the present study FGF17 was more potent at suppressing P_4 secretion; E_2 secretion was inhibited only at higher doses of FGF17. This suggests that while all FGFs studied to date inhibit steroidogenesis in bovine granulosa cells, they display different specificities for estrogenic and progestagenic pathways. Potential mechanisms of this differential action have not been investigated, but are likely related to mitogen-activated protein kinase (MAPK) activity; FGFRs activate the MAPK pathway (Powers *et al.* 2000), and MAPK activity suppresses aromatase (*CYP19A1*) mRNA expression (Silva & Price 2000, Fan *et al.* 2009). As it would be critical to keep FGF17 levels low to allow E_2 secretion and continued follicle growth, it is possible that FSH and IGF-I may stimulate E_2 production in part through the inhibition of FGF expression.

In conclusion, the present study suggests a physiological role for FGF17 in the control of granulosa cell differentiation. FGF17 suppressed steroid secretion from granulosa cells, and mRNA abundance was decreased by FSH and IGF-I. This is consistent with lower *FGF17* mRNA levels in healthy compared with atretic follicles. One potential role for FGF17 may be to inhibit E_2 secretion and follicle growth in regressing follicles; therefore, it may be critical for FSH and IGF-I to suppress *FGF17* expression during growth of the dominant follicle.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil, and the Natural Sciences and Engineering Research Council of Canada (NSERC).

Acknowledgements

We thank Dr J F Garcia (Universidade Estadual Paulista, Araçatuba) for sequencing the *FGF17* PCR products, and Drs A C S Castilho, P Pinczowski, and D M Guerra for their technical assistance.

References

- Bé langer A, Couture J, Caron S & Roy R 1990 Determination of nonconjugated and conjugated steroid levels in plasma and prostate after separation on C-18 columns. *Annals of the New York Academy of Sciences* **595** 251–259.
- Belleudi F, Ceridono M, Capone A, Serafino A, Marchese C, Picardo M, Frati L & Torrisi M 2002 The endocytic pathway followed by the keratinocyte growth factor receptor. *Histochemistry and Cell Biology* **118** 1–10.
- Berisha B, Schams D, Kosmann M, Amselgruber W & Einspanier R 2000 Expression and localisation of vascular endothelial growth factor and basic fibroblast growth factor during the final growth of bovine ovarian follicles. *Journal of Endocrinology* **167** 371–382.

- Berisha B, Sinowatz F & Schams D 2004 Expression and localization of fibroblast growth factor (FGF) family members during the final growth of bovine ovarian follicles. *Molecular Reproduction and Development* **67** 162–171.
- Buratini J Jr, Glapinski VF, Giometti IC, Teixeira AB, Costa IB, Avellar MC, Barros CM & Price CA 2005a Expression of fibroblast growth factor-8 and its cognate receptors, fibroblast growth factor receptor (FGFR)-3c and -4, in fetal bovine preantral follicles. *Molecular Reproduction and Development* **70** 255–261.
- Buratini J Jr, Teixeira AB, Costa IB, Glapinski VF, Pinto MGL, Giometti IC, Barros CM, Cao M, Nicola ES & Price CA 2005b Expression of fibroblast growth factor-8 and regulation of cognate receptors, fibroblast growth factor receptor (FGFR)-3c and -4, in bovine antral follicles. *Reproduction* **130** 343–350.
- Buratini J Jr, Pinto MG, Castilho AC, Amorim RL, Giometti IC, Portela VM, Nicola ES & Price CA 2007 Expression and function of fibroblast growth factor 10 and its receptor, fibroblast growth factor receptor 2B, in bovine follicles. *Biology of Reproduction* **77** 743–750.
- Cao M, Nicola E, Portela VM & Price CA 2006 Regulation of serine protease inhibitor-E₂ and plasminogen activator expression and secretion by follicle stimulating hormone and growth factors in non-luteinizing bovine granulosa cells *in vitro*. *Matrix Biology* **25** 342–354.
- Fan H-Y, Liu Z, Shimada M, Sterneck E, Johnson PF, Hedrick SM & Richards JS 2009 MAPK3/1(ERK1/2) in ovarian granulosa cells are essential for female fertility. *Science* **324** 938–941.
- Ford-Perriss M, Abud H & Murphy M 2001 Fibroblast growth factors in the developing central nervous system. *Clinical and Experimental Pharmacology and Physiology* **28** 493–503.
- Grimes RW & Ireland JJ 1986 Relationship of macroscopic appearance of the surface of bovine ovarian follicles concentrations of steroids in follicular fluid, and maturation of oocytes *in vitro*. *Biology of Reproduction* **35** 725–732.
- Gutiérrez CG, Campbell BK & Webb R 1997 Development of a long-term bovine granulosa cell culture system: induction and maintenance of estradiol production, response to follicle-stimulating hormone, and morphological characteristics. *Biology of Reproduction* **56** 608–616.
- Ireland J, Good T, Knight P & Ireland J 1994 Alterations in amounts of different forms of inhibin during follicular atresia. *Biology of Reproduction* **50** 1265–1276.
- Itoh N & Ornitz DM 2004 Evolution of the *Fgf* and *Fgfr* gene families. *Trends in Genetics* **20** 563–569.
- Jeong YJ, Choi HW, Shin HS, Cui XS, Kim NH, Gerton GL & Jun JH 2005 Optimization of real time RT-PCR methods for the analysis of gene expression in mouse eggs and preimplantation embryos. *Molecular Reproduction and Development* **71** 284–289.
- Ji Q, Liu PI, Chen PK & Aoyama C 2004 Follicle stimulating hormone-induced growth promotion and gene expression profiles on ovarian surface epithelial cells. *International Journal of Cancer* **112** 803–814.
- Kiefer P & Dickson C 1995 Nucleolar association of fibroblast growth factor 3 via specific sequence motifs has inhibitory effects on cell growth. *Molecular and Cellular Biology* **15** 4364–4374.
- Krejci P, Krakow D, Mekikian PB & Wilcox WR 2007 Fibroblast growth factors 1, 2, 17, and 19 are the predominant FGF ligands expressed in human fetal growth plate cartilage. *Pediatric Research* **61** 267–272.
- Lafrance M & Goff AK 1985 Effect of pregnancy on oxytocin-induced release of prostaglandin F_{2α} in heifers. *Biology of Reproduction* **33** 1113–1119.
- Lavranos TC, Rodgers HF, Bertonecello I & Rodgers RJ 1994 Anchorage-independent culture of bovine granulosa cells: the effects of basic fibroblast growth factor and dibutyryl cAMP on cell division and differentiation. *Experimental Cell Research* **211** 245–251.
- Leibfried L & First NL 1979 Characterization of bovine follicular oocytes and their ability to mature *in vitro*. *Journal of Animal Science* **48** 76–86.
- Silva JM & Price CA 2000 Effect of follicle-stimulating hormone on steroid secretion and messenger ribonucleic acids encoding cytochromes P450 aromatase and cholesterol side-chain cleavage in bovine granulosa cells *in vitro*. *Biology of Reproduction* **62** 186–191.
- Nilsson E, Parrott JA & Skinner MK 2001 Basic fibroblast growth factor induces primordial follicle development and initiates folliculogenesis. *Molecular and Cellular Endocrinology* **175** 123–130.
- O'Leary DD, Chou SJ & Sahara S 2007 Area patterning of the mammalian cortex. *Neuron* **56** 252–269.
- Olsnes S, Klingenberg O & Wiedlocha A 2003 Transport of exogenous growth factors and cytokines to the cytosol and to the nucleus. *Physiological Reviews* **83** 163–182.
- Ornitz DM, Xu J, Colvin JS, McEwen DG, MacArthur CA, Coulier F, Gao G & Goldfarb M 1996 Receptor specificity of the fibroblast growth factor family. *Journal of Biological Chemistry* **271** 15292–15297.
- Parrott JA & Skinner MK 1998 Developmental and hormonal regulation of keratinocyte growth factor expression and action in the ovarian follicle. *Endocrinology* **139** 228–235.
- Pfaff MW 2001 A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* **29** e45.
- Polnaszek N, Kwabi-Addo B, Wang J & Ittmann M 2004 FGF17 is an autocrine prostatic epithelial growth factor and is upregulated in benign prostatic hyperplasia. *Prostate* **60** 18–24.
- Powers CJ, McLeskey SW & Wellstein A 2000 Fibroblast growth factors, their receptors and signaling. *Endocrine-Related Cancer* **7** 165–197.
- Ramakers C, Ruijter JM, Deprez RH & Moorman AF 2003 Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neuroscience Letters* **339** 62–66.
- Sasson R, Dantes A, Tajima K & Amsterdam A 2003 Novel genes modulated by FSH in normal and immortalized FSH-responsive cells: new insights into the mechanism of FSH action. *FASEB Journal* **17** 1256–1266.
- Sheng Z, Lewis JA & Chirico WJ 2004 Nuclear and nucleolar localization of 18-kDa fibroblast growth factor-2 is controlled by C-terminal signals. *Journal of Biological Chemistry* **279** 40153–40160.
- Spicer LJ & Stewart RE 1996 Interactions among basic fibroblast growth factor, epidermal growth factor, insulin, and insulin-like growth factor-I (IGF-I) on cell numbers and steroidogenesis of bovine thecal cells: role of IGF-I receptors. *Biology of Reproduction* **54** 255–263.
- Sugiura K, Su YQ, Diaz FJ, Pangas SA, Sharma S, Wigglesworth K, O'Brien MJ, Matzuk MM, Shimasaki S & Eppig JJ 2007 Oocyte-derived BMP15 and FGFs cooperate to promote glycolysis in cumulus cells. *Development* **134** 2593–2603.
- Valve E, Penttilä TL, Paranko J & Harkonen P 1997 FGF-8 is expressed during specific phases of rodent oocyte and spermatogonium development. *Biochemical and Biophysical Research Communications* **232** 173–177.
- Vernon RK & Spicer LJ 1994 Effects of basic fibroblast growth factor and heparin on follicle-stimulating hormone-induced steroidogenesis by bovine granulosa cells. *Journal of Animal Science* **72** 2696–2702.
- Webb R, Nicholas B, Gong JG, Campbell BK, Gutierrez CG, Garverick HA & Armstrong DG 2003 Mechanisms regulating follicular development and selection of the dominant follicle. *Reproduction* **61** 71–90.
- Wesche J, Malecki J, Wiedlocha A, Skjerpen CS, Claus P & Olsnes S 2006 FGF-1 and FGF-2 require the cytosolic chaperone Hsp90 for translocation into the cytosol and the cell nucleus. *Journal of Biological Chemistry* **281** 11405–11412.
- van Wezel IL, Umaphysivam K, Tilley WD & Rodgers RJ 1995 Immunohistochemical localization of basic fibroblast growth factor in bovine ovarian follicles. *Molecular and Cellular Endocrinology* **115** 133–140.
- Zhan X, Hu X, Friedman S & Maciag T 1992 Analysis of endogenous and exogenous nuclear translocation of fibroblast growth factor-1 in NIH 3T3 cells. *Biochemical and Biophysical Research Communications* **188** 982–991.
- Zhang X, Ibrahim OA, Olsen SK, Umemori H, Mohammadi M & Ornitz DM 2006 Receptor specificity of the fibroblast growth factor family. The complete mammalian FGF family. *Journal of Biological Chemistry* **281** 15694–15700.
- Zhong W, Wang QT, Sun T, Wang F, Liu J, Leach R, Johnson A, Puscheck EE & Rappolee DA 2006 FGF ligand family mRNA expression profile for mouse preimplantation embryos, early gestation human placenta, and mouse trophoblast stem cells. *Molecular Reproduction and Development* **73** 540–550.

Received in final form 5 June 2009

Accepted 17 June 2009

Made available online as an Accepted Preprint
17 June 2009