Acute cytokine-mediated downregulation of the zinc transporter ZnT8 alters pancreatic β -cell function

Malek El Muayed*, Liana K Billings^{1,*}, Meera R Raja^{2,*}, Xiaomin Zhang³, Paul J Park⁴, Marsha V Newman, Dixon B Kaufman³, Thomas V O'Halloran² and William L Lowe Jr

Division of Endocrinology, Metabolism and Molecular Medicine, Feinberg School of Medicine, Northwestern University, 303 East Chicago Avenue, Tarry 15-755, Chicago, Illinois 60611, USA

¹Massachusetts General Hospital, Harvard University, 55 Fruit Street, Boston, Massachusetts 02114, USA

²The Chemistry of Life Processes Institute and Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208-3113, USA

³Division of Transplant Surgery, Feinberg School of Medicine, Northwestern University, 303 East Chicago Avenue, Chicago, Illinois 60611, USA

⁴Northwestern University, 2145 Sheridan Road, Evanston, Illinois 60208, USA

(Correspondence should be addressed to M El Muayed; Email: m-muayed@northwestern.edu)

*(M El Muayed, L K Billings and M R Raja contributed equally to this work)

Abstract

Genetic studies suggest that Zn transporters such as ZnT8 play a role in insulin secretion by pancreatic β -cells; however, little is known about the dynamic roles of Zn trafficking pathways on β -cell physiology. To test the acute effects of the inflammatory cytokines interleukin 1 β (IL1 β) and tumor necrosis factor α (TNF α) on Zn homeostasis, the mRNA expression profile of Zn transporters of the ZnT and ZIP families was examined. Exposure of MIN6 cells or primary murine islets to IL1 β or TNF α altered the mRNA expression profile of Zn transporters; most notable was decreased ZnT8 mRNA levels. siRNA-mediated gene knockdown was used to examine the effects of decreased ZnT8 expression in primary

dispersed murine islet cells from C57/BL6 mice and MIN6 cells. ZnT8 knockdown in these murine islets led to reduced glucose stimulated insulin secretion without altering the total cellular insulin content or cell viability at normal or supraphysiological Zn concentrations. The labile Zn content determined by flow cytometry after loading with the Zn-specific sensor FluoZin-3 AM was decreased in MIN6 cells following ZnT8 knockdown or IL1 β treatment. These results suggest that an acute decrease in ZnT8 levels impairs β -cell function and Zn homeostasis, and may contribute to inflammatory cytokine-induced alterations in β -cell function. *Journal of Endocrinology* (2010) **206**, 159–169

Introduction

A low level chronic inflammatory state with elevated circulating inflammatory cytokine levels is thought to be an important contributor to the development of type 2 diabetes mellitus (T2DM). Impaired function of insulin secreting β -cells after exposure to the inflammatory cytokines interleukin 1 β (IL1 β) or tumor necrosis factor α (TNF α) has been shown in human and rodent β -cells as well as in MIN6 cells, a β -cell line. Several mechanisms for this effect have been implicated (Xenos *et al.* 1992, 1993, 1994, Corbett & McDaniel 1995, Rabinovitch & Suarez-Pinzon 1998, Wu *et al.* 2001, Donath *et al.* 2008). Altered expression of several genes involved in β -cell function is thought to be one of the mechanisms underlying this effect of cytokines (Donath *et al.* 2008).

Zn is the second most abundant transition metal in mammals (Wallwork *et al.* 1983, Milne *et al.* 1985, Ward & Mason 1987) and is utilized in a number of biological

processes (Cousins *et al.* 2006). Zn homeostasis is tightly regulated on the cellular and systemic level through various mechanisms (Outten & O'Halloran 2001, Cousins *et al.* 2006) and is of particular importance to pancreatic β -cell physiology (Chausmer 1998, Dodson & Steiner 1998, Chimienti *et al.* 2005, 2006, Lemaire *et al.* 2009). β -Cells have exceedingly high intracellular concentrations of Zn, particularly within secretory vesicles, where Zn facilitates the packaging of insulin into hexamers (Figlewicz *et al.* 1984, Dodson & Steiner 1998, Chimienti *et al.* 2006, Lemaire *et al.* 2009). In order to utilize Zn, the β -cells require the trafficking of Zn across cell and vesicular membranes, mainly through members of the Zn transporter (ZnT, *SLC30A*) and Zrt-Irt-like protein (ZIP, *SLC39A*) transporter families. These facilitate Zn diffusion out of or into the cytoplasm respectively. The Zn transporter encoded by *SLC30A8*, ZnT8, plays a particularly important role in Zn homeostasis in β -cells (Chimienti *et al.* 2005, 2006, Lemaire *et al.* 2009). ZnT8 is highly expressed in β -cells (Chimienti *et al.* 2005, Mocchegiani *et al.* 2008), and

localizes mainly to the membranes of secretory vesicles in β -cells (Chimienti *et al.* 2006). Other genetic variations in *SLC30A8* have been found to be associated with fasting glucose levels (Dupuis *et al.* 2010). Strong association in humans has been demonstrated between a nonsynonymous single-nucleotide polymorphism rs1326634 in *SLC30A8*, and increased risk for the development of T2DM or β -cell dysfunction (Saxena *et al.* 2007, Scott *et al.* 2007, Sladek *et al.* 2007, Zeggini *et al.* 2007, Boesgaard *et al.* 2008, Hertel *et al.* 2008, van Hoek *et al.* 2008, Horikawa *et al.* 2008, Omori *et al.* 2008, Wu *et al.* 2008, Xiang *et al.* 2008). Other genetic variations in *SLC30A8* have been found to be associated with fasting glucose levels (Dupuis *et al.* 2010). Recently, three studies described impaired Zn accumulation in secretory vesicles of islets isolated from ZnT8 knockout mice. Impaired β -cell function was present under certain age and high-fat dietary conditions (Lemaire *et al.* 2009, Nicolson *et al.* 2009, Pound *et al.* 2009). This highlighted the deleterious effects of a chronic and complete loss of ZnT8 function on islet cell physiology, and the likely role of compensatory and environmental factors in this process.

Given the importance of well-calibrated Zn homeostasis for β -cell physiology, we examined whether the effects of the cytokines IL1 β and TNF α on β -cell function may be mediated, in part, through altered expression of Zn transporters. We examined the effect of treating MIN6 cells with these cytokines on the expression of Zn transporters in the ZIP and ZnT families. Additionally, we examined the consequences of the most significant cytokine-induced alteration in Zn transporter expression, decreased expression of ZnT8, in MIN6 cells and primary murine islets.

Materials and Methods

Islet purification

Islets from 10 to 14 weeks old C57BL/6 male mice (Jackson Laboratories, Bar Harbor, ME, USA) were isolated as described previously (Blomeier *et al.* 2006). After washing, the islets were handpicked and counted under microscopic guidance. All studies were approved by the Northwestern University Animal Care and Use Committee.

MIN6 cell culture

MIN6 cells were a gift from Dr Junichi Miyazaki (Osaka University Medical School, Osaka, Japan), and were grown and maintained as described previously (Miyazaki *et al.* 1990). Cells between passages 28 and 35 were used.

Cytokine treatment

Cytokine treatment was performed as previously reported with modifications (Chin-Chance *et al.* 2006). Briefly, MIN6 cells between passages 28 and 34 were plated at a density

of 2×10^6 cells per 60-mm dish. Forty-eight hours after plating, the culture medium was exchanged for serum reduced culture medium containing 1% heat inactivated fetal bovine serum (FBS) supplemented with either no cytokines, 5 ng/ml murine IL1 β , 10 ng/ml murine TNF α , or a combination of both (R&D Systems, Minneapolis, MN, USA). The cells were treated with cytokines for 6, 24, or 48 h. For studies with murine islets, purified islets were incubated for 24 h in RPMI 1640 (ATCC, Manassas, VA, USA) supplemented with 1% non-heat inactivated FBS. The media were supplemented with either no cytokines, 5 ng/ml IL1 β , or 10 ng/ml TNF α .

mRNA extraction and real-time PCR

RNA was extracted from MIN6 and dispersed murine islet cells using Trizol reagent (Invitrogen) according to the manufacturer's instructions. RNA was extracted from intact murine islets using the RNeasy Mini kit (Qiagen) according to the manufacturer's recommendations. The iScript cDNA synthesis kit (Bio-Rad) was used for reverse transcription of 0.5 μ g RNA following treatment with DNase (Turbo DNase kit, Ambion, Austin, TX, USA) according to the manufacturer's instructions. Real-time PCR was performed using the SYBR-Green super mix reagent (Bio-Rad) and the Bio-Rad iCycler PCR system. Primers for ZIP1 through 14, ZnT1 through 10, and β -actin, and 18S (idtDNA, Coralville, IA, USA) were used in the PCRs. The sequences for the real-time PCR primers are listed in [Supplementary Table 1](#), see section on [supplementary data](#) given at the end of this article. For real-time PCR, an initial denaturation step at 95 °C $\times$ 5 min was followed by 40 cycles at 95 °C $\times$ 15 s and 60 °C $\times$ 30 s. A melt curve analysis was performed by incubating the reaction at 95 °C $\times$ 1 min, cooling to 60 °C for 1 min, and then increasing the temperature 0.5 °C every 10 s up to 100 °C. All primers were validated using mRNA from mouse islets, MIN6 cells, and embryoid bodies prepared from murine embryonic stem cells. 2% agarose gel electrophoresis demonstrated a product of the expected size for all the ZnTs and ZIPs. Melt curve analyses demonstrated a single PCR product for each reaction. The $\Delta - \Delta C_t$ method was used to calculate the relative level of mRNA of the gene of interest compared with the control condition after normalizing using the level of β -actin mRNA (Livak & Schmittgen 2001). 18S was used as an additional normalizer for experiments comparing MIN6 and primary islet Zn transporter expression profiles.

MIN6 cell imaging

Untreated live MIN6 cells were imaged following incubation for 1 h in phenol red-free DMEM media (Invitrogen) supplemented with the labile Zn-specific fluorescent sensor FluoZin-3-AM (Invitrogen) at a concentration of 5 μ M. The images were obtained on an IX2-DSU/BX-DSU spinning disc confocal microscope (Olympus, Center Valley, PA, USA)

in the Northwestern University cell imaging facility. Images were analyzed with Metamorph software (Universal Imaging Corp., Downingtown, PA, USA).

Flow cytometry

Flow cytometry was carried out 72 h after transfection with siRNA or after treatment with IL1 β for 24 h. Cells were washed twice with MIN6 cell culture medium containing 1 or 15% heat-inactivated FBS for IL1 β or ZnT8 knockdown experiments respectively. Cells were subsequently incubated for 1 h in basic media supplemented with 2 mM FluoZin-3 AM (Invitrogen) and 1 mM CellTrace Red (Invitrogen) at 37 °C in 5% CO₂. Cells were then washed twice with cell culture medium and once with PBS. The cells were dispersed using 0.05% trypsin, collected by centrifugation, resuspended in PBS, and passed through a 70- μ m Cell Strainer (BD Biosciences, Bedford, MA, USA). Cells were analyzed on an LSR II system (BD, Franklin Lakes, NH, USA). Only the Zn fluorescence of live, single cells, as determined by the CellTrace side scatter and forward scatter fluorescence patterns, was analyzed.

ZnT8 knockdown in dispersed islet cells or MIN6 cells

Islets harvested from C57BL/6 mice were washed with PBS (Invitrogen), pelleted by centrifugation at 0.5 relative centrifugal force (RCF) $\times$ 1 min, resuspended in 0.5% trypsin (Invitrogen), and incubated at 37 °C for 4–5 min. The cells were dispersed by pipetting at the 2-min time point and end of the incubation period. The incubation was terminated by adding culture medium (RPMI 1640 supplemented with 10% non-heat inactivated FBS (Hyclone, Logan, UT, USA), 100 U/ml penicillin, 100 μ g/ml streptomycin, and 2 mmol/l glutamine). The dispersed cells were collected by centrifugation (0.5 RCF $\times$ 5 min), resuspended in culture medium, and plated onto tissue culture dishes (BD-Falcon, San Jose, CA, USA). MIN6 cells were cultured as previously described (Miyazaki *et al.* 1990).

MIN6 cells or dispersed murine islet cells were plated in a cell media volume of 1.5 ml at a density of 1.0×10^5 or 2.0×10^5 cells per 35-mm tissue culture dish respectively. Transfection with siRNA was carried out 5 or 24 h after plating for dispersed murine islets or MIN6 cells respectively. Cells were transfected with control (negative control siRNA #3, Dharmacon, Lafayette, CO, USA) or ZnT8 siRNA (Nicolson *et al.* 2009; see Supplementary Table 1 for sequence of the ZnT8 siRNA) by adding 500 μ l of transfection mix (500 μ l Opti-MEM-I transfection medium (Invitrogen), 150 pmol siRNA (final concentration 75 nM, Dharmacon), and 7 μ l Lipofectamin RNAimax (Invitrogen), prepared according to the manufacturer's instructions). In MIN6 cell experiments, the cell culture medium was exchanged for antibiotic-free standard cell culture medium immediately prior to transfection. Cells were maintained in this mix of culture and transfection medium until experiments were

carried out 72 h later. In each experiment, cells were transfected with ZnT8 siRNA or non-targeting control siRNA in parallel. Decreased ZnT8 mRNA and protein levels were verified using real-time PCR and western blot analysis (see below).

Western blot analysis

Western blot analysis of cell lysates was carried out as described previously (Liu *et al.* 2001), with modifications as described herein. Samples (20 μ g of protein) were suspended in β -mercaptoethanol (β -ME)-containing laemmli buffer without heating and subjected to 10% SDS-PAGE followed by semi dry transfer for 2 h to a polyvinylidene fluoride membrane (Immobilon-P, Millipore, Bedford, MA, USA). The membranes were blocked for 2 h and then hybridized overnight with a 1:2000 dilution of polyclonal rabbit-anti-rat ZnT8 antibody (Mellitech, Grenoble, France). The membranes were then washed $\times$ 3 for 10 min in Tris-Buffered Saline Tween-20 (TBS-T) and hybridized with a 1:7500 dilution of secondary antibody (HRP-conjugated goat anti-rabbit IgG, Promega) for 1 h. The membrane was washed $\times$ 3 for 10 min in TBS-T. Immunoreactive bands were detected using an ECL detection kit (Amersham). All of the above steps were carried out at 4 °C on ice. The blocking and antibody incubation steps were carried out in TBS-T containing 5% w/v non-fat milk powder (Carnation, Wilkes-Barre, PA, USA). To document equivalent loading of protein, the immunoblots were stripped in 62.5 mM Tris-HCl (pH 6.7), 20% SDS, and 100 mM β -ME for 30 min at 50 °C, rehybridized with a monoclonal mouse anti-mouse GAPDH antibody (Cell Signalling, Danvers, MA, USA), and an HRP-coupled secondary goat anti-mouse antibody.

Glucose-stimulated insulin secretion and total cellular insulin content

To measure glucose-stimulated insulin secretion (GSIS), dispersed murine islet cells were washed three times with PBS 72 h after transfection. Subsequently, the cells were equilibrated in HEPES-buffered Krebs solution (130 mM NaCl, 3.6 mM KCl, 0.5 mM NaH₂PO₄, 0.5 mM MgSO₄, 1.5 mM CaCl₂, 1% HEPES, 0.1% BSA, pH 7.4) containing 5.6 mM glucose for 30 min. This was followed by incubation in HEPES-buffered Krebs solution with 5.6 mM glucose for 30 min, and then HEPES-buffered Krebs solution with 16.7 mM glucose for 30 min. After each period of incubation, 200 μ l of the supernatant was collected. Samples were stored at -80 °C. In separate experiments, cells were incubated for 72 h following transfection and then lysed after washing three times with PBS by adding 100 μ l of RIPA lysis buffer supplemented with a cocktail of protease and phosphatase inhibitors (Calbiochem, Madison, WI, USA). Total protein concentration in the lysate was measured using the Micro BCA Protein Assay kit (Pierce, Thermo Fisher Scientific, Waltham, MA, USA) according to the

manufacturer's recommendations. Measurement of insulin was carried out by ELISA (Alpco Diagnostics, San Francisco, CA, USA) according to the manufacturer's instructions.

Cell viability

Cell viability experiments were performed in black, opaque 96-well tissue culture plates (BD-Falcon). Dispersed murine islet cells were cultured and transfected with either control or ZnT8 siRNA as described above with proportional adjustment for the decreased culture area. The medium was supplemented with ZnCl₂ to final concentrations of Zn of 5, 19, 34, 48, 62, 76, 91, or 105 μ M at the time of transfection. Cell viability was assessed 72 h post transfection using the double fluorescence Multitox assay (Promega) according to the manufacturer's instructions. Briefly, 100 μ l of the assay mix was added to each well and incubated at 37 °C for 2 h. Fluorescence was measured on a fluorescence plate reader at excitation wavelength (Ex) 400 nm/emission wavelength (Em) 505 nm and Ex 485/Em 520, which measured fluorescence from proteases in viable and dead cells respectively. Fluorescence values were transformed to dead and live cell readout values using a standard curve obtained using a standardized range of cell densities for the live cell signal and digitonin-induced cell death for the dead cell signal. Readout values were normalized to values obtained from untreated cells. Results were reported as a cell viability index calculated as the ratio of live cell to dead cell readouts.

Measurement of Zn concentration in the media

Inductively coupled plasma mass spectrometry (ICP-MS) was used to measure Zn content in the culture media. The media were diluted with ultra-pure laboratory grade water (Millipore, Billerica, MA, USA), and an internal standard mixture of Sc, Tb, Y, In, and Bi (CPI International, Santa Rosa, CA, USA) was added to this solution. Standards between 0 and 90 ppb were made using a mixed element solution (CPI International). 2% v/v trace metal grade nitric acid (Fisher Scientific, Pittsburgh, PA, USA) was added to the standards and samples. All measurements were performed on an XSeries II ICP-MS instrument (Thermo Scientific, Waltham, MA, USA).

Statistical analysis

Results are reported as means $\pm$ S.E.M. unless otherwise stated. The 'n' represents the number of independent experiments performed. Changes in parameters normalized to the control condition were tested for statistical significance using a two-sided *t*-test with unequal variance to account for the normalization to the control. In all other experiments comparing the effects of various treatments on physiological parameters not normalized to the control, groups were compared directly to the control condition

using a two-sided *t*-test with equal variance. *P* values ≤ 0.05 and 0.005 were considered statistically significant and highly significant respectively (STATA-IC V.10.1, College Station, TX, USA).

Results

Cytokine-induced alterations in expression of Zn transporters

Initial studies compared the expression profile of members of the ZnT and ZIP families in MIN6 cells and primary murine islets. As shown in Fig. 1, mRNAs encoding 14 members of the ZIP family and nine members of the ZnT family are expressed in murine islets. The expression profile of the ZnTs and ZIPs in MIN6 cells was comparable to that in islets with the exception of a markedly lower expression of ZnT2, ZnT3, ZIP4, and ZIP5.

Having established comparable expression profiles of Zn transporters in islets and MIN6 cells, MIN6 cells were used as a model to examine the impact of cytokines on the level of mRNAs encoding the different Zn transporters. Treatment of MIN6 cells with either 5 ng/ml IL1 β or 10 ng/ml TNF α for 24 h altered the expression of several Zn transporters (Fig. 2). The transporters with the most marked change in expression were ZnT8 and ZIP4. The level of ZnT8 mRNA was decreased by 54.6 ± 6.9 and $44.7 \pm 5.4\%$ by IL1 β and TNF α respectively (*P* = 0.004 and 0.004, Fig. 2C). The level of ZIP4 mRNA was increased 14 ± 4.2 -fold after IL1 β treatment

Figure 1 Relative level of mRNAs encoding ZIP and ZnT transporters in MIN6 cells and murine islets. mRNA was extracted from untreated murine islets and MIN6 cells, and real-time PCR was performed. mRNA levels were normalized to both β -actin and 18S. Values represent the relative level of ZIP or ZnT mRNA normalized to the level of both β -actin and 18S mRNA, and are the mean $\pm$ S.E.M. of the results of three independent experiment.

Figure 2 Effect of cytokines on ZIP and ZnT mRNA levels in MIN6 cells. MIN6 cells were treated for 24 h with 5 ng/ml IL1 β or 10 ng/ml TNF α , and the level of mRNAs encoding ZIPs (A and B) and ZnTs (C) was determined. Values represent the relative level of mRNA compared with the level in control cells, which was defined as 1.0 and represent the mean $\pm$ S.E.M. (n = 4–5 for the different transporters). * P < 0.05 compared with control; ** P < 0.005 compared with control.

(P < 0.05, Fig. 2B). There was also a trend, albeit not significant, toward increased expression of ZIP4 mRNA following treatment with TNF α (4.1 ± 1.2 -fold, P = 0.089, Fig. 2B). Treatment for 48 h with a combination of 5 ng/ml IL1 β and 10 ng/ml TNF α decreased ZnT8 mRNA by $50 \pm 14\%$ (P = 0.024, Fig. 3C), while ZIP4 mRNA levels showed a clear but statistically non-significant trend toward increased expression (21 ± 7.6 -fold, P = 0.057, Fig. 3D). In time course experiments, IL1 β treatment decreased ZnT8 mRNA levels by $51 \pm 4.2\%$, $54.6 \pm 6.9\%$, and $64.1 \pm 0.6\%$ following treatment for 6, 24, and 48 h respectively (Fig. 3A). Treatment with 10 ng/ml TNF α for 6 h had no effect on ZnT8 mRNA levels but decreased ZnT8 mRNA levels by 44.7 ± 5.4 and $50 \pm 5.2\%$ after treatment for 24 and 48 h respectively (Fig. 3A). IL1 β treatment for 6, 24, and 48 h increased ZIP4 mRNA levels by 16 ± 1.2 -, 14 ± 4.2 -, and 47 ± 4.1 -fold respectively (Fig. 3B). Treatment with TNF α for 6, 24, and 48 h stimulated a nonsignificant trend toward increased ZIP4 mRNA levels with increases of 6.8 ± 1.2 -, 4.1 ± 1.2 -, and 13 ± 3.5 -fold respectively (Fig. 3B).

Subsequent experiments determined whether cytokines had a similar effect on ZnT8 and ZIP4 mRNA levels in murine islets. Treatment of islets with 5 ng/ml IL1 β for 24 h decreased ZnT8 mRNA levels by $81 \pm 7\%$, while treatment for 24 h with 10 ng/ml TNF α decreased ZnT8 mRNA levels by $68 \pm 11\%$ (P = 0.0012 and 0.009, Fig. 4A). Treatment of islets with 5 ng/ml IL1 β for 24 h showed a trend toward increased ZIP4 mRNA levels (3.81 ± 0.75 -fold, P = 0.064, Fig. 4B). Treatment with 10 ng/ml TNF α for 24 h resulted in a similar nonsignificant trend toward increased ZIP4 mRNA levels (2.62 ± 0.50 -fold, P = 0.083, Fig. 4B).

β -Cell labile Zn distribution and cytokine-induced changes of labile Zn

Laser scanning confocal imaging of live MIN6 cells following incubation with FluoZin-3 AM showed a high concentration of labile Zn (unbound or weakly bound intracellular Zn), which appeared to be distributed primarily in secretory vesicles (Fig. 5A). Given the changes in Zn transporter expression induced by cytokine treatment, we examined the changes in labile, FluoZin-3 AM-detectable Zn in β -cells after cytokine treatment which, based upon the live cell imaging, represents primarily Zn present in secretory vesicles. In order to avoid interference by other cell types contained in islets, we used the pancreatic β -cell line MIN6. After 24 h treatment with 5 ng/ml IL1 β , the average Zn-related FluoZin-3 AM fluorescence decreased by $22.2 \pm 8.8\%$ compared with control cells (P = 0.0012, Fig. 5B).

Effect of an acute decrease in ZnT8 expression on GSIS

Given the high baseline expression of ZnT8, its localization to insulin secretory granules, and the marked cytokine-induced changes in ZnT8 mRNA levels, we investigated the effect of an acute downregulation of ZnT8 mRNA levels in β -cells using siRNA. Initial studies demonstrated the ability of ZnT8 siRNA to decrease the level of ZnT8 mRNA. Following transfection of MIN6 or primary dispersed murine islet cells with ZnT8 siRNA, the level of ZnT8 mRNA was 23 ± 1.4 and $54.4 \pm 4.4\%$ (mean $\pm$ S.E.M.) of the level in cells

Figure 3 (A and B) Time course of effect of cytokines on ZnT8 and ZIP4 mRNA levels. The level of ZnT8 (A) and ZIP4 (B) mRNA was determined in MIN6 cells after treatment with 5 ng/ml IL1 β or 10 ng/ml TNF α for 6 (n = 2), 24 (n = 4), or 48 (n = 2) h. Values represent the relative level of ZnT8 and ZIP4 mRNA compared with the level in control cells, and are the mean $\pm$ S.E.M. (C and D) Effect of a combination of cytokines on ZnT8 and ZIP4 mRNA levels. The level of ZnT8 (C) and ZIP4 (D) mRNA was determined in MIN6 cells after treatment with a combination of 5 ng/ml IL1 β and 10 ng/ml TNF α for 48 h. Values represent the relative level of ZnT8 and ZIP4 mRNA compared with the level in control cells, and are the mean $\pm$ S.E.M. (n = 5). * P < 0.05.

Figure 4 Effect of cytokines on ZnT8 and ZIP4 mRNA levels in intact murine islets. ZnT8 (A) and ZIP4 (B) mRNA levels were determined in intact murine islets after treatment for 24 h with 5 ng/ml IL1 β or 10 ng/ml TNF α . Values are the mean level of ZnT8 and ZIP4 mRNA compared with the level in control cells, which was defined as 1.0 and are the mean $\pm$ s.e.m. ($n=4$ for ZnT8, $n=3$ for ZIP4). * $P<0.05$, ** $P<0.005$ compared with control.

transfected with control siRNA respectively (Fig. 6A). A decrease in ZnT8 protein level in dispersed primary murine islets following treatment with ZnT8 siRNA was verified by western blotting (Fig. 6B).

Having established that ZnT8 siRNA was able to decrease ZnT8 expression, GSIS in dispersed murine islet cells following transfection with ZnT8 or control siRNA was determined. The relative ratio of insulin concentration in medium following incubation in 16.7 compared with 5.6 mM glucose was 2.0 ± 0.25 in cells treated with control siRNA (Fig. 7A), while in islet cells treated with ZnT8 siRNA, the ratio was significantly decreased to 1.2 ± 0.21 ($P<0.05$ compared with the ratio in cells transfected with control siRNA). In contrast to the effect on GSIS, transfection with ZnT8 siRNA did not affect total cellular insulin content. Total cellular insulin content in islet cell lysates was 17.1 ± 1.5 ng insulin/ μ g protein following transfection with ZnT8 siRNA compared with 15.3 ± 1.6 ng insulin/ μ g protein following transfection with control siRNA ($P=0.41$, Fig. 7B).

Effects of decreased ZnT8 mRNA levels on labile Zn

We next examined whether acute downregulation of ZnT8 expression altered Zn homeostasis similar to treatment with IL1 β . For these studies, MIN6 cells were transfected with either ZnT8 or control siRNA, and labile Zn was determined using flow cytometry after staining with the Zn-specific fluorescent probe FluoZin-3 AM and the live-cell dye CellTrace calcein red-orange AM. The relative mean labile Zn-related fluorescence from live cells decreased by $26.5 \pm 17.6\%$ following transfection with ZnT8 siRNA compared with control siRNA ($P<0.05$, Fig. 8A).

Effect of decreased ZnT8 mRNA levels on islet cell viability in various Zn concentrations

Subsequent studies examined the impact of decreased ZnT8 expression and the resulting changes in Zn homeostasis on islet cell viability. Initial studies using ICP-MS demonstrated that the concentration of Zn in the culture medium was 5 μ M. Following transfection of dispersed murine islet cells with

Figure 5 (A and B) The distribution of labile Zn in single untreated live MIN6 cells. Labile Zn was imaged following incubation in FluoZin-3 AM in phenol red-free DMEM for 30 min using confocal spinning disc microscopy (100 $\times$ magnification, A, FluoZin-3 AM fluorescence alone; B, phase contrast + fluorescence overlay). (C) Effect of IL1 β on labile Zn in MIN6 cells. Cells were analyzed using flow cytometry after staining with the Zn-specific fluorescent probe FluoZin-3 AM and live-cell dye CellTrace calcein red-orange, AM. FluoZin-3 AM fluorescence was analyzed in live cells only. In total, 20 000–30 000 live cell events were recorded in each replicate. The change in FluoZin-3 AM detectable Zn was determined in MIN6 cells after treatment or not with 5 ng/ml IL1 β for 24 h. Values represent the relative FluoZin-3 AM-related fluorescence in live cells normalized to the control condition, and are the mean $\pm$ s.e.m. ($n=3$, each in triplicate). ** $P<0.005$ compared with control.

Figure 6 Effect of ZnT8 siRNA on ZnT8 expression. (A) ZnT8 mRNA levels were determined in MIN6 and dispersed murine islet cells following transfection with either ZnT8 or control siRNA. The values represent the relative level of ZnT8 mRNA in cells treated with ZnT8 siRNA compared with cells treated with control siRNA, which was defined as 1.0, and are the mean $\pm$ S.E.M. ($n=3$ for MIN6 cells, $n=4$ for islet cells). The level of ZnT8 mRNA was normalized to the level of β -actin mRNA. ** $P<0.005$ compared with control siRNA. (B) Western blot analysis comparing ZnT8 in dispersed murine islet cells treated with ZnT8 siRNA and control siRNA. The autoradiograms are representative of the results of three and five independent experiments for MIN6 and dispersed islets respectively.

either ZnT8 or control siRNA, cell viability was determined using the Multitox assay. The viability indices, which are proportional to the ratio of live to dead cells each normalized to the values of control cells with no added Zn, were 1.0002 ± 0.0004 and 0.9937 ± 0.0070 following transfection with control and ZnT8 siRNAs respectively ($P=0.386$, $n=4$, each performed in duplicate or triplicate, Fig. 8B). Supplementing the medium with ZnCl_2 to final Zn concentrations of 19, 34, 48, 62, 76, 91, or 105 μM enhanced cell viability at all concentrations. The viability index increased to 1.5803 ± 0.0608 and 1.5493 ± 0.1017 at 105 μM Zn in cells transfected with control and ZnT8 siRNAs respectively ($P=0.802$ for the knockdown versus control comparison, $n=4$, each performed in duplicate or triplicate).

Discussion

Impaired function of human and rodent β -cells as well as MIN6 cells after exposure to $\text{IL1}\beta$ or $\text{TNF}\alpha$ is well established (Donath *et al.* 2008). Several potential mechanisms have been suggested for this effect (Xenos *et al.* 1992, 1993, 1994, Corbett & McDaniel 1995, Rabinovitch & Suarez-Pinzon

1998, Wu *et al.* 2001, Donath *et al.* 2008). These include activation of the mitogen-activated protein kinases, c-jun N-terminal kinase, p38, and Extracellular Signal-Regulated-Kinase (ERK) with subsequent activation of the nuclear factor kappa-light-chain-enhancer of activated β -cells (NF- κB) pathway and altered gene expression (Donath *et al.* 2008). In this study, we examined whether cytokines affect Zn transporter expression as a possible contributing factor to cytokine-induced β -cell dysfunction. We demonstrated that treatment of murine islets or MIN6 cells with $\text{IL1}\beta$ or $\text{TNF}\alpha$ alters the expression profile of Zn transporters of the ZnT and ZIP families. One of the observed changes which may be of particular significance is the cytokine-induced decrease in ZnT8 expression. ZnT8 has a high baseline expression level in β -cells, and plays an important role in β -cell function. Moreover, we observed similar alterations in cellular labile

Figure 7 Glucose-stimulated insulin secretion and cellular insulin content in dispersed murine islet cells following transfection with ZnT8 siRNA. (A) 72 h after transfection of dispersed murine islet cells with either ZnT8 or control siRNA, the insulin concentration in the medium was determined following incubation in 5.6 and 16.7 mM glucose. Values represent the ratio of the insulin concentration in medium after incubation for 30 min in 16.7 mM glucose compared with 5.6 mM glucose, and are the mean $\pm$ S.E.M. ($n=4$). * $P<0.05$ compared with cells treated with control siRNA. (B) Total cellular content of insulin in dispersed murine islet cells 72 h after transfection with either ZnT8 or control siRNA. Values represent the total insulin content in cell lysates which was determined as described in the Materials and Methods, and are the mean $\pm$ S.E.M. ($n=3$).

Figure 8 Effect of decreased ZnT8 expression on MIN6 cell labile Zn content and dispersed islet cell viability. (A) MIN6 cells were analyzed using flow cytometry after staining with the Zn-specific fluorescent probe FluoZin-3 AM and live-cell dye CellTrace calcein red-orange. FluoZin-3 AM fluorescence was analyzed in live cells only. In total, 20 000–30 000 live cell events were recorded in each replicate. FluoZin-3-chelatable Zn in MIN6 cells transfected with control siRNA or ZnT8 siRNA was then determined. Values represent the relative FluoZin-3-related fluorescence in live cells normalized to the control condition, and are the mean $\pm$ s.e.m. ($n=3$, each in duplicate or triplicate), * $P<0.05$. (B) Dispersed islet cells were transfected with either ZnT8 or control siRNA, and then incubated in the presence of increasing concentrations of Zn. Cell viability was determined using the Multitox assay. Changes in the cell viability index correspond to changes in the ratio of the live to dead cell readouts. Values represent the mean cell viability index, and are the mean $\pm$ s.d. of the results ($n=4$, each in duplicate or triplicate).

Zn content following IL1 β exposure and ZnT8 knockdown. This finding and the decrease in GSIS observed following ZnT8 knockdown are consistent with alterations in Zn homeostasis due to decreased ZnT8 expression contributing to cytokine-induced impairment of β -cell function.

Our finding of a cytokine-induced decrease in ZnT8 expression generally agrees with a recent report by Egefjord *et al.* (2009) which described an IL1 β -induced decrease in ZnT8 mRNA levels in INS-1 cells, a rat β -cell line, as well as in rat islets. In contrast to their findings, we did not observe

reduced ZnT3, ZnT6, and ZIP6 mRNA levels; however, we do find a significant cytokine-induced increase in ZIP4 mRNA levels. It is likely that these differences relate to differences in the models used.

Zn is critical for the function of mammalian cells. The vast majority of the intracellular Zn is bound to proteins or is accrued in specialized compartments (Suhly *et al.* 1999, Hambidge 2000, Colvin *et al.* 2003, Lemire *et al.* 2007). A vanishingly small amount of free Zn is typically available in the cytosol of most cells (Suhly *et al.* 1999, Outten & O'Halloran 2001, Bozym *et al.* 2006). In most cells, Zn is important for the structural integrity and function of transcription factors in the Zn finger family of proteins and for maintaining proper function of Zn-dependent enzymes that protect cells against oxidative stress and apoptosis (Truong-Tran *et al.* 2001, Prasad *et al.* 2004). The cellular content of Zn in β -cells is markedly higher than most mammalian cells, which is likely related to its unique roles in β -cells (Chausmer 1998, Dodson & Steiner 1998, Chimienti *et al.* 2005, Cousins *et al.* 2006). Estimates suggest that $\sim 50\%$ of Zn in β -cells is present within secretory vesicles (Zalewski *et al.* 1994), where it has been postulated to play a role in the packaging of proinsulin and insulin as hexamers. In this model, Zn enables storage of insulin at high concentrations and facilitates adequate insulin release in response to a glucose load (Huber & Gershoff 1973, Figlewicz *et al.* 1984, Huang & Arvan 1995, Dodson & Steiner 1998, Chimienti *et al.* 2005, 2006).

ZnT8 is highly expressed in β -cells (Chimienti *et al.* 2005, 2006), and localizes mainly to membranes of secretory vesicles, suggesting that it plays an important role in Zn transport into β -cell secretory vesicles (Chimienti *et al.* 2006, Mocchegiani *et al.* 2008). As noted earlier, overexpression of ZnT8 enhanced the insulin secretory capacity of a β -cell line (Chimienti *et al.* 2006). On the other hand, an R325W mutation in *SLC30A8* is associated with type 2 diabetes as well as decreased first phase insulin secretion in non-diabetic subjects bearing at least one copy of the risk allele (Saxena *et al.* 2007, Scott *et al.* 2007, Sladek *et al.* 2007, Zeggini *et al.* 2007, Boesgaard *et al.* 2008, Hertel *et al.* 2008, van Hoek *et al.* 2008, Horikawa *et al.* 2008, Omori *et al.* 2008, Wu *et al.* 2008, Xiang *et al.* 2008). Complete loss of ZnT8 expression in mice homozygous for a null mutation of *SLC30A8* led to decreased Zn accumulation in islets (Lemaire *et al.* 2009, Nicolson *et al.* 2009, Pound *et al.* 2009). ZnT8 null mice also exhibited decreased GSIS and altered glucose homeostasis (Pound *et al.* 2009). However, ZnT8 null mice exhibited altered glucose tolerance as well as *in vivo* and *in vitro* disturbances in GSIS only under certain age, gender, and dietary conditions (Lemaire *et al.* 2009, Nicolson *et al.* 2009). Furthermore, an unexpected increase in weight gain was observed in ZnT8 null mice fed a high fat diet (Nicolson *et al.* 2009). Given these observations, secondary effects during various developmental stages as well as environmental interactions are likely to have influenced the findings in this global knockout model. In the present study, we have now demonstrated that an acute decrease in ZnT8 expression in

dispersed murine islet cells decreases GSIS in cells without altering the total cellular insulin content. In contrast to the global ZnT8 knockout mouse model, our model demonstrates the impact of an isolated, acute decrease in ZnT8 function prior to the induction of compensatory changes during development. This approach furthers our understanding of β -cell physiology by examining the impact of decreased ZnT8 expression independent of potential developmental changes or secondary compensatory mechanisms present in the ZnT8 knockout model.

ZnT8 knockout models demonstrated decreased GSIS under certain age and dietary conditions without changes in total cellular insulin content (Lemaire *et al.* 2009, Nicolson *et al.* 2009, Pound *et al.* 2009). This similarity between the effects of acutely decreased ZnT8 expression and chronic, complete loss of ZnT8 expression in the knockout mice indicates that β -cells have a limited capacity to compensate for changes in Zn homeostasis. Our findings and those in ZnT8 knockout mice in which decreased or absent ZnT8 expression did not impact total cellular insulin content differ from data obtained using the INS-1 β -cell line after an siRNA-mediated decrease in ZnT8 expression (Fu *et al.* 2009). In that model, both the number of insulin-containing secretory vesicles and total cellular insulin content were decreased after ZnT8 knockdown. These differences may be due to differences in the physiology of primary islet cells and INS cells.

The effect of decreased ZnT8 expression on GSIS in murine islet cells could be due to one or more possible mechanisms. Decreased ZnT8 expression likely results in reduced transport of Zn into secretory vesicles, which would impair the packaging of insulin as hexamers around Zn cores and thereby alter insulin secretion (Huang & Arvan 1995, Dodson & Steiner 1998). Another possible mechanism would be a toxic effect of Zn accumulation in the cytoplasm exceeding the metal-buffering capacity of cytoplasmic proteins (Frederickson *et al.* 1989, Koh *et al.* 1996, Kim *et al.* 1999a,b, Lobner *et al.* 2000, Frederickson *et al.* 2005, Lecane *et al.* 2005, Lemire *et al.* 2007, Oyama *et al.* 2007, Park *et al.* 2007, Zhang *et al.* 2007, Matsui *et al.* 2008). To examine this, we studied the changes in labile Zn with decreased ZnT8 expression in MIN6 cells using the fluorescent Zn probe FluoZin-3 AM. This approach has been previously validated for Zn in MIN6 cells (Gee *et al.* 2002). A significant decrease in intracellular FluoZin-3 AM-detectable Zn, which correlated with decreased ZnT8 expression, was observed. FluoZin-3 AM is reported to have a moderate binding affinity for Zn (K_d of 15 nM per the manufacturer) which is a significantly lower affinity for Zn than that of typical Zn-binding enzymes such as carbonic anhydrase (Bozym *et al.* 2006). Thus, Zn detected by FluoZin-3 AM represents the fraction of cellular Zn that is loosely bound to assorted intracellular ligands that have a lower affinity for Zn. Most cytoplasmic Zn is reported to be bound to metallothioneins, which have a much higher affinity for Zn than FluoZin-3 AM (Jacob *et al.* 1998). Therefore, it is unlikely that the

FluoZin-3 AM signal reflects cytosolic Zn unless labile Zn concentrations exceed the buffering capacity of the abundant metallothioneins present in the cytoplasm of β -cells (Laychock *et al.* 2000, Li *et al.* 2004). FluoZin-3 AM is known to penetrate intracellular vesicles, and, consistent with our imaging of FluoZin-3 AM loaded live β -cells, it is likely that the majority of the labile Zn detected using FluoZin-3 AM is present within secretory vesicles. Taken together, our results suggest that acute downregulation of ZnT8 reduces the capacity of β -cells to accumulate Zn in β -cell insulin secretory vesicles.

The present studies also demonstrated that decreased ZnT8 expression did not have a significant impact on islet cell viability at the physiological Zn concentration of 5 $\mu\text{mol/l}$ or in the presence of supraphysiological concentrations of Zn (up to 105 $\mu\text{mol/l}$) in the growth medium (Isbir *et al.* 1994, Sampson *et al.* 1997, Chausmer 1998, Maret & Sandstead 2006). In contrast to other cell types, β -cells have been shown to tolerate supraphysiological ambient Zn concentrations <200 $\mu\text{mol/l}$ (Huber & Gershoff 1973, Figlewicz *et al.* 1981, Frederickson *et al.* 1989, Koh *et al.* 1996, Kim *et al.* 1999a,b, 2000, Lobner *et al.* 2000, Chang *et al.* 2003, Frederickson *et al.* 2005, Lecane *et al.* 2005, Lemire *et al.* 2007, Oyama *et al.* 2007, Park *et al.* 2007, Zhang *et al.* 2007, Matsui *et al.* 2008). The absence of toxic effects following ZnT8 knockdown in cells exposed to 105 $\mu\text{mol/l}$ Zn in our study suggests that impaired GSIS with decreased ZnT8 expression is not likely due to toxic effects secondary to free Zn accumulation in the cytoplasm.

Given the low level of ZIP4 mRNA, the physiologic relevance of the increase in ZIP4 expression following exposure to IL1 β is uncertain. ZIP4 regulates Zn influx into the cytoplasm of various mammalian cell types. In intestinal enterocytes, ZIP4 is located in the apical cell membrane and facilitates the uptake of sufficient quantities of dietary Zn. ZIP4 expression is reported to increase in Zn-deficient intestinal cells through posttranscriptional mRNA stabilization as well as decreased protein degradation (Dufner-Beattie *et al.* 2003, Weaver *et al.* 2007, Andrews 2008). Little is known about the role of ZIP4 in the pancreas, although ZIP4 was reported to be aberrantly expressed in pancreatic cancer (Li *et al.* 2007). Based upon the real-time PCR assay, the expression of ZIP4 mRNA was relatively low compared with other Zn transporters but increased following cytokine exposure. It is possible that this represents a compensatory response to the IL1 β -induced decrease in labile Zn (Fig. 5B). Future studies will be needed to address that issue; specifically, the level of ZIP4 protein and changes in response to cytokine exposure and/or ZnT8 knockdown will need to be examined.

This study highlights the importance of the Zn transporter ZnT8 in pancreatic β -cell physiology. Beyond demonstrating a critical role for ZnT8 in GSIS, our findings suggest that cytokine-mediated β -cell dysfunction may occur, in part, through disruption of Zn metabolism via decreased ZnT8 expression.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1677/JOE-09-0420>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was funded by a Northwestern Memorial Foundation MD-Scientist Fellowship in Genetic Medicine award for ME. The study was also funded by two NIH grants: R37GM038784 (TVO) and U42RR023245 (TVO) and T32 DK007169-27 NRSA Institutional Training Grant (LKB). Imaging was performed at the Northwestern University Cell Imaging Facility generously supported by CCSG P30 CA060553 awarded to the Robert H Lurie Comprehensive Cancer Center. ICP-MS Metal analysis was performed at the Northwestern University Quantitative Bioelemental Imaging Center generously supported by NASA Ames Research Center NNA04CC36G.

References

- Andrews GK 2008 Regulation and function of Zip4, the acrodermatitis enteropathica gene. *Biochemical Society Transactions* **36** 1242–1246.
- Blomeier H, Zhang X, Rives C, Brissova M, Hughes E, Baker M, Powers AC, Kaufman DB, Shea LD & Lowe WL Jr 2006 Polymer scaffolds as synthetic microenvironments for extrahepatic islet transplantation. *Transplantation* **82** 452–459.
- Boesgaard TW, Zilinskaite J, Vanttinen M, Laakso M, Jansson PA, Hammarstedt A, Smith U, Stefan N, Fritsche A, Haring H *et al.* 2008 The common SLC30A8 Arg325Trp variant is associated with reduced first-phase insulin release in 846 non-diabetic offspring of type 2 diabetes patients – the EUGENE2 study. *Diabetologia* **51** 816–820.
- Bozym RA, Thompson RB, Stoddard AK & Fierke CA 2006 Measuring picomolar intracellular exchangeable zinc in PC-12 cells using a ratiometric fluorescence biosensor. *ACS Chemical Biology* **1** 103–111.
- Chang I, Cho N, Koh JY & Lee MS 2003 Pyruvate inhibits zinc-mediated pancreatic islet cell death and diabetes. *Diabetologia* **46** 1220–1227.
- Chausmer AB 1998 Zinc, insulin and diabetes. *Journal of the American College of Nutrition* **17** 109–115.
- Chimienti F, Favier A & Seve M 2005 ZnT-8, a pancreatic β -cell-specific zinc transporter. *Biomaterials* **18** 313–317.
- Chimienti F, Devergnas S, Pattou F, Schuit F, Garcia-Cuenca R, Vandewalle B, Kerr-Conte J, Van Lommel L, Grunwald D, Favier A *et al.* 2006 *In vivo* expression and functional characterization of the zinc transporter ZnT8 in glucose-induced insulin secretion. *Journal of Cell Science* **119** 4199–4206.
- Chin-Chance CV, Newman MV, Aronovitz A, Blomeier H, Kruger J, Lee EJ & Lowe WL Jr 2006 Role of the mitogen-activated protein kinases in cytokine-mediated inhibition of insulin gene expression. *Journal of Investigative Medicine* **54** 132–142.
- Colvin RA, Fontaine CP, Laskowski M & Thomas D 2003 Zn²⁺ transporters and Zn²⁺ homeostasis in neurons. *European Journal of Pharmacology* **479** 171–185.
- Corbett JA & McDaniel ML 1995 Intra-islet release of interleukin 1 inhibits β cell function by inducing β cell expression of inducible nitric oxide synthase. *Journal of Experimental Medicine* **181** 559–568.
- Cousins RJ, Liuzzi JP & Lichten LA 2006 Mammalian zinc transport, trafficking, and signals. *Journal of Biological Chemistry* **281** 24085–24089.
- Dodson G & Steiner D 1998 The role of assembly in insulin's biosynthesis. *Current Opinion in Structural Biology* **8** 189–194.
- Donath MY, Storling J, Berchtold LA, Billestrup N & Mandrup-Poulsen T 2008 Cytokines and β -cell biology: from concept to clinical translation. *Endocrine Reviews* **29** 334–350.
- Dufner-Beattie J, Wang F, Kuo YM, Gitschier J, Eide D & Andrews GK 2003 The acrodermatitis enteropathica gene ZIP4 encodes a tissue-specific, zinc-regulated zinc transporter in mice. *Journal of Biological Chemistry* **278** 33474–33481.
- Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, Jackson AU, Wheeler E, Glazer NL, Bouatia-Naji N, Gloyn AL *et al.* 2010 New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nature Genetics* **42** 105–116.
- Egefjord L, Jensen JL, Bang-Berthelsen CH, Petersen AB, Smidt K, Schmitz O, Karlsen AE, Pociot F, Chimienti F, Rungby J *et al.* 2009 Zinc transporter gene expression is regulated by pro-inflammatory cytokines: a potential role for zinc transporters in β -cell apoptosis? *BMC Endocrine Disorder* **9** 7.
- Figlewicz DP, Heldt A, Forhan SE & Grodsky GM 1981 Effect of exogenous zinc on insulin secretion *in vitro*. *Endocrinology* **108** 730–732.
- Figlewicz DP, Forhan SE, Hodgson AT & Grodsky GM 1984 65Zinc and endogenous zinc content and distribution in islets in relationship to insulin content. *Endocrinology* **115** 877–881.
- Frederickson CJ, Hernandez MD & McGinty JF 1989 Translocation of zinc may contribute to seizure-induced death of neurons. *Brain Research* **480** 317–321.
- Frederickson CJ, Koh JY & Bush AI 2005 The neurobiology of zinc in health and disease. *Nature Reviews. Neuroscience* **6** 449–462.
- Fu Y, Tian W, Pratt EB, Dirling LB, Shyng SL, Meshul CK & Cohen DM 2009 Down-regulation of ZnT8 expression in INS-1 rat pancreatic β cells reduces insulin content and glucose-inducible insulin secretion. *PLoS ONE* **4** e5679.
- Gee KR, Zhou ZL, Qian WJ & Kennedy R 2002 Detection and imaging of zinc secretion from pancreatic β -cells using a new fluorescent zinc indicator. *Journal of the American Chemical Society* **124** 776–778.
- Hambidge M 2000 Human zinc deficiency. *Journal of Nutrition* **130** 1344S–1349S.
- Hertel JK, Johansson S, Raeder H, Midthjell K, Lyssenko V, Groop L, Molven A & Njolstad PR 2008 Genetic analysis of recently identified type 2 diabetes loci in 1,638 unselected patients with type 2 diabetes and 1,858 control participants from a Norwegian population-based cohort (the HUNT study). *Diabetologia* **51** 971–977.
- van Hoek M, Dehgan A, Witteman JC, van Duijn CM, Uitterlinden AG, Oostra BA, Hofman A, Sijbrands EJ & Janssens AC 2008 Predicting type 2 diabetes based on polymorphisms from genome wide association studies: a population-based study. *Diabetes* **57** 3122–3128.
- Horikawa Y, Miyake K, Yasuda K, Enya M, Hirota Y, Yamagata K, Hinokio Y, Oka Y, Iwasaki N, Iwamoto Y *et al.* 2008 Replication of genome-wide association studies of type 2 diabetes susceptibility in Japan. *Journal of Clinical Endocrinology and Metabolism* **93** 3136–3141.
- Huang XF & Arvan P 1995 Intracellular transport of proinsulin in pancreatic β -cells. Structural maturation probed by disulfide accessibility. *Journal of Biological Chemistry* **270** 20417–20423.
- Huber AM & Gershoff SN 1973 Effect of zinc deficiency in rats on insulin release from the pancreas. *Journal of Nutrition* **103** 1739–1744.
- Isbir T, Tamer L, Taylor A & Isbir M 1994 Zinc, copper and magnesium status in insulin-dependent diabetes. *Diabetes Research* **26** 41–45.
- Jacob C, Maret W & Vallee BL 1998 Control of zinc transfer between thionein, metallothionein, and zinc proteins. *PNAS* **95** 3489–3494.
- Kim EY, Koh JY, Kim YH, Sohn S, Joe E & Gwag BJ 1999a Zn²⁺ entry produces oxidative neuronal necrosis in cortical cell cultures. *European Journal of Neuroscience* **11** 327–334.
- Kim YH, Kim EY, Gwag BJ, Sohn S & Koh JY 1999b Zinc-induced cortical neuronal death with features of apoptosis and necrosis: mediation by free radicals. *Neuroscience* **89** 175–182.
- Kim BJ, Kim YH, Kim S, Kim JW, Koh JY, Oh SH, Lee MK, Kim KW & Lee MS 2000 Zinc as a paracrine effector in pancreatic islet cell death. *Diabetes* **49** 367–372.
- Koh JY, Suh SW, Gwag BJ, He YY, Hsu CY & Choi DW 1996 The role of zinc in selective neuronal death after transient global cerebral ischemia. *Science* **272** 1013–1016.
- Laychock SG, Duzen J & Simpkins CO 2000 Metallothionein induction in islets of Langerhans and insulinoma cells. *Molecular and Cellular Endocrinology* **165** 179–187.

- Lecane PS, Karaman MW, Sirisawad M, Naumovski L, Miller RA, Hacia JG & Magda D 2005 Motexafin gadolinium and zinc induce oxidative stress responses and apoptosis in B-cell lymphoma lines. *Cancer Research* **65** 11676–11688.
- Lemaire K, Ravier MA, Schraenen A, Creemers JW, Van de Plas R, Granvik M, Van Lommel L, Waelkens E, Chimienti F, Rutter GA *et al.* 2009 Insulin crystallization depends on zinc transporter ZnT8 expression, but is not required for normal glucose homeostasis in mice. *PNAS* **106** 14872–14877.
- Lemire J, Mailloux R & Appanna VD 2007 Zinc toxicity alters mitochondrial metabolism and leads to decreased ATP production in hepatocytes. *Journal of Applied Toxicology* **28** 175–182.
- Li X, Chen H & Epstein PN 2004 Metallothionein protects islets from hypoxia and extends islet graft survival by scavenging most kinds of reactive oxygen species. *Journal of Biological Chemistry* **279** 765–771.
- Li M, Zhang Y, Liu Z, Bharadwaj U, Wang H, Wang X, Zhang S, Liuzzi JP, Chang SM, Cousins RJ *et al.* 2007 Aberrant expression of zinc transporter ZIP4 (SLC39A4) significantly contributes to human pancreatic cancer pathogenesis and progression. *PNAS* **104** 18636–18641.
- Liu W, Liu Y & Lowe WL Jr 2001 The role of phosphatidylinositol 3-kinase and the mitogen-activated protein kinases in insulin-like growth factor-I-mediated effects in vascular endothelial cells. *Endocrinology* **142** 1710–1719.
- Livak KJ & Schmittgen TD 2001 Analysis of relative gene expression data using real-time quantitative PCR and the $2(-\Delta\Delta C(T))$ method. *Methods* **25** 402–408.
- Loebner D, Canzoniero LM, Manzerra P, Gotttron F, Ying H, Knudson M, Tian M, Dugan LL, Kerchner GA, Sheline CT *et al.* 2000 Zinc-induced neuronal death in cortical neurons. *Cellular and Molecular Biology* **46** 797–806.
- Maret W & Sandstead HH 2006 Zinc requirements and the risks and benefits of zinc supplementation. *Journal of Trace Elements in Medicine and Biology* **20** 3–18.
- Matsui H, Sakanashi Y, Oyama TM, Oyama Y, Yokota S, Ishida S, Okano Y, Oyama TB & Nishimura Y 2008 Imidazole antifungals, but not triazole antifungals, increase membrane Zn^{2+} permeability in rat thymocytes possible contribution to their cytotoxicity. *Toxicology* **248** 142–150.
- Milne DB, Ralston NV & Wallwork JC 1985 Zinc content of blood cellular components and lymph node and spleen lymphocytes in severely zinc-deficient rats. *Journal of Nutrition* **115** 1073–1078.
- Miyazaki J, Araki K, Yamato E, Ikegami H, Asano T, Shibasaki Y, Oka Y & Yamamura K 1990 Establishment of a pancreatic β cell line that retains glucose-inducible insulin secretion: special reference to expression of glucose transporter isoforms. *Endocrinology* **127** 126–132.
- Mocchegiani E, Giacconi R & Malavolta M 2008 Zinc signalling and subcellular distribution: emerging targets in type 2 diabetes. *Trends in Molecular Medicine* **14** 419–428.
- Nicolson TJ, Bellomo EA, Wijesekara N, Loder MK, Baldwin JM, Gylkhandanyan AV, Koshkin V, Tarasov AI, Carzaniga R, Kronenberger K *et al.* 2009 Insulin storage and glucose homeostasis in mice null for the granule zinc transporter ZnT8 and studies of the type 2 diabetes-associated variants. *Diabetes* **58** 2070–2083.
- Omori S, Tanaka Y, Takahashi A, Hirose H, Kashiwagi A, Kaku K, Kawamori R, Nakamura Y & Maeda S 2008 Association of CDKAL1, IGF2BP2, CDKN2A/B, HHEX, SLC30A8, and KCNJ11 with susceptibility to type 2 diabetes in a Japanese population. *Diabetes* **57** 791–795.
- Outten CE & O'Halloran TV 2001 Femtomolar sensitivity of metallo-regulatory proteins controlling zinc homeostasis. *Science* **292** 2488–2492.
- Oyama Y, Matsui H, Morimoto M, Sakanashi Y, Nishimura Y, Ishida S & Okano Y 2007 Synergic cytotoxic action induced by simultaneous application of zinc and clotrimazole in rat thymocytes. *Toxicology Letters* **171** 138–145.
- Park SE, Park JW, Cho YS, Ryu JH, Paick JS & Chun YS 2007 HIF-1 α promotes survival of prostate cells at a high zinc environment. *Prostate* **67** 1514–1523.
- Pound LD, Sarkar S, Benninger RK, Wang Y, Suwanichkul A, Shadoan MK, Printz RL, Oeser JK, Lee CE, Piston DW *et al.* 2009 Deletion of the mouse Slc30a8 gene encoding zinc transporter-8 results in impaired insulin secretion. *Biochemical Journal* **421** 371–376.
- Prasad AS, Bao B, Beck FW, Kucuk O & Sarkar FH 2004 Antioxidant effect of zinc in humans. *Free Radical Biology & Medicine* **37** 1182–1190.
- Rabinovitch A & Suarez-Pinzon WL 1998 Cytokines and their roles in pancreatic islet β -cell destruction and insulin-dependent diabetes mellitus. *Biochemical Pharmacology* **55** 1139–1149.
- Sampson B, Kovar IZ, Rauscher A, Fairweather-Tait S, Beattie J, McArdle HJ, Ahmed R & Green C 1997 A case of hyperzincemia with functional zinc depletion: a new disorder? *Pediatric Research* **42** 219–225.
- Saxena R, Voight BF, Lyssenko V, Burt NP, de Bakker PI, Chen H, Roix JJ, Kathiresan S, Hirschhorn JN, Daly MJ *et al.* 2007 Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* **316** 1331–1336.
- Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, Erdos MR, Stringham HM, Chines PS, Jackson AU *et al.* 2007 A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* **316** 1341–1345.
- Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin P, Vincent D, Belisle A, Hadjadj S *et al.* 2007 A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* **445** 881–885.
- Suh Y, Simon KD, Linzer DI & O'Halloran TV 1999 Metallothionein is part of a zinc-scavenging mechanism for cell survival under conditions of extreme zinc deprivation. *Journal of Biological Chemistry* **274** 9183–9192.
- Truong-Tran AQ, Carter J, Ruffin RE & Zalewski PD 2001 The role of zinc in caspase activation and apoptotic cell death. *Biometals* **14** 315–330.
- Wallwork JC, Johnson LK, Milne DB & Sandstead HH 1983 The effect of interactions between dietary egg white protein and zinc on body weight, bone growth and tissue trace metals in the 30-day-old rat. *Journal of Nutrition* **113** 1307–1320.
- Ward NI & Mason JA 1987 Neuron activation analysis techniques for identifying elemental status in Alzheimer's disease. *Journal of Radioanalytical and Nuclear Chemistry* **113** 515–526.
- Weaver BP, Dufner-Beattie J, Kambe T & Andrews GK 2007 Novel zinc-responsive post-transcriptional mechanisms reciprocally regulate expression of the mouse Slc39a4 and Slc39a5 zinc transporters (Zip4 and Zip5). *Biological Chemistry* **388** 1301–1312.
- Wu JJ, Chen X, Cao XC, Baker MS & Kaufman DB 2001 Cytokine-induced metabolic dysfunction of MIN6 β cells is nitric oxide independent. *Journal of Surgical Research* **101** 190–195.
- Wu Y, Li H, Loos RJ, Yu Z, Ye X, Chen L, Pan A, Hu FB & Lin X 2008 Common variants in CDKAL1, CDKN2A/B, IGF2BP2, SLC30A8 and HHEX/IDE genes are associated with type 2 diabetes and impaired fasting glucose in a Chinese Han population. *Diabetes* **57** 2834–2842.
- Xenos ES, Farney AC, Widmer MB, Casanova D, Stevens RB, Blazar BR, Sutherland DE & Gores PF 1992 Effect of tumor necrosis factor α and of the soluble tumor necrosis factor receptor on insulin secretion of isolated islets of Langerhans. *Transplantation Proceedings* **24** 2863–2864.
- Xenos ES, Stevens RB, Gores PF, Casanova D, Farney AC, Sutherland DE & Platt JL 1993 IL-1 β -induced inhibition of β -cell function is mediated through nitric oxide. *Transplantation Proceedings* **25** 994.
- Xenos ES, Stevens RB, Sutherland DE, Lokesh A, Ansit JD, Casanova D, Gores PF & Platt JL 1994 The role of nitric oxide in IL-1 β -mediated dysfunction of rodent islets of Langerhans. Implications for the function of intrahepatic islet grafts. *Transplantation* **57** 1208–1212.
- Xiang J, Li XY, Xu M, Hong J, Huang Y, Tan JR, Lu X, Dai M, Yu B & Ning G 2008 Zinc transporter-8 gene (SLC30A8) is associated with type 2 diabetes in Chinese. *Journal of Clinical Endocrinology and Metabolism* **93** 4107–4112.
- Zalewski PD, Millard SH, Forbes IJ, Kapaniris O, Slavotinek A, Betts WH, Ward AD, Lincoln SF & Mahadevan I 1994 Video image analysis of labile zinc in viable pancreatic islet cells using a specific fluorescent probe for zinc. *Journal of Histochemistry and Cytochemistry* **42** 877–884.
- Zeggini E, Weedon MN, Lindgren CM, Frayling TM, Elliott KS, Lango H, Timpson NJ, Perry JR, Rayner NW, Freathy RM *et al.* 2007 Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science* **316** 1336–1341.
- Zhang Y, Aizenman E, Defranco DB & Rosenberg PA 2007 Intracellular zinc release, 12-lipoxygenase activation and MAPK dependent neuronal and oligodendroglial death. *Molecular Medicine* **13** 350–355.

Received in final form 10 May 2010

Accepted 27 May 2010

Made available online as an Accepted Preprint
27 May 2010