REVIEW

Kisspeptin and fertility

Saira Hameed, Channa N Jayasena and Waljit S Dhillon

Division of Diabetes, Endocrinology and Metabolism, Section of Investigative Medicine, Imperial College London, Hammersmith Hospital, 6th Floor
Commonwealth Building, Du Cane Road, London W12 0NN, UK

(Correspondence should be addressed to W S Dhillon; Email: w.dhillon@imperial.ac.uk)

Abstract

The kisspeptins are a family of peptide hormones, which in recent years have been shown to play a critical role in the regulation of the hypothalamic–pituitary–gonadal axis, thus in turn influencing fertility and reproduction. This review examines the physiological role of kisspeptin and the kisspeptin receptor in

the control of gonadotrophin and gonadal steroid hormone secretion and the implications of these findings with respect to fertility. In addition, the potential therapeutic use of kisspeptin in the treatment of reproductive disorders will be examined.

Journal of Endocrinology (2011) **208**, 97–105

Introduction

Infertility affects up to one in six couples in the United Kingdom (Human Fertilisation and Embryology Authority 2009, <http://www.hfea.gov.uk/infertility-facts.html#1248>). The currently available hormone-based treatments for infertility act through the manipulation of the hypothalamic–pituitary–gonadal (HPG) axis at the level of GnRH or below. Although effective, these treatments also have a significant failure rate as well as an associated morbidity. The discovery of kisspeptin in 1996 and the subsequent identification of the kisspeptin receptor (previously known as G-protein-coupled receptor 54, GPR54) have added a new critical dimension to our understanding of the physiology of the HPG axis, reproduction and fertility (Lee *et al.* 1996, Clements *et al.* 2001). Mice and humans lacking kisspeptin receptor expression show a phenotype of hypogonadotrophic hypogonadism and consequent infertility (de Roux *et al.* 2003, Seminara *et al.* 2003). However, kisspeptin receptor knockout mice have normal levels of hypothalamic GnRH expression (Seminara *et al.* 2003) and normal GnRH neuronal morphology (Messenger *et al.* 2005). These findings have been pivotal in the emergence of kisspeptin signalling as critical regulator of normal fertility, and future work may lead to the development of therapeutic use of kisspeptin in the treatment of reproductive disorders.

Kisspeptin structure and distribution

The kisspeptins are a family of RF peptide hormones, so named as the arginine–phenylalanine residues are present at the amino terminal (Arg-Phe-NH₂; Clements *et al.* 2001,

Kotani *et al.* 2001). The kisspeptins are products of the *KISS1* gene derived from the plasma proteolytic cleavage of the 145-amino acid gene product, the suffix denoting the number of amino acids. All of the kisspeptin fragments have a C-terminal decapeptide that is critical for biological activity, and all of the kisspeptin forms show similar agonist activity for kisspeptin receptor (Clements *et al.* 2001, Kotani *et al.* 2001, Ohtaki *et al.* 2001).

Kisspeptin is found in both the peripheral and the central nervous system (CNS). In the periphery, kisspeptin has been identified in the testis, ovary, anterior pituitary gonadotrophs, pancreas and small intestine (Ohtaki *et al.* 2001, Richard *et al.* 2008, Gaytan *et al.* 2009). However, peripheral expression of kisspeptin is highest in the placenta with maternal plasma levels of kisspeptin in the third trimester of pregnancy rising to 7000-fold greater than in the non-pregnant state (Muir *et al.* 2001, Ohtaki *et al.* 2001, Horikoshi *et al.* 2003). It has been postulated that the function of kisspeptin production by the placenta may be to down-regulate the HPG axis during pregnancy. However, this is not in keeping with the finding that when kisspeptin-10 is centrally administered to pregnant rats, stimulation of the HPG axis is found to be preserved (Roa *et al.* 2006). Kisspeptin-10 inhibits trophoblast migration in human placental explants (Bilban *et al.* 2004). Plasma kisspeptin IR is elevated in patients with gestational trophoblastic neoplasia when compared with non-pregnant controls and falls during and after chemotherapy (Dhillon *et al.* 2006). Furthermore, levels of kisspeptin and kisspeptin receptor mRNA in placental tissue are increased in cases of gestational trophoblastic disease when compared with normal placental tissue (Janneau *et al.* 2002). Such observations have

led to speculation that kisspeptin may act to regulate trophoblastic invasion of uterine tissue. However, patients with inactivating mutations of the kisspeptin receptor are able to undergo normal pregnancy after treatment with GnRH or gonadotrophin (Pallais *et al.* 2006). The function of placental kisspeptin signalling is therefore currently unknown; however, its presence may not be essential for placental function in humans.

Within the rodent CNS, both *Kiss1* mRNA and kisspeptin protein are particularly highly expressed within the hypothalamus in the arcuate nucleus (ARC), anteroventral periventricular nucleus (AVPV) and periventricular nucleus (Gottsch *et al.* 2004). In primates including humans, hypothalamic *KISS1* mRNA is predominantly found within the infundibular nucleus, which is the equivalent of the ARC in this order of mammals (Rometo *et al.* 2007).

The kisspeptin receptor

The kisspeptin receptor is a member of the rhodopsin family of seven transmembrane GPRs with structural similarities to the galanin receptor. However, it does not appear to bind galanin *in vitro* (Lee *et al.* 1999, Muir *et al.* 2001). Initially classified as an orphan receptor, the kisspeptin receptor (previously known as GPR54) was found to be the cognate receptor for kisspeptin 5 years after the hormone was first described (Kotani *et al.* 2001, Muir *et al.* 2001, Ohtaki *et al.* 2001).

Within the CNS, the kisspeptin receptor is found within the hypothalamus but is also widely expressed within both cortical and subcortical regions (Lee *et al.* 1999). In the periphery, it is notably expressed in the placenta and by pituitary gonadotrophs (Muir *et al.* 2001, Richard *et al.* 2008).

Kisspeptin stimulates the release of GnRH through action at the kisspeptin receptor, which in turn stimulates gonadotrophin release

The GnRH neurons of primates, rodents and sheep are found in close apposition with kisspeptin neurons (Silverman *et al.* 1977, Barry 1979, Rance *et al.* 1994, Clarkson & Herbison 2006). GnRH neurons express the kisspeptin receptor, and when kisspeptin is incubated with hypothalamic explants, it stimulates the release of GnRH. This effect is not observed in kisspeptin receptor knockout mice (*kiss1r*^{-/-}; d'Anglemont de Tassigny *et al.* 2008). The pivotal role of the kisspeptin receptor in kisspeptin-stimulated GnRH release is further demonstrated by the attenuation of the rise in plasma LH following kisspeptin administration in male mice and rats pre-treated with i.c.v. injections of a kisspeptin receptor (Roseweir *et al.* 2009).

I.c.v. administration of kisspeptin-52 to male rodents results in the expression of c-Fos (a well-established marker of neuronal activation) within the cell bodies of GnRH neurons.

In addition, GnRH neurons show an increase in firing rate *in vitro* following kisspeptin treatment. This effect is attenuated by the application of a kisspeptin receptor antagonist (Irwig *et al.* 2004, Liu *et al.* 2008, Roseweir *et al.* 2009). Furthermore, kisspeptin treatment results in a dose- and time-dependent increase in *GnRH* mRNA levels in GnRH-secreting neuronal cell lines (Novaira *et al.* 2009).

Both i.c.v. and peripheral administration of kisspeptin result in a marked rise in plasma LH and to a lesser extent FSH in several mammalian species including rats, mice, sheep, monkeys and humans. This effect is abolished in *GPR54*^{-/-} mice (Gottsch *et al.* 2004, Thompson *et al.* 2004, Dhillo *et al.* 2005, 2007, Messenger *et al.* 2005, Seminara *et al.* 2006, Caraty *et al.* 2007). The effect of kisspeptin on gonadotrophin release is likely to be due to kisspeptin stimulation of GnRH release into the portal circulation, which in turn stimulates the release of LH and FSH from the gonadotrophs of the anterior pituitary gland. Evidence for this includes the abolition of the kisspeptin-induced gonadotrophin rise following pre-treatment with a GnRH antagonist.

These findings are consistent with the phenotype of hypogonadotrophic hypogonadism of *kiss1r*^{-/-} and *kiss1*^{-/-} null mice (de Roux *et al.* 2003, Seminara *et al.* 2003, d'Anglemont de Tassigny *et al.* 2007). These mice have normal levels of hypothalamic GnRH. As expected, kisspeptin administration to *Kiss1*^{-/-} mice results in gonadotrophin release, but this effect is not observed in *Gpr54*^{-/-} mice (de Roux *et al.* 2003, Seminara *et al.* 2003, Messenger *et al.* 2005, d'Anglemont de Tassigny *et al.* 2007). Thus, evidence suggests that kisspeptin is the major ligand for the kisspeptin receptor.

Kisspeptin may also act directly on pituitary gonadotrophs to stimulate gonadotrophin release

Some lines of evidence point towards a direct effect of kisspeptin on the gonadotrophs of the anterior pituitary gland, stimulating the release of LH and FSH. Both the *Kiss1* and *Kiss1R* genes are expressed in pituitary gonadotrophs, and *in vitro* exposure of pituitary cells and tissue explants to kisspeptin results in the dose-dependent release of LH (Kotani *et al.* 2001, Navarro *et al.* 2005, Gutierrez-Pascual *et al.* 2007, Richard *et al.* 2008). Furthermore, in ovariectomised female rats, there is a fall in the expression of *Kiss1* mRNA within gonadotrophs, which can be prevented by the administration of oestradiol (E₂; Richard *et al.* 2008). These data suggest that circulating sex steroids may be required for pituitary *Kiss1* expression. This is consistent with the observation that circulating levels of E₂ modulate responsiveness of pituitary gonadotrophs to GnRH (Knobil *et al.* 1980).

Nevertheless, *kiss1r*^{-/-} mice that show a phenotype of hypogonadotrophic hypogonadism have preserved gonadotroph function as evidenced by an appropriate response to exogenous GnRH (Seminara *et al.* 2003). In addition, i.v. administration of kisspeptin fails to stimulate LH release in

GnRH-replaced, ovariectomised, hypothalamic–pituitary-disconnected ewes. Furthermore, the pre-treatment of male monkeys with a GnRH antagonist results in the failure of administered kisspeptin to stimulate an LH rise (Plant *et al.* 2006, Smith *et al.* 2008).

The role of kisspeptin within the pituitary is currently unknown. *In vivo* and *ex vivo* evidences suggest a stimulatory role of kisspeptin within the anterior pituitary. However, pituitary kisspeptin signalling does not appear critical for the robust stimulation of gonadotrophin release observed following kisspeptin administration *in vivo*. Hence, kisspeptin signalling within the pituitary may have a regulatory role on gonadotrophin function, which is distinct to its better characterised role within the hypothalamus.

Kisspeptin plays a key role in mediating gonadal steroid feedback to the hypothalamus

Steroid hormones produced by the gonads feed back to the hypothalamus exerting a positive or a negative regulatory effect on GnRH production and release. The oestrogen receptors (ERs) are transcription factors, which exist as two isoforms, ER α and ER β . These receptors bind to specific DNA sequences known as oestrogen-response elements found in the promoter region of oestrogen-responsive genes, resulting in the activation or suppression of gene transcription.

During most of the menstrual cycle, oestrogen suppresses gonadotrophin secretion, but at mid-cycle, the effect of oestrogen on the HPG axis changes to a potent positive feedback effect, leading to a surge in the plasma LH and ovulation. Increasing evidence suggests that hypothalamic kisspeptin signalling plays a critical role in the generation of the pre-ovulatory LH surge, which is necessary for normal fertility. Evidence for this comes from ovariectomised *GPR54* and *Kiss1* null mice treated with oestrogen and progesterone, which fail to mount an LH surge unlike their wild-type litter mates (Dungan *et al.* 2007, Clarkson *et al.* 2008). In addition, infusion of a monoclonal anti-rat kisspeptin antibody into the pre-optic area (POA), which contains GnRH cell bodies, leads to the complete blockade of the pre-ovulatory LH surge in oestrogen-treated ovariectomised female rats (Kinoshita *et al.* 2005). Lastly, continuous i.c.v. administration of a kisspeptin antagonist to female rats in the morning of oestrous until the afternoon of the following pro-oestrous prevents the pre-ovulatory LH surge (Pineda *et al.* 2010).

ER α has been shown to be critical for the positive feedback effect of oestrogen, which results in the female mid-cycle LH surge. Adult female rodents pre-administered with an ER α antagonist fail to ovulate or undergo a pre-ovulatory LH surge and have blunted LH response following administration of kisspeptin-10 (Roa *et al.* 2008). Furthermore, ovariectomised, oestrogen-replaced, neuron-specific ER α null mice are infertile. It is not known whether such mice have altered levels of *Kiss1* expression, but they fail to generate

a pre-ovulatory LH peak despite having normal basal levels of LH (Wintermantel *et al.* 2006). Interestingly, the pre-ovulatory LH surge is preserved in ER β knockouts (Wintermantel *et al.* 2006). GnRH neurons express ER β but not ER α , which implies that the mid-cycle positive regulation of GnRH release by oestrogen is mediated by a separate population of ER α -expressing neurons with afferent input to GnRH neurons (Hrabovszky *et al.* 2000). Dual *in situ* hybridisation studies have revealed the co-expression of *Kiss1* and ER α mRNA in the AVPV and the ARC. In addition, work using neuronal viral retrograde tracing in mice has demonstrated ER α -expressing neurons in the AVPV and ARC with afferent input to GnRH neurons (Smith *et al.* 2005, Wintermantel *et al.* 2006). In female rats, *Kiss1* mRNA expression in the AVPV peaks during the evening of pro-oestrous but falls to a nadir in the ARC (Smith *et al.* 2006b). *Kiss1* neurons in the AVPV show high levels of c-Fos expression during the LH surge, while c-Fos expression at dioestrous is virtually absent (Smith *et al.* 2006b). Conversely, c-Fos is almost undetectable in the ARC at pro-oestrous (Smith *et al.* 2006b). Lastly, in female mice following ovariectomy, *Kiss1* expression in the AVPV falls but is increased following oestrogen replacement in these animals. Interestingly, the opposite pattern is observed in the ARC (Smith *et al.* 2005).

Taken altogether, these findings suggest that E₂ within ER α -expressing kisspeptin neurons in the AVPV positively regulates GnRH neurons culminating in the pre-ovulatory LH surge. However, ER α -expressing kisspeptin neurons in the ARC respond to oestrogen stimulation by inhibiting GnRH production and release (Fig. 1). This feed forward-feedback control on GnRH release by gonadal steroids is critical for normal fertility.

Kisspeptin neurons in the ARC co-express other neuropeptides, which regulate the HPG axis

Kisspeptin neurons in the ARC co-express the neuropeptides neurokinin B (NKB) and dynorphin, and this co-localisation is highly conserved in several mammalian species including humans (Goodman *et al.* 2007, Rance 2009, Hrabovszky *et al.* 2010). Dynorphin is an endogenous opioid peptide, which plays a role in the progesterone-mediated negative feedback control of GnRH release, while NKB is a member of the substance P-related tachykinin family (Goodman *et al.* 2004, Krajewski *et al.* 2005). The receptor for NKB is tachykinin neurokinin 3 receptor (NK3R), which is expressed on GnRH neurons (Todman *et al.* 2005).

Recently, it has been reported in humans that loss-of-function mutations in the gene encoding NKB (*TAC3*) or the gene encoding NK3R (*TAC3R*) result in normosmic hypogonadotropic hypogonadism and pubertal failure. This phenotype is remarkably similar to rodent and human models of defective kisspeptin signalling (de Roux *et al.* 2003,

Figure 1 Diagram of kisspeptin signalling within the female central nervous system. KISS1-expressing neurons within the hypothalamic arcuate nucleus (ARC; equivalent to the infundibular nucleus in primates) are negatively regulated by oestrogen and inhibit GnRH release. KISS1-expressing neurons within the hypothalamic anteroventral periventricular nucleus (AVPV) are positively regulated by oestrogen and stimulate GnRH during the pre-ovulatory surge. Kisspeptin may also have a direct modulatory effect on pituitary LH and FSH release. KISS1 neurons in the ARC co-express neurokinin B (NKB), which stimulates LH and FSH release in a GnRH-dependent manner. Leptin signalling has a permissive effect on kisspeptin signalling within the ARC, which may be mediated by the mammalian target of rapamycin (mTOR; also known as mechanistic target of rapamycin, MTOR) pathway.

Seminara *et al.* 2003, d'Anglemont de Tassigny *et al.* 2007, Guran *et al.* 2009, Topaloglu *et al.* 2009).

Kisspeptin neurons in the ARC project to GnRH neurons. The association of NKB and kisspeptin within kisspeptin neurons may be suggestive of a synergy between the two neuropeptides in the regulation of the HPG axis. For example, it is known that the expression of both peptides is inhibited by E₂. In addition, in post-menopausal women, the distribution and morphology of the hypertrophied *KISS1*-positive neurons in the infundibular nucleus are similar to that of *NKB*-expressing cells (Romero *et al.* 2007, Navarro *et al.* 2009).

Until recently, the effect of NKB on reproductive hormone release had been inconclusive with conflicting results from studies in sheep and rodents (Jayasena & Dhillon 2010). However, recently it has been reported that i.v. administration of NKB or an NKB agonist to male monkeys results in a potent stimulation of LH release, an effect that is abolished by pre-treatment with a GnRH antagonist (Ramaswamy *et al.* 2010). Interestingly, repetitive administration of NKB is

not associated with a sustained pattern of LH release (Ramaswamy *et al.* 2010). Furthermore, monkeys remain responsive to kisspeptin injection despite losing responsiveness to NKB following repetitive administration. This suggests that NKB stimulates GnRH release in a kisspeptin-independent manner. It is not known whether animals may respond to NKB despite losing responsiveness to kisspeptin following chronic administration; such an observation would suggest NKB and kisspeptin to have parallel influences of GnRH release.

A role for kisspeptin signalling in pubertal development and seasonal reproductive activity

During puberty, the immature mammal develops adult physical and hormonal characteristics rendering it fertile and capable of reproduction. Juvenile *kiss1*^{-/-} and *kiss1*^{-/-} mice and humans with inactivating kisspeptin receptor mutations fail to enter puberty (de Roux *et al.* 2003,

Seminara *et al.* 2003, d'Anglemont de Tassigny *et al.* 2007); these observations therefore suggest that kisspeptin signalling plays a critical role in the onset of puberty.

Further evidence for the importance of kisspeptin signalling in the onset of puberty comes from studies of immature female rats administered twice daily i.c.v. injections of kisspeptin from postnatal day 26 to day 31 (Navarro *et al.* 2004). This results in precocious vaginal opening, increased uterine weight and raised plasma LH and E₂ levels relative to vehicle-treated controls. These findings complement the report of central precocious puberty in an 8-year-old girl due to an activating mutation of KISS1R (Teles *et al.* 2008).

At postnatal day 10, kisspeptin-expressing neurons in the AVPV are not detectable in either male or female mice. However, these become apparent, lying in close apposition to GnRH neurons, from postnatal day 25, reaching adult levels by the onset of puberty at postnatal day 31 (Clarkson & Herbison 2006). In support of these findings, the expression of *KISS1* mRNA in the mediobasal hypothalamus (MBH) of agonadal male rhesus monkeys has been found to be significantly greater in the pubertal group than in the juvenile cohort (Shahab *et al.* 2005). Similarly, in female monkeys with intact gonads, a threefold increase in *KISS1* mRNA expression is observed in the MBH in early pubertal animals that in a juvenile group (Shahab *et al.* 2005).

Kisspeptin is also implicated in seasonal regulation of reproductive activity in seasonal breeders. In Syrian hamsters, hypothalamic *Kiss1* expression is high during long day conditions (associated with sexual activity) and reduced during short day conditions (associated with a quiescence of sexual activity); furthermore, administration of kisspeptin-10 to these animals restores testicular weight and testosterone release (Revel *et al.* 2006, 2007). Sheep display seasonal sexual activity during short day conditions. *KISS1* expression within the ARC is increased under short day conditions in ovariectomised ewes, but no seasonal change in *KISS1* expression is detected within the POA (Clarke *et al.* 2009).

These findings suggest that kisspeptin acts as an essential gatekeeper to the onset of puberty and consequent fertility in several mammalian species. Furthermore, kisspeptin may play a role in the seasonal regulation of reproductive activity in Syrian hamsters and sheep.

Kisspeptin provides a link between nutritional status and fertility

It has long been recognised that an intimate relationship exists between nutritional status and fertility. Leptin is a peptide hormone synthesised and secreted by adipocytes conveying information about body energy stores and nutritional status (Zhang *et al.* 1994, Pelleymounter *et al.* 1995). The level of circulating leptin is proportional to fat mass and falls in both mice and humans following weight loss (Maffei *et al.* 1995). The leptin-deficient *ob/ob* mouse is a well-established model of hypoleptinaemia (Zhang *et al.* 1994). These mice have

delayed puberty and are infertile as a consequence of hypogonadotrophic hypogonadism. Interestingly, a similar phenotype of hypogonadotrophic hypogonadism is also present in humans with mutations of leptin or its receptor and in women with hypoleptinaemia as a consequence of low body weight (Coleman 1978, Clement *et al.* 1998, Farooqi *et al.* 1999, Welt *et al.* 2004). Leptin therapy in hypoleptinaemic mice and humans reverses these reproductive abnormalities. In addition, when leptin is administered to juvenile wild-type mice, this accelerates the onset of puberty (Chehab *et al.* 1996, 1997, Ahima *et al.* 1997, Farooqi *et al.* 1999, Welt *et al.* 2004).

The leptin receptor (Ob-Rb) is not expressed by GnRH neurons of the hypothalamus. However *Ob-Rb* mRNA is found in 40% of *Kiss1* mRNA-expressing cells of the ARC (Smith *et al.* 2006a). The expression of *Kiss1* mRNA in the ARC of *ob/ob* mice is reduced in comparison with wild-type mice. Interestingly, although kisspeptin mRNA in the ARC of *ob/ob* mice is significantly increased following leptin treatment, its expression is not restored to that of the wild-type controls (Smith *et al.* 2006a).

The expression of *Kiss1* mRNA has been shown to be influenced by nutritional status. In pre-pubertal rats, which have been food deprived for 72 h, the hypothalamic expression of *Kiss1* mRNA is markedly reduced (Castellano *et al.* 2005). In a model of chronic undernutrition in pre-pubertal rats, daily i.c.v. administration of kisspeptin from postnatal day 30 to 37 restores the delayed vaginal opening of these animals and increases the suppressed levels of plasma LH, FSH and E₂ (Castellano *et al.* 2005).

The ARC contains discrete subpopulations of first-order leptin-responsive neurons. One subpopulation expresses the orexigenic peptide neuropeptide Y (NPY), while another expresses the anorectic peptide α -melanocyte-stimulating hormone (α -MSH), which is derived from pro-opiomelanocortin (POMC; Cheung *et al.* 1997, Broberger *et al.* 1998). Using double-label fluorescent immunohistochemistry, kisspeptin fibres in ewes have been shown to be in close apposition with ARC NPY and POMC neurons (Backholer *et al.* 2010). I.c.v. administration of an α -MSH-like agonist to ewes increases *KISS1* mRNA in the POA and results in a rise in plasma LH (Backholer *et al.* 2009). Conversely, the orexigenic peptide melanin-concentrating hormone, which is highly expressed in the lateral hypothalamus, inhibits the stimulatory effects of kisspeptin on GnRH neurons (Wu *et al.* 2009).

Mammalian target of rapamycin protein (mTOR; also known as mechanistic target of rapamycin, MTOR) is another regulator of energy homeostasis, which has been shown to influence kisspeptin signalling. mTOR is a ubiquitously expressed serine-threonine protein kinase, which plays a vital role in the regulation of cell growth and differentiation (Schmelzle & Hall 2000). When nutrient availability is low, mTOR activity falls thus inhibiting the high-energy demand cell cycle. Interestingly, i.c.v. administration of leucine (which stimulates mTOR signalling) leads to an increase in plasma LH in female peri-pubertal rats

(Cota *et al.* 2006, Roa *et al.* 2009). Conversely, when mTOR activation is blocked (by rapamycin), the expression of *Kiss1* mRNA in the ARC becomes almost undetectable, and plasma levels of LH fall (Roa *et al.* 2009).

Reproduction is a highly energy demanding process, and during food deprivation the HPG axis is down-regulated in order to conserve energy. In summary, the findings presented above demonstrate a key role for hypothalamic kisspeptin as a link between nutritional status and fertility.

Could kisspeptin offer a novel therapy for the treatment of fertility disorders?

As our understanding of the biology of the kisspeptin signalling system grows, it is becoming increasingly tempting to speculate on the possible therapeutic use of kisspeptin in the treatment of fertility disorders. Infertility affects up to one in six couples in the United Kingdom, and although effective, both the hormonal and surgical therapies, which are currently employed are not without significant side effects and failure rates (National Institute for Health and Clinical Excellence 2004, Human Fertilisation and Embryology Authority 2009, <http://www.hfea.gov.uk/infertility-facts.html#1248>).

When administered to healthy male subjects, i.v. infusion of kisspeptin results in a significant increase in plasma LH, FSH and testosterone (Dhillon *et al.* 2005). S.c. injection of kisspeptin to healthy pre-menopausal female subjects elicits a marked rise in LH, which is most pronounced in the pre-ovulatory phase of the menstrual cycle (Dhillon *et al.* 2007). This finding sits well with the animal studies reviewed above, which suggest an essential role for kisspeptin in the generation of the LH surge.

Kisspeptin has been administered to humans by s.c. and i.v. injection without any observed adverse effects (Dhillon *et al.* 2005, 2007). In particular, kisspeptin has not been shown to cause changes in heart rate or blood pressure, which is important in view of the vasoconstrictor properties of kisspeptin that have been demonstrated *in vitro* (Mead *et al.* 2007, Nijher *et al.* 2010).

When kisspeptin is administered twice daily s.c. injection to infertile women with functional hypothalamic amenorrhoea (HA) due to low body weight, it effectively stimulates a rise in plasma gonadotrophins (Jayasena *et al.* 2009). This effect is most marked following the first injection of kisspeptin and is significantly diminished after 2 weeks of treatment. However, biweekly administration of kisspeptin results in a sustained gonadotrophin response to kisspeptin (Jayasena *et al.* 2010). Furthermore, it has been shown that in women with HA, the LH response to injected kisspeptin is fourfold greater than that of healthy female subjects studied in the follicular phase (Jayasena *et al.* 2009). This may be due to an enhanced responsiveness to kisspeptin in women with HA or increased pituitary sensitivity to the effects of GnRH.

Gonadotrophin injections provide the current mainstay of infertility therapy (Elchalal & Schenker 1997). It is therefore

interesting to consider if kisspeptin-based therapies would offer any potential advantages over existing therapies. Kisspeptin acts by stimulating endogenous hypothalamic GnRH release, which in turn triggers endogenous pituitary gonadotrophin release (Irwig *et al.* 2004, Thompson *et al.* 2004, Shahab *et al.* 2005). Kisspeptin therapy might therefore stimulate a more natural pattern of reproductive hormone release than existing therapies. Furthermore, the stimulation of endogenous gonadotrophin release by kisspeptin may be predicted to confer a lower risk of ovarian hyperstimulation syndrome associated with exogenous gonadotrophin injections (Elchalal & Schenker 1997). More data are needed to test if such potential benefits would exist over current therapies for infertility.

Concluding remarks

Over the past decade, our increasing insight into the biology of the kisspeptin pathway has significantly added to our understanding of the physiology and pathophysiology of the HPG axis. Kisspeptin has now been safely and successfully used in both healthy and infertile human subjects, and it is possible that in the future the manipulation of kisspeptin signalling may be used in the treatment of reproductive disorders.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

SH is supported by a Medical Research Council Clinical Research Training Fellowship. CNJ is supported by an NIHR Clinical Lectureship and a Wellcome Trust Clinical Research Training Fellowship. WSD is supported by a HEFCE Clinical Senior Lecturer Award. The department is funded by an Integrative Mammalian Biology (IMB) Capacity Building Award and the NIHR Biomedical Research Centre Funding Scheme.

References

- Ahima RS, Dushay J, Flier SN, Prabakaran D & Flier JS 1997 Leptin accelerates the onset of puberty in normal female mice. *Journal of Clinical Investigation* **99** 391–395. (doi:10.1172/JCI119172)
- Backholer K, Smith J & Clarke IJ 2009 Melanocortins may stimulate reproduction by activating orexin neurons in the dorsomedial hypothalamus and kisspeptin neurons in the preoptic area of the ewe. *Endocrinology* **150** 5488–5497. (doi:10.1210/en.2009-0604)
- Backholer K, Smith JT, Rao A, Pereira A, Iqbal J, Ogawa S, Li Q & Clarke IJ 2010 Kisspeptin cells in the ewe brain respond to leptin and communicate with neuropeptide Y and proopiomelanocortin cells. *Endocrinology* **151** 2233–2243. (doi:10.1210/en.2009-1190)
- Barry J 1979 Immunohistochemistry of luteinizing hormone-releasing hormone-producing neurons of the vertebrates. *International Review of Cytology* **60** 179–221. (doi:10.1016/S0074-7696(08)61263-2)
- Bilban M, Ghaffari-Tabrizi N, Hintermann E, Bauer S, Molzer S, Zoratti C, Malli R, Sharabi A, Hiden U, Graier W *et al.* 2004 Kisspeptin-10, a

- KiSS-1/metastatin-derived decapeptide, is a physiological invasion inhibitor of primary human trophoblasts. *Journal of Cell Science* **117** 1319–1328. (doi:10.1242/jcs.00971)
- Broberger C, Johansen J, Johansson C, Schalling M & Hokfelt T 1998 The neuropeptide Y/agouti gene-related protein (AGRP) brain circuitry in normal, anorectic, and monosodium glutamate-treated mice. *PNAS* **95** 15043–15048. (doi:10.1073/pnas.95.25.15043)
- Caraty A, Smith JT, Lomet D, Ben SS, Morrissey A, Cagnie J, Doughton B, Baril G, Briant C & Clarke IJ 2007 Kisspeptin synchronizes preovulatory surges in cyclical ewes and causes ovulation in seasonally acyclic ewes. *Endocrinology* **148** 5258–5267. (doi:10.1210/en.2007-0554)
- Castellano JM, Navarro VM, Fernandez-Fernandez R, Nogueiras R, Tovar S, Roa J, Vazquez MJ, Vigo E, Casanueva FF, Aguilar E *et al.* 2005 Changes in hypothalamic KiSS-1 system and restoration of pubertal activation of the reproductive axis by kisspeptin in undernutrition. *Endocrinology* **146** 3917–3925. (doi:10.1210/en.2005-0337)
- Chehab FF, Lim ME & Lu R 1996 Correction of the sterility defect in homozygous obese female mice by treatment with the human recombinant leptin. *Nature Genetics* **12** 318–320. (doi:10.1038/ng0396-318)
- Chehab FF, Mounzih K, Lu R & Lim ME 1997 Early onset of reproductive function in normal female mice treated with leptin. *Science* **275** 88–90. (doi:10.1126/science.275.5296.88)
- Cheung CC, Clifton DK & Steiner RA 1997 Proopiomelanocortin neurons are direct targets for leptin in the hypothalamus. *Endocrinology* **138** 4489–4492. (doi:10.1210/en.138.10.4489)
- Clarke IJ, Smith JT, Caraty A, Goodman RL & Lehman MN 2009 Kisspeptin and seasonality in sheep. *Peptides* **30** 154–163. (doi:10.1016/j.peptides.2008.08.022)
- Clarkson J & Herbison AE 2006 Postnatal development of kisspeptin neurons in mouse hypothalamus; sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinology* **147** 5817–5825. (doi:10.1210/en.2006-0787)
- Clarkson J, d'Anglemont de Tassigny X, Moreno AS, Colledge WH & Herbison AE 2008 Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. *Journal of Neuroscience* **28** 8691–8697. (doi:10.1523/JNEUROSCI.1775-08.2008)
- Clement K, Vaisse C, Lahlou N, Cabrol S, Pelloux V, Casuto D, Gourmelen M, Dina C, Chambaz J, Lacorte JM *et al.* 1998 A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature* **392** 398–401. (doi:10.1038/32911)
- Clements MK, McDonald TP, Wang R, Xie G, O'Dowd BF, George SR, Austin CP & Liu Q 2001 FMRamide-related neuropeptides are agonists of the orphan G-protein-coupled receptor GPR54. *Biochemical and Biophysical Research Communications* **284** 1189–1193. (doi:10.1006/bbrc.2001.5098)
- Coleman DL 1978 Obese and diabetes: two mutant genes causing diabetes-obesity syndromes in mice. *Diabetologia* **14** 141–148. (doi:10.1007/BF00429772)
- Cota D, Proulx K, Smith KA, Kozma SC, Thomas G, Woods SC & Seeley RJ 2006 Hypothalamic mTOR signaling regulates food intake. *Science* **312** 927–930. (doi:10.1126/science.1124147)
- d'Anglemont de Tassigny X, Fagg LA, Dixon JP, Day K, Leitch HG, Hendrick AG, Zahn D, Franceschini I, Caraty A, Carlton MB *et al.* 2007 Hypogonadotropic hypogonadism in mice lacking a functional Kiss1 gene. *PNAS* **104** 10714–10719. (doi:10.1073/pnas.0704114104)
- d'Anglemont de Tassigny X, Fagg LA, Carlton MB & Colledge WH 2008 Kisspeptin can stimulate gonadotropin-releasing hormone (GnRH) release by a direct action at GnRH nerve terminals. *Endocrinology* **149** 3926–3932. (doi:10.1210/en.2007-1487)
- Dhillon WS, Chaudhri OB, Patterson M, Thompson EL, Murphy KG, Badman MK, McGowan BM, Amber V, Patel S, Ghatei MA *et al.* 2005 Kisspeptin-54 stimulates the hypothalamic-pituitary-gonadal axis in human males. *Journal of Clinical Endocrinology and Metabolism* **90** 6609–6615. (doi:10.1210/jc.2005-1468)
- Dhillon WS, Savage P, Murphy KG, Chaudhri OB, Patterson M, Nijher GM, Foggo VM, Dancey GS, Mitchell H, Seckl MJ *et al.* 2006 Plasma kisspeptin is raised in patients with gestational trophoblastic neoplasia and falls during treatment. *American Journal of Physiology. Endocrinology and Metabolism* **291** E878–E884. (doi:10.1152/ajpendo.00555.2005)
- Dhillon WS, Chaudhri OB, Thompson EL, Murphy KG, Patterson M, Ramachandran R, Nijher GK, Amber V, Kokkinos A, Donaldson M *et al.* 2007 Kisspeptin-54 stimulates gonadotropin release most potently during the preovulatory phase of the menstrual cycle in women. *Journal of Clinical Endocrinology and Metabolism* **92** 3958–3966. (doi:10.1210/jc.2007-1116)
- Dungan HM, Gottsch ML, Zeng H, Gragerov A, Bergmann JE, Vassilatis DK, Clifton DK & Steiner RA 2007 The role of kisspeptin-GPR54 signaling in the tonic regulation and surge release of gonadotropin-releasing hormone/luteinizing hormone. *Journal of Neuroscience* **27** 12088–12095. (doi:10.1523/JNEUROSCI.2748-07.2007)
- Elchahal U & Schenker JG 1997 The pathophysiology of ovarian hyperstimulation syndrome – views and ideas. *Human Reproduction* **12** 1129–1137. (doi:10.1093/humrep/12.6.1129)
- Farooqi IS, Jebb SA, Langmack G, Lawrence E, Cheetham CH, Prentice AM, Hughes IA, McCamish MA & O'Rahilly S 1999 Effects of recombinant leptin therapy in a child with congenital leptin deficiency. *New England Journal of Medicine* **341** 879–884. (doi:10.1056/NEJM199909163411204)
- Gaytan F, Gaytan M, Castellano JM, Romero M, Roa J, Aparicio B, Garrido N, Sanchez-Criado JE, Millar RP, Pellicer A *et al.* 2009 KiSS-1 in the mammalian ovary: distribution of kisspeptin in human and marmoset and alterations in KiSS-1 mRNA levels in a rat model of ovulatory dysfunction. *American Journal of Physiology. Endocrinology and Metabolism* **296** E520–E531. (doi:10.1152/ajpendo.90895.2008)
- Goodman RL, Coolen LM, Anderson GM, Hardy SL, Valent M, Connors JM, Fitzgerald ME & Lehman MN 2004 Evidence that dynorphin plays a major role in mediating progesterone negative feedback on gonadotropin-releasing hormone neurons in sheep. *Endocrinology* **145** 2959–2967. (doi:10.1210/en.2003-1305)
- Goodman RL, Lehman MN, Smith JT, Coolen LM, de Oliveira CV, Jafarzadehshirazi MR, Pereira A, Iqbal J, Caraty A, Ciofi P *et al.* 2007 Kisspeptin neurons in the arcuate nucleus of the ewe express both dynorphin A and neurokinin B. *Endocrinology* **148** 5752–5760. (doi:10.1210/en.2007-0961)
- Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, Crowley WF, Seminara S, Clifton DK & Steiner RA 2004 A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology* **145** 4073–4077. (doi:10.1210/en.2004-0431)
- Guran T, Tolhurst G, Bereket A, Rocha N, Porter K, Turan S, Gribble FM, Kotan LD, Akcay T, Atay Z *et al.* 2009 Hypogonadotropic hypogonadism due to a novel missense mutation in the first extracellular loop of the neurokinin B receptor. *Journal of Clinical Endocrinology and Metabolism* **94** 3633–3639. (doi:10.1210/jc.2009-0551)
- Gutierrez-Pascual E, Martinez-Fuentes AJ, Pinilla L, Tena-Sempere M, Malagon MM & Castano JP 2007 Direct pituitary effects of kisspeptin: activation of gonadotrophs and somatotrophs and stimulation of luteinizing hormone and growth hormone secretion. *Journal of Neuroendocrinology* **19** 521–530. (doi:10.1111/j.1365-2826.2007.01558.x)
- Horikoshi Y, Matsumoto H, Takatsu Y, Ohtaki T, Kitada C, Usuki S & Fujino M 2003 Dramatic elevation of plasma metastin concentrations in human pregnancy: metastin as a novel placenta-derived hormone in humans. *Journal of Clinical Endocrinology and Metabolism* **88** 914–919. (doi:10.1210/jc.2002-021235)
- Hrabovszky E, Shughrue PJ, Merchenthaler I, Hajszan T, Carpenter CD, Liposits Z & Petersen SL 2000 Detection of estrogen receptor- β messenger ribonucleic acid and 125I-estrogen binding sites in luteinizing hormone-releasing hormone neurons of the rat brain. *Endocrinology* **141** 3506–3509. (doi:10.1210/en.141.9.3506)
- Hrabovszky E, Ciofi P, Vida B, Horvath MC, Keller E, Caraty A, Bloom SR, Ghatei MA, Dhillon WS, Liposits Z *et al.* 2010 The kisspeptin system of the human hypothalamus: sexual dimorphism and relationship with gonadotropin-releasing hormone and neurokinin B neurons. *European Journal of Neuroscience* **31** 1984–1998. (doi:10.1111/j.1460-9568.2010.07239.x)

- Irwig MS, Fraley GS, Smith JT, Acohido BV, Popa SM, Cunningham MJ, Gottsch ML, Clifton DK & Steiner RA 2004 Kisspeptin activation of gonadotropin releasing hormone neurons and regulation of KiSS-1 mRNA in the male rat. *Neuroendocrinology* **80** 264–272. (doi:10.1159/000083140)
- Janneau JL, Maldonado-Estrada J, Tachdjian G, Miran I, Motte N, Saulnier P, Sabourin JC, Cote JF, Simon B, Frydman R *et al.* 2002 Transcriptional Expression of genes involved in cell invasion and migration by normal and tumoral trophoblast cells. *Journal of Clinical Endocrinology and Metabolism* **87** 5336–5339. (doi:10.1210/jc.2002-021093)
- Jayasena CN, Nijher GM, Chaudhri OB, Murphy KG, Ranger A, Lim A, Patel D, Mehta A, Todd C, Ramachandran R *et al.* 2009 Subcutaneous injection of kisspeptin-54 acutely stimulates gonadotropin secretion in women with hypothalamic amenorrhea, but chronic administration causes tachyphylaxis. *Journal of Clinical Endocrinology and Metabolism* **94** 4315–4323. (doi:10.1210/jc.2009-0406)
- Jayasena CN & Dhillon WS 2010 Neurokinin B and Kisspeptin – sexual partners or single agents? *Endocrinology* **151** 4090–4091. (doi:10.1210/en.2010-0696)
- Jayasena CN, Nijher GMK, Abbara A, Murphy KG, Lim A, Patel DA, Mehta A, Todd C, Donaldson M, Trew G *et al.* 2010 Twice-weekly administration of kisspeptin-54 for eight weeks stimulates reproductive hormone release in women with hypothalamic amenorrhea, Endo 2010 meeting, San Diego, CA, USA.
- Kinoshita M, Tsukamura H, Adachi S, Matsui H, Uenoyama Y, Iwata K, Yamada S, Inoue K, Ohtaki T, Matsumoto H *et al.* 2005 Involvement of central metastin in the regulation of preovulatory luteinizing hormone surge and estrous cyclicity in female rats. *Endocrinology* **146** 4431–4436. (doi:10.1210/en.2005-0195)
- Knobil E, Plant TM, Wildt L, Belchetz PE & Marshall G 1980 Control of the rhesus monkey menstrual cycle: permissive role of hypothalamic gonadotropin-releasing hormone. *Science* **207** 1371–1373. (doi:10.1126/science.6766566)
- Kotani M, Detheux M, Vandenbogaerde A, Communi D, Vanderwinden JM, Le PE, Brezillon S, Tyldesley R, Suarez-Huerta N, Vandeput F *et al.* 2001 The metastasis suppressor gene KiSS-1 encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor GPR54. *Journal of Biological Chemistry* **276** 34631–34636. (doi:10.1074/jbc.M104847200)
- Krajewski SJ, Anderson MJ, Iles-Shih L, Chen KJ, Urbanski HF & Rance NE 2005 Morphologic evidence that neurokinin B modulates gonadotropin-releasing hormone secretion via neurokinin 3 receptors in the rat median eminence. *Journal of Comparative Neurology* **489** 372–386. (doi:10.1002/cne.20626)
- Lee JH, Miele ME, Hicks DJ, Phillips KK, Trent JM, Weissman BE & Welch DR 1996 KiSS-1, a novel human malignant melanoma metastasis-suppressor gene. *Journal of the National Cancer Institute* **88** 1731–1737. (doi:10.1093/jnci/88.23.1731)
- Lee DK, Nguyen T, O'Neill GP, Cheng R, Liu Y, Howard AD, Coulombe N, Tan CP, Tang-Nguyen AT, George SR *et al.* 1999 Discovery of a receptor related to the galanin receptors. *FEBS Letters* **446** 103–107. (doi:10.1016/S0014-5793(99)00009-5)
- Liu X, Lee K & Herbison AE 2008 Kisspeptin excites gonadotropin-releasing hormone neurons through a phospholipase C/calcium-dependent pathway regulating multiple ion channels. *Endocrinology* **149** 4605–4614. (doi:10.1210/en.2008-0321)
- Maffei M, Halaas J, Ravussin E, Pratley RE, Lee GH, Zhang Y, Fei H, Kim S, Lallone R, Ranganathan S *et al.* 1995 Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nature Medicine* **1** 1155–1161. (doi:10.1038/nm1195-1155)
- Mead EJ, Maguire JJ, Kuc RE & Davenport AP 2007 Kisspeptins are novel potent vasoconstrictors in humans, with a discrete localization of their receptor, G protein-coupled receptor 54, to atherosclerosis-prone vessels. *Endocrinology* **148** 140–147. (doi:10.1210/en.2006-0818)
- Message S, Chatzidakis EE, Ma D, Hendrick AG, Zahn D, Dixon J, Thresher RR, Malinge I, Lomet D, Carlton MB *et al.* 2005 Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *PNAS* **102** 1761–1766. (doi:10.1073/pnas.0409330102)
- Muir AI, Chamberlain L, Elshourbagy NA, Michalovich D, Moore DJ, Calamari A, Szekeres PG, Sarau HM, Chambers JK, Murdock P *et al.* 2001 AXOR12, a novel human G protein-coupled receptor, activated by the peptide KiSS-1. *Journal of Biological Chemistry* **276** 28969–28975. (doi:10.1074/jbc.M102743200)
- National Institute for Health and Clinical Excellence 2004 *Fertility Assessment and Treatment for People with Fertility Problems*. <http://www.nice.org.uk/nice/media/live/10936/29269/29269.pdf>.
- Navarro VM, Fernandez-Fernandez R, Castellano JM, Roa J, Mayen A, Barreiro ML, Gaytan F, Aguilar E, Pinilla L, Dieguez C *et al.* 2004 Advanced vaginal opening and precocious activation of the reproductive axis by KiSS-1 peptide, the endogenous ligand of GPR54. *Journal of Physiology* **561** 379–386. (doi:10.1113/jphysiol.2004.072298)
- Navarro VM, Castellano JM, Fernandez-Fernandez R, Tovar S, Roa J, Mayen A, Nogueiras R, Vazquez MJ, Barreiro ML, Magni P *et al.* 2005 Characterization of the potent luteinizing hormone-releasing activity of KiSS-1 peptide, the natural ligand of GPR54. *Endocrinology* **146** 156–163. (doi:10.1210/en.2004-0836)
- Navarro VM, Gottsch ML, Chavkin C, Okamura H, Clifton DK & Steiner RA 2009 Regulation of gonadotropin-releasing hormone secretion by kisspeptin/dynorphin/neurokinin B neurons in the arcuate nucleus of the mouse. *Journal of Neuroscience* **29** 11859–11866. (doi:10.1523/JNEUROSCI.1569-09.2009)
- Nijher G, Chaudhri O, Ramachandran R, Murphy K, Zac-Varghese S, Fowler A, Chinthapalli K, Patterson M, Thompson E, Williamson C *et al.* 2010 The effects of kisspeptin-54 on blood pressure in humans and plasma kisspeptin levels in hypertensive diseases of pregnancy. *British Journal of Clinical Pharmacology* **70** 674–681. (doi:10.1111/j.1365-2125.2010.03746.x)
- Novaira HJ, Ng Y, Wolfe A & Radovick S 2009 Kisspeptin increases GnRH mRNA expression and secretion in GnRH secreting neuronal cell lines. *Molecular and Cellular Endocrinology* **311** 126–134. (doi:10.1016/j.mce.2009.06.011)
- Ohtaki T, Shintani Y, Honda S, Matsumoto H, Hori A, Kanehashi K, Terao Y, Kumano S, Takatsu Y, Masuda Y *et al.* 2001 Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-coupled receptor. *Nature* **411** 613–617. (doi:10.1038/35079135)
- Pallais JC, Bo-Abbas Y, Pitteloud N, Crowley WF Jr & Seminara SB 2006 Neuroendocrine, gonadal, placental, and obstetric phenotypes in patients with IHH and mutations in the G-protein coupled receptor, GPR54. *Molecular and Cellular Endocrinology* **254–255** 70–77. (doi:10.1016/j.mce.2006.04.019)
- Pelleymounter MA, Cullen MJ, Baker MB, Hecht R, Winters D, Boone T & Collins F 1995 Effects of the obese gene product on body weight regulation in ob/ob mice. *Science* **269** 540–543. (doi:10.1126/science.7624776)
- Pineda R, Garcia-Galiano D, Roseweir A, Romero M, Sanchez-Garrido MA, Ruiz-Pino F, Morgan K, Pinilla L, Millar RP & Tena-Sempere M 2010 Critical roles of kisspeptins in female puberty and preovulatory gonadotropin surges as revealed by a novel antagonist. *Endocrinology* **151** 722–730. (doi:10.1210/en.2009-0803)
- Plant TM, Ramaswamy S & Dipietro MJ 2006 Repetitive activation of hypothalamic G protein-coupled receptor 54 with intravenous pulses of kisspeptin in the juvenile monkey (*Macaca mulatta*) elicits a sustained train of gonadotropin-releasing hormone discharges. *Endocrinology* **147** 1007–1013. (doi:10.1210/en.2005-1261)
- Ramaswamy S, Seminara SB, Ali B, Ciofi P, Amin NA & Plant TM 2010 Neurokinin B stimulates GnRH release in the male monkey (*Macaca mulatta*) and is colocalized with kisspeptin in the arcuate nucleus. *Endocrinology* **151** 4494–4503. (doi:10.1210/en.2010-0223)
- Rance NE 2009 Menopause and the human hypothalamus: evidence for the role of kisspeptin/neurokinin B neurons in the regulation of estrogen negative feedback. *Peptides* **30** 111–122. (doi:10.1016/j.peptides.2008.05.016)
- Rance NE, Young WS III & McMullen NT 1994 Topography of neurons expressing luteinizing hormone-releasing hormone gene transcripts in the human hypothalamus and basal forebrain. *Journal of Comparative Neurology* **339** 573–586. (doi:10.1002/cne.903390408)
- Revel FG, Saboureaux M, Masson-Pévet M, Pévet P, Mikkelsen JD & Simonneaux V 2006 Kisspeptin mediates the photoperiodic control of reproduction in hamsters. *Current Biology* **16** 1730–1735. (doi:10.1016/j.cub.2006.07.025)

- Revel FG, Ansel L, Klosen P, Saboureaux M, Pévet P, Mikkelsen JD & Simonneaux V 2007 Kisspeptin: a key link to seasonal breeding. *Reviews in Endocrine and Metabolic Disorders* **8** 57–65. (doi:10.1007/s11154-007-9031-7)
- Richard N, Galmiche G, Corvaisier S, Caraty A & Kottler ML 2008 KiSS-1 and GPR54 genes are co-expressed in rat gonadotrophs and differentially regulated *in vivo* by oestradiol and gonadotrophin-releasing hormone. *Journal of Neuroendocrinology* **20** 381–393. (doi:10.1111/j.1365-2826.2008.01653.x)
- Roa J, Vigo E, Castellano JM, Navarro VM, Fernandez-Fernandez R, Casanueva FF, Dieguez C, Aguilar E, Pinilla L & Tena-Sempere M 2006 Hypothalamic expression of KiSS-1 system and gonadotropin-releasing effects of kisspeptin in different reproductive states of the female rat. *Endocrinology* **147** 2864–2878. (doi:10.1210/en.2005-1463)
- Roa J, Vigo E, Castellano JM, Gaytan F, Navarro VM, Aguilar E, Dijkens FA, Ederveen AG, Pinilla L, van Noort PI *et al.* 2008 Opposite roles of estrogen receptor (ER)- α and ER β in the modulation of luteinizing hormone responses to kisspeptin in the female rat: implications for the generation of the preovulatory surge. *Endocrinology* **149** 1627–1637. (doi:10.1210/en.2007-1540)
- Roa J, Garcia-Galiano D, Varela L, Sanchez-Garrido MA, Pineda R, Castellano JM, Ruiz-Pino F, Romero M, Aguilar E, Lopez M *et al.* 2009 The mammalian target of rapamycin as novel central regulator of puberty onset via modulation of hypothalamic Kiss1 system. *Endocrinology* **150** 5016–5026. (doi:10.1210/en.2009-0096)
- Rometo AM, Krajewski SJ, Voytko ML & Rance NE 2007 Hypertrophy and increased kisspeptin gene expression in the hypothalamic infundibular nucleus of postmenopausal women and ovariectomized monkeys. *Journal of Clinical Endocrinology and Metabolism* **92** 2744–2750. (doi:10.1210/jc.2007-0553)
- Roseweir AK, Kauffman AS, Smith JT, Guerriero KA, Morgan K, Pielecka-Fortuna J, Pineda R, Gottsch ML, Tena-Sempere M, Moenter SM *et al.* 2009 Discovery of potent kisspeptin antagonists delineate physiological mechanisms of gonadotropin regulation. *Journal of Neuroscience* **29** 3920–3929. (doi:10.1523/JNEUROSCI.5740-08.2009)
- de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL & Milgrom E 2003 Hypogonadotropic hypogonadism due to loss of function of the KiSS1-derived peptide receptor GPR54. *PNAS* **100** 10972–10976. (doi:10.1073/pnas.1834399100)
- Schmelzle T & Hall MN 2000 TOR, a central controller of cell growth. *Cell* **103** 253–262.
- Seminara SB, Messager S, Chatzidaki EE, Thresher RR, Acierno JS Jr, Shagoury JK, Bo-Abbas Y, Kuohung W, Schwino KM, Hendrick AG *et al.* 2003 The GPR54 gene as a regulator of puberty. *New England Journal of Medicine* **349** 1614–1627. (doi:10.1056/NEJMoa035322)
- Seminara SB, Dipietro MJ, Ramaswamy S, Crowley WF Jr & Plant TM 2006 Continuous human metastatin 45–54 infusion desensitizes G protein-coupled receptor 54-induced gonadotropin-releasing hormone release monitored indirectly in the juvenile male Rhesus monkey (*Macaca mulatta*): a finding with therapeutic implications. *Endocrinology* **147** 2122–2126. (doi:10.1210/en.2005-1550)
- Shahab M, Mastronardi C, Seminara SB, Crowley WF, Ojeda SR & Plant TM 2005 Increased hypothalamic GPR54 signaling: a potential mechanism for initiation of puberty in primates. *PNAS* **102** 2129–2134. (doi:10.1073/pnas.0409822102)
- Silverman AJ, Antunes JL, Ferin M & Zimmerman EA 1977 The distribution of luteinizing hormone-releasing hormone (LHRH) in the hypothalamus of the rhesus monkey. Light microscopic studies using immunoperoxidase technique. *Endocrinology* **101** 134–142. (doi:10.1210/endo-101-1-134)
- Smith JT, Cunningham MJ, Rissman EF, Clifton DK & Steiner RA 2005 Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* **146** 3686–3692. (doi:10.1210/en.2005-0488)
- Smith JT, Acohido BV, Clifton DK & Steiner RA 2006a KiSS-1 neurones are direct targets for leptin in the ob/ob mouse. *Journal of Neuroendocrinology* **18** 298–303. (doi:10.1111/j.1365-2826.2006.01417.x)
- Smith JT, Popa SM, Clifton DK, Hoffman GE & Steiner RA 2006b Kiss1 neurons in the forebrain as central processors for generating the preovulatory luteinizing hormone surge. *Journal of Neuroscience* **26** 6687–6694. (doi:10.1523/JNEUROSCI.1618-06.2006)
- Smith JT, Rao A, Pereira A, Caraty A, Millar RP & Clarke IJ 2008 Kisspeptin is present in ovine hypophysial portal blood but does not increase during the preovulatory luteinizing hormone surge: evidence that gonadotropes are not direct targets of kisspeptin *in vivo*. *Endocrinology* **149** 1951–1959. (doi:10.1210/en.2007-1425)
- Teles MG, Bianco SD, Brito VN, Trarbach EB, Kuohung W, Xu S, Seminara SB, Mendonca BB, Kaiser UB & Latronico AC 2008 A GPR54-activating mutation in a patient with central precocious puberty. *New England Journal of Medicine* **358** 709–715. (doi:10.1056/NEJMoa073443)
- Thompson EL, Patterson M, Murphy KG, Smith KL, Dhillo WS, Todd JF, Ghatei MA & Bloom SR 2004 Central and peripheral administration of kisspeptin-10 stimulates the hypothalamic–pituitary–gonadal axis. *Journal of Neuroendocrinology* **16** 850–858. (doi:10.1111/j.1365-2826.2004.01240.x)
- Todman MG, Han SK & Herbison AE 2005 Profiling neurotransmitter receptor expression in mouse gonadotropin-releasing hormone neurons using green fluorescent protein-promoter transgenics and microarrays. *Neuroscience* **132** 703–712. (doi:10.1016/j.neuroscience.2005.01.035)
- Topaloglu AK, Reimann F, Guclu M, Yalin AS, Kotan LD, Porter KM, Serin A, Mungan NO, Cook JR, Ozbek MN *et al.* 2009 TAC3 and TACR3 mutations in familial hypogonadotropic hypogonadism reveal a key role for neurokinin B in the central control of reproduction. *Nature Genetics* **41** 354–358. (doi:10.1038/ng.306)
- Welt CK, Chan JL, Bullen J, Murphy R, Smith P, DePaoli AM, Karalis A & Mantzoros CS 2004 Recombinant human leptin in women with hypothalamic amenorrhea. *New England Journal of Medicine* **351** 987–997. (doi:10.1056/NEJMoa040388)
- Wintermantel TM, Campbell RE, Porteous R, Bock D, Grone HJ, Todman MG, Korach KS, Greiner E, Perez CA, Schutz G *et al.* 2006 Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. *Neuron* **52** 271–280. (doi:10.1016/j.neuron.2006.07.023)
- Wu M, Dumalska I, Morozova E, van den PA & Alreja M 2009 Melanin-concentrating hormone directly inhibits GnRH neurons and blocks kisspeptin activation, linking energy balance to reproduction. *PNAS* **106** 17217–17222. (doi:10.1073/pnas.0908200106)
- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L & Friedman JM 1994 Positional cloning of the mouse obese gene and its human homologue. *Nature* **372** 425–432. (doi:10.1038/372425a0)

Received in final form 8 November 2010

Accepted 17 November 2010

Made available online as an Accepted Preprint
17 November 2010