Maternal protein restriction leads to enhanced hepatic gluconeogenic gene expression in adult male rat offspring due to impaired expression of the liver X receptor

Thin Xuan Vo^{1,3,4}, Andrew Revesz^{1,3,4}, Gurjeev Sohi^{2,3}, Noelle Ma^{1,2,3} and Daniel B Hardy^{1,2,3,4}

Departments of ¹Physiology and Pharmacology and ²Obstetrics and Gynecology, The University of Western Ontario, London, Ontario, Canada N6A 5C1

³The Children's Health Research Institute, London, Ontario, Canada N6A 5C1

⁴The Lawson Health Research Institute, London, Ontario, Canada N6A 5C1

Correspondence
should be addressed
to D B Hardy
Email
daniel.hardy@schulich.uwo.ca

Abstract

Epidemiological studies demonstrate that the link between impaired fetal development and glucose intolerance in later life is exacerbated by postnatal catch-up growth. Maternal protein restriction (MPR) during pregnancy and lactation in the rat has been previously demonstrated to lead to impaired glucose tolerance in adulthood, however the effects of protein restoration during weaning on glucose homeostasis are largely unknown. Recent *in vitro* studies have identified that the liver X receptor α (LXR α) maintains glucose homeostasis by inhibiting critical genes involved in gluconeogenesis including *G6pase* (*G6pc*), *11 β -Hsd1* (*Hsd11b1*) and *Pepck* (*Pck1*). Therefore, we hypothesized that MPR with postnatal catch-up growth would impair LXR α *in vivo*, which in turn would lead to augmented gluconeogenic LXR α -target gene expression and glucose intolerance. To examine this hypothesis, pregnant Wistar rats were fed a control (20%) protein diet (C) or a low (8%) protein diet during pregnancy and switched to a control diet at birth (LP). At 4 months, the LP offspring had impaired glucose tolerance. In addition, LP offspring had decreased LXR α expression, while hepatic expression of 11 β -HSD1 and G6Pase was significantly higher. This was concomitant with decreased binding of LXR α to the putative LXRE on *11 β -Hsd1* and *G6pase*. Finally, we demonstrated that the acetylation of histone H3 (K9,14) surrounding the transcriptional start site of hepatic *Lxr α* (*Nr1h3*) was decreased in LP offspring, suggesting MPR-induced epigenetic silencing of the *Lxr α* promoter. In summary, our study demonstrates for the first time the important role of LXR α in mediating enhanced hepatic gluconeogenic gene expression and consequent glucose intolerance in adult MPR offspring.

Key Words

- ▶ fetal programming
- ▶ liver X receptor
- ▶ gluconeogenesis
- ▶ nuclear receptors
- ▶ maternal protein restriction

Journal of Endocrinology
(2013) 218, 85–97

Introduction

Epidemiological evidence suggests that adverse events *in utero* (e.g. placental insufficiency-induced intrauterine growth restriction (PI-IUGR)) can permanently alter

physiological processes leading to hypertension and type 2 diabetes (Barker *et al.* 1993, Nilsson *et al.* 1997, Jaquet *et al.* 2000, Levitt *et al.* 2000, Huxley *et al.* 2007).

Previous animal models of maternal protein restriction (MPR) have consistently linked asymmetric IUGR (Desai & Hales 1997) with symptoms of type 2 diabetes long-term in the offspring. For example, Petrik *et al.* (1999) demonstrated a low protein diet during pregnancy and weaning induced a decrease in birth weight and disrupted pancreatic β -cell proliferation in the adult offspring. Other studies have found altered glucagon-stimulated and insulin-stimulated hepatic glucose output as well as reduced glucokinase expression and structural modifications in the livers of low protein offspring (Ozanne *et al.* 1996, Burns *et al.* 1997). In addition, Chamson-Reig *et al.* (2009) have demonstrated that low protein offspring have impaired glucose tolerance as early as 130 days of age in rat offspring. Thus, the evidence strongly suggests that maternal low protein-mediated IUGR in the rat predisposes the offspring to impaired glucose tolerance and a type 2 diabetes-like phenotype. However, the molecular mechanisms underlying these low protein-induced alterations in the output of hepatic glucose are not completely understood.

The liver X receptor (LXR) is a transcription factor belonging to the 1H subfamily of nuclear receptors. LXR exists as two isoforms: LXR α and LXR β . LXR α is mainly expressed in the liver, adipose tissue, macrophages and intestines (Apfel *et al.* 1994, Willy *et al.* 1995), while LXR β is ubiquitously expressed (Song *et al.* 1994). Endogenous ligands for LXR are mainly derivatives of cholesterol (i.e. oxysterols; Lehmann *et al.* 1997, Song & Liao 2000). Consequently, LXR has principally been implicated in regulating genes involved in the metabolism and transport of cholesterol (Lehmann *et al.* 1997, Venkateswaran *et al.* 2000) and in enhancing the expression of lipogenic enzymes (Repa *et al.* 2000). Recent studies have also demonstrated that LXR can silence genes involved in glucose production including phosphoenolpyruvate kinase (PEPCK) and glucose-6-phosphatase (G6Pase), both critical enzymes involved in the gluconeogenic pathway (Stulnig *et al.* 2002a,b, Cao *et al.* 2003). In addition, LXR has also been found to indirectly down-regulate hepatic glucose production through inhibition of the enzyme 11 β -hydroxysteroid dehydrogenase-1 (11 β -HSD1; Stulnig *et al.* 2002a). 11 β -HSD1 reduces inactive corticosteroids to their active form (e.g. 11-dehydrocorticosterone to corticosterone in the rodent). Since active corticosteroids are responsible for increased glucose production, LXR-mediated inhibition of 11 β -HSD1 would indirectly decrease glucose production.

Previous studies from our own laboratory have found that MPR leads to decreases in the expression of the LXR-target gene, *Cyp7a1*, the critical enzyme involved with

cholesterol catabolism. The decrease in *Cyp7a1* leads to hypercholesterolemia in male offspring by 4 months (Sohi *et al.* 2011). This was found to be due, in part, to repressive changes in histone modifications at the LXRE site of the *Cyp7a1* promoter. Other studies in mice have demonstrated that MPR leads to hypermethylation of the *Lxr α* (*Nr1h3*) promoter in association with decreased *Lxr α* mRNA in the liver tissue of embryonic day 19.5 fetuses, however the effect on posttranslational histone modifications surrounding *Lxr α* remains elusive (van Straten *et al.* 2010). While others and we have demonstrated that MPR can lead to long-term epigenetic alterations of LXR-target genes involved in cholesterol and lipid homeostasis, it is not known if LXR-target genes impairing hepatic gluconeogenesis are altered.

The aims of the present study were to examine whether MPR alters LXR α -mediated gluconeogenesis in the liver. Given the role of LXR α in lipid, glucose and cholesterol homeostasis, it is an attractive candidate in elucidating the molecular mechanisms underlying IUGR-related fetal programming. We hypothesized that decreased maternal protein availability during gestation would impair hepatic gluconeogenesis in the adult offspring through decreases in LXR α and aberrant activity of its target genes (*G6pase* (*G6pc*), *Pepck* (*Pck1*) and *11 β -Hsd1* (*Hsd11b1*)). Using a well-established model of MPR in rat pregnancy, we assessed the effects of a low protein diet in gestation on long-term glucose handling, LXR α activity, and the expression of hepatic LXR-target genes involved in gluconeogenesis. In the control group, dams were fed a 20% protein diet throughout life. Low protein dams received an 8% protein diet until birth of the offspring, followed by a 20% protein diet during the weaning period (until postnatal day 21). We decided to examine the effects of restoring protein immediately after birth as opposed to waiting until after the weaning period because we have already demonstrated that earlier restoration of protein promotes accelerated catch-up growth (Sohi *et al.* 2011). Moreover, postnatal accelerated growth of IUGR offspring has been demonstrated to exacerbate the effects of IUGR-related programming and reduce the lifespans of these offspring (Ozanne & Hales 2004, Chen *et al.* 2009, Tarry-Adkins *et al.* 2013).

Materials and methods

Animal experiments and dietary regime

All procedures were performed in accordance with the guidelines set by the Canadian Council of Animal Care and upon approval of the Animal Care Committee of the

University of Western Ontario. Male and female Wistar rats at breeding age (250 g) were purchased from Charles River (La Salle, St-Constant, QC, Canada) and were allowed to acclimatize to their new environment for 2 weeks. Rats were housed at room temperature on a 12 h light:12 h darkness cycle. Females were housed in separate cages and were cohabitated with a male for mating upon entering proestrus. Conception was confirmed by the presence of sperm in the vaginal smear the following day.

Dams and offspring received isocaloric diets (Bio-Serv, Frenchtown, NJ, USA) varying in protein composition, depending on their experimental group. Briefly, the control offspring and dams received 20% protein throughout life. Protein-restricted dams received low protein chow (8%) throughout gestation and then restored on a 20% protein chow immediately after birth (herein termed 'LP'). All diets and water were administered *ad libitum*. Previous studies by our laboratory have demonstrated that the food intake between both offspring groups is practically identical (Sohi *et al.* 2011).

At embryonic day 19, a subset of dams (three control dams; four LP dams) was sacrificed and livers from the fetuses were extracted. The livers were flash frozen for further molecular analysis. The other subset of dams (four control dams; four LP dams) delivered spontaneously. All litters with less than ten pups were arbitrarily culled down to 9–10 pups to ensure a consistent litter size per dam.

After their intraperitoneal glucose tolerance tests (IPGTT) at postnatal days 120–125, all offspring were sacrificed using a lethal dose (50 mg/kg) of Euthanyl forte pentobarbital sodium (Bimeda-MTC, Cambridge, ON, Canada) at postnatal day 130. This age was chosen because previous studies have demonstrated that in other models of protein restriction, impaired glucose tolerance was not observed earlier than 4 months (Chamson-Reig *et al.* 2009). Following sacrifice, liver and blood were immediately extracted and flash frozen at -80°C for molecular analysis. We did not examine the female offspring in this study to prevent confounding factors related to their estrus cycle and hormone profile. More importantly, the maternal low protein model has been demonstrated to exhibit early life programming effects in a sexually dimorphic manner, which was not the focus of this investigation (Guan *et al.* 2005, Chamson-Reig *et al.* 2009, Sohi *et al.* 2011). For molecular analysis, one-to-two male pups from each of four separate dams were arbitrarily chosen. All available male pups were used for the IPGTT.

Glucose tolerance tests

At postnatal days 120–125, male offspring were subject to an IPGTT. Prior to the IPGTT, the animals were fasted overnight for 14–16 h. Animals were awake throughout the experiment. Blood glucose measurements were obtained using a Bayer Breeze 2 Blood Glucose Meter (Bayer). Fasted blood glucose levels were obtained prior to the glucose injection. Animals then received 2 g/kg of glucose via injection into the i.p. cavity. Blood glucose was sampled at the tail vein at $t=0, 5, 10, 15, 30, 60$ and 120 min. Area under the curve of each animal was calculated using (GraphPad Prism Software, San Diego, California, USA, www.graphpad.com). IPGTTs were performed on six control males and ten LP males.

Quantitative real-time PCR for gene expression analysis

Total RNA was extracted from the medial lobe of offspring livers at embryonic day 19 and postnatal day 130 as previously described, using the one-step TRIzol (Invitrogen) method (Sohi *et al.* 2011). Total RNA was subsequently treated with deoxyribonuclease to eliminate contaminating DNA. Four micrograms of total RNA were then reverse transcribed to cDNA using random primers and Superscript II RNase H-reverse transcriptase (Invitrogen). TaqMan probes and sequences for the genes of interest (*11 β -Hsd1*, *G6pase*, *Lxr α* , *Pepck* and *β -actin (Actb)*) and TaqMan Universal Master Mix were obtained from Invitrogen. Quantitative analysis of mRNA expression was measured using the Bio-Rad CFX384 Real Time System. The cycling conditions were as follows: polymerase activation (95°C for 10 min) followed by 40 cycles of denaturing (95°C for 15 s) and annealing (60°C for 1 min). The cycle threshold was set where the exponential increase in amplification was equivalent between all samples. Relative fold changes were calculated using the comparative cycle times (Ct) method with β -actin as the reference gene. ΔCt values for each probe set were standardized to the experimental samples with the lowest transcript abundance (highest Ct value). The relative abundance of each primer set compared with calibrator was determined by the formula, $2^{\Delta\Delta\text{Ct}}$, where $\Delta\Delta\text{Ct}$ was the standardized Ct value.

Tissue protein extraction and western immunoblotting

Tissue protein was extracted from the medial lobe of snap-frozen offspring livers using a lysis buffer solution (pH 7.4, Tris-HCl 50 mM, NP-40 1%, sodium deoxycholate 0.25%,

NaCl 150 mM, EDTA 1 mM, NaF 50 mM, Na₃VO₄ 1 mM and β-glycerophosphate 25 mM). Prior to tissue homogenization, a mini protease inhibitor tablet was added to the lysis buffer.

First, a small chunk of snap-frozen liver was added to 600 μl of RIPA buffer. The tissue was then homogenized with the IKA T10 Basic S1 Dispersing Tool (IKA Works, Inc., Wilmington, NC, USA) for 10–15 s at speed 6. After letting the homogenized tissue sit on ice for 5 min, the tissue was then sonicated. Following sonication, the tissue was rotated at 4 °C for 5 min and then centrifuged for 15 min at 300 *g* and 4 °C. The supernatant was retained for further centrifugation at 20 000 *g* for 20 min at 4 °C. The final supernatant was retained for protein quantification and western immunoblotting.

Equal concentrations of total protein were normalized using a colorimetric BCA Protein Assay (Pierce Corp., Madison, WI, USA). Proteins were then fractionated in 17-well gradient polyacrylamide gels (Invitrogen) and transferred onto PVDF membrane (Millipore, Etobicoke, ON, Canada). Amido black and Coomassie brilliant blue staining confirmed sufficient transfer of proteins onto the membrane.

Immunoblots were probed using LXRα (LXR; 1:1000; cat no. sc-13068), PEPCK (1:2000; cat no. sc-32879), G6Pase-α (1:1000; cat no. sc-25840), PI3-kinase p85α (Z-8; 1:1000; cat no. sc-423) and 11β-HSD1 (1:800; cat no. sc-20175) all from Santa Cruz Biotechnology. In addition, p-Akt1 (Serine 473; 1:1000; cat no. ab66138), p-Akt (Threonine 308; 1:500; cat no. 4796) and Akt (1:125; cat no. ab6076) antibodies used to assess hepatic insulin sensitivity were purchased from Abcam, Inc. (Cambridge, MA, USA). In addition, we also assessed insulin sensitivity by using antibodies against p-IRS-1 (Serine 302; 1:500; cat no. 2384), p-IRS-1 (Serine 1101; 1:500; cat no. 2385) and IRS-1 (1:500; cat no. 2382) all purchased from Cell Signaling (Danvers, MA, USA). Monoclonal HRP conjugated β-actin (1:50 000; cat no. A3854; Sigma-Aldrich) diluted in 5% milk-TBS-Tween-20 (0.1%) buffer and HRP conjugated donkey anti-rabbit IgG (1:10 000, cat no. 711-035-152; Jackson ImmunoResearch Laboratories, West Grove, PA, USA) diluted in 5% milk-TBS-Tween-20 (0.1%) buffer were used as the secondary antibodies. Finally, immunostained bands were then visualized using an ECL detection system (Thermo Scientific, Waltham, MA, USA).

Chromatin immunoprecipitation

Chromatin was extracted from the medial lobe of offspring livers as previously described (Sohi *et al.* 2011). Briefly, a

small piece of snap-frozen liver was homogenized and incubated with 1% formaldehyde for 10 min at room temperature to cross-link proteins and DNA. Cross-linking was terminated by the addition of glycine (0.125 M, final concentration). The liver tissue was washed once with cold PBS and placed in 500 μl of SDS lysis buffer (Millipore) with a protease inhibitor cocktail (Roche). The lysates were sonicated on ice to produce sheared, soluble chromatin. The lysates were diluted 10 times with the addition of chromatin immunoprecipitation (ChIP) dilution buffer (Millipore) and aliquoted to 400 μl amounts. Each of the aliquots was precleared with protein A/G Plus agarose beads (40 μl; Millipore) at 4 °C for 30 min. The samples were centrifuged at 20 000 *g* to pellet the beads, and the supernatant containing the sheared chromatin was placed in new tubes. The aliquots were incubated with 4 μg of antibodies against RNA polymerase II (cat no. 05-623B; Millipore), trimethylated histone H3 lysine 4 (K4; cat no. ab1012; Abcam, Inc.), acetylated histone H3 lysine 9,14 (K9,14; cat no. 05-399; Millipore), trimethylated histone H3 lysine 9 (K9; cat no. 07-442; Millipore) and ChIP-grade LXRα (cat no. sc-13068x; Santa Cruz Biotechnology) at 4 °C overnight. Two aliquots were reserved as 'controls' – one incubated without antibody and the other with nonimmune IgG (Millipore). Protein A/G Plus agarose beads (60 μl) were added to each tube, the mixtures incubated for 1 h at 4 °C and the immune complexes collected by centrifugation. The beads containing the immunoprecipitated complexes were washed sequentially for 5 min in wash buffer I (20 mM Tris-HCl, pH 8.1, 2 mM EDTA, 0.1% SDS, 1% Triton X-100 and 150 mM NaCl), wash buffer II (same as I, except containing 500 mM NaCl), wash buffer III (10 mM Tris-HCl, pH 8.1, 1 mM EDTA, 1% NP-40, 1% deoxycholate and 0.25 M LiCl), and in 2× TE buffer. The beads were eluted with 250 μl elution buffer (1% SDS, 0.1 mM NaHCO₃ + 20 g salmon sperm DNA (Sigma-Aldrich)) at room temperature. This was repeated once and eluates were then combined. Cross-linking of the immunoprecipitated chromatin complexes and 'input controls' (10% of the total soluble chromatin) was reversed by heating the samples at 65 °C for 4 h. Proteinase K (15 μg; Invitrogen) was added to each sample in buffer (50 mM Tris-HCl, pH 8.5, 1% SDS and 10 mM EDTA) and incubated for 1 h at 45 °C. The DNA was purified by phenol-chloroform extraction and precipitated in EtOH overnight at 20 °C. Samples and 'input controls' were diluted in 10–100 μl TE buffer just prior to quantitative real-time PCR (qRT-PCR).

Putative LXR-binding sites (threshold of 0.7) on the promoters of *G6pase* and *11β-Hsd1* were determined using

the MatInspector Software (Genomatix, Munich, Germany). The MatInspector Software was used to match the LXR consensus binding site (AGGTCA_DR-4_AGGTCA; Willy *et al.* 1995) with putative transcription factor binding sequences based on algorithms as described by Cartharius *et al.* (2005). qRT-PCR was employed using forward (5'-GGTCACTGCATGATCACAGG-3') and reverse (5'-CCTTGAATCCAGAATGCTC-3') primers that amplify a -35 to +92 bp region encompassing the rat *G6pase* LXRE site (+22 to +46 bp), and forward (5'-TTCGCCAACTCTGACCTCT-3') and reverse (5'-ACAGGTTTGGCCTGAT-GT-3') primers that amplify a -115 to -7 bp region encompassing the rat *11 β -Hsd1* LXRE site (-114 to -90 bp) (PE Applied Biosystems). The LXR α (gene: *Nr1h3*) transcriptional start site (TSS) was found using the Ensembl Genome Browser (<http://www.ensembl.org>). Forward (5'-GGCTTCACTGGTTGATCCAT-3') and reverse (5'-AGGGGGTTGATTCTTGAGGT-3') primers were designed to amplify the -135 to +144 bp region surrounding the +1 bp TSS of Lxr α . Recent evidence indicates that there is epigenetic regulation in the CG-rich regions of the Lxr α promoter around the TSS in another rodent model of MPR (van Straten *et al.* 2010). Thus, primers around the promoter were used to examine the binding of RNA polymerase II, acetylation of histone H3 (K9,14), methylation of histone H3 (K4) and trimethylation of histone H3 (K9) at the TSS of Lxr α .

The aforementioned constructed ChIP primers were then used in conjunction with Sso-Fast EvaGreen Supermix (Bio-Rad) to perform qRT-PCR. Similar to the gene expression assays, the relative abundance of the immunoprecipitated chromatin compared with input chromatin was determined using the $2^{\Delta\Delta Ct}$ method.

Statistical analysis

All data are represented as a mean of an arbitrary value $\pm$ S.E.M. IPGTT, areas under the curve, qRT-PCR (including ChIP) and quantified western immunoblot bands were analyzed using the unpaired Student's *t*-test. All data with a *P* value <0.05 were considered statistically significant.

Results

MPR leads to impaired glucose tolerance at 4 months of age in male offspring

At 4 months of age, all of the male offspring underwent an IPGTT to assess fasted glucose tolerance after an administered glucose load. Resting levels of glucose were not

significantly different between control and LP animals. After administration of the glucose (2 g/kg), measured blood glucose levels were significantly elevated ($P<0.05$) in LP animals at the 10-, 15-, 30- and 60-min time points (Fig. 1A). Blood glucose levels in the LP animals returned to the same levels as the control animals by the 120-min time point. At the end of the experiment, both control and LP animals had similar blood glucose levels. The area under the curve for the LP animals was increased by 32.8% ($P<0.05$) compared with the control animals (Fig. 1B), further indicating impaired glucose tolerance at 4 months of age. Although we did not perform insulin tolerance tests, hepatic insulin sensitivity was assessed through western immunoblot detection of phosphorylated-Akt1 (S473 and T308), the p85 subunit of phosphoinositide 3-kinase (p85), and the phosphorylated IRS-1 (S302 and S1101), all markers of insulin sensitivity (Guo *et al.* 2009,

Figure 1

(A) Intraperitoneal GTT (2 g/kg) administered to fasted male offspring at 4 months of age. Control and LP animals were analyzed together at each time point ($t=0, 5, 10, 15, 30, 60$ and 120 min) using the Student's unpaired *t*-test. (B) Area under the curve of control and LP animals. Area under the curve was calculated using GraphPad Prism Software. (Control $n=6$, LP $n=10$). Results are expressed as the mean $\pm$ S.E.M. *Statistically significant ($P<0.05$).

Valverde & Gonzalez-Rodriguez 2011). Protein expression of these proteins was unchanged between control and LP animals, suggesting no difference in insulin sensitivity between the experimental groups at 4 months of age (Supplementary Figure 1A and B, see section on supplementary data given at the end of this article).

The steady-state levels of hepatic *Lxrα* mRNA are decreased, concomitant with an increase in *G6pase* and *11β-Hsd1* mRNA in LP animals by 4 months of age

Given the LP animals exhibited glucose intolerance at 4 months of age, we subsequently investigated the expression of hepatic LXRα and its target genes involved in gluconeogenesis. To determine differences in *in vivo* hepatic mRNA levels of *Lxrα*, *G6pase*, *11β-Hsd1* and *Pepck* at 4 months of age in the male offspring, qRT-PCR was employed with TaqMan probes for each gene. *Lxrα* mRNA was significantly decreased by 45% ($P < 0.05$) in the LP offspring, while *Pepck* mRNA was unchanged between groups (Fig. 2). Hepatic *G6pase* and *11β-Hsd1* mRNA were significantly increased ($P < 0.05$, 1.6-fold) in the LP offspring (Fig. 2).

The levels of hepatic LXRα protein are decreased, concomitant with an increase in G6PASE and 11β-HSD1 protein levels in LP animals by 4 months of age

To assess the effect of a maternal low protein diet on the protein levels of LXRα and LXR-target genes in 4-month-old offspring, we performed western immunoblotting to determine if there would be similar trends to what was observed in the steady-state mRNA levels. At 4 months of age, LXRα protein expression was decreased by 40% ($P < 0.05$), while both G6PASE and 11β-HSD1 protein levels were increased ($P < 0.05$, 1.5- and 1.6-fold respectively) in the LP animals compared with the control animals (Fig. 3). PEPCK protein expression was not different between the two groups.

LXRα binding to the LXRE on the promoters of *G6pase* and *11β-Hsd1* is decreased by 4 months of age in the LP offspring

To investigate whether the changes in the expression of G6Pase and 11β-HSD1 between the control and LP offspring were due to the decreased binding of LXRα to the promoters of *G6pase* and *11β-Hsd1*, we employed ChIP

Figure 2

The effect of LP on *in vivo* hepatic levels of (A) *Lxrα* mRNA, (B) *Pepck* mRNA, (C) *G6pase* mRNA and (D) *11β-Hsd1* mRNA in control and LP offspring at 4 months of age. Data were quantified from qRT-PCR (TaqMan) and the

ΔΔCt method. Data are represented as arbitrary values and were analyzed using the Student's unpaired *t*-test. Results are expressed as the mean ± S.E.M. *Statistically significant. $n = 4-6$ per experimental group.

Figure 3

The effect of LP on the *in vivo* hepatic levels of (A) LXRα protein (50 kDa), (B) PEPCK protein (62 kDa), (C) G6PASE protein (36 kDa) and (D) 11β-HSD1 (34 kDa) protein in control and LP offspring at 4 months of age. Data were obtained from western immunoblotting experiments. Immunoblots were

quantified using densitometry and normalized to β-ACTIN (42 kDa) protein expression. Data are represented as arbitrary values. Results are expressed as the mean ± s.e.m. *Statistically significant. *n* = 4–6 per experimental group.

to immunoprecipitate LXRα. After using MatInspector (Genomatix) to find putative LXREs on the promoters of *G6pase* and *11β-Hsd1*, qRT-PCR was employed to examine LXRα binding at these putative LXRE sites. By 4 months of age, the LP animals exhibited a marked decrease in the binding of LXRα to the promoter of *11β-Hsd1* (45% decrease) and *G6pase* (50% decrease) compared with the control animals ($P < 0.05$) (Fig. 4). The nonspecific binding of IgG was tested and found to be minimal (Ct value > 34, data not shown).

Acetylation of lysine residues 9 and 14 on histone H3 is decreased surrounding the TSS of *Lxrα* in LP offspring by 4 months of age

We further employed ChIP to examine the epigenetic regulation of *Lxrα* at its TSS. By immunoprecipitating chromatin with antibodies specific to RNA polymerase II, trimethylated histone H3 (K9), acetylated histone H3 (K9,14) and trimethylated histone H3 (K4) we were able to examine the transcriptional and epigenetic regulation of *Lxrα* in our model of MPR. Using primers specific to the –144 to +134 region of the *Lxrα* gene promoter and qRT-PCR, we found a significant 45% reduction ($P < 0.05$)

in the acetylation of histone H3 (K9,14), a hallmark of chromatin silencing, near the TSS of *Lxrα* (Fig. 5C). While not significant, we also found a decreasing trend in the recruitment of RNA polymerase II binding and histone H3 trimethylation (K4) at the same site (Fig. 5A and B). Again, the nonspecific binding of IgG was tested and found to be minimal (Ct value > 34, data not shown). These results, in combination, support the notion that *Lxrα* is transcriptionally and epigenetically silenced long-term in our MPR model of IUGR.

The steady-state levels of hepatic *Lxrα* mRNA are unchanged between control and LP offspring concomitant with a decrease in *G6Pase* and *11β-Hsd1* mRNA in LP animals at embryonic day 19

To assess the direct effects of the LP diet on *Lxrα* and LXR-target gene expression during gestation and prior to birth, we analyzed the livers of fetuses sacrificed at embryonic day 19. qRT-PCR was employed with TaqMan probes for (*Lxrα*, *G6pase* and *11β-HSD1* (*Pepck* could not be detected in our embryonic liver tissue). At embryonic day 19, there were no differences in *Lxrα* mRNA expression between the control and LP animals. However,

Figure 4

The effect of LP on the *in vivo* hepatic binding of LXRα to the promoters of (A) *G6pase* (+22 to +46 bp) and (B) *11β-Hsd1* (-114 to -90 bp) in control and LP offspring at 4 months of age. Putative LXRE sites were determined using the MatInspector Software from Genomatix. Livers were immunoprecipitated with antibodies specific to LXRα. Quantification was performed using qRT-PCR (Sso-Fast EvaGreen) with primers specific to the proposed LXRE sites. The relative amount of immunoprecipitated genomic DNA was normalized to total genomic DNA. Data are represented as arbitrary values using the $\Delta\Delta C_t$ method. Results are expressed as the mean $\pm$ S.E.M. *Statistically significant. $n=4-6$ per experimental group.

both *G6pase* and *11β-Hsd1* mRNA expression were decreased in LP offspring compared with control offspring ($P<0.05$; Fig. 6).

Discussion

Our present study demonstrates that male offspring of LP dams exhibit increased expression of hepatic gluconeogenic genes due to aberrant expression of hepatic LXRα. This is of great interest considering that previous studies have indicated that MPR leads to glucose dysregulation (Ozanne *et al.* 1996, Burns *et al.* 1997, Chamson-Reig *et al.* 2009). We present evidence for the first time that suppressed expression of LXRα may mediate the enhanced transcription of the gluconeogenic genes *G6pase* and *11β-Hsd1* due to its decreased binding on these promoters, ultimately removing its ability to suppress hepatic gluconeogenesis (Stulnig *et al.* 2002a, Cao *et al.* 2003, Laffitte *et al.* 2003, Mitro *et al.* 2007).

Given placental insufficiency in humans can produce protein deficiency in the fetus (Crosby 1991), this LP model shares features in common with PI-IUGR (Ross & Beall 2008). Previous studies done in our laboratory with the same cohort of animals have already demonstrated that LP offspring exhibit a 15% lower fetal to placental weight ratio and a 40% decreased fetal liver to body weight ratio at embryonic day 19 (Sohi *et al.* 2011). While switching the low protein offspring to a control diet at weaning led to glucose intolerance (Chamson-Reig *et al.* 2009), little is known about how catch-up growth due to early restoration of protein (Sohi *et al.* 2011) influences their hepatic glucose handling by adulthood. In our LP model, after switching to a control (20% protein) diet at birth, the animals exhibited full catch-up growth by 3 weeks of age (Sohi *et al.* 2011). Moreover, by 4 months, these offspring exhibited impaired glucose tolerance with no evidence of hepatic insulin insensitivity. Given that glucose intolerance precedes insulin resistance, it is likely that these MPR offspring will develop insulin resistance at a later time point. Interestingly, the impaired glucose tolerance was similar, not worse, to low protein offspring at 4 months whereby their diet was switched to 20% at weaning (Chamson-Reig *et al.* 2009). Collectively, both studies further support of the main tenets of the Thrifty Phenotype hypothesis (Hales & Barker 1992).

We previously found that LXRα expression and binding could be influenced by maternal diet by 3 weeks of age in the offspring (Sohi *et al.* 2011), but the expression of LXRα at 4 months was unknown. In this study we demonstrated that in LP offspring, hepatic *Lxrα* mRNA and

Figure 5

The effect of LP on the *in vivo* transcriptional and epigenetic regulation of the *Lxrα* TSS (–135 to +144 bp) at 4 months of age. (A) Binding of RNA polymerase II to the *Lxrα* TSS, (B) trimethylation of histone H3 lysine 4, (C) acetylation of histone H3 lysine 9 and 14 and (D) trimethylation of histone H3 lysine 9. Primers were designed based on sequencing from Ensembl. Livers were immunoprecipitated with antibodies specific to RNA polymerase II, trimethylated histone H3 (K4), acetylated histone H3 (K9,14),

and trimethylated histone H3 (K9). Quantification was performed using qRT-PCR (Sso-Fast EvaGreen) with primers specific to the proposed LXR element sites. The relative amount of immunoprecipitated genomic DNA was normalized to total genomic DNA. Data are represented as arbitrary values using the $\Delta\Delta C_t$ method. Results are expressed as the mean $\pm$ S.E.M. *Statistically significant. $n=4-6$ per experimental group.

protein were decreased at 4 months of age compared with control offspring. Given aberrant LXR α expression and activity can alter the expression of genes involved in hepatic gluconeogenesis (e.g. *Pepck*, *G6pase* and *11 β -Hsd1*; Stulnig *et al.* 2002a,b, Cao *et al.* 2003, Laffitte *et al.* 2003), we next examined whether the protein expression of these LXR-target genes was altered in LP offspring. At 4 months, we found increases in the steady-state mRNA and protein levels of G6PASE in LP male rats. This is of great interest considering that this LXR-target gene is responsible for the final catalytic step of gluconeogenesis, the conversion of glucose-6-phosphate to glucose. Moreover, overproduction of G6Pase does not necessarily lead to increases in fasting glucose levels, which we also observed, but it does lead to an enhanced glucose response (e.g. a greater area under the curve during GTT; Trinh *et al.* 1998). To directly

implicate whether alterations in LXR α expression influenced the binding of LXR α to the promoter of *G6pase*, we then employed ChIP to examine the *in vivo* binding of LXR α to its putative LXRE on the promoter of *G6pase*. At 4 months of age, we observed a decrease in the binding of LXR α to the LXRE site (+22 to +46) of the *G6pase* promoter. These data suggests that the increase of *G6pase* expression seen in protein restricted offspring is at least partly due to the decreased binding of LXR α to the putative *G6pase* promoter. Overexpression of PEPCCK has also been demonstrated to impair glucose tolerance and leads to noninsulin-dependent diabetes (Valera *et al.* 1994), however we did not find any significant alterations in *Pepck* mRNA or protein. This is in contrast to other studies whereby hepatic PEPCCK activity increased in 3-week-old and 11-month-old offspring fed a low protein

Figure 6

The effect of LP on *in vivo* hepatic levels of (A) *Lxrα* mRNA, (B) *G6pase* mRNA and (C) *11β-Hsd1* mRNA in control and LP offspring at embryonic day 19. Data were quantified from qRT-PCR (TaqMan) and the $\Delta\Delta C_t$ method. Data are represented as arbitrary values and were analyzed using Student's unpaired *t*-test. Results are expressed as the mean $\pm$ S.E.M. *Statistically significant. *n* = 4–6 per experimental group.

diet during gestation (Desai *et al.* 1995), and low protein offspring fed a high sucrose diet (500 g/kg) *postpartum* (Burdge *et al.* 2007). The difference in the former study may be due to the fact that the offspring were cross-fostered to dams not subjected to a low protein diet, potentially leading to even greater catch-up growth.

11 β -HSD1 may also play a role in impairing glucose homeostasis in our LP model. Our study demonstrated an

increase in 11 β -Hsd1 mRNA along with elevated 11 β -HSD1 protein levels in the LP offspring. Interestingly, while previous nutrient restriction models have demonstrated no change in 11 β -HSD1 expression in the adipose tissue of adult rat offspring (Dutriez-Casteloot *et al.* 2008), its expression in the liver was not examined. Similar to *G6pase*, we proposed that a decrease in LXR α expression and binding would lead to the loss of inhibitory action on the 11 β -Hsd1 promoter and a subsequent increase in 11 β -Hsd1 gene expression. Our ChIP experiments confirmed our speculation by demonstrating a decrease in LXR α binding to the putative LXRE (–114 to –90) on 11 β -Hsd1. With increased expression of 11 β -HSD1, it is conceivable that there would be enhanced conversion of inactive glucocorticoids to active glucocorticoids. Since glucocorticoids have stimulatory effects on the expression of gluconeogenic genes such as *G6pase* and *Pepck* (Sasaki *et al.* 1984, Lange *et al.* 1994, Lin *et al.* 1998), the sustained overproduction of glucocorticoids would lead to an augmented glucose response, as observed in the IPGTTs. Collectively, it is likely that the overproduction of G6Pase may not only occur due to the direct actions of LXR α , but also indirectly through enhanced 11 β -HSD1 expression.

Previous studies strongly suggest the role of epigenetics in mediating the effects of fetal programming long-term into adulthood (Rees *et al.* 2000, Lillycrop *et al.* 2005, Burdge *et al.* 2007, Park *et al.* 2008, van Straten *et al.* 2010). By 4 months of age, we demonstrated that LP males exhibited significantly decreased acetylation of histone H3 (K9,14) associated with trends of decreased RNA polymerase II recruitment and decreased methylation of histone H3 (K4) surrounding the promoter of *Lxrα*. Considering acetylation of histone H3 (K9,14) and methylation of histone H3 (K4) are both known to be hallmarks of chromatin opening (Jenuwein & Allis 2001, Santos-Rosa *et al.* 2002, Yan & Boyd 2006), our findings suggest that the *Lxrα* TSS is silenced through epigenetic mechanisms by adulthood. These findings are congruent with the decreased levels of *Lxrα* mRNA and protein expression observed. We previously demonstrated that MPR leads to long-term decreases in histone H3 acetylation (K9,14) surrounding the promoter of the LXR-target gene *Cyp7a1* resulting in hypercholesterolemia in these offspring (Sohi *et al.* 2011). To address whether the low protein diet itself directly alters hepatic LXR α and LXR-target genes *in vivo*, we measured their fetal expression (embryonic day 19) during the low protein insult. Interestingly, the low protein diet impaired these hepatic genes involved in gluconeogenesis without changes to LXR α expression. This suggests that the augmented expression of LXR α , G6Pase

and 11 β -HSD1 observed in adulthood is more likely due to the indirect actions of the low protein diet, namely, a protein mismatch in postnatal life associated with rapid catch-up growth.

In view of the fact that LXR α suppresses glucose production, it may be a suitable target as a therapeutic intervention to prevent glucose intolerance. Animal studies have widely demonstrated that the administration of LXR agonists (e.g. GW3965 and T0901317) leads to improved glucose tolerance (Cao *et al.* 2003, Laffitte *et al.* 2003). Given that during the newborn period in the rat there is a high rate of replication, neogenesis and apoptosis leading to extensive liver remodeling (Cascio & Zaret 1991), this period represents a critical, but opportune window for therapy designed to improve hepatic growth and function long-term. For example, it has been demonstrated in IUGR rats derived from uterine artery-ligated dams that neonatal administration of Exendin-4 (a GLP-1 analog) prevented the development of hepatic oxidative stress and insulin resistance (Raab *et al.* 2009). Therefore it is plausible that LXR α agonists, administered in neonatal life, a period of liver plasticity, may prevent the development of glucose intolerance long-term through activation of hepatic LXR α . While LXR agonists appear promising, the negative effects of LXR α activation must also be considered, given that it can activate lipogenesis through increased expression of fatty acid synthase, acetyl-CoA carboxylase and the master lipid regulator, sterol regulatory element binding protein-1c (Steffensen & Gustafsson 2004).

In summary, our study demonstrates for the first time the role of LXR α in mediating the transcriptional regulation of hepatic gluconeogenic genes in our rat LP model. In these offspring, decreased expression of hepatic LXR α reduced the transcriptional inhibition of hepatic G6Pase and 11 β -HSD1. This increased expression of G6Pase and 11 β -HSD1 in LP offspring would contribute, in part, to the aberrant glucose handling observed in these animals. Given the role of hepatic LXR α in reducing glucose production, it serves as a possible therapeutic target of intervention due to its anti-diabetic properties. Further studies will be necessary to find a suitable balance between anti-diabetic and lipogenic actions of LXR α before it could be considered as an ideal candidate for preventing glucose intolerance in IUGR offspring.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/JOE-13-0055>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Canadian Institutes of Health Research (grant number R4228A10) and Natural Sciences and Engineering Research Council of Canada.

Author contribution statement

T X V designed and performed all the experiments in this study and was the primary author in writing this manuscript. A R and N M assisted in the glucose tolerance tests and tissue collection, while G S assisted in the molecular analyses experiments (qRT-PCR and western immunoblotting). D B H assisted in the designing of the experiments and in the preparation of this manuscript.

Acknowledgements

We thank Dr Lin Zhao (Department of Obstetrics and Gynaecology, Western University) for his technical assistance.

References

- Apfel R, Benbrook D, Lernhardt E, Ortiz MA, Salbert G & Pfahl M 1994 A novel orphan receptor specific for a subset of thyroid hormone-responsive elements and its interaction with the retinoid/thyroid hormone receptor subfamily. *Molecular and Cellular Biology* **14** 7025–7035.
- Barker DJ, Hales CN, Fall CH, Osmond C, Phipps K & Clark PM 1993 Type 2 (non-insulin-dependent) diabetes mellitus, hypertension and hyperlipidaemia (syndrome X): relation to reduced fetal growth. *Diabetologia* **36** 62–67. (doi:10.1007/BF00399095)
- Burdge GC, Slater-Jefferies J, Torrens C, Phillips ES, Hanson MA & Lillycrop KA 2007 Dietary protein restriction of pregnant rats in the F0 generation induces altered methylation of hepatic gene promoters in the adult male offspring in the F1 and F2 generations. *British Journal of Nutrition* **97** 435–439. (doi:10.1017/S0007114507352392)
- Burns SP, Desai M, Cohen RD, Hales CN, Iles RA, Germain JP, Going TC & Bailey RA 1997 Gluconeogenesis, glucose handling, and structural changes in livers of the adult offspring of rats partially deprived of protein during pregnancy and lactation. *Journal of Clinical Investigation* **100** 1768–1774. (doi:10.1172/JCI119703)
- Cao G, Liang Y, Broderick CL, Oldham BA, Beyer TP, Schmidt RJ, Zhang Y, Staybrook KR, Suen C, Otto KA *et al.* 2003 Antidiabetic action of a liver X receptor agonist mediated by inhibition of hepatic gluconeogenesis. *Journal of Biological Chemistry* **278** 1131–1136. (doi:10.1074/jbc.M210208200)
- Cartharius K, Frech K, Grote K, Klocke B, Haltmeier M, Klingenhoff A, Frisch M, Bayerlein M & Werner T 2005 MatInspector and beyond: promoter analysis based on transcription factor binding sites. *Bioinformatics* **21** 2933–2942. (doi:10.1093/bioinformatics/bti473)
- Cascio S & Zaret KS 1991 Hepatocyte differentiation initiates during endodermal–mesenchymal interactions prior to liver formation. *Development* **113** 217–225.
- Chamson-Reig A, Thyssen SM, Hill DJ & Arany E 2009 Exposure of the pregnant rat to low protein diet causes impaired glucose homeostasis in the young adult offspring by different mechanisms in males and

- females. *Experimental Biology and Medicine* **234** 1425–1436. (doi:10.3181/0902-RM-69)
- Chen JH, Martin-Gronert MS, Tarry-Adkins J & Ozanne SE 2009 Maternal protein restriction affects postnatal growth and the expression of key proteins involved in lifespan regulation in mice. *PLoS ONE* **4** e4950. (doi:10.1371/journal.pone.0004950)
- Crosby WM 1991 Studies in fetal malnutrition. *American Journal of Diseases of Children* **145** 871–876.
- Desai M & Hales CN 1997 Role of fetal and infant growth in programming metabolism in later life. *Biological Reviews of the Cambridge Philosophical Society* **72** 329–348. (doi:10.1017/S0006323196005026)
- Desai M, Crowther NJ, Ozanne SE, Lucas A & Hales CN 1995 Adult glucose and lipid metabolism may be programmed during fetal life. *Biochemical Society Transactions* **23** 331–335.
- Dutriez-Casteloot I, Breton C, Coupe B, Hawchar O, Enache M, Dickes-Coopman A, Keyzer Y, Deloof S, Lesage J & Vieau D 2008 Tissue-specific programming expression of glucocorticoid receptors and 11 β -HSDs by maternal perinatal undernutrition in the HPA axis of adult male rats. *Hormone and Metabolic Research* **40** 257–261. (doi:10.1055/s-2008-1058064)
- Guan H, Arany E, van Beek JP, Chamson-Reig A, Thyssen S, Hill DJ & Yang K 2005 Adipose tissue gene expression profiling reveals distinct molecular pathways that define visceral adiposity in offspring of maternal protein-restricted rats. *American Journal of Physiology. Endocrinology and Metabolism* **288** E663–E673. (doi:10.1152/ajpendo.00461.2004)
- Guo S, Copps KD, Dong X, Park S, Cheng Z, Pocai A, Rossetti L, Sajan M, Farese RV & White MF 2009 The Irs1 branch of the insulin signaling cascade plays a dominant role in hepatic nutrient homeostasis. *Molecular and Cellular Biology* **29** 5070–5083. (doi:10.1128/MCB.00138-09)
- Hales CN & Barker DJ 1992 Type 2 (non-insulin-dependent) diabetes mellitus: the Thrifty Phenotype hypothesis. *Diabetologia* **35** 595–601. (doi:10.1007/BF00400248)
- Huxley R, Owen CG, Whincup PH, Cook DG, Rich-Edwards J, Smith GD & Collins R 2007 Is birth weight a risk factor for ischemic heart disease in later life? *American Journal of Clinical Nutrition* **85** 1244–1250.
- Jaquet D, Gaboriau A, Czernichow P & Levy-Marchal C 2000 Insulin resistance early in adulthood in subjects born with intrauterine growth retardation. *Journal of Clinical Endocrinology and Metabolism* **85** 1401–1406. (doi:10.1210/jc.85.4.1401)
- Jenuwein T & Allis CD 2001 Translating the histone code. *Science* **293** 1074–1080. (doi:10.1126/science.1063127)
- Laffitte BA, Chao LC, Li J, Walczak R, Hummasti S, Joseph SB, Castrillo A, Wilpitz DC, Mangelsdorf DJ, Collins JL *et al.* 2003 Activation of liver X receptor improves glucose tolerance through coordinate regulation of glucose metabolism in liver and adipose tissue. *PNAS* **100** 5419–5424. (doi:10.1073/pnas.0830671100)
- Lange AJ, Argaud D, el-Maghrabi MR, Pan W, Maitra SR & Pilks SJ 1994 Isolation of a cDNA for the catalytic subunit of rat liver glucose-6-phosphatase: Regulation of gene expression in FAO hepatoma cells by insulin, dexamethasone and cAMP. *Biochemical and Biophysical Research Communications* **201** 302–309. (doi:10.1006/bbrc.1994.1702)
- Lehmann JM, Kliewer SA, Moore LB, Smith-Oliver TA, Oliver BB, Su JL, Sundseth SS, Winegar DA, Blanchard DE, Spencer TA *et al.* 1997 Activation of the nuclear receptor LXR by oxysterols defines a new hormone response pathway. *Journal of Biological Chemistry* **272** 3137–3140. (doi:10.1074/jbc.272.6.3137)
- Levitt NS, Lambert EV, Woods D, Hales CN, Andrew R & Seckl JR 2000 Impaired glucose tolerance and elevated blood pressure in low birth weight, nonobese, young South African adults: early programming of cortisol axis. *Journal of Clinical Endocrinology and Metabolism* **85** 4611–4618. (doi:10.1210/jc.85.12.4611)
- Lillicrop KA, Phillips ES, Jackson AA, Hanson MA & Burdge GC 2005 Dietary protein restriction of pregnant rats induces and folic acid supplementation prevents epigenetic modification of hepatic gene expression in the offspring. *Journal of Nutrition* **135** 1382–1386.
- Lin B, Morris DW & Chou JY 1998 Hepatocyte nuclear factor 1 α is an accessory factor required for activation of glucose-6-phosphatase gene transcription by glucocorticoids. *DNA and Cell Biology* **17** 967–974. (doi:10.1089/dna.1998.17.967)
- Mitro N, Mak PA, Vargas L, Godio C, Hampton E, Molteni V, Kreusch A & Saez E 2007 The nuclear receptor LXR is a glucose sensor. *Nature* **445** 219–223. (doi:10.1038/nature05449)
- Nilsson PM, Ostergren PO, Nyberg P, Soderstrom M & Allebeck P 1997 Low birth weight is associated with elevated systolic blood pressure in adolescence: a prospective study of a birth cohort of 149378 Swedish boys. *Journal of Hypertension* **15** 1627–1631. (doi:10.1097/00004872-199715120-00064)
- Ozanne SE & Hales CN 2004 Lifespan: catch-up growth and obesity in male mice. *Nature* **427** 411–412. (doi:10.1038/427411b)
- Ozanne SE, Smith GD, Tikerpae J & Hales CN 1996 Altered regulation of hepatic glucose output in the male offspring of protein-malnourished rat dams. *American Journal of Physiology* **270** E559–E564.
- Park JH, Stoffers DA, Nicholls RD & Simmons RA 2008 Development of type 2 diabetes following intrauterine growth retardation in rats is associated with progressive epigenetic silencing of Pdx1. *Journal of Clinical Investigation* **118** 2316–2324. (doi:10.1172/JCI32011)
- Petrik J, Reusens B, Arany E, Remacle C, Coelho C, Hoet JJ & Hill DJ 1999 A low protein diet alters the balance of islet cell replication and apoptosis in the fetal and neonatal rat and is associated with a reduced pancreatic expression of insulin-like growth factor-II. *Endocrinology* **140** 4861–4873. (doi:10.1210/en.140.10.4861)
- Raab EL, Vuguin PM, Stoffers DA & Simmons RA 2009 Neonatal Exendin-4 treatment reduces oxidative stress and prevents hepatic insulin resistance in intrauterine growth-retarded rats. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **297** R1785–R1794. (doi:10.1152/ajpregu.00519.2009)
- Rees WD, Hay SM, Brown DS, Antipatis C & Palmer RM 2000 Maternal protein deficiency causes hypermethylation of DNA in the livers of rat fetuses. *Journal of Nutrition* **130** 1821–1826.
- Repa JJ, Liang G, Ou J, Bashmakov Y, Lobaccaro JM, Shimomura I, Shan B, Brown MS, Goldstein JL & Mangelsdorf DJ 2000 Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXR α and LXR β . *Genes and Development* **14** 2819–2830. (doi:10.1101/gad.844900)
- Ross MG & Beall MH 2008 Adult sequelae of intrauterine growth restriction. *Seminars in Perinatology* **32** 213–218. (doi:10.1053/j.semperi.2007.11.005)
- Santos-Rosa H, Schneider R, Bannister AJ, Sherriff J, Bernstein BE, Emre NC, Schreiber SL, Mellor J & Kouzarides T 2002 Active genes are tri-methylated at K4 of histone H3. *Nature* **419** 407–411. (doi:10.1038/nature01080)
- Sasaki K, Cripe TP, Koch SR, Andreone TL, Petersen DD, Beale EG & Granner DK 1984 Multihormonal regulation of phosphoenolpyruvate carboxykinase gene transcription the dominant role of insulin. *Journal of Biological Chemistry* **259** 15242–15251.
- Sohi G, Marchand K, Revesz A, Arany E & Hardy DB 2011 Maternal protein restriction elevates cholesterol in adult rat offspring due to repressive changes in histone modifications at the cholesterol 7 α -hydroxylase promoter. *Molecular Endocrinology* **25** 785–798. (doi:10.1210/me.2010-0395)
- Song C & Liao S 2000 Cholestenoic acid is a naturally occurring ligand for liver X receptor α . *Endocrinology* **141** 4180–4184. (doi:10.1210/en.141.11.4180)
- Song C, Kokontis JM, Hiipakka RA & Liao S 1994 Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors. *PNAS* **91** 10809–10813. (doi:10.1073/pnas.91.23.10809)

- Steffensen KR & Gustafsson JA 2004 Putative metabolic effects of the liver X receptor (LXR). *Diabetes* **53**(Suppl 1) S36–S42. (doi:10.2337/diabetes.53.2007.S36)
- van Straten EM, Bloks VW, Huijkman NC, Baller JF, Meer H, Lutjohann D, Kuipers F & Plosch T 2010 The liver X-receptor gene promoter is hypermethylated in a mouse model of prenatal protein restriction. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **298** R275–R282. (doi:10.1152/ajpregu.00413.2009)
- Stulnig TM, Oppermann U, Steffensen KR, Schuster GU & Gustafsson JA 2002a Liver X receptors downregulate 11 β -hydroxysteroid dehydrogenase type 1 expression and activity. *Diabetes* **51** 2426–2433. (doi:10.2337/diabetes.51.8.2426)
- Stulnig TM, Steffensen KR, Gao H, Reimers M, Dahlman-Wright K, Schuster GU & Gustafsson JA 2002b Novel roles of liver X receptors exposed by gene expression profiling in liver and adipose tissue. *Molecular Pharmacology* **62** 1299–1305. (doi:10.1124/mol.62.6.1299)
- Tarry-Adkins JL, Martin-Gronert MS, Fernandez-Twinn DS, Hargreaves I, Alfaradhi MZ, Land JM, Aiken CE & Ozanne SE 2013 Poor maternal nutrition followed by accelerated postnatal growth leads to alterations in DNA damage and repair, oxidative and nitrosative stress, and oxidative defense capacity in rat heart. *FASEB J* **27** 379–390. (doi:10.1096/fj.12-218685)
- Trinh KY, O'Doherty RM, Anderson P, Lange AJ & Newgard CB 1998 Perturbation of fuel homeostasis caused by overexpression of the glucose-6-phosphatase catalytic subunit in liver of normal rats. *Journal of Biological Chemistry* **273** 31615–31620. (doi:10.1074/jbc.273.47.31615)
- Valera A, Pujol A, Pelegrin M & Bosch F 1994 Transgenic mice overexpressing phosphoenolpyruvate carboxykinase develop non-insulin-dependent diabetes mellitus. *PNAS* **91** 9151–9154. (doi:10.1073/pnas.91.19.9151)
- Valverde AM & Gonzalez-Rodriguez A 2011 IRS2 and PTP1B: two opposite modulators of hepatic insulin signalling. *Archives of Physiology and Biochemistry* **117** 105–115. (doi:10.3109/13813455.2011.557386)
- Venkateswaran A, Laffitte BA, Joseph SB, Mak PA, Wilpitz DC, Edwards PA & Tontonoz P 2000 Control of cellular cholesterol efflux by the nuclear oxysterol receptor LXR α . *PNAS* **97** 12097–12102. (doi:10.1073/pnas.200367697)
- Willy PJ, Umesono K, Ong ES, Evans RM, Heyman RA & Mangelsdorf DJ 1995 LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes and Development* **9** 1033–1045. (doi:10.1101/gad.9.9.1033)
- Yan C & Boyd DD 2006 Histone H3 acetylation and H3 K4 methylation define distinct chromatin regions permissive for transgene expression. *Molecular and Cellular Biology* **26** 6357–6371. (doi:10.1128/MCB.00311-06)

Received in final form 25 April 2013

Accepted 30 April 2013

Accepted Preprint published online 30 April 2013