Vitamin D receptor-mediated control of *Soggy*, *Wise*, and *Hairless* gene expression in keratinocytes

Jui-Cheng Hsieh¹, Rudolf C Estess¹, Ichiro Kaneko^{1,2}, G Kerr Whitfield¹, Peter W Jurutka^{1,2} and Mark R Haussler¹

¹Department of Basic Medical Sciences, University of Arizona College of Medicine, 425 North 5th Street, Phoenix, Arizona 85004-2157, USA

²School of Mathematical and Natural Sciences, Arizona State University, Phoenix, Arizona 85306, USA

Correspondence should be addressed to M R Haussler
Email
haussler@email.arizona.edu

Abstract

The vitamin D receptor (VDR), but not its hormonal ligand, 1,25-dihydroxyvitamin D₃ (1,25D), is required for the progression of the mammalian hair cycle. We studied three genes relevant to hair cycle signaling, *DKKL1* (*Soggy*), *SOSTDC1* (*Wise*), and *HR* (*Hairless*), to determine whether their expression is regulated by VDR and/or its 1,25D ligand. *DKKL1* mRNA was repressed 49–72% by 1,25D in primary human and CCD-1106 KERTr keratinocytes; a functional vitamin D responsive element (VDRE) was identified at –9590 bp in murine *Soggy*. Similarly, *SOSTDC1* mRNA was repressed 41–59% by 1,25D in KERTr and primary human keratinocytes; a functional VDRE was located at –6215 bp in human *Wise*. In contrast, *HR* mRNA was upregulated 1.56- to 2.77-fold by 1,25D in primary human and KERTr keratinocytes; a VDRE (TGGTGAggtgAGGACA) consisting of an imperfect direct repeat separated by three nucleotides (DR3) was identified at –7269 bp in the human *Hairless* gene that mediated dramatic induction, even in the absence of 1,25D ligand. In parallel, a DR4 thyroid hormone responsive element, TGGTGAggccAGGACA, was identified at +1304 bp in the human *HR* gene that conferred tri-iodothyronine (T₃)-independent transcriptional activation. Because the thyroid hormone receptor controls *HR* expression in the CNS, whereas VDR functions in concert with the HR corepressor specifically in skin, a model is proposed wherein unliganded VDR upregulates the expression of *HR*, the gene product of which acts as a downstream comodulator to feedback-repress *DKKL1* and *SOSTDC1*, resulting in integration of bone morphogenic protein and Wnt signaling to drive the mammalian hair cycle and/or influencing epidermal function.

Key Words

- ▶ gene regulation
- ▶ vitamin D
- ▶ transcription factors
- ▶ hormone receptors
- ▶ skin

Journal of Endocrinology
(2014) 220, 165–178

Introduction

Molecular control of the mammalian hair cycle is incompletely characterized. Three gene products that are involved in this process are hairless (HR), the vitamin D receptor (VDR), and retinoid X receptor- α (RXR α). Loss-of-function mutations in any of the genes encoding these proteins in mammals result in failure to reinitiate the hair

cycle after the loss of the first coat of hair, leading to alopecia and dermal cysts (Bergman *et al.* 2005). It has therefore been proposed that these three gene products function together in a single pathway to initiate a new hair cycle (Wang *et al.* 2007, Hsieh *et al.* 2010). However, further details about this pathway and the other gene products

that might be included have not been elucidated, although there is evidence that multiple signaling pathways are involved, including Wnt proteins (Fuchs *et al.* 2001), sonic hedgehog (Teichert *et al.* 2010), and bone morphogenic proteins (BMPs; O'Shaughnessy *et al.* 2004).

Several clues have emerged from mouse gene knockout experiments, including a *hairless* gene ablation study by Thompson and colleagues (Beaudoin *et al.* 2005) that showed an inverse relationship between the expression of *hairless* and sclerostin domain-containing protein 1 (*SOSTDC1*) genes, the latter also known as Wnt modulator in surface ectoderm (Wise). *SOSTDC1*-encoded Wise inhibits both the Wnt and BMP signaling pathways in the course of the hair cycle (Lintern *et al.* 2009). Thompson and colleagues proposed that Hr suppression of *Sostdc1* expression is important in triggering reinitiation of the anagen phase of the hair cycle by allowing keratinocytes to respond to an undetermined signal that presumably impinges on the hair-cycle-controlling pathways, including the Wnt- β -catenin pathway (Beaudoin *et al.* 2005).

Another gene with a possible role in the hair cycle is dickkopf-like 1 (*DkkL1*), encoding soggy. It has been reported that mRNA levels of both *DkkL1* and *Sostdc1* are upregulated in *hr* null mice, downregulated in *hr*-overexpressing transgenic mice, and repressed in keratinocytes that have been 'rescued' by expression of the *hr* gene under the control of a keratin-14 promoter (Zarach *et al.* 2004, Thompson *et al.* 2006). *DkkL1* is a member of the Dickkopf family of secreted proteins, several of which regulate signaling by the canonical Wnt pathway (Niehrs 2006). Thompson *et al.* (2006) have shown that *DkkL1* expression is measurable in hair follicles of mice and peaks during late anagen and early catagen. In contrast, the mammalian Hr protein, which has been shown to be essential for progression of the hair cycle, exhibits a complementary expression pattern, i.e. strong expression during early anagen, with expression declining sharply at the anagen-catagen transition (Panteleyev *et al.* 2000). These authors put forth a model wherein the Hr protein suppresses the expression of *DkkL1* (Thompson *et al.* 2006) and also a second protein, Wise (Beaudoin *et al.* 2005). The suppression of both proteins is proposed to play a permissive role in allowing Wnt signaling to initiate a new anagen phase of the hair cycle.

Mammalian Hr is a highly conserved 130 kDa transcription factor that, according to *in situ* hybridization analysis in mice (Cachon-Gonzalez *et al.* 1999), is chiefly expressed in skin, cartilage, retina, inner ear, brain, colon, and oral/tongue/nasal/bladder/urethral epithelia. Loss-of-function mutations in the human *HR* gene cause atrichia

with papular lesions (Ahmad *et al.* 1999, Klein *et al.* 2002) and mutations leading to overexpression of HR protein also result in a hair-skin phenotype known as Marie Unna hereditary hypotrichosis (Wen *et al.* 2009, Ramot *et al.* 2010). In the *hr/hr* mouse (Cachon-Gonzalez *et al.* 1994), a mutation leading to partial loss of Hr protein causes disappearance of most hair after completion of the first hair cycle, along with dermal cysts, whereas a total loss of HR function, such as from a premature stop codon (Cachon-Gonzalez *et al.* 1999), results in complete alopecia after 3–4 weeks, but also thickened and wrinkled skin with many dermal cysts. Hr and VDR have been shown to physically and functionally interact (Hsieh *et al.* 2003a). VDR activates transcription in response to 1,25-dihydroxy-vitamin D₃ (1,25D) by forming a heterodimer with one of the RXRs and binding to a vitamin D responsive element (VDRE) in or near each target gene (Whitfield *et al.* 2005). One consequence of a VDR/Hr interaction is that Hr inhibits the ability of VDR to activate transcription of its target genes in response to the 1,25D ligand (Hsieh *et al.* 2003a, Xie *et al.* 2006). Hr also has been shown to attenuate transactivation by the thyroid hormone receptor (TR; Potter *et al.* 2001), as well as the retinoic acid receptor-related orphan receptor- α (ROR α ; Moraitis & Giguere 2003).

The molecular basis of Hr-mediated transrepression of VDR, TR, and ROR α signaling is not well characterized. It has been proposed that Hr exerts its repression via its Jumonji C-like domain to recruit histone deacetylases, which modify chromatin structure to silence gene transcription (Wang *et al.* 2007, Hsieh *et al.* 2010). It has also been reported that HR possesses intrinsic histone 3 lysine 9 demethylase activity, possibly controlling transcription catalytically via the histone code as an epigenetic 'eraser' (Liu *et al.* 2011). An interesting feature of human HR, as opposed to homologs reported from other mammalian species, is the presence of two isoforms, generated via alternative mRNA splicing. The α isoform (HR α) is produced from a complete mRNA transcript, whereas the β isoform (HR β) is translated from a transcript that lacks exon 17 (Malloy *et al.* 2009).

Based upon its functioning as a corepressor of VDR, we tested the hypothesis that HR may target VDR–VDRE signaling and subsequently modulate downstream *DKKL1* and *SOSTDC1* expression. *In silico* analysis of the regions surrounding the *DkkL1* and *SOSTDC1* genes revealed candidate VDREs, two of which are shown herein to be functionally active. We further hypothesized that *HR* may be transcriptionally activated by VDR, reasoning that the resulting HR corepressor could reciprocally suppress *HR* expression and *VDR* mRNA, thus establishing a novel

inhibitory feedback loop to control the level of both proteins. In this study, an *in silico* search for potential hormone responsive elements in human *HR* led to the identification of an apparent constitutive VDRE 5' of the promoter, as well as a novel, ligand-independent thyroid hormone responsive element (TRE) in the first intron.

Materials and methods

Cell culture

Cell lines were cultured in the following media: human CCD-1106 KERTr (KERTr) in keratinocyte serum-free medium (Invitrogen) with the recommended additives; human Caco-2 colorectal adenocarcinoma and green monkey COS7 kidney, human HEK-293 embryonic kidney, and human HaCaT keratinocyte cells were all cultured in DMEM supplemented with 10% fetal bovine serum (FBS), with the latter two lines receiving supplements of 4 and 2 mM L-glutamine respectively. Normal primary neonatal human epidermal keratinocytes (HEKn) were purchased from Invitrogen and were cultured in serum-free EpiLife medium containing the recommended HKGS supplement kit reagents. Primary keratinocytes were incubated for 24 h in 1.2 mM calcium to elicit differentiation, which was evidenced by a clear change in morphology, with the uniform monolayer at 60 μ M calcium transforming to dense/compacted, differentiated keratinocytes. Thus, for the present experiments, KERTr and HaCaT keratinocytes were undifferentiated, whereas HEKn cells were used in the differentiated state.

Plasmid constructs

Oligonucleotides corresponding to four copies of candidate VDREs along with four bases of flanking sequence on either side were separately annealed and inserted into the HindIII and BglII sites of pLuc-MCS (Stratagene Corp., La Jolla, CA, USA). Positive control reporter plasmids were similarly constructed with four copies of the rat osteocalcin VDRE (rOC-Luc) or 5500 bp from the promoter of the human *CYP24A1* gene (p24OHaseLuc). The integrity of each of these reporter vectors was confirmed by DNA sequencing. The expression vector for WT rat Hr (pRK5myc-rhr), which contains the CMV promoter, has been described previously (Potter *et al.* 2001). An expression vector for full-length human HR (*HR α*) was kindly provided by Dr. A Hillmer (Rheinische Friedrich-Wilhelms-Universität, Germany). *HR α* is cloned into the mammalian expression vector p3xFLAG-CMV-7.1 with

expression driven by the CMV promoter, yielding *HR α* with a triple FLAG-tag at the N-terminus. An expression vector for *HR β* (HR Δ 1072-1126) was derived from the parent *HR α* vector using the Quickchange XL Mutagenesis Kit (Stratagene). The expression plasmids for human VDR (pSG5hVDR) and for human TR β 1 have been described previously (Hsieh *et al.* 1991). The reporter construct used for assaying TR signaling contained two copies of the TRE from the rat myosin heavy chain gene (Hsieh *et al.* 1991).

Transcriptional activity assays

Cultured cells were plated in 24-well plates at a density of 80 000 cells/well in 1 ml of the appropriate medium. Six or more hours following plating (when the cells had become attached), wells were cotransfected using Lipofectamine and PLUS Transfection Reagents (Invitrogen) with 250 ng/well of reporter plasmid (either the rOC-Luc reporter plasmid or one of the VDRE-Luc or TRE-Luc constructs), along with pRL-CMV (1 ng/well), pSG5hVDR (250 ng/well), and 499 ng of pTZ18U plasmid as carrier DNA. For each well, plasmids (total 1.0 μ g) were diluted into 25 μ l serum-free medium. PLUS reagent (4 μ l) was added and incubated at room temperature for 15 min. Lipofectamine Reagent (2 μ l) was diluted into 25 μ l serum-free medium in a second tube. The pre-complex DNA and Lipofectamine tubes were combined and incubated for an additional 15 min. The final DNA-Plus-Lipofectamine complexes (54 μ l/well) were added to the cultured cells and incubated for 48 h with or without 1,25D. The wells were washed twice with PBS and lysed with 150 μ l passive lysis buffer (from DLR Assay Kit, Promega). Firefly and Renilla luciferase activities were measured sequentially from each lysate using a Sirius Luminometer (Titertek Berthold, Pforzheim, Germany) and a Dual Luciferase Assay Kit (Promega) as per the manufacturer's instructions. The ratio of firefly:Renilla luciferase activity was calculated to normalize for transfection efficiency.

In silico, bioinformatic search for candidate VDREs

The search criteria were based on the following direct repeat 3 (DR3) degenerate sequence, with hexanucleotide repeats in upper case and the three nucleotide spacer in lower case: RGKDBRnnrRGKDBR, where R=A or G, K=G or T, D=G, T or A, B=G, T or C, and *n*=any base.

Electrophoretic mobility shift assay

Annealed, ³²P-labeled oligonucleotides were used in an electrophoretic mobility shift assay (EMSA) as follows.

Double-stranded oligonucleotides encoding two tandem copies of the candidate VDREs plus four bases of flanking sequences and four-base overhangs were labeled with [α - 32 P]dCTP and used along with *Escherichia coli*-expressed, partially purified human VDR and RXR α as described previously (Hsieh *et al.* 2003b). A double-stranded oligonucleotide, containing a dual copy of the VDRE sequence from the rat osteocalcin gene, served as a positive control that binds VDR and RXR. Reaction mixtures were resolved on 4% nondenaturing polyacrylamide gels, dried, and exposed to Amersham Hyperfilm at room temperature.

Quantitative real-time PCR assays

KERTr and HEKn were plated at 800 000 cells/100 mm plate in 10 ml of DMEM with 10% FBS. The cells were treated with 10^{-8} M 1,25D for 24 h and harvested by trypsinization. Total RNA was extracted using a High Pure RNA isolation kit (Roche Applied Science) according to the manufacturer's instructions. First-strand cDNA was synthesized from 1.0 μ g total RNA using an iScript cDNA synthesis kit (Bio-Rad Laboratories). The cDNA was used in 20 μ l PCRs containing 10 μ l FastStart Universal SYBR Green Master Mix (Roche Applied Science) with primers. Reactions were performed on an ABI 7500 Fast instrument (Life Technologies). Data were analyzed using the comparative Ct method as a means of relative quantitation, normalized to an endogenous reference (glyceraldehyde phosphate dehydrogenase (GAPDH)) and relative to a calibrator (normalized Ct value from vehicle-treated cells), and expressed as $2^{-\Delta\Delta C_t}$ according to Applied Biosystems' User Bulletin 2: Rev B, 'Relative Quantitation of Gene Expression'. The primer sets for real-time PCR were as follows: Human *HR* primers were 5'-GGGACACATCGATAGGGAACAAGGAT-3' (forward primer) and 5'-TATGTCCTGAAGTCCCGGTCC-3' (reverse primer). Primers for human GAPDH expression were 5'-ACAACCTTGGTATCGTGGAAGGAC-3' (forward primer) and 5'-CAGGGATGATGTTCTGGAGAGC-3' (reverse primer). Primers for human *DKKL1* were 5'-GACAACAAGACAGGAGAGGTG-3' (forward primer) and 5'-TCAAATCACCCTCGAAGCTC-3' (reverse primer). Primers for human *SOSTDC1* were 5'-TGTTCCATAGCCTCCTCCAATCCAGTTA-3' (forward primer) and 5'-AACTGCGTTCCACCAAATACATCTCTGAT-3' (reverse primer). Human *CYP24A1* was detected using forward primer 5'-CAGCGAACTGAACAAATGGTCG-3' and reverse primer 5'-TCTCTTCTCATACAACACGAGGCAG-3'.

Results

The 1,25D hormone regulates the expression of genes relevant to skin/hair cycle

Assuming Hr functions as a corepressor in cooperation with VDR to modulate genes encoding factors that ensure hair cycle progression, we initially examined several genes observed to be overexpressed in keratinocytes of *hr*-ablated mice. As illustrated in Fig. 1, we evaluated the effect of 10^{-8} M 1,25D for 24 h on the expression of *DKKL1*, *SOSTDC1*, and *HR* in cultured human keratinocytes using qRT-PCR. *CYP24A1* was employed as a highly induced positive control for VDR-mediated 1,25D action to ensure that the tested cells expressed VDR and thereby responded to the 1,25D ligand in the expected fashion. Indeed, Fig. 1A illustrates the dramatic 89- and 124-fold enhancement of *CYP24A1* mRNA concentrations elicited by the vitamin D hormone in undifferentiated KERTr and differentiated primary human keratinocytes, respectively, indicating that both systems express approximately equal VDR levels and represent valid models to probe VDR-targeted gene expression. Furthermore, we observed that CASP14, which is a nonapoptotic caspase family

Figure 1

1,25D effect on *CYP24A1*, *DKKL1*, *SOSTDC1*, and *HR* expression in keratinocytes. A (*CYP24A1*), B (*DKKL1*), C (*SOSTDC1*), D (*HR*). Real-time PCR was performed on total RNA prepared from: KERTr (center black bars in each panel) and primary normal human epidermal keratinocytes (HEKn; right black bars in each panel) using specific primers for each gene, as described in Materials and methods. All results were normalized to GAPDH, and cells treated with 10^{-8} M 1,25D for 24 h were compared with vehicle ethanol (EtOH) control, which was set to 1.0-fold. Each bar represents the average of three independent experiments $\pm$ s.d. *All 1,25D treated groups were statistically significantly different from EtOH control ($P < 0.05$), with the exception of *HR* in HEKn cells ($P = 0.08$), although this latter induction by 1,25D has been observed in three additional HEKn primary cultures to average 1.40-fold (data not shown).

member that is essential for keratinocyte differentiation (Rendl *et al.* 2002), is induced 2.11-fold by 1,25D in KERTr cells (data not shown), a finding verified by cDNA microarray analysis (Haussler *et al.* 2013). Thus, despite their immortality through transformation by human papillomavirus 16 E6/E7, KERTr cells apparently retain the cadre of comodulators required for 1,25D/VDR influence on skin/hair cycle relevant genes.

In accordance with this, the data in Fig. 1B reveal that 1,25D represses *DKKL1* mRNA concentrations by 72 and 49% in KERTr and normal primary human keratinocytes respectively. Thus, *DKKL1* fits the concept of repression by the VDR–RXR complex in cooperation with Hr, with relieving of this repression (Beaudoin *et al.* 2005) in the keratinocytes from *hr*-null mice. As depicted in Fig. 1C, 1,25D also represses *SOSTDC1* mRNA levels by 41 and 59% in KERTr and primary human keratinocytes respectively. Therefore, like *DKKL1*, *SOSTDC1* fits the concept of repression by the VDR–RXR complex in cooperation with Hr, with relieving of this repression (Beaudoin *et al.* 2005) in the keratinocytes from *hr*-null mice. With respect to regulation of human *HR* expression by 1,25D, qRT-PCR results illustrated in Fig. 1D reveal a 2.77- and 1.56-fold enhancement of *HR* mRNA expression by 1,25D treatment of KERTr and primary human keratinocytes respectively. Thus, we hypothesized that VDR binds to VDREs in the *HR* gene to activate transcription, and in *DKKL1* and *SOSTDC1* to repress transcription.

Regulation of *DKKL1* and *SOSTDC1* by 1,25D

We first investigated the mechanism whereby 1,25D controls *DKKL1* and *SOSTDC1*, noting that VDREs have yet to be identified in the vicinity of either gene. Consequently, *in silico* analysis was performed on both mouse and human *DKKL1* genes in order to locate potential VDREs. As shown in Fig. 2A for mouse *DkkL1*, a single candidate VDRE, GGGTCAtggAGGGCA, was located at –9590 bp relative to the transcriptional start site. A single candidate VDRE was also found in the human *DKKL1* gene at –9830 bp (Fig. 2B). Both candidates were analyzed via an EMSA to test their *in vitro* VDR-binding capabilities. As shown in Fig. 2C, only the mouse candidate VDRE was capable of binding VDR–RXR complex when compared with the rOC VDRE positive control. Transcriptional activity of the mouse VDRE was examined using the dual luciferase assay procedure. As shown in Fig. 2D, the –9590 VDRE is activated by 1,25D, driving liganded, VDR-mediated transcriptional activation with a 12-fold effect over the vehicle control, approaching the activity of

the human *CYP24A1* natural promoter fragment (5.5 kb) containing a classic dual VDRE (Fig. 2D), and exceeding the activity of the rat osteocalcin VDRE (Fig. 2E). As shown in Fig. 2E, activation of the –9590 VDRE by 1,25D is reduced from 16.7- to 2.5-fold by rat Hr, similar to VDREs in well-characterized, 1,25D-regulated genes. Curiously, the basal activity of the –9590 VDRE is increased by approximately a factor of two in the presence of the rHr, a phenomenon not usually observed for VDREs, wherein both basal and 1,25D-stimulated transcriptions are suppressed. The mechanistic caveats regarding the murine –9590 VDRE in *DkkL1* are that it is: i) specific to the mouse, as a conserved homolog has yet to be located in the human *DKKL1* gene and ii) functioning as a positive VDRE out of the keratinocyte setting, whereas 1,25D represses *DKKL1* expression in intact keratinocytes (Fig. 1B).

Similar to *DKKL1*, *SOSTDC1* expression is repressed by 1,25D in keratinocytes. (Fig. 1C), and both genes are overexpressed in keratinocytes derived from *hr* knockout mice. Given these observations, it is evident that both genes are regulated by 1,25D/VDR and by Hr, probably involving a pattern of comodulators specific to differentiated keratinocytes. Further experiments probing the human *SOSTDC1* gene were conducted via *in silico* analysis, yielding the identification of two potential human *SOSTDC1* VDREs (Fig. 3A). An EMSA revealed that both the –6215 and –5857 bp responsive elements in human *SOSTDC1* were capable of binding a VDR–RXR complex, *in vitro* (Fig. 3B). However, as shown in Fig. 3C, only the –6215 element was capable of mediating repression of transcription by 1,25D even though, in the same experiment, 1,25D dramatically induced transcription directed by the mouse *DkkL1* VDRE located at –9590 as well as by the human *CYP24A1* promoter fragment. As depicted in Fig. 3D, in a repeat experiment, the –6215 bp VDRE mediated significant repression of reporter transcription in the presence of 1,25D, a phenomenon that is best shown in the left panel of Fig. 3D. This repression is reproducible in the presence of endogenous levels of VDR in HEK-293 cells (bars in right panel of Fig. 3D, ‘no added VDR’). Thus, the data in Fig. 3 indicate that the molecular mechanism whereby 1,25D represses *SOSTDC1* expression in human cells probably involves a negative VDRE at –6215 bp in the *SOSTDC1* gene.

Identification and functional analysis of a putative VDRE in the human *HR* gene

In silico searches revealed six candidate VDREs, located at –7269, –7410, –8171, –9087, –23 029, and

Figure 2

In vitro characterization and functional analysis of an *in silico*-identified VDRE in the mouse *Dkk1* gene. (A and B) Bioinformatic identification of candidate VDREs in the vicinity of the mouse and human *DKKL1* genes respectively. (C) EMSA of VDR/RXR binding to candidate VDREs as described in Materials and methods. Lane 1 contains the rat osteocalcin (roC) VDRE probe plus added RXR α (50 ng) and VDR (100 ng); the shifted complex is indicated by an arrow at the left. Lanes 2 and 5 contain labeled probes with no protein added to the candidate mouse and human VDREs respectively. Lanes 3 and 6 contain probes plus added (50 ng) RXR α only. Lanes 4 and 7 each contain respective probes plus added RXR α (50 ng) and VDR (100 ng). (D) Reporter gene assays testing the functionality of the mouse *Dkk1* candidate VDRE in COS7 cells. A double-stranded oligonucleotide corresponding to the -9590 VDRE was inserted as a four copy tandem repeat upstream of a luciferase reporter gene and cotransfected into COS7 cells with a pSG5VDR expression vector to supply exogenous VDR. The human *CYP24A1* 5.5 kb natural promoter-luciferase

construct was used as a positive control. Indicated wells received 1,25D (10^{-8} M final concentration) at 18 h post-transfection. After an additional 24 h, cell lysates were harvested and luciferase activity was assayed as described in Materials and methods. Each bar represents the average of three independent experiments performed in triplicate $\pm$ s.d.

*1,25D-treated groups statistically significantly different from the EtOH control ($P < 0.05$). (E) Rat Hr inhibits transactivation by VDR in the -9590 *Dkk1* VDRE-luciferase construct. COS7 cells were cotransfected with the following plasmids as indicated: pSG5hVDR (250 ng/well), pRK5rHr (200 ng/well), and the -9590 *Dkk1* VDRE luciferase reporter (250 ng/well) or a positive control plasmid containing four copies of the rat osteocalcin VDRE. Transcriptional activities were quantified by luciferase assay and all values were normalized to the expression of Renilla luciferase. Each bar represents the average of three independent experiments performed in triplicate $\pm$ s.d. *1,25D-treated groups statistically significantly different from the EtOH control ($P < 0.05$).

-24 780 bp relative to the transcriptional start site of the *HR* gene (Ahmad *et al.* 1999; Fig. 4A, upper). It was next determined whether any of these candidate VDREs could bind VDR, *in vitro*, via EMSA. As shown in Fig. 4A (lower), VDR-RXR α complexes were formed in the absence of 1,25D ligand on four of the six tested VDREs: the -7269 element (lane 12), the -7410 element (lane 10), the -8171 element (lane 8), and a weak complex on the -24 780 element (lane 2). Luciferase assays revealed that only the -7269 VDRE is capable of conferring 1,25D induction onto the reporter gene in HEK-293 cells (Fig. 4B). It is noteworthy that transcription of the -7269 VDRE-linked reporter gene is significantly upregulated even in the absence of 1,25D, although the addition

of ligand results in a further increase by 3.5-fold (Fig. 4B). The -7269 VDRE performs similarly in HaCaT keratinocytes (Fig. 4C), with significant ligand-independent activity and approximately a twofold augmentation of transcription by 1,25D. In COS7, however, the VDRE is completely ligand-independent in the mediation of transactivation (Fig. 4D). These data indicate that VDR-mediated transactivation driven by the *HR* -7269 VDRE is cell-type-specific in terms of degree of ligand autonomy. An unusual feature of this VDRE is the presence of a T nucleotide at the first (underlined) position of the VDRE (TGGTGAggtAGGTCA). We suggest that the presence of a thymidine in this position might confer ligand-independent regulation onto a reporter gene.

Figure 3

In vitro characterization and functional analysis of *in silico*-identified VDREs in the *SOSTDC1* gene. (A) Bioinformatic identification of candidate VDREs in the vicinity of the human *SOSTDC1* gene. Binding sites for the CTCF insulator protein were determined in normal human epidermal keratinocytes (Mikkelsen *et al.* 2007); these sites were used to delineate the genomic region that was searched. (B) Annealed, 32 P-labeled oligonucleotides were used in EMSA as described in Materials and methods and the legend to Fig. 2. (C) Reporter gene assays testing the functionality of the human *SOSTDC1* candidate VDREs in COS7 cells. Double-stranded oligonucleotides corresponding to the -5857 and -6215 VDREs were evaluated by luciferase assay as described in Materials and methods and the legend to Fig. 2; data are expressed as fold-stimulation of transcription by 1,25D treatment on the ordinate. The luciferase constructs containing the human *CYP24A1* natural promoter and the -9590 mouse *DkkL1* VDRE were used as positive controls. Mock represents empty reporter vector

transfections. Each bar represents the average of three independent experiments performed in triplicate $\pm$ s.d. *1,25D-treated groups statistically significantly different from the EtOH control ($P < 0.05$). (D) Reporter gene assays testing the functionality of the human *SOSTDC1* candidate VDREs in COS7 and HEK293 cells. Experiments were carried out as described in (C) above, except that the data for the *CYP24A1* natural promoter positive control are not shown. The left panel represents a magnification of the data obtained in COS7 in an independent experiment, excluding *CYP24A1*, allowing visualization of the repressive effect of the -6215 VDRE. The right panel illustrates an independent experiment in HEK293 cells, wherein no exogenous VDR was included as indicated. Each bar represents the average of three independent experiments performed in triplicate $\pm$ s.d. *1,25D-treated groups statistically significantly different from EtOH control ($P < 0.05$).

Functional interaction of -7269 VDRE with human HR isoforms and rat Hr protein

To examine whether transactivation mediated by the -7269 VDRE is repressed by Hr, and whether this repression affects 1,25D-dependent transcription, ligand-independent transcription, or both, HEK-293 cells were cotransfected with hVDR and Hr expression plasmids along with the -7269 VDRE-reporter construct, and 1,25D-induced transcriptional activity was measured in the absence and presence of 10^{-8} M of 1,25D (Fig. 4E). Cotransfection of either HR α or HR β resulted in a dramatic repression of ligand-independent VDRE-VDR

transcription ($\sim 70\%$) and a milder reduction in transcriptional activation by 1,25D-liganded VDR-VDRE ($\sim 30\%$), with no significant differences between the two human HR isoforms. In contrast, rat Hr sharply repressed both ligand-independent and 1,25D-dependent transactivation, almost completely blunting any effect of 1,25D (Fig. 4E, right two bars). Although rat Hr resides in the pRK5-myc vector, whereas human HR isoforms lie in the p3xFLAG-CMV-7.1 vector, both vectors employ the strong CMV promoter, yielding efficient expression. Western blotting (data not shown) analysis confirmed that human HR α and HR β are equally expressed, but lack

Figure 4

In vitro characterization and functional analysis of the *in silico*-identified VDREs in the human *HR* gene. (A, upper panel) Bioinformatic identification of candidate VDREs in the vicinity of the human *HR* gene. Locations and sequences of candidate VDREs relative to the *HR* transcriptional start site (*HR* start) are depicted employing an image from the UC Santa Cruz web browser (hg18 version; Kent *et al.* 2002). As in Fig. 3 (legend), CTCF binding sites were used as boundaries for the genomic interval to be searched. Orientation of each candidate element is indicated as sense (S) or antisense (AS). Neighboring genes include receptor accessory protein 4 (REEP4). (A, lower panel) Annealed, ³²P-labeled oligonucleotides were used in an EMSA as described in Materials and methods and the legend to Fig. 2. Even numbered lanes contained partially purified human VDR and RXRα proteins; odd numbered lanes contained probe only. (B) Reporter gene assays testing the functionality of human *HR* candidate VDREs in HEK-293 cells. Double-stranded oligonucleotides corresponding to each *HR* VDRE that tested positively in the EMSA were separately evaluated by luciferase assay as described in Materials and methods and in the legend to Fig. 2.

Each bar represents the average of three independent experiments performed in triplicate ± s.d. *1,25D-treated groups statistically significantly different from ETOH control ($P < 0.05$). (C and D) The -7269 VDRE-luciferase construct was tested in HaCaT (C) and COS7 (D) cell lines. Each bar represents the average of three independent experiments performed in triplicate ± s.d. *1,25D-treated groups statistically significantly different from ETOH control ($P < 0.05$). (E) The ability of human *HR* isoforms to inhibit transactivation by VDR. HEK-293 cells were cotransfected with the following plasmids as indicated: pSG5hVDR (250 ng/well), p3xflagCMVhHrα (200 ng/well), p3xflagCMVhHrβ (200 ng/well), pRK5hR (200 ng/well) and the -7269 VDRE luciferase reporter (250 ng/well). Transcriptional activities were quantified by luciferase assay and all values were normalized to the expression of Renilla luciferase. Each bar represents the average of three independent experiments performed in triplicate ± s.d. *1,25D-treated groups statistically significantly different from ETOH control ($P < 0.05$). #Hairless-treated groups statistically significantly different from untreated control ($P < 0.05$).

of an effective antibody prevented the quantitation of rat Hr expression. Nevertheless, because rat Hr proved to be more potent than the human isoforms in suppressing VDR-mediated transactivation, there is little doubt about the activity of rat Hr as a VDR corepressor.

Characterization of a novel TRE in the human *HR* gene

A 3 kb portion of the human *HR* promoter (-2902 to +102) was previously shown to be differentially regulated

by T_3 in neuroblastoma cells and keratinocytes, and a TRE (AGGGCATctgAGGACA) was localized -2632 to -2647 bp upstream of the human *HR* gene (Engelhard & Christiano 2004). Because functional VDREs apparently accommodate a thymidine in the first position (such as the -7269 element), the search for TRE sequences was expanded to include such half-sites to determine if a novel TRE might be found in the human *HR* gene. A candidate TRE with the sequence TGGTGAggccAGGACA was indeed identified at +1304 to +1319 in the

first intron of the human *HR* gene (Fig. 5A). Evaluation of the transcriptional enhancer capacity of the -2632 and $+1304$ TREs revealed that transcription of the -2632 TRE-linked reporter is activated only in the presence of T_3 (10^{-7} and 10^{-8} M), with very modest ligand-independent activity (Fig. 5B). In contrast, the $+1304$ TRE conferred significant luciferase expression in the absence of T_3 that was not further enhanced by the addition of T_3 (Fig. 5C), unveiling a ligand-independent TRE analogous to the constitutive -7269 VDRE in the *HR* gene (Figs 4D and 5D).

Figure 5

TREs in the human *HR* locus. (A) A previously reported TRE at -2632 along with a novel TRE at $+1304$ bp in the context of the human *HR* gene. (B and C) Reporter gene assays of each TRE in the presence of either TR β alone or TR β + T_3 (two concentrations as indicated) for 24 h. Mock transfected wells that received empty reporter plasmid are shown for comparison. Each bar is the compilation of three independent experiments performed in triplicate $\pm$ s.d. * T_3 -treated groups statistically significantly different from EtOH control ($P < 0.05$). Ligand independency is demonstrated in (C) by the fact that $-T_3$ and $+T_3$ values are not statistically different ($P > 0.1$). (D) Sequence comparison of the novel ligand-independent TRE and VDRE sequences, the thymidine residue at the first (n1) position of each responsive element are highlighted with a box. Half-element sequences are identical between the VDRE and TRE, with differences only in the three and four nucleotide spacers.

Discussion

Control of the hair cycle by VDR is crucial in pathobiology because hair serves a critical function in terrestrial mammals by protecting skin from DNA-damaging u.v. irradiation. Rachitic, VDR-null mice display the phenotype of alopecia and dermal cysts, which is not ameliorated by a high calcium, lactose, and phosphate rescue diet that reverses the calcium and bone mineral defects (Sakai *et al.* 2001). However, there exists no corresponding pathological phenotype in the skin of mice that unable to synthesize 1,25D, suggesting that at least a part of the action of VDR in skin is independent of the 1,25D ligand. The hair and skin abnormalities observed in mice with *hr* loss-of-function mutations are largely recapitulated in VDR knockout mice (Miller *et al.* 2001), an observation that is consistent with a functional interaction between the VDR and Hr nuclear proteins in signal transduction pathways that drive the hair cycle. Because Hr is a nuclear receptor corepressor, it is presumably this action of Hr-VDR in mammals that is required for the progression of the hair cycle. Following this reasoning, Thompson and colleagues (Beaudoin *et al.* 2005) have defined *Sostdc1*, *Dkk11*, and *Casp14* as genes overexpressed in keratinocytes from *hr*-null mice, and Kato and colleagues (Yamamoto *et al.* 2009) characterized *S100A8* and *CASP14* as two genes overexpressed in VDR-null keratinocytes. It was previously observed (Haussler *et al.* 2010) that *S100A8* is rapidly repressed by 1,25D in human keratinocytes. Similar to *S100A8*, *SOSTDC1* is significantly repressed by 1,25D-activated VDR in KERTr and primary human keratinocytes (Fig. 1C). Suppression of *SOSTDC1* mRNA by 1,25D was verified using reverse transcriptase PCR in human keratinocytes (Haussler *et al.* 2010), and cDNA microarray analysis of Caco2 cells (data not shown). The present results demonstrate the interaction of VDR with a VDRE in the human *SOSTDC1* gene at -6215 bp (Fig. 3B), and this VDRE elicited 1,25D-dependent repression when linked in a reporter construct, indicating that it acts as a negative VDRE (Fig. 6, upper right). Because *SOSTDC1*-encoded Wise not only antagonizes the Wnt pathway by binding to lipoprotein-receptor-related protein (LRP) but also inhibits the BMP pathway through neutralization of BMP4 (Lintern *et al.* 2009), repression of *SOSTDC1* by VDR-Hr could constitute a major event in initiation of the mammalian hair cycle (Fig. 6). However, the relationship between *SOSTDC1* repression (as well as *HR* induction) by 1,25D and progression of the hair cycle is unclear, because post-morphogenic responses to VDR in the mammalian hair follicle are vitamin D ligand-independent.

Figure 6

Model for VDR and Hr action in keratinocytes. Regulation of *hr*, *Sostdc1*, and *DkkL1* expression by 1,25D/VDR may contribute to epidermal cell functions as depicted in the central and right portions of the model. However, along with VDR (Cianferotti *et al.* 2007), and Hr (Cachon-Gonzalez *et al.* 1999), β -catenin is absolutely required in keratinocytes (Huelsen *et al.* 2001) to permit mammalian hair cycling, and the ligand-independent action of VDR to drive the hair cycle is thought to involve Wnt signaling in skin stem cells (Beaudoin *et al.* 2005, Cianferotti *et al.* 2007). Thus, as illustrated in the left portion of the figure, by upregulating *HR*, and in turn repressing *Sostdc1* (*Wise*) and *DkkL1* (*Soggy*) expression, unliganded VDR could conceivably drive the hair cycle. Signaling in the mammalian hair cycling is complex, consisting of the convergence of two signaling pathways, BMP and Wnt. Noggin from the dermal papilla initially antagonizes BMP4 signaling in bulb (or bulge) keratinocytes, allowing for the accumulation of Lef1/TCF, a transcriptional coactivator that targets genes via DNA-binding partners such as β -catenin. Cessation of Noggin signaling reinstates BMP signal transduction via SMADs provided that *Wise* (encoded by *Sostdc1*), which antagonizes both Wnt and BMP pathways

(Lintern *et al.* 2009), has also been repressed, either directly by VDR (upper right), indirectly through VDR induction of Hr (right center), or by a combination thereof. Wnt ligand (e.g. Wnt 10b) signaling leads to accumulation of β -catenin, which cooperates with Lef1/TCF to induce genes encoding factors such as sonic hedgehog (*Shh*), that trigger the hair cycle to transition from telogen (resting) to anagen (growth). Finally, the figure is not meant to imply that keratinocytes and hair follicle bulge stem cells are the same population of cells. Because mature keratinocytes rather than hair follicle stem cells that express Sox9, K15, Cd34, and alpha 6, were used for the present experiments, one must be cautious in applying this model to control of the hair cycle over regulation of more straightforward mature epidermal cell functions. See text for additional discussion. Abbreviations not defined in the text are: HDMe, histone demethylase; Wnt, ortholog of *Drosophila* wingless and mouse int-1; Lef1, lymphoid enhancer factor-1; TCF, T cell-specific factor; *msx-1* and *msx-2*, orthologs of *Drosophila* muscle-specific homeobox protein. Factors that are membrane receptors or transporters are boxed. Solid arrows indicate activation and dotted lines ending in a continuous perpendicular line denote inhibition.

As illustrated in Fig. 1B, *DKKL1* mRNA also is repressed by 1,25D in KERTr and primary human keratinocytes. *DkkL1* is expressed in the hair follicle in a manner inversely related to the expression of *Hr* (Thompson *et al.* 2006), leading to the suggestion that *DkkL1* may be a Wnt inhibitor. Should this be the case, *DkkL1* suppression by VDR-Hr would potentiate the action of the Wnt signaling pathway to initiate a new cycle of hair growth. Indeed, Wnt is the major inductive signaling pathway activated in hair follicle stem cells (HFSCs) during the onset of a new hair cycle, and the VDR does play a role in HFSC function, although in an apparently unliganded fashion, and perhaps in cooperation with Hr to repress *DkkL1* expression. One cautionary note this hypothesis is that *DkkL1* expression does not affect Wnt signaling in the testes (Kohn *et al.* 2005). However, it is possible that *DkkL1* lacks the activity to inhibit Wnt signaling in the testes but

possesses such activity in the context of the hair cycle. Alternatively, other major Wnt regulators such as *wif1*, *Dkk2*, *msx*, etc., may instead replace *DkkL1* as a pivotal Wnt regulator modulated by VDR-Hr. A second cautionary note is that the data in Fig. 2D are seemingly in conflict with those in Fig. 1B showing repression, because 1,25D is observed to upregulate transcription driven by a murine *DkkL1*-linked reporter gene in Fig. 2D. Thus, the murine –9590 *DkkL1* VDRE performs like a classic ‘positive’ VDRE in isolation (Fig. 2C and D), but considering that 1,25D represses *DKKL1* expression in intact keratinocytes (Fig. 1B), this VDRE apparently adopts a repressive character in its natural context of DNA and associated comodulators in the differentiated keratinocyte. We term such VDREs ‘conditional’, rather than attempting to classify them as ‘positive’ or ‘negative’. Indeed, the direction and/or magnitude of *DKKL1* gene control may

be cell-context specific as well as differentiation stage selective, with repression mediated by the VDR–RXR–HR–HDAC complex bound to the VDRE (corepressors shown at lower right; Fig. 6; Skoriya *et al.* 2005). In isolation, outside the context of the differentiated keratinocyte, this repression of *DKKL1* is postulated to be relieved by VDR liganding with 1,25D in conjunction with recruitment of coactivators to the VDRE, effectively replacing HR/corepressors as depicted in Fig. 6 (lower right object). Although it is clear that *DKKL1* expression is regulated by VDR, the mechanism via the murine –9590 *Dkk1* VDRE appears to be species-specific, as we have yet to identify an active VDRE in the human *DKKL1* gene to account for the repression by 1,25D shown in Fig. 1B. Moreover, we have on rare occasions observed in some preparations of calcium-differentiated primary human keratinocytes, but never in KERTr cells, that 1,25D induces rather than represses both *DKKL1* and *SOSTDC1* expression (data not shown), suggesting that both the *DKKL1* and *SOSTDC1* VDREs should perhaps be referred to as ‘conditional’. This observation, probably accounted for by incomplete differentiation of occasional primary human keratinocyte lots, is also consistent with the recent finding that HR elicits coactivation of the cathelicidin gene while corepressing *CYP24A1* mRNA in the context of HaCaT keratinocytes (Chuma *et al.* 2012). Finally, although no dramatic epidermal or hair cycle phenotype exists in either *Dkk1*- or *Sostdc1*-knockout mice, the two gene products could redundantly regulate the hair cycle, requiring a double knockout to generate a hair cycle phenotype, or both gene products may coordinate with other genes within the skin compartment to control hair growth.

A major goal of the present experiments was to characterize the regulation and functions of the Hr comodulator. It has been demonstrated previously that Hr functions as a VDR corepressor (Hsieh *et al.* 2003a), raising the question as to whether Hr and VDR might reciprocally control one another via a counterregulatory feedback loop. Indeed as shown herein, activated VDR is a bona fide positive regulator of *HR* mRNA expression (Fig. 1D). It has been reported that 816 bp of 5' flanking sequence derived from the human *HR* gene could support VDR-dependent transrepression that was amplified by 1,25D whereas, paradoxically, the ‘full promoter’ supported ligand-independent transactivation by VDR in a cell-specific fashion (Engelhard *et al.* 2008), thus suggesting the existence of both negative and positive feedback loops whereby VDR modulates *HR* expression. In this study, we report that human *HR* gene expression is augmented by 1,25D in both keratinocyte (Fig. 1D) and

enterocyte (Caco-2, data not shown) cell lines. It is therefore concluded that the induction of *HR* by 1,25D-liganded VDR represents a feedforward counter-regulatory action to curtail the effects of the 1,25D hormone–VDR complex through the synthesis of its corepressor, HR.

In a quest to seek functional VDREs in the vicinity of the human *HR* gene, we identified, *in silico*, the existence of a novel VDRE that is proposed to account for at least part of the ability of VDR to regulate *HR* gene expression. This candidate VDRE, located 7269 bp 5' of the transcription start site in the human *HR* gene, confers varying degrees of ligand-dependent transactivation in HEK-293 (Fig. 4B), HaCaT (Fig. 4C), and COS7 (Fig. 4D) cells. The ability of the –7269 VDRE to activate luciferase also contains a significant ligand-independent component in HaCaT cells (Fig. 4C) and especially in COS7 cells (Fig. 4D). The molecular basis of this cell-type-specific activity is not defined as yet, but it may be generated by differential expression of transcriptional comodulators which are then recruited when RXR–VDR is conformed by docking on this particular responsive element, which we have designated as a ‘constitutive’ VDRE (Fig. 6, central right object). The phenomenon of vitamin D ligand-independent transactivation mediated by a VDRE has been reported by MacDonald and colleagues (Ellison *et al.* 2007) for the human *CYP24A1* promoter, but only when it is transfected into normal human keratinocytes, and not into either transformed keratinocytes or normal fibroblasts. A foundation for ligand independence may reside primarily in the precise sequence of the VDRE, providing there exists support for VDR–RXR heterodimerization other than via ligand binding, as there apparently does in the case of the *CYP24A1* promoter in normal human keratinocytes (Ellison *et al.* 2007). The conclusion herein is that vitamin D ligand-unoccupied VDR appears to be capable of transactivation of *HR* via its constitutive VDRE (Fig. 6, centre right), triggering the switch for progression of the hair cycle through subsequent repression of *SOSTDC1* (Fig. 6, upper right object) and possibly of *DKKL1* (Fig. 6, lower right object). Alternative mechanisms explaining vitamin D ligand-independent transactivation by VDR include: VDR activation via phosphorylation by a cell-selective kinase, cell- or promoter-specific coactivation of unoccupied VDR–RXR, and endogenous synthesis of an alternative lipophilic, nonvitamin D VDR ligand.

The role of thyroid hormone and its receptor, TR β , in the hair follicle (Billoni *et al.* 2000) does not appear to be as crucial as the roles of VDR and Hr. However, human patients who are hypothyroid or hyperthyroid (van Beek

et al. 2008) or who have mutations in TR β (Guran *et al.* 2009) show a diffuse hair loss or hair thinning, although neither perturbations in thyroid hormone status nor receptor ablation fully mimic the *hr* knockout phenotype (van Beek *et al.* 2008). In this study, we identified an atypical TRE, TGGTGAggccAGGACA, at +1304 bp within the first intron of the human HR gene (Fig. 5A). This novel TRE confers ligand-independent transactivation onto a heterologous reporter gene in COS7 cells (Fig. 5C). This study thus reveals the existence of two ligand-independent elements, a TRE and a VDRE, both of which contain a thymidine residue in the first position of the first half-element (Fig. 5D). A thymidine in this position is neither found in previously characterized DR3 VDREs (Whitfield *et al.* 2005) nor is it observed in consensus DR4 TREs (Umesono *et al.* 1991). We propose that a thymidine in this position, corresponding to the binding site for the RXR heterodimeric partner of either VDR (Jin & Pike 1996) or TR (Perlmann *et al.* 1993), may cause a conformational change in RXR that is then transferred to the primary receptor (VDR or TR), resulting in the ability of the heterodimer to attract coactivators and thereby function predominantly as a ligand-independent transactivator of the HR gene. The plausibility of such a mechanism was demonstrated in a recent study of VDR–RXR binding to two different VDRE sequences (Zhang *et al.* 2011), in which the point was made that VDR–RXR heterodimer binding to differing DNA sequences can relay information to the ligand-binding domain and specifically to the AF-2 C-terminal helix domain of VDR that makes contact with coactivators. It is therefore implied that unique nuclear receptor responsive elements may exist in genes which are regulated by unliganded nuclear receptor complexes, many of which are indeed observed bound to DNA in the human genome (Meyer *et al.* 2012). In conclusion, although the three novel VDREs identified herein exist in genes that are well established to affect the hair cycle, further research including chromatin immunoprecipitation-sequencing (ChIP-seq) experiments is required to prove that the present findings on soggy, Wise, and hairless close the gap in our biological understanding of control of the hair cycle and/or epidermal keratinocyte function.

Importantly, although the present communication emphasizes the findings as they may relate to the mammalian hair cycle, no actual hair cycle analyses were carried out. The current experiments used keratinocytes, not purified bulge stem cells, and no investigation of the functioning of bulge stem cells or their progeny after 1,25D treatment, *in vivo*, has been performed.

The observed regulation by 1,25D of HR, *SOSTDC1*, and *DKKL1* could constitute epidermal keratinocyte phenomena. Thus, an alternative interpretation of the results herein is that they are more pertinent to epidermal keratinocyte function, than to regulation of the hair cycle, *per se*. Indeed, 1,25D induces the expression of a number of genes in cultured keratinocytes, the products of which are potential prodifferentiative and structural components, as well as detoxification, immunomodulation, and anti-inflammatory/anti-oxidation principles (Bikle 2012). For example, 1,25D induces caspase-14 in keratinocytes (Haussler *et al.* 2013). This nonapoptotic caspase is crucial for keratinocyte differentiation (Rendl *et al.* 2002). 1,25D induces cathelicidin and several defensins in keratinocytes (Bikle 2012), indicating that vitamin D modulates the immune complement in skin. Also, late cornified envelope (*LCE-1D*, *-1F*, *-2B*) genes in the epidermal differentiation complex are induced by 1,25D in human keratinocytes (Haussler *et al.* 2013). Finally, 1,25D increases the expression of a number of keratin-related gene products (Haussler *et al.* 2013) which, when considered along with the LCE proteins, indicates that vitamin D signaling supports the skin structurally and mediates barrier function development. Therefore, inductive epidermal and mesenchyme signaling after 1,25D treatment is more likely to influence specific epidermal cell properties than to play a part in the hair cycle as shown by previous studies (e.g. hair reconstitution assays). In summary, VDR functions to drive the mammalian hair cycle in cooperation with Hr, primarily via the repression of gene expression, whereas 1,25D acts via VDR binding to signal transcription of genes for which the products support the development and barrier function of the skin.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was supported by National Institutes of Health grant NIH DK033351 to M R H, NIH CA140285 to P W J, and a grant from the Arizona Biomedical Research Commission to J-C H.

Author contribution statement

Conception and design: J-C H, G K W, P W J, and M R H. Development of methodology: J-C H, R C E, G K W, and I K. Acquisition of data: J-C H, R C E, G K W, and I K. Writing, review, and/or revision of the manuscript: J-C H, G K W, P W J, and M R H. Administrative, technical, or material support: J-C H, G K W, P W J, and M R H. Study supervision: J-C H and M R H.

Acknowledgements

The authors are grateful to Dr Yifei Wu for technical assistance.

References

- Ahmad W, Zlotogorski A, Panteleyev AA, Lam H, Ahmad M, ul Haque MF, Abdallah HM, Dragan L & Christiano AM 1999 Genomic organization of the human hairless gene (HR) and identification of a mutation underlying congenital atrichia in an Arab Palestinian family. *Genomics* **56** 141–148. (doi:10.1006/geno.1998.5699)
- Beaudoin GM III, Sisk JM, Coulombe PA & Thompson CC 2005 Hairless triggers reactivation of hair growth by promoting Wnt signaling. *Proceedings of the National Academy of Sciences USA* **102** 14653–14658. (doi:10.1073/pnas.0507609102)
- van Beek N, Bodo E, Kromminga A, Gaspar E, Meyer K, Zmijewski MA, Slominski A, Wenzel BE & Paus R 2008 Thyroid hormones directly alter human hair follicle functions: anagen prolongation and stimulation of both hair matrix keratinocyte proliferation and hair pigmentation. *Journal of Clinical Endocrinology and Metabolism* **93** 4381–4388. (doi:10.1210/jc.2008-0283)
- Bergman R, Schein-Goldshmid R, Hochberg Z, Ben-Izhak O & Sprecher E 2005 The alopecias associated with vitamin D-dependent rickets type IIA and with hairless gene mutations: a comparative clinical, histologic, and immunohistochemical study. *Archives of Dermatology* **141** 343–351. (doi:10.1001/archderm.141.3.343)
- Bikle DD 2012 Vitamin D and the skin: Physiology and pathophysiology. *Reviews in Endocrine and Metabolic Disorders* **13** 3–19. (doi:10.1007/s11154-011-9194-0)
- Billoni N, Buan B, Gautier B, Gaillard O, Mahe YF & Bernard BA 2000 Thyroid hormone receptor β 1 is expressed in the human hair follicle. *British Journal of Dermatology* **142** 645–652. (doi:10.1046/j.1365-2133.2000.03408.x)
- Cachon-Gonzalez MB, Fenner S, Coffin JM, Moran C, Best S & Stoye JP 1994 Structure and expression of the hairless gene of mice. *Proceedings of the National Academy of Sciences USA* **91** 7717–7721. (doi:10.1073/pnas.91.16.7717)
- Cachon-Gonzalez MB, San-Jose I, Cano A, Vega JA, Garcia N, Freeman T, Schimmang T & Stoye JP 1999 The hairless gene of the mouse: relationship of phenotypic effects with expression profile and genotype. *Developmental Dynamics* **216** 113–126. (doi:10.1002/(SICI)1097-0177(199910)216:2<113::AID-DVDY3>3.0.CO;2-M)
- Chuma M, Endo-Umeda K, Shimba S, Yamada S & Makishima M 2012 Hairless modulates ligand-dependent activation of the vitamin D receptor–retinoid X receptor heterodimer. *Biological & Pharmaceutical Bulletin* **35** 582–587. (doi:10.1248/bpb.35.582)
- Cianferotti L, Cox M, Skoriya K & Demay MB 2007 Vitamin D receptor is essential for normal keratinocyte stem cell function. *Proceedings of the National Academy of Sciences USA* **104** 9428–9433. (doi:10.1073/pnas.0702884104)
- Ellison TI, Eckert RL & MacDonald PN 2007 Evidence for 1,25-dihydroxyvitamin D₃-independent transactivation by the vitamin D receptor: uncoupling the receptor and ligand in keratinocytes. *Journal of Biological Chemistry* **282** 10953–10962. (doi:10.1074/jbc.M609717200)
- Engelhard A & Christiano AM 2004 The hairless promoter is differentially regulated by thyroid hormone in keratinocytes and neuroblastoma cells. *Experimental Dermatology* **13** 257–264. (doi:10.1111/j.0906-6705.2004.00175.x)
- Engelhard A, Bauer RC, Casta A, Djabali K & Christiano AM 2008 Ligand-independent regulation of the hairless promoter by vitamin D receptor. *Photochemistry and Photobiology* **84** 515–521. (doi:10.1111/j.1751-1097.2008.00301.x)
- Fuchs E, Merrill BJ, Jamora C & DasGupta R 2001 At the roots of a never-ending cycle. *Developmental Cell* **1** 13–25. (doi:10.1016/S1534-5807(01)00022-3)
- Guran T, Bircan R, Turan S & Bereket A 2009 Alopecia: association with resistance to thyroid hormones. *Journal of Pediatric Endocrinology & Metabolism* **22** 1075–1081. (doi:10.1515/JPEM.2009.22.11.1075)
- Haussler MR, Haussler CA, Whitfield GK, Hsieh JC, Thompson PD, Barthel TK, Bartik L, Egan JB, Wu Y, Kubicek JL *et al.* 2010 The nuclear vitamin D receptor controls the expression of genes encoding factors which feed the “Fountain of Youth” to mediate healthful aging. *Journal of Steroid Biochemistry and Molecular Biology* **121** 88–97. (doi:10.1016/j.jsbmb.2010.03.019)
- Haussler MR, Whitfield GK, Kaneko I, Haussler CA, Hsieh D, Hsieh JC & Jurutka PW 2013 Molecular mechanisms of vitamin D action. *Calcified Tissue International* **92** 77–98. (doi:10.1007/s00223-012-9619-0)
- Hsieh J-C, Jurutka PW, Galligan MA, Terpening CM, Haussler CA, Samuels DS, Shimizu Y, Shimizu N & Haussler MR 1991 Human vitamin D receptor is selectively phosphorylated by protein kinase C on serine S1, a residue crucial to its trans-activation function. *Proceedings of the National Academy of Sciences USA* **88** 9315–9319. (doi:10.1073/pnas.88.20.9315)
- Hsieh J-C, Sisk JM, Jurutka PW, Haussler CA, Slater SA, Haussler MR & Thompson CC 2003a Physical and functional interaction between the vitamin D receptor and hairless corepressor, two proteins required for hair cycling. *Journal of Biological Chemistry* **278** 38665–38674. (doi:10.1074/jbc.M304886200)
- Hsieh J-C, Whitfield GK, Jurutka PW, Haussler CA, Thatcher ML, Thompson PD, Dang HTL, Galligan MA, Oza AK & Haussler MR 2003b Two basic amino acids C-terminal of the P-box specify functional binding of the vitamin D receptor to its rat osteocalcin DNA responsive element. *Endocrinology* **144** 5065–5080. (doi:10.1210/en.2003-0635)
- Hsieh JC, Slater SA, Whitfield GK, Dawson JL, Hsieh G, Sheedy C, Haussler CA & Haussler MR 2010 Analysis of hairless corepressor mutants to characterize molecular cooperation with the vitamin D receptor in promoting the mammalian hair cycle. *Journal of Cellular Biochemistry* **110** 671–686. (doi:10.1002/jcb.22578)
- Huelsken J, Vogel R, Erdmann B, Cotsarelis G & Birchmeier W 2001 β -Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. *Cell* **105** 533–545. (doi:10.1016/j.bbr.2011.03.031)
- Jin CH & Pike JW 1996 Human vitamin D receptor-dependent transactivation in *Saccharomyces cerevisiae* requires retinoid X receptor. *Molecular Endocrinology* **10** 196–205. (doi:10.1210/me.10.2.196)
- Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM & Haussler D 2002 The human genome browser at UCSC. *Genome Research* **12** 996–1006. (doi:10.1101/gr.229102)
- Klein I, Bergman R, Indelman M & Sprecher E 2002 A novel missense mutation affecting the human hairless thyroid receptor interacting domain 2 causes congenital atrichia. *Journal of Investigative Dermatology* **119** 920–922. (doi:10.1046/j.1523-1747.2002.00268.x)
- Kohn MJ, Kaneko KJ & DePamphilis ML 2005 DkkL1 (Soggy), a Dickkopf family member, localizes to the acrosome during mammalian spermatogenesis. *Molecular Reproduction and Development* **71** 516–522. (doi:10.1002/mrd.20314)
- Lintern KB, Guidato S, Rowe A, Saldanha JW & Itasaki N 2009 Characterization of wise protein and its molecular mechanism to interact with both Wnt and BMP signals. *Journal of Biological Chemistry* **284** 23159–23168. (doi:10.1074/jbc.M109.025478)
- Liu L, Kim H, Casta LC, Kobayashi Y, Shapiro LS & Christiano AM 2011 Hairless is a H3K9 histone demethylase. *Journal of Investigative Dermatology* **131** S69. (doi:10.1038/jid.2011.75)
- Malloy PJ, Wang J, Jensen K & Feldman D 2009 Modulation of vitamin D receptor activity by the corepressor hairless: differential effects of hairless isoforms. *Endocrinology* **150** 4950–4957. (doi:10.1210/en.2009-0358)
- Meyer MB, Goetsch PD & Pike JW 2012 VDR/RXR and TCF4/ β -catenin cistromes in colonic cells of colorectal tumor origin: impact on c-FOS

- and c-MYC gene expression. *Molecular Endocrinology* **26** 37–51. (doi:10.1210/me.2011-1109)
- Mikkelsen TS, Ku M, Jaffe DB, Issac B, Lieberman E, Giannoukos G, Alvarez P, Brockman W, Kim TK, Koche RP *et al.* 2007 Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* **448** 553–560. (doi:10.1038/nature06008)
- Miller J, Djabali K, Chen T, Liu Y, Ioffreda M, Lyle S, Christiano AM, Holick M & Cotsarelis G 2001 Atrichia caused by mutations in the vitamin D receptor gene is a phenocopy of generalized atrichia caused by mutations in the hairless gene. *Journal of Investigative Dermatology* **117** 612–617. (doi:10.1046/j.0022-202x.2001.01438.x)
- Moraitis AN & Giguere V 2003 The co-repressor hairless protects ROR α orphan nuclear receptor from proteasome-mediated degradation. *Journal of Biological Chemistry* **278** 52511–52518. (doi:10.1074/jbc.M308152200)
- Niehrs C 2006 Function and biological roles of the Dickkopf family of Wnt modulators. *Oncogene* **25** 7469–7481. (doi:10.1038/sj.onc.1210054)
- O'Shaughnessy RF, Christiano AM & Jahoda CA 2004 The role of BMP signalling in the control of ID3 expression in the hair follicle. *Experimental Dermatology* **13** 621–629. (doi:10.1111/j.0906-6705.2004.00206.x)
- Panteleyev AA, Paus R & Christiano AM 2000 Patterns of hairless (*hr*) gene expression in mouse hair follicle morphogenesis and cycling. *American Journal of Pathology* **157** 1071–1079. (doi:10.1016/S0002-9440(10)64621-4)
- Perlmann T, Rangarajan PN, Umesono K & Evans RM 1993 Determinants for selective RAR and TR recognition of direct repeat HREs. *Genes and Development* **7** 1411–1422. (doi:10.1101/gad.7.7b.1411)
- Potter GB, Beaudoin GM III, DeRenzo CL, Zarach JM, Chen SH & Thompson CC 2001 The *hairless* gene mutated in congenital hair loss disorders encodes a novel nuclear receptor corepressor. *Genes and Development* **15** 2687–2701. (doi:10.1101/gad.916701)
- Ramot Y, Horev L, Smolovich I, Molho-Pessach V & Zlotogorski A 2010 Marie Unna hereditary hypotrichosis caused by a novel mutation in the human hairless transcript. *Experimental Dermatology* **19** e320–e322. (doi:10.1111/j.1600-0625.2009.01042.x)
- Rendl M, Ban J, Mrass P, Mayer C, Lengauer B, Eckhart L, Declerq W & Tschachler E 2002 Caspase-14 expression by epidermal keratinocytes is regulated by retinoids in a differentiation-associated manner. *Journal of Investigative Dermatology* **119** 1150–1155. (doi:10.1046/j.1523-1747.2002.19532.x)
- Sakai Y, Kishimoto J & Demay MB 2001 Metabolic and cellular analysis of alopecia in vitamin D receptor knockout mice. *Journal of Clinical Investigation* **107** 961–966. (doi:10.1172/JCI11676)
- Skorija K, Cox M, Sisk JM, Dowd DR, MacDonald PN, Thompson CC & Demay MB 2005 Ligand-independent actions of the vitamin D receptor maintain hair follicle homeostasis. *Molecular Endocrinology* **19** 855–862. (doi:10.1210/me.2004-0415)
- Teichert A, Elalieh H & Bikle D 2010 Disruption of the hedgehog signaling pathway contributes to the hair follicle cycling deficiency in Vdr knockout mice. *Journal of Cellular Physiology* **225** 482–489. (doi:10.1002/jcp.22227)
- Thompson CC, Sisk JM & Beaudoin GM III 2006 Hairless and Wnt signaling: allies in epithelial stem cell differentiation. *Cell Cycle* **5** 1913–1917. (doi:10.4161/cc.5.17.3189)
- Umesono K, Murakami KK, Thompson CC & Evans RM 1991 Direct repeats as selective response elements for the thyroid hormone, retinoic acid, and vitamin D₃ receptors. *Cell* **65** 1255–1266. (doi:10.1016/0092-8674(91)90020-Y)
- Wang J, Malloy PJ & Feldman D 2007 Interactions of the vitamin D receptor with the corepressor hairless: analysis of hairless mutants in atrichia with papular lesions. *Journal of Biological Chemistry* **282** 25231–25239. (doi:10.1074/jbc.M702939200)
- Wen Y, Liu Y, Xu Y, Zhao Y, Hua R, Wang K, Sun M, Li Y, Yang S, Zhang XJ *et al.* 2009 Loss-of-function mutations of an inhibitory upstream ORF in the human hairless transcript cause Marie Unna hereditary hypotrichosis. *Nature Genetics* **41** 228–233. (doi:10.1038/ng.276)
- Whitfield GK, Jurutka PW, Haussler CA, Hsieh JC, Barthel TK, Jacobs ET, Encinas Dominguez C, Thatcher ML & Haussler MR 2005 Nuclear vitamin D receptor: structure–function, molecular control of gene transcription, and novel bioactions. In *Vitamin D*, 2nd edn, pp 219–261. Eds D Feldman, JW Pike & FH Glorieux. Oxford, UK: Elsevier Academic Press.
- Xie Z, Chang S, Oda Y & Bikle DD 2006 Hairless suppresses vitamin D receptor transactivation in human keratinocytes. *Endocrinology* **147** 314–323. (doi:10.1210/en.2005-1111)
- Yamamoto Y, Memezawa A, Takagi K, Ochiai S, Shindo M & Kato S 2009 A tissue-specific function by unliganded VDR. Abstracts from the 14th Workshop on Vitamin D, Brugge, Belgium (October 4–8, 2009 66).
- Zarach JM, Beaudoin GM III, Coulombe PA & Thompson CC 2004 The co-repressor hairless has a role in epithelial cell differentiation in the skin. *Development* **131** 4189–4200. (doi:10.1242/dev.01303)
- Zhang J, Chalmers MJ, Stayrook KR, Burris LL, Wang Y, Busby SA, Pascal BD, Garcia-Ordóñez RD, Bruning JB, Istrate MA *et al.* 2011 DNA binding alters coactivator interaction surfaces of the intact VDR–RXR complex. *Nature Structural & Molecular Biology* **18** 556–563. (doi:10.1038/nsmb.2046)

Received in final form 30 October 2013

Accepted 4 November 2013

Accepted Preprint published online 4 November 2013