Determinants of GH resistance in malnutrition

Pouneh K Fazeli and Anne Klibanski

Neuroendocrine Unit, Massachusetts General Hospital, 55 Fruit Street, Bulfinch 457, Boston, Massachusetts 02114, USA

Correspondence should be addressed to P K Fazeli
Email
pkfazeli@partners.org

Abstract

States of undernutrition are characterized by GH resistance. Decreased total energy intake, as well as isolated protein–calorie malnutrition and isolated nutrient deficiencies, result in elevated GH levels and low levels of IGF1. We review various states of malnutrition and a disease state characterized by chronic undernutrition – anorexia nervosa – and discuss possible mechanisms contributing to the state of GH resistance, including fibroblast growth factor 21 and Sirtuin 1. We conclude by examining the hypothesis that GH resistance is an adaptive response to states of undernutrition, in order to maintain euglycemia and preserve energy.

Key Words

- ▶ growth hormone
- ▶ IGF
- ▶ neuroendocrinology
- ▶ pituitary

Journal of Endocrinology
(2014) **220**, R57–R65

Introduction

Insulin-like growth factor 1 (IGF1) is a hormone produced primarily by the liver and regulated by growth hormone (GH) secretion by the somatotroph cells of the anterior pituitary gland. Many, but not all, of the actions of GH are mediated by IGF1. In addition to regulation by GH, IGF1 secretion is also responsive to nutritional cues. In obesity, a state of overnutrition, GH secretion is decreased and may result in IGF1 levels that are lower than those in normal-weight individuals (Maccario *et al.* 1997), although bioactive IGF1 levels may be comparable to those of lean controls (Frydryk *et al.* 2009). In states of undernutrition, GH levels are normal or elevated in the setting of low IGF1 levels; therefore, there is a state of GH resistance, with an inappropriate response to GH at the level of the liver. This state of acquired GH resistance is likely an adaptive response to decreased energy intake. A number of potential mechanisms of GH resistance in malnutrition have been elucidated, including possible hormonal determinants. This review will examine the current

evidence of determinants of GH resistance in states of malnutrition (Fig. 1).

States of malnutrition

A number of models of malnutrition have been studied to better understand the state of GH resistance in undernutrition. These models include total energy deficiency, isolated protein deficiency (kwashiorkor) – a state in which individuals consume an appropriate number of calories but have profoundly low levels of protein intake – and anorexia nervosa, a psychiatric disease that is characterized by low body weight and self-induced starvation. There are a number of other types of malnutrition as well, such as isolated nutrient deficiencies, which may also result in GH resistance. These various states of malnutrition will be discussed in detail below.

Figure 1

Potential causes and mediators of growth hormone (GH) resistance. Starvation, protein-calorie malnutrition, and isolated vitamin deficiencies can lead to a state of GH resistance, with normal or elevated levels of GH coincident with low levels of insulin-like growth factor 1 (IGF1).

Recent evidence suggests that the state of GH resistance in starvation may result from a decrease in STAT5 phosphorylation, mediated by fibroblast growth factor 21 (FGF21) and/or Sirtuin 1 (SIRT1).

Starvation

Malnutrition may be due to isolated energy deficiencies – such as protein-calorie deficiency – or total energy deficiency, the extreme case of which is starvation. States of total energy deficiency result in a GH resistance state. IGF1 levels decrease by ~50% after only 4 days of fasting (Grinspoon *et al.* 1995). Fasting individuals have elevated GH levels coupled with the low IGF1 levels (Fichter *et al.* 1986, Ho *et al.* 1988) and may also have hyperresponsive GH secretion after thyrotropin-releasing hormone stimulation (Shimizu *et al.* 1991). The fact that endogenous hypersecretion of GH does not result in elevated IGF1 levels suggests that fasting either decreases GH receptors in the liver or there is a post-receptor defect resulting in an inability of GH to stimulate IGF1 production. In animal models of starvation, low IGF1 levels are coupled with decreased GH receptor mRNA (Straus & Takemoto 1990) and decreased GH binding (Baxter *et al.* 1981), suggesting that GH receptor downregulation in the liver contributes to the state of GH resistance.

Exogenous GH is also unable to stimulate IGF1 production during fasting. The effects of exogenous GH in the fed vs the fasted state were investigated in subjects with isolated GH deficiency (Merimee *et al.* 1982). Subjects were treated with 5 mg GH twice/day for 5 days, either while consuming an *ad libitum* diet or during a 5-day fast (Merimee *et al.* 1982). In response to GH, IGF1 levels increased tenfold in the fed state but only doubled in the

fasted state, demonstrating resistance to exogenous GH during fasting (Merimee *et al.* 1982). Interestingly, in a study of acromegals undergoing a 36-h fast, serum IGF1 levels did not decrease in the acromegals whereas levels decreased in the fasted control patients (Grottoli *et al.* 2008). Therefore, it is likely that adaptations to fasting are abnormal in individuals with baseline GH hypersecretion, although the effects of a longer term fast in individuals with acromegaly are unknown.

Protein deficiency

Isolated protein deficiency, with otherwise normal energy intake, also results in a state of GH resistance. Studies in children with protein-calorie malnutrition demonstrate elevated GH levels and low IGF1 levels with normalization after nutritional recovery (Olusi *et al.* 1977, Robinson & Picou 1977, Hintz *et al.* 1978, Soliman *et al.* 1986). In adults who were fasted and then re-fed a protein-deficient diet, IGF1 levels increased significantly less than in adults fed an iso-caloric, protein-sufficient diet (Isley *et al.* 1983).

Importantly, in animal models with low IGF1 levels due to protein deficiency, pharmacological doses of GH do not increase IGF1 levels to the normal range or normalize growth (Thissen *et al.* 1990, 1991a), but increasing dietary protein does increase IGF1 levels. In animal models, varying the protein content of food did not affect plasma GH levels, whereas IGF1 levels increased significantly with increasing protein content (Reeves *et al.* 1979). Similarly, in a

population of elderly individuals, 6 months of protein supplementation increased IGF1 levels by 85% (Schurch *et al.* 1998) and in individuals with type 2 diabetes mellitus, IGF1 levels increased by 30% in those fed a diet consisting of 30% protein for 5 weeks compared with those fed a diet consisting of 15% protein (Gannon & Nuttall 2011).

Unlike in starvation, in which a decrease in GH receptors is likely contributing to the GH-resistant state, in protein deficiency, the GH resistance state is likely due to a post-receptor defect. In animal models of protein deficiency, GH binding is not reduced (Maes *et al.* 1988, Thissen *et al.* 1990) and *Igf1* mRNA production in response to exogenous GH is similar in hypophysectomized rats on a protein-deficient vs protein-sufficient diet (Thissen *et al.* 1991a). Protein deficiency not only results in GH resistance but also likely results in a state of end-organ resistance to IGF1. Rats with low IGF1 levels due to protein restriction and a control population of rats with low IGF1 levels due to hypophysectomy were treated with recombinant human IGF1 at a dose of 300 µg/day (Thissen *et al.* 1991b). Serum IGF1 levels normalized in both groups but increases in body weight, tibial epiphyseal widening, and tail growth only occurred in the hypophysectomized rats, not in the protein-restricted rats (Thissen *et al.* 1991b). Therefore, in addition to a state of GH resistance, there may also be a state of IGF1 resistance in protein restriction.

Anorexia nervosa

Anorexia nervosa is a psychiatric disorder, predominantly affecting women, with a lifetime prevalence of 2.2% (Keski-Rahkonen *et al.* 2007). The disease is characterized by self-imposed starvation and an inability to maintain a normal weight for height (American Psychiatric Association 1994). Individuals with anorexia nervosa are in a state of chronic starvation. Like individuals with total energy deficiency and isolated protein deficiency, individuals with anorexia nervosa are GH resistant. In girls and women with anorexia nervosa, basal and pulsatile GH secretion is increased and IGF1 levels are low (Garfinkel *et al.* 1975, Scacchi *et al.* 1997, Stoving *et al.* 1999, Misra *et al.* 2003). Systemic IGF1 levels in anorexia nervosa are ~50% of those of normal weight women and levels of IGF-binding protein 3, the predominant binding protein for IGF1, are low (Counts *et al.* 1992). Yet, bioactive IGF1 levels are also low (Stoving *et al.* 2007), suggesting that the low IGF1 levels in anorexia nervosa are not simply due to decreased levels of binding proteins but due to decreased bioactive IGF1. The low IGF1 levels in women with anorexia nervosa are exquisitely sensitive to nutritional repletion – after

only 3 days of hyperalimentation therapy, levels have been shown to increase by ~50% (Hotta *et al.* 2000).

Individuals with anorexia nervosa also have an exaggerated response to GH-releasing hormone (GHRH) (Rolla *et al.* 1990) and abnormal GH suppressibility after a glucose load (Tamai *et al.* 1991). In healthy individuals, pretreatment with a cholinergic muscarinic antagonist blocks GH secretion in response to GH-releasing factor (Massara *et al.* 1984). The mechanism by which this occurs may be an increase in the release of somatostatin – a potent inhibitor of GH secretion – by the hypothalamus. By contrast, in girls with anorexia nervosa, pretreatment with a cholinergic muscarinic antagonist does not block the exaggerated GH response to GHRH (Rolla *et al.* 1990, Tamai *et al.* 1990), suggesting that there may be an abnormality in the release of or response to somatostatin.

We investigated whether supraphysiological doses of GH can overcome the state of GH resistance in women with anorexia nervosa (Fazeli *et al.* 2010a). Subjects ($n=21$) were randomized to either placebo or recombinant human GH treatment (mean maximum daily dose: 1.4 mg/day) for 12 weeks (Fazeli *et al.* 2010a). At the conclusion of the study, IGF1 levels did not differ between the groups, demonstrating that even very high levels of exogenous GH cannot overcome the insensitivity to GH, at the level of the liver, in anorexia nervosa (Fazeli *et al.* 2010a). One purported mechanism for the GH resistance state in anorexia nervosa is the downregulation of GH receptors in the liver (Counts *et al.* 1992). The failure of supraphysiological doses of GH to increase IGF1 levels supports this hypothesis.

Isolated nutrient deficiencies

Isolated vitamin deficiencies may also cause a state of GH resistance. Vitamin A-deficient rats have lower IGF1 levels compared with pair-fed controls despite comparable GH levels (Mohan & Jaya Rao 1980). Compared with similar weight controls, children with vitamin A deficiency have normal GH levels coincident with lower IGF1 levels (Mohan & Jaya Rao 1979). Similarly, rats fed a diet deficient in vitamin B6 for 4 weeks were found to have low IGF1 levels and normal GH levels (Rao & Mohan 1982).

In a rodent model, both zinc deficiency and magnesium deficiency are associated with decreased IGF1 levels in the setting of normal GH levels (Dorup *et al.* 1991). Young rats fed a magnesium-deficient diet were found to have significantly lower serum IGF1 levels compared with pair-fed controls, whereas basal GH levels and GH levels after stimulation with GH-releasing factor were similar in the magnesium-deficient group compared with an *ad libitum*-fed

group (Dorup *et al.* 1991). Similarly, young rats fed a zinc-deficient diet were also found to have significantly lower serum IGF1 levels compared with pair-fed controls but levels of GH were similar in the zinc-deficient rats and the *ad libitum*-fed zinc-sufficient rats after treatment with GH-releasing factor, suggesting a state of GH resistance (Dorup *et al.* 1991). In postmenopausal women, zinc intake is positively associated with IGF1 levels, even after adjustment for possible confounders, but because GH levels were not measured in this study, it is not known whether GH resistance contributes to the lower IGF1 levels (Devine *et al.* 1998).

Potassium-deficient diets have also been studied in an animal model (Flyvbjerg *et al.* 1991). Rats consuming a potassium-deficient fodder had significantly lower serum IGF1 levels compared with pair-fed controls, but in this case, the response to GH-releasing factor stimulation was significantly less in the potassium-deficient group compared with the controls (Flyvbjerg *et al.* 1991). Therefore, the low IGF1 levels in potassium-deficient rodents are not due to GH resistance but likely due to downregulation of an appropriately functioning GH-IGF1 axis.

Effects of dietary composition

In *in vitro* studies, saturated fatty acids, but not unsaturated fatty acids, inhibit promoter activity at the GH receptor gene and thereby decrease GH receptor mRNA and protein levels (Thimmarayappa *et al.* 2006), suggesting that saturated fatty acids may contribute to GH resistance. In a large cohort of healthy women ($n=1000$), a cross-sectional examination of plasma IGF1 levels and various dietary components demonstrated a positive association between IGF1 levels and total energy intake (Holmes *et al.* 2002). There was also a positive association between IGF1 levels and increasing levels of protein intake (Holmes *et al.* 2002). Smaller human studies ($n=115$) have also investigated the cross-sectional relationship between IGF1 and dietary composition (Kaklamani *et al.* 1999). Positive associations between IGF1 and fats, red meat, and oil consumption have been reported as well as an inverse association between IGF1 levels and carbohydrate intake (Kaklamani *et al.* 1999). Importantly, GH levels were not measured in these studies and therefore we do not know whether these differences in IGF1 levels are GH mediated or due to GH insensitivity.

Mechanisms of GH resistance

GH is secreted by somatotroph cells in the anterior pituitary gland and binds to GH receptors in many tissues,

including hepatocytes. The binding of GH to its receptor activates JAK2, which leads to phosphorylation of STAT5. STAT5 is subsequently translocated into the nucleus where it is able to bind to regulatory elements of target genes including IGF1. Recently, two proteins have been shown to be important regulators of GH resistance in states of nutritional deprivation: fibroblast growth factor 21 (FGF21) and Sirtuin 1 (SIRT1). Both FGF21 and SIRT1 induce GH resistance via STAT5 inhibition.

Fibroblast growth factor 21

FGF21, a member of the FGF family of proteins, is a hormone produced in the liver (Nishimura *et al.* 2000) and adipocytes (Zhang *et al.* 2008). Serum FGF21 levels correlate with BMI and also with *FGF21* mRNA expression in subcutaneous fat (Zhang *et al.* 2008), suggesting that serum FGF21 levels primarily originate in adipose tissue. FGF21 also stimulates insulin-independent glucose uptake in adipocytes via the glucose transporter-1 (Kharitonov *et al.* 2005) and is currently being investigated as a therapeutic agent in individuals with type 2 diabetes mellitus (Gaich *et al.* 2013). In animal models, starvation induces FGF21 production in the liver via a mechanism requiring peroxisome proliferator-activated receptor α (Inagaki *et al.* 2007, Lundasen *et al.* 2007). FGF21 subsequently induces peroxisome proliferator-activated receptor γ coactivator protein 1 α (PGC1 α) expression, leading to fatty acid oxidation and increased gluconeogenesis (Potthoff *et al.* 2009). In humans, a very low-calorie diet and prolonged fasting result in increased levels of FGF21 (Galman *et al.* 2008, Mraz *et al.* 2009). Therefore, FGF21 may act differently in states of nutritional sufficiency compared with states of nutritional deprivation.

In a transgenic animal model, *Fgf21* transgenic mice have high levels of GH and significantly lower levels of IGF1 compared with WT mice (Inagaki *et al.* 2008), consistent with GH resistance. The mechanism of GH resistance in *Fgf21* transgenic mice is a decrease in STAT5 phosphorylation (Inagaki *et al.* 2008). Similarly, we have shown that FGF21 levels are significantly higher in adolescent girls with anorexia nervosa compared with normal-weight girls after controlling for percent body fat and insulin resistance, suggesting that the increased levels of FGF21 in anorexia nervosa are due to increased levels of liver-derived FGF21 (Fazeli *et al.* 2010b). We have also demonstrated a significant positive relationship between GH area under the curve and FGF21 and an inverse relationship between IGF1 and elevated levels of FGF21,

after controlling for percent body fat and insulin resistance (Fazeli *et al.* 2010b). These data suggest that FGF21 may be a mediator of GH resistance in the human model of chronic starvation – anorexia nervosa.

Sirtuin 1

SIRT1, a class III histone deacetylase, is regulated by nutrient availability and promotes gluconeogenesis and fatty acid oxidation during periods of fasting (Gillum *et al.* 2010). Recently, SIRT1 has been shown to play a role in GH resistance in states of starvation (Yamamoto *et al.* 2013). SIRT1 knockdown mice that underwent a 48-h fast had higher serum levels of IGF1 and higher *Igf1* mRNA levels in the liver when compared with WT fasted mice, suggesting that SIRT1 mediates GH resistance in states of undernutrition (Yamamoto *et al.* 2013). Like FGF21, the mechanism by which SIRT1 acts is a reduction in STAT5 phosphorylation; this decrease in STAT5 phosphorylation is a result of SIRT1 deacetylating lysine residues on STAT5 (Yamamoto *et al.* 2013).

Insulin

Type 1 diabetes mellitus is also a state of GH resistance (Yde 1969, Lundbaek *et al.* 1970, Horner *et al.* 1981). Therefore, insulin has also been investigated as a possible determinant of GH resistance. GH receptors in the liver are reduced in diabetic female rats and this decrease correlates with the decreased serum insulin levels (Baxter & Turtle 1978). GH receptors also increase in response to treatment with insulin (Baxter & Turtle 1978) and *in vitro* models demonstrate upregulation of GH receptors by insulin (Leung *et al.* 2000). Therefore, insulin levels, which are low in states of nutritional deprivation such as fasting, may in part mediate GH resistance by downregulation of GH receptors in the liver. Although, this is unlikely to be the only mechanism by which GH resistance is induced in states of undernutrition, as studies in diabetic rats have demonstrated independent effects of insulin and a low-protein diet (Maiter *et al.* 1989).

Insulin-like growth factor 1

The low IGF1 levels in the GH resistance state – and the resultant decrease in negative feedback at the level of the pituitary – contribute to the elevated GH levels characteristic of GH resistance. In a study of healthy, fasting men, a recombinant human IGF1 infusion resulted in a significant decrease in serum GH levels (Hartman *et al.* 1993).

Similarly, in a study of women with anorexia nervosa, administration of recombinant human IGF1 (20 µg/kg) significantly reduced mean basal GH concentrations, although levels remained higher than those in normal weight controls (Gianotti *et al.* 2000).

Other hormonal factors

Triiodothyronine Triiodothyronine (T₃) may play a role in the GH resistance state induced by fasting (Ikeda *et al.* 1990). Both plasma IGF1 and T₃ levels are low in fasted rats (Ikeda *et al.* 1990). S.c. injections of T₃ (5 µg/kg) normalized T₃ levels in fasted rats and plasma IGF1 levels also increased significantly, suggesting that T₃ may play a role in mediating GH resistance in states of undernutrition (Ikeda *et al.* 1990).

Leptin Levels of leptin, an anorexigenic hormone secreted by adipocytes, are low in states of undernutrition (Frederich *et al.* 1995), and these low levels may mediate the GH resistance state. In sheep, an animal model that also demonstrates GH resistance in response to fasting, ovariectomized ewes that were fasted for 72 h and infused with leptin centrally had levels of GH, which were lower than those of the vehicle-treated, fasted, and ovariectomized ewes (Henry *et al.* 2004). GH levels in the leptin-treated ewes were comparable to levels in *ad libitum*-fed ewes (Henry *et al.* 2004). Therefore, central leptin may reduce GH secretion and the low levels of leptin during starvation may result in an increase in GH levels, contributing to the GH resistance state.

In humans, children with protein–energy malnutrition have significantly lower levels of leptin and IGF1 compared with healthy controls and significantly higher levels of GH (Soliman *et al.* 2000, Kilic *et al.* 2004). Leptin levels are also positively associated with IGF1 and inversely associated with GH (Soliman *et al.* 2000, Kilic *et al.* 2004). When women with hypothalamic amenorrhea, a state of negative energy balance, were treated with recombinant human methionyl leptin for 3 months – resulting in a normalization of leptin levels – IGF1 levels increased to levels comparable to euleptinemic controls, even after controlling for estradiol (E₂) levels and changes in weight (Chan *et al.* 2008). Therefore, low leptin levels may be a mediator of GH resistance in states of nutritional deprivation.

Testosterone Testosterone has also been shown to be a significant predictor of IGF1 levels in women with anorexia nervosa (Brick *et al.* 2010). In a group of women

with anorexia nervosa and normal weight controls, free testosterone was found to be a significant and independent predictor of IGF1 levels and was able to account for 36% of the variability in IGF1 levels after controlling for other possible hormonal predictors including E₂ (Brick *et al.* 2010). Therefore, the low testosterone levels in anorexia nervosa (Miller *et al.* 2007) and other states of starvation (Klibanski *et al.* 1981) may contribute to the state of GH resistance.

GH resistance is an adaptive response to undernutrition

In states of undernutrition, GH resistance may be an adaptive response. Elevated GH levels may be necessary to maintain euglycemia, whereas decreased levels of IGF1 help conserve energy during periods of nutritional deprivation.

Mobilization of fat stores

Elevated GH levels may be important for mobilization of fat stores in states of malnutrition. Mice with adult-onset, isolated GH deficiency and their GH-replete littermates were calorie restricted for 11 days (Gahete *et al.* 2013); GH levels increased in the GH-replete mice but not in the GH-deficient mice in response to the calorie deprivation (Gahete *et al.* 2013). Importantly, the loss of fat mass and subsequent increase in free fatty acid levels were significantly greater in the GH-replete mice compared with the GH-deficient mice (Gahete *et al.* 2013), suggesting that mobilization of fat stores during starvation is a significant, and likely adaptive, consequence of the elevated GH levels. In humans, GH has also been shown to be a critical factor in increasing the rate of lipolysis during fasting (Sakharova *et al.* 2008).

Maintenance of euglycemia

Elevated GH levels may also be important for maintaining euglycemia in states of starvation. Ghrelin is an orexigenic hormone produced primarily in the fundus of the stomach. Importantly, there are ghrelin receptors – called GH secretagogue receptors – in the anterior pituitary gland and octanoylated ghrelin acts as a potent stimulator of GH secretion (Kojima & Kangawa 2005). In an animal model, elimination of ghrelin O-acyltransferase (GOAT) – which attaches octanoate to proghrelin – results in mice that lack octanoylated ghrelin or acylated ghrelin (Zhao *et al.* 2010). *Goat* knockout mice on normal or high-fat diets grow and

gain weight normally but when calorie restricted, the *Goat* knockout mice become profoundly hypoglycemic in contrast to the WT, calorie-restricted mice (Zhao *et al.* 2010). GH levels were also found to be twofold higher in the WT mice compared with the *Goat* knockout mice and infusion of GH was able to rescue the *Goat* knockout mice from hypoglycemia and normalize blood sugars, suggesting that an important role of elevated GH levels in states of malnutrition is to maintain euglycemia (Zhao *et al.* 2010). Although there is recent controversy regarding the role of ghrelin and GH in protecting against hypoglycemia during periods of calorie restriction (Yi *et al.* 2012, Gahete *et al.* 2013), it is clear that the increasing levels of GH serve to mobilize energy during periods of malnutrition and are an adaptive response to undernutrition.

Decreased energy expenditure

Elevated GH levels are potentially an important means of mobilizing fat stores and maintaining euglycemia in states of undernutrition. Yet, if the GH-IGF1 axis remained intact during states of nutritional deprivation, the elevated levels of GH would result in elevated IGF1 levels, leading to increased energy expenditure on growth and therefore incur a survival disadvantage in states of malnutrition. Therefore, GH resistance, with an inability of GH to appropriately stimulate IGF1 production, is likely an adaptive mechanism to preserve energy during periods of undernutrition.

Conclusions

Our evolutionary past was marked by periods of undernutrition. These periods of undernutrition were characterized by isolated protein-calorie deficiency, isolated vitamin deficiencies, and even starvation and survival depended on the body's ability to mobilize energy stores and prevent hypoglycemia. GH plays a key role in mobilizing energy and therefore elevated GH levels confer a survival advantage during periods of undernutrition. Yet, increasing energy expenditure for growth during these periods – which would result if the GH-IGF1 axis remained intact and IGF1 levels increased – would be disadvantageous in states of undernutrition. Therefore, the elevated GH levels and low IGF1 levels characteristic of GH resistance are an important adaptive response to calorie and nutrient deprivation and an important mechanism for survival.

Declaration of interest

A K receives support for an investigator-initiated study from Rhythm Pharmaceuticals.

Funding

NIH grants K23 DK094820 (P K F) and R01 DK052625 (A K).

References

- American Psychiatric Association 1994 Diagnostic and statistical manual of mental disorders (DSM-IV), 4th edn. Washington, DC.
- Baxter RC & Turtle JR 1978 Regulation of hepatic growth hormone receptors by insulin. *Biochemical and Biophysical Research Communications* **84** 350–357. (doi:10.1016/0006-291X(78)90177-8)
- Baxter RC, Bryson JM & Turtle JR 1981 The effect of fasting on liver receptors for prolactin and growth hormone. *Metabolism* **30** 1086–1090. (doi:10.1016/0026-0495(81)90052-4)
- Brick DJ, Gerweck AV, Meenaghan E, Lawson EA, Misra M, Fazeli P, Johnson W, Klibanski A & Miller KK 2010 Determinants of IGF1 and GH across the weight spectrum: from anorexia nervosa to obesity. *European Journal of Endocrinology* **163** 185–191. (doi:10.1530/EJE-10-0365)
- Chan JL, Williams CJ, Raciti P, Blakeman J, Kelesidis T, Kelesidis I, Johnson ML, Thorner MO & Mantzoros CS 2008 Leptin does not mediate short-term fasting-induced changes in growth hormone pulsatility but increases IGF-I in leptin deficiency states. *Journal of Clinical Endocrinology and Metabolism* **93** 2819–2827. (doi:10.1210/jc.2008-0056)
- Counts DR, Gwirtsman H, Carlsson LM, Lesem M & Cutler GB Jr 1992 The effect of anorexia nervosa and refeeding on growth hormone-binding protein, the insulin-like growth factors (IGFs), and the IGF-binding proteins. *Journal of Clinical Endocrinology and Metabolism* **75** 762–767. (doi:10.1210/jc.75.3.762)
- Devine A, Rosen C, Mohan S, Baylink D & Prince RL 1998 Effects of zinc and other nutritional factors on insulin-like growth factor I and insulin-like growth factor binding proteins in postmenopausal women. *American Journal of Clinical Nutrition* **68** 200–206.
- Dorup I, Flyvbjerg A, Everts ME & Clausen T 1991 Role of insulin-like growth factor-1 and growth hormone in growth inhibition induced by magnesium and zinc deficiencies. *British Journal of Nutrition* **66** 505–521. (doi:10.1079/BJN19910051)
- Fazeli PK, Lawson EA, Prabhakaran R, Miller KK, Donoho DA, Clemmons DR, Herzog DB, Misra M & Klibanski A 2010a Effects of recombinant human growth hormone in anorexia nervosa: a randomized, placebo-controlled study. *Journal of Clinical Endocrinology and Metabolism* **95** 4889–4897. (doi:10.1210/jc.2010-0493)
- Fazeli PK, Misra M, Goldstein M, Miller KK & Klibanski A 2010b Fibroblast growth factor-21 may mediate growth hormone resistance in anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **95** 369–374. (doi:10.1210/jc.2009-1730)
- Fichter MM, Pirke KM & Holsboer F 1986 Weight loss causes neuroendocrine disturbances: experimental study in healthy starving subjects. *Psychiatry Research* **17** 61–72. (doi:10.1016/0165-1781(86)90042-9)
- Flyvbjerg A, Dorup I, Everts ME & Orskov H 1991 Evidence that potassium deficiency induces growth retardation through reduced circulating levels of growth hormone and insulin-like growth factor I. *Metabolism* **40** 769–775. (doi:10.1016/0026-0495(91)90001-D)
- Frederich RC, Lollmann B, Hamann A, Napolitano-Rosen A, Kahn BB, Lowell BB & Flier JS 1995 Expression of *ob* mRNA and its encoded protein in rodents. Impact of nutrition and obesity. *Journal of Clinical Investigation* **96** 1658–1663. (doi:10.1172/JCI118206)
- Frystyk J, Brick DJ, Gerweck AV, Utz AL & Miller KK 2009 Bioactive insulin-like growth factor-I in obesity. *Journal of Clinical Endocrinology and Metabolism* **94** 3093–3097. (doi:10.1210/jc.2009-0614)
- Gahete MD, Cordoba-Chacon J, Luque RM & Kineman RD 2013 The rise in growth hormone during starvation does not serve to maintain glucose levels or lean mass but is required for appropriate adipose tissue response in female mice. *Endocrinology* **154** 263–269. (doi:10.1210/en.2012-1849)
- Gaich G, Chien JY, Fu H, Glass LC, Deeg MA, Holland WL, Kharitonov A, Bumol T, Schilske HK & Moller DE 2013 The effects of LY2405319, an FGF21 analog, in obese human subjects with type 2 diabetes. *Cell Metabolism* **18** 333–340. (doi:10.1016/j.cmet.2013.08.005)
- Galman C, Lundasen T, Kharitonov A, Bina HA, Eriksson M, Hafstrom I, Dahlin M, Amark P, Angelin B & Rudling M 2008 The circulating metabolic regulator FGF21 is induced by prolonged fasting and PPAR α activation in man. *Cell Metabolism* **8** 169–174. (doi:10.1016/j.cmet.2008.06.014)
- Gannon MC & Nuttall FQ 2011 Effect of a high-protein diet on ghrelin, growth hormone, and insulin-like growth factor-I and binding proteins 1 and 3 in subjects with type 2 diabetes mellitus. *Metabolism* **60** 1300–1311. (doi:10.1016/j.metabol.2011.01.016)
- Garfinkel PE, Brown GM, Stancer HC & Moldofsky H 1975 Hypothalamic-pituitary function in anorexia nervosa. *Archives of General Psychiatry* **32** 739–744. (doi:10.1001/archpsyc.1975.01760240067005)
- Gianotti L, Pincelli AI, Scacchi M, Rolla M, Bellitti D, Arvat E, Lanfranco F, Torsello A, Ghigo E, Cavagnini F *et al.* 2000 Effects of recombinant human insulin-like growth factor I administration on spontaneous and growth hormone (GH)-releasing hormone-stimulated GH secretion in anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **85** 2805–2809. (doi:10.1210/jc.85.8.2805)
- Gillum MP, Erion DM & Shulman GI 2010 Sirtuin-1 regulation of mammalian metabolism. *Trends in Molecular Medicine* **17** 8–13. (doi:10.1016/j.molmed.2010.09.005)
- Grinspoon SK, Baum HB, Peterson S & Klibanski A 1995 Effects of rhIGF-I administration on bone turnover during short-term fasting. *Journal of Clinical Investigation* **96** 900–906. (doi:10.1172/JCI118137)
- Grottoli S, Gasco V, Mainolfi A, Beccuti G, Corneli G, Aimaretti G, Dieguez C, Casanueva F & Ghigo E 2008 Growth hormone/insulin-like growth factor I axis, glucose metabolism, and lipolysis but not leptin show some degree of refractoriness to short-term fasting in acromegaly. *Journal of Endocrinological Investigation* **31** 1103–1109.
- Hartman ML, Clayton PE, Johnson ML, Celniker A, Perlman AJ, Alberti KG & Thorner MO 1993 A low dose euglycemic infusion of recombinant human insulin-like growth factor I rapidly suppresses fasting-enhanced pulsatile growth hormone secretion in humans. *Journal of Clinical Investigation* **91** 2453–2462. (doi:10.1172/JCI116480)
- Henry BA, Goding JW, Tilbrook AJ, Dunshea FR, Blache D & Clarke IJ 2004 Leptin-mediated effects of undernutrition or fasting on luteinizing hormone and growth hormone secretion in ovariectomized ewes depend on the duration of metabolic perturbation. *Journal of Neuroendocrinology* **16** 244–255. (doi:10.1111/j.0953-8194.2004.01157.x)
- Hintz RL, Suskind R, Amatayakul K, Thanangkul O & Olson R 1978 Plasma somatomedin and growth hormone values in children with protein-calorie malnutrition. *Journal of Pediatrics* **92** 153–156. (doi:10.1016/S0022-3476(78)80099-7)
- Ho KY, Veldhuis JD, Johnson ML, Furlanetto R, Evans WS, Alberti KG & Thorner MO 1988 Fasting enhances growth hormone secretion and amplifies the complex rhythms of growth hormone secretion in man. *Journal of Clinical Investigation* **81** 968–975. (doi:10.1172/JCI113450)
- Holmes MD, Pollak MN, Willett WC & Hankinson SE 2002 Dietary correlates of plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 concentrations. *Cancer Epidemiology, Biomarkers & Prevention* **11** 852–861.
- Horner JM, Kemp SF & Hintz RL 1981 Growth hormone and somatomedin in insulin-dependent diabetes mellitus. *Journal of Clinical Endocrinology and Metabolism* **53** 1148–1153. (doi:10.1210/jcem-53-6-1148)

- Hotta M, Fukuda I, Sato K, Hizuka N, Shibasaki T & Takano K 2000 The relationship between bone turnover and body weight, serum insulin-like growth factor (IGF) I, and serum IGF-binding protein levels in patients with anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **85** 200–206. (doi:10.1210/jc.85.1.200)
- Ikedo T, Fujiyama K, Hoshino T, Takeuchi T, Mashiba H & Tominaga M 1990 Possible role of thyroid hormone in decreased somatomedin-C levels in diabetic and starved rats. *Annals of Nutrition & Metabolism* **34** 8–12. (doi:10.1159/000177564)
- Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, Parameswara V, Li Y, Goetz R, Mohammadi M, Esser V *et al.* 2007 Endocrine regulation of the fasting response by PPAR α -mediated induction of fibroblast growth factor 21. *Cell Metabolism* **5** 415–425. (doi:10.1016/j.cmet.2007.05.003)
- Inagaki T, Lin VY, Goetz R, Mohammadi M, Mangelsdorf DJ & Kliewer SA 2008 Inhibition of growth hormone signaling by the fasting-induced hormone FGF21. *Cell Metabolism* **8** 77–83. (doi:10.1016/j.cmet.2008.05.006)
- Isley WL, Underwood LE & Clemmons DR 1983 Dietary components that regulate serum somatomedin-C concentrations in humans. *Journal of Clinical Investigation* **71** 175–182. (doi:10.1172/JCI110757)
- Kaklamani VG, Linos A, Kaklamani E, Markaki I, Koumantaki Y & Mantzoros CS 1999 Dietary fat and carbohydrates are independently associated with circulating insulin-like growth factor 1 and insulin-like growth factor-binding protein 3 concentrations in healthy adults. *Journal of Clinical Oncology* **17** 3291–3298.
- Keski-Rahkonen A, Hoek HW, Susser ES, Linna MS, Sihvola E, Raevuori A, Bulik CM, Kaprio J & Rissanen A 2007 Epidemiology and course of anorexia nervosa in the community. *American Journal of Psychiatry* **164** 1259–1265. (doi:10.1176/appi.ajp.2007.06081388)
- Kharitonov A, Shiyanova TL, Koester A, Ford AM, Micanovic R, Galbreath EJ, Sandusky GE, Hammond LJ, Moyers JS, Owens RA *et al.* 2005 FGF-21 as a novel metabolic regulator. *Journal of Clinical Investigation* **115** 1627–1635. (doi:10.1172/JCI23606)
- Kilic M, Taskin E, Ustundag B & Aygun AD 2004 The evaluation of serum leptin level and other hormonal parameters in children with severe malnutrition. *Clinical Biochemistry* **37** 382–387. (doi:10.1016/j.clinbiochem.2003.12.010)
- Klibanski A, Beitins IZ, Badger T, Little R & McArthur JW 1981 Reproductive function during fasting in men. *Journal of Clinical Endocrinology and Metabolism* **53** 258–263. (doi:10.1210/jcem-53-2-258)
- Kojima M & Kangawa K 2005 Ghrelin: structure and function. *Physiological Reviews* **85** 495–522. (doi:10.1152/physrev.00012.2004)
- Leung KC, Doyle N, Ballesteros M, Waters MJ & Ho KK 2000 Insulin regulation of human hepatic growth hormone receptors: divergent effects on biosynthesis and surface translocation. *Journal of Clinical Endocrinology and Metabolism* **85** 4712–4720. (doi:10.1210/jc.85.12.4712)
- Lundasen T, Hunt MC, Nilsson LM, Sanyal S, Angelin B, Alexson SE & Rudling MP 2007 PPAR α is a key regulator of hepatic FGF21. *Biochemical and Biophysical Research Communications* **360** 437–440. (doi:10.1016/j.bbrc.2007.06.068)
- Lundbaek K, Christensen NJ, Jensen VA, Johansen K, Olsen TS, Hansen AP, Orskov H & Osterby R 1970 Diabetes, diabetic angiopathy, and growth hormone. *Lancet* **2** 131–133. (doi:10.1016/S0140-6736(70)92706-6)
- Maccario M, Valetto MR, Savio P, Aimaretti G, Baffoni C, Procopio M, Grottoli S, Olandri SE, Arvat E & Ghigo E 1997 Maximal secretory capacity of somatotrope cells in obesity: comparison with GH deficiency. *International Journal of Obesity and Related Metabolic Disorders* **21** 27–32. (doi:10.1038/sj.ijo.0800356)
- Maes M, Amand Y, Underwood LE, Maiter D & Ketelslegers JM 1988 Decreased serum insulin-like growth factor I response to growth hormone in hypophysectomized rats fed a low protein diet: evidence for a postreceptor defect. *Acta Endocrinologica* **117** 320–326.
- Maiter D, Fliesen T, Underwood LE, Maes M, Gerard G, Davenport ML & Ketelslegers JM 1989 Dietary protein restriction decreases insulin-like growth factor I independent of insulin and liver growth hormone binding. *Endocrinology* **124** 2604–2611. (doi:10.1210/endo-124-5-2604)
- Massara F, Ghigo E, Goffi S, Molinatti GM, Muller EE & Camanni F 1984 Blockade of hp-GRF-40-induced GH release in normal men by a cholinergic muscarinic antagonist. *Journal of Clinical Endocrinology and Metabolism* **59** 1025–1026. (doi:10.1210/jcem-59-5-1025)
- Merimee TJ, Zapf J & Froesch ER 1982 Insulin-like growth factors in the fed and fasted states. *Journal of Clinical Endocrinology and Metabolism* **55** 999–1002. (doi:10.1210/jcem-55-5-999)
- Miller KK, Lawson EA, Mathur V, Wexler TL, Meenaghan E, Misra M, Herzog DB & Klibanski A 2007 Androgens in women with anorexia nervosa and normal-weight women with hypothalamic amenorrhea. *Journal of Clinical Endocrinology and Metabolism* **92** 1334–1339. (doi:10.1210/jc.2006-2501)
- Misra M, Miller KK, Bjornson J, Hackman A, Aggarwal A, Chung J, Ott M, Herzog DB, Johnson ML & Klibanski A 2003 Alterations in growth hormone secretory dynamics in adolescent girls with anorexia nervosa and effects on bone metabolism. *Journal of Clinical Endocrinology and Metabolism* **88** 5615–5623. (doi:10.1210/jc.2003-030532)
- Mohan PS & Jaya Rao KS 1979 Plasma somatomedin activity in vitamin A deficient children. *Clinica Chimica Acta* **96** 241–246. (doi:10.1016/0009-8981(79)90434-0)
- Mohan PS & Jaya Rao KS 1980 Sulfate metabolism in vitamin A-deficient rats. *Journal of Nutrition* **110** 868–875.
- Mraz M, Bartlova M, Lacinova Z, Michalsky D, Kasalicky M, Haluzikova D, Matoulek M, Dostalova I, Humenanska V & Haluzik M 2009 Serum concentrations and tissue expression of a novel endocrine regulator fibroblast growth factor-21 in patients with type 2 diabetes and obesity. *Clinical Endocrinology* **71** 369–375. (doi:10.1111/j.1365-2265.2008.03502.x)
- Nishimura T, Nakatake Y, Konishi M & Itoh N 2000 Identification of a novel FGF, FGF-21, preferentially expressed in the liver. *Biochimica et Biophysica Acta* **1492** 203–206. (doi:10.1016/S0167-4781(00)00067-1)
- Olusi SO, Orrell DH, Morris PM & McFarlane H 1977 A study of endocrine function in protein-energy malnutrition. *Clinica Chimica Acta* **74** 261–269. (doi:10.1016/0009-8981(77)90293-5)
- Potthoff MJ, Inagaki T, Satapati S, Ding X, He T, Goetz R, Mohammadi M, Finck BN, Mangelsdorf DJ, Kliewer SA *et al.* 2009 FGF21 induces PGC-1 α and regulates carbohydrate and fatty acid metabolism during the adaptive starvation response. *PNAS* **106** 10853–10858. (doi:10.1073/pnas.0904187106)
- Rao KS & Mohan PS 1982 Plasma somatomedin activity, growth-hormone and insulin levels in vitamin B6 deficient rats. *Hormone and Metabolic Research* **14** 580–582. (doi:10.1055/s-2007-1019086)
- Reeves RD, Dickinson L, Lee J, Kilgore B, Branham B & Elders MJ 1979 Effects of dietary composition on somatomedin activity in growing rats. *Journal of Nutrition* **109** 613–620.
- Robinson H & Picou D 1977 A comparison of fasting plasma insulin and growth hormone concentrations in marasmic, kwashiorkor, marasmic-kwashiorkor and underweight children. *Pediatric Research* **11** 637–640. (doi:10.1203/00006450-197705000-00003)
- Rolla M, Andreoni A, Belliti D, Ferdeghini M, Cristofani R & Muller EE 1990 Effects of cholinergic muscarinic antagonist pirenzepine on GH response to GHRH 1–40 in patients with anorexia nervosa. *Endocrinologia Experimentalis* **24** 195–204.
- Sakharova AA, Horowitz JF, Surya S, Goldenberg N, Harber MP, Symons K & Barkan A 2008 Role of growth hormone in regulating lipolysis, proteolysis, and hepatic glucose production during fasting. *Journal of Clinical Endocrinology and Metabolism* **93** 2755–2759. (doi:10.1210/jc.2008-0079)
- Scacchi M, Pincelli AI, Caumo A, Tomasi P, Delitala G, Baldi G & Cavagnini F 1997 Spontaneous nocturnal growth hormone secretion in anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **82** 3225–3229. (doi:10.1210/jc.82.10.3225)
- Schurck MA, Rizzoli R, Slosman D, Vadas L, Vergnaud P & Bonjour JP 1998 Protein supplements increase serum insulin-like growth factor-I levels and attenuate proximal femur bone loss in patients with recent hip fracture. A randomized, double-blind, placebo-controlled trial. *Annals of Internal Medicine* **128** 801–809. (doi:10.7326/0003-4819-128-10-199805150-00002)

- Shimizu H, Miyazaki M, Shimomura Y & Kobayashi I 1991 Altered hormonal status in a female deprived of food for 18 days. *Journal of Medicine* **22** 201–210.
- Soliman AT, Hassan AE, Aref MK, Hintz RL, Rosenfeld RG & Rogol AD 1986 Serum insulin-like growth factors I and II concentrations and growth hormone and insulin responses to arginine infusion in children with protein–energy malnutrition before and after nutritional rehabilitation. *Pediatric Research* **20** 1122–1130. (doi:10.1203/00006450-198611000-00012)
- Soliman AT, ElZalabany MM, Salama M & Ansari BM 2000 Serum leptin concentrations during severe protein–energy malnutrition: correlation with growth parameters and endocrine function. *Metabolism* **49** 819–825. (doi:10.1053/meta.2000.6745)
- Stoving RK, Veldhuis JD, Flyvbjerg A, Vinten J, Hangaard J, Koldkjaer OG, Kristiansen J & Hagen C 1999 Jointly amplified basal and pulsatile growth hormone (GH) secretion and increased process irregularity in women with anorexia nervosa: indirect evidence for disruption of feedback regulation within the GH–insulin-like growth factor I axis. *Journal of Clinical Endocrinology and Metabolism* **84** 2056–2063. (doi:10.1210/jc.84.6.2056)
- Stoving RK, Chen JW, Glinborg D, Brixen K, Flyvbjerg A, Horder K & Frydlyk J 2007 Bioactive insulin-like growth factor (IGF) I and IGF-binding protein-1 in anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **92** 2323–2329. (doi:10.1210/jc.2006-1926)
- Straus DS & Takemoto CD 1990 Effect of fasting on insulin-like growth factor-I (IGF-I) and growth hormone receptor mRNA levels and IGF-I gene transcription in rat liver. *Molecular Endocrinology* **4** 91–100. (doi:10.1210/mend-4-1-91)
- Tamai H, Komaki G, Matsubayashi S, Kobayashi N, Mori K, Nakagawa T, Truong MP, Walter RM Jr & Kumagai LF 1990 Effect of cholinergic muscarinic receptor blockade on human growth hormone (GH)-releasing hormone-(1–44)-induced GH secretion in anorexia nervosa. *Journal of Clinical Endocrinology and Metabolism* **70** 738–741. (doi:10.1210/jcem-70-3-738)
- Tamai H, Kiyohara K, Mukuta T, Kobayashi N, Komaki G, Nakagawa T, Kumagai LF & Aoki TT 1991 Responses of growth hormone and cortisol to intravenous glucose loading test in patients with anorexia nervosa. *Metabolism* **40** 31–34. (doi:10.1016/0026-0495(91)90188-3)
- Thimmarayappa J, Sun J, Schultz LE, Dejjamron P, Lu C, Giallongo A, Merchant JL & Menon RK 2006 Inhibition of growth hormone receptor gene expression by saturated fatty acids: role of Kruppel-like zinc finger factor, ZBP-89. *Molecular Endocrinology* **20** 2747–2760. (doi:10.1210/me.2006-0128)
- Thissen JP, Triest S, Underwood LE, Maes M & Ketelslegers JM 1990 Divergent responses of serum insulin-like growth factor-I and liver growth hormone (GH) receptors to exogenous GH in protein-restricted rats. *Endocrinology* **126** 908–913. (doi:10.1210/endo-126-2-908)
- Thissen JP, Triest S, Moats-Staats BM, Underwood LE, Mauerhoff T, Maiter D & Ketelslegers JM 1991a Evidence that pretranslational and translational defects decrease serum insulin-like growth factor-I concentrations during dietary protein restriction. *Endocrinology* **129** 429–435. (doi:10.1210/endo-129-1-429)
- Thissen JP, Underwood LE, Maiter D, Maes M, Clemmons DR & Ketelslegers JM 1991b Failure of insulin-like growth factor-I (IGF-I) infusion to promote growth in protein-restricted rats despite normalization of serum IGF-I concentrations. *Endocrinology* **128** 885–890. (doi:10.1210/endo-128-2-885)
- Yamamoto M, Iguchi G, Fukuoka H, Suda K, Bando H, Takahashi M, Nishizawa H, Seino S & Takahashi Y 2013 SIRT1 regulates adaptive response of the growth hormone – insulin-like growth factor-I axis under fasting conditions in liver. *PNAS* **110** 14948–14953. (doi:10.1073/pnas.1220606110)
- Yde H 1969 The growth hormone dependent sulfation factor in serum from patients with various types of diabetes. *Acta Medica Scandinavica* **186** 293–297. (doi:10.1111/j.0954-6820.1969.tb01479.x)
- Yi CX, Heppner KM, Kirchner H, Tong J, Bielohuby M, Gaylinn BD, Muller TD, Bartley E, Davis HW, Zhao Y *et al.* 2012 The GOAT–ghrelin system is not essential for hypoglycemia prevention during prolonged calorie restriction. *PLoS ONE* **7** e32100. (doi:10.1371/journal.pone.0032100)
- Zhang X, Yeung DC, Karpisek M, Stejskal D, Zhou ZG, Liu F, Wong RL, Chow WS, Tso AW, Lam KS *et al.* 2008 Serum FGF21 levels are increased in obesity and are independently associated with the metabolic syndrome in humans. *Diabetes* **57** 1246–1253. (doi:10.2337/db07-1476)
- Zhao TJ, Liang G, Li RL, Xie X, Sleeman MW, Murphy AJ, Valenzuela DM, Yancopoulos GD, Goldstein JL & Brown MS 2010 Ghrelin O-acyltransferase (GOAT) is essential for growth hormone-mediated survival of calorie-restricted mice. *PNAS* **107** 7467–7472. (doi:10.1073/pnas.1002271107)

Received in final form 10 December 2013

Accepted 19 December 2013

Accepted Preprint published online 20 December 2013