Protocatechuic acid exerts a cardioprotective effect in type 1 diabetic rats

Yoswaris Semaming^{1,2}, Sirinart Kumfu^{1,3}, Patchareewan Pannangpetch^{1,2}, Siriporn C Chattipakorn^{1,4} and Nipon Chattipakorn^{1,3}

¹Cardiac Electrophysiology Research and Training Center, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand

²Department of Pharmacology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002, Thailand

³Cardiac Electrophysiology Unit, Department of Physiology, Faculty of Medicine and ⁴Department of Oral Biology and Diagnostic Sciences, Faculty of Dentistry, Chiang Mai University, Chiang Mai 50200, Thailand

Correspondence
should be addressed
to N Chattipakorn
Email
nchattip@gmail.com

Abstract

Oxidative stress has been shown to play an important role in the pathogenesis of diabetes-induced cardiac dysfunction. Protocatechuic acid (PCA) is a phenolic compound, a main metabolite of anthocyanin, which has been reported to display various pharmacological properties. We proposed the hypothesis that PCA exerts cardioprotection in type 1 diabetic (T1DM) rats. T1DM was induced in male Sprague–Dawley rats by a single i.p. injection of 50 mg/kg streptozotocin (STZ) and groups of these animals received the following treatments for 12 weeks: i) oral administration of vehicle, ii) oral administration of PCA at a dose of 50 mg/kg per day, iii) oral administration of PCA at a dose of 100 mg/kg per day, iv) s.c. injection of insulin at a dose of 4 U/kg per day, and v) a combination of PCA, 100 mg/kg per day and insulin, 4 U/kg per day. Metabolic parameters, results from echocardiography, and heart rate variability were monitored every 4 weeks, and the HbA1c, cardiac malondialdehyde (MDA), cardiac mitochondrial function, and cardiac BAX/BCL2 expression were evaluated at the end of treatment. PCA, insulin, and combined drug treatments significantly improved metabolic parameters and cardiac function as shown by increased percentage fractional shortening and percentage left ventricular ejection fraction and decreased low-frequency:high-frequency ratio in T1DM rats. Moreover, all treatments significantly decreased plasma HbA1c and cardiac MDA levels, improved cardiac mitochondrial function, and increased BCL2 expression. Our results demonstrated for the first time, to our knowledge, the efficacy of PCA in improving cardiac function and cardiac autonomic balance, preventing cardiac mitochondrial dysfunction, and increasing anti-apoptotic protein in STZ-induced T1DM rats. Thus, PCA possesses a potential cardioprotective effect and could restore cardiac function when combined with insulin treatment. These findings indicated that supplementation with PCA might be helpful for the prevention and alleviation of cardiovascular complications in T1DM.

Key Words

- ▶ type 1 diabetes
- ▶ oxidative stress
- ▶ cardiac mitochondrial dysfunction
- ▶ protocatechuic acid

Journal of Endocrinology
(2014) 223, 13–23

Introduction

Diabetes and cardiac complications have become a public health problem of considerable magnitude. Cardiac dysfunction is the most common serious complication of diabetes mellitus and has become one of the leading causes of the increased mortality in both Type 1 and Type 2 diabetes. A streptozotocin (STZ)-induced type 1 diabetic rat has been used as an animal model for the investigation of pharmacological interventions and also for the study of diabetic complications including diabetes-induced cardiac dysfunction (Krishna *et al.* 2005, Naowaboot *et al.* 2009). Clinically, cardiac dysfunction may occur without major vascular lesions, indicating a primary role for the direct effects of diabetes on cardiomyocytes (Trost & LeWinter 2001, Dyntar *et al.* 2006). These characteristics further result in diabetes-induced cardiac dysfunction through cellular pathological changes, especially increased oxidative stress, mitochondrial dysfunction, and cardiac autonomic neuropathy (An & Rodrigues 2006, Boudina & Abel 2007). Oxidative damage plays an important role in the occurrence and development of cardiac dysfunction in this case (Asbun & Villarreal 2006, Kumar & Sitasawad 2009). Oxidative stress is known to potentially lead to biological damage to macromolecules and activates maladaptive signaling pathways, which may cause cell dysfunction and cell death (Sies 1991).

Protocatechuic acid (PCA, 3,4-dihydroxybenzoic acid), a phenolic compound, is isolated from the dried flowers of *Hibiscus sabdariffa* Linn. It is a main metabolite of complex polyphenols, especially anthocyanins, which are converted to PCA and also abundantly formed and absorbed in the large intestine because of microbial metabolism (Kay *et al.* 2005, Vitaglione *et al.* 2007). Previous reports of both *in vitro* and *in vivo* studies have demonstrated that PCA has anti-carcinogen (Tseng *et al.* 2000, Yip *et al.* 2006, Lin *et al.* 2007, Yin *et al.* 2009), anti-hyperglycemia (Lin *et al.* 2009, 2011, Harini & Pugalendi 2010, Scazzocchio *et al.* 2011), antioxidant (Liu *et al.* 2002, Sroka & Cisowski 2003, Masella *et al.* 2004, Shi *et al.* 2006, Tarozzi *et al.* 2007, Lin *et al.* 2009, Chou *et al.* 2010, Vari *et al.* 2011, Zhang *et al.* 2011), and anti-inflammatory properties (Yan *et al.* 2004, Min *et al.* 2010, Wang *et al.* 2010, 2011). However, no scientific investigation has yet been conducted on the effect of PCA on diabetes-induced cardiac dysfunction, which is also caused by oxidative stress, and thus this investigation was conducted to study the anti-hyperglycemic activity of PCA and the effect of PCA on diabetes-induced cardiac dysfunction and other related metabolic parameters in STZ-induced diabetic rats.

We tested the hypothesis that PCA could attenuate cardiac complications by reducing oxidative stress, ameliorating cardiomyocyte apoptosis, and improving the cardiac autonomic balance and cardiac mitochondrial function in STZ-induced diabetic rats.

Materials and methods

Animals

A total of 36 male Sprague–Dawley rats with body weights of 250–280 g were acquired from the National Laboratory Animal Center, Mahidol University, Salaya, Nakornpathom. All experiments complied with the standards for the care and use of experimental animals and were approved by the Faculty of Medicine, Chiang Mai University Institutional Animal Care and Use Committee. After 7 days of acclimatization, diabetes was induced in 30 of the rats by a single i.p. injection of 50 mg/kg body weight of STZ dissolved in 0.1 M citric buffer (pH 4.5). Normal control rats ($n=6$) received injections of citric buffer (vehicle). Ten days after vehicle/STZ injection, only diabetic rats with fasting blood glucose (FBG) over 300 mg/dl were included in the experiments (Jiang *et al.* 2011, Zhao *et al.* 2014). Rats were assigned to one of six groups with six rats in each group and were treated as follows: normal rats treated with vehicle (NMV), diabetic rats treated with vehicle (DMV), diabetic rats treated with insulin at a dose of 4 U/kg (DMI), diabetic rats treated with PCA at a dose of 50 mg/kg (DML), diabetic rats treated with PCA at a dose of 100 mg/kg (DMH) (Harini & Pugalendi 2010), and diabetic rats treated with DMH and insulin at a dose of 4 U/kg (DMHI). All treatments were administered daily for 12 weeks. Fifty percent propylene glycol (vehicle) and PCA (Sigma Chemical Co. Ltd) were administered orally using gavage feeding, and insulin was injected subcutaneously. The animals were given free access to feed and drinking water. The body weight and food intake were recorded weekly. Echocardiography was performed and heart rate variability (HRV) was measured every 4 weeks and blood sampling from tail vein was performed after fasting for 12 h to measure plasma glucose, insulin, and malondialdehyde (MDA) levels. At the end of a 12-week treatment, animals were deeply anesthetized and then killed; the hearts were collected for investigation of mitochondrial function, MDA, and BAX and BCL2 protein levels, and blood was collected for investigation of HbA1c.

FBG, plasma insulin, and HbA1c level determination

FBG level was determined using a glucometer (Accu-Chek Advantage II; Roche Diagnostics). Plasma insulin levels were measured by Sandwich ELISA (LINCO Research, Saint Charles, MO, USA; Pratchayasakul *et al.* 2011). Plasma HbA1c was determined based on the competitive turbidimetric inhibition immunoassay (Roche Diagnostics Ltd).

Plasma and cardiac MDA level determination

Plasma and cardiac MDA level was determined using a HPLC-based assay (Thermo scientific, Bangkok, Thailand). Cardiac tissue was homogenized in phosphate buffer, pH 2.8. Plasma and cardiac tissue were mixed with H_3PO_4 and thiobarbituric acid (TBA) to produce TBA-reactive substances (TBARSs). The plasma and cardiac TBARS concentrations were determined from a standard curve and reported as being equivalent to the MDA concentration (Apaijai *et al.* 2013).

Echocardiography

The echocardiographic parameters were measured in each rat using an HP/Agilent Philips Sonos 4500 (Agilent Technologies, Santa Clara, CA, USA). An echocardiography probe was placed in gentle contact with the chest, and images were collected along the parasternal short axis of the heart (Lekawanvijit *et al.* 2012). M-mode echocardiography was performed at the level of the papillary muscles. Percentage fractional shortening (%FS) and percentage left ventricular ejection fraction (%LVEF) were determined.

HRV analysis

The output from electrocardiogram (ECG) lead II was recorded in each rat using the PowerLab (ADInstruments, Sydney, NSW, Australia) and the Chart 5.0 programs (Raheer *et al.* 2008). During ECG recording, rats were placed in a restraint and prohibited from movement (Incharoen *et al.* 2007, Raheer *et al.* 2008, Kumfu *et al.* 2012). The high-frequency (HF, 0.6–3.0 Hz) component, representing cardiac parasympathetic activity, and low-frequency (LF, 0.2–0.6 Hz) component, representing cardiac sympathetic and parasympathetic activity, were determined using the MATLAB program (Raheer *et al.* 2008). The LF:HF ratio was considered to be an indicator of cardiac sympathetic/parasympathetic tone balance (Ohuchi *et al.* 2000). An increased LF:HF ratio (i.e. depressed HRV) indicated a cardiac sympathovagal imbalance (Incharoen *et al.* 2007).

Cardiac mitochondrial function determination

Cardiac mitochondrial isolation was performed as described previously (Thummasorn *et al.* 2011). Cardiac mitochondrial function was assessed by determining production of cardiac mitochondrial reactive oxygen species (ROS), cardiac mitochondrial membrane potential change, and cardiac mitochondrial swelling (Thummasorn *et al.* 2011, Apaijai *et al.* 2012, Chinda *et al.* 2013). The components of the buffers used in the analysis of cardiac mitochondrial function consisted of KCl, sucrose, HEPES, KH_2PO_4 , and pyruvate/malate that have no ADP/ATP. The measurements of mitochondrial function in each group were made during State IV respiration.

Cardiac mitochondrial ROS production Cardiac mitochondria were incubated with 2- μM DCFH-DA dye at 25 °C for 20 min. The dye was excited at λ_{ex} 485 nm and detected at λ_{em} 530 nm using a fluorescent microplate reader (BioTek Instruments, Winooski, VT, USA). An increase in the fluorescence intensity indicated increased mitochondrial ROS production (Thummasorn *et al.* 2011, Apaijai *et al.* 2013, Chinda *et al.* 2013).

Cardiac mitochondrial membrane potential changes

Cardiac mitochondrial membrane potential changes were determined using 5- μM JC-1 dye as described previously (Thummasorn *et al.* 2011, Apaijai *et al.* 2013). In brief, cardiac mitochondria were incubated with JC-1 at 37 °C for 30 min. JC-1 monomer form (green-fluorescent) was excited at λ_{ex} 485 nm and detected at λ_{em} 590 nm, and JC-1 aggregate form (red-fluorescent) was excited at λ_{ex} 485 nm and detected at λ_{em} 530 nm using a fluorescent microplate reader. A decrease in the red:green fluorescence intensity ratio indicated cardiac mitochondrial membrane depolarization (Thummasorn *et al.* 2011, Apaijai *et al.* 2013, Chinda *et al.* 2013).

Cardiac mitochondrial swelling Cardiac mitochondria were incubated with 1.5-mM respiration buffer containing 100 mM KCl, 10 mM HEPES, and 5 mM KH_2PO_4 , and the absorbance was determined using a spectrophotometer as described previously (Thummasorn *et al.* 2011, Apaijai *et al.* 2013). Cardiac mitochondrial swelling was indicated by a decrease in the detected absorbance (Thummasorn *et al.* 2011, Apaijai *et al.* 2013, Chinda *et al.* 2013).

Cardiac mitochondrial morphology Cardiac mitochondria were fixed with 2.5% glutaraldehyde in 0.1-M

phosphate buffer overnight and post-fixed in 1% cacodylate buffer osmium tetroxide for 2 h, and then dehydrated with a graded ethanol series (Thummasorn *et al.* 2011). Cardiac mitochondria were embedded in Epon-Araldite, cut with a diamond knife, and stained with uranyl acetate and lead acetate (Thummasorn *et al.* 2011). Cardiac mitochondrial morphology was detected using a transmission electron microscope (TEM; Thummasorn *et al.* 2011).

Western blot analysis of BAX and BCL2 protein expression levels

Myocardial protein extracts were prepared by homogenization of nitrogen-frozen myocardial tissues in a 300-ml extraction buffer containing 20-mM Tris-HCl (pH 6.8), 1-mM sodium orthovanadate, 5-mM sodium fluoride, and a protease inhibitor. Total protein concentration was determined using a Bio-Rad protein assay kit (Bio-Rad Laboratories). Samples of 50–80 µg of total protein were mixed with a loading buffer (5% betamercaptoethanol, 0.05% bromophenol blue, 75 mM Tris-HCl (pH 6.8), 2% SDS, and 10% glycerol), and loaded on 10% SDS-acrylamide gels. Proteins were transferred onto PVDF membrane in a glycine/methanol-transfer buffer (Palee *et al.* 2013, Surinkaew *et al.* 2013) in a Wet/Tank blotting system (Bio-Rad Laboratories). Membranes were blocked in 5% skim milk in Tris-buffered saline and Tween (TBST) buffer. Western blot analysis for BAX and BCL2 was performed using myocardial tissues. Membranes were exposed to mouse monoclonal anti-rat Bax and Bcl2 (1:1000 dilution, Santa Cruz Biotechnology). Bound antibody was detected by HRP conjugated with an anti-rabbit IgG (1:2000 dilution, Cell

Signaling Technology, Danvers, MA, USA). The membranes were developed using the Amersham ECL Prime Western Blotting Detection Reagent (GE Healthcare, Buckinghamshire, UK) and densitometric analysis was carried out (Surinkaew *et al.* 2013).

Statistical analysis

Metabolic parameters, echocardiography, and HRV data are expressed as mean $\pm$ s.d., and others are expressed as mean $\pm$ s.e.m. One-way ANOVA followed by the least significant difference (LSD) *post hoc* test was used to test the differences among the groups. $P < 0.05$ was considered statistically significant.

Results

Effects of pharmacological interventions on body weight, food intake, FBG, and plasma insulin in diabetic rats

At 10 days after STZ injection, diabetic rats were characterized by body weight loss, polyphagia, hyperglycemia, and insulin deficiency (Tables 1 and 2). Beginning at week 4 of insulin and combined drug treatments in diabetic rats, there were significant increases in body weight and plasma insulin levels, and decreases in food intake and FBG levels, compared with DMV rats (Tables 3, 4 and 5). Beginning at week 4 for both doses of PCA treatment in diabetic rats, the FBG level significantly decreased when compared with DMV rats, whereas there was no alteration in body weight, food intake, and plasma insulin level in these rats (Tables 3, 4 and 5).

Table 1 Metabolic parameters, echocardiography, and heart rate variability (HRV) in normal and diabetic rats at baseline and 10 days after vehicle/STZ injection. Values are expressed as mean $\pm$ s.d.

Parameters	Baseline		10 days after vehicle/STZ injection	
	NM	DM	NM	DM
Metabolic parameters				
Body weight (g)	274 $\pm$ 9	274 $\pm$ 11	316 $\pm$ 9*	261 $\pm$ 17 [†]
Food intake (g/day)	20 $\pm$ 2	21 $\pm$ 2	22 $\pm$ 2	24 $\pm$ 2 [†]
Fasting blood glucose (mg/dl)	94 $\pm$ 3	95 $\pm$ 6	100 $\pm$ 5	400 $\pm$ 36 [†]
Plasma insulin (ng/ml)	1.49 $\pm$ 0.37	1.48 $\pm$ 0.38	1.39 $\pm$ 0.44	0.56 $\pm$ 0.25 [†]
Plasma MDA (µmol/ml)	1.05 $\pm$ 0.03	1.03 $\pm$ 0.03	1.04 $\pm$ 0.02	1.04 $\pm$ 0.02
Echocardiography				
Fractional shortening (%)	55 $\pm$ 3	54 $\pm$ 4	55 $\pm$ 2	53 $\pm$ 3
LV ejection fraction (%)	81 $\pm$ 3	82 $\pm$ 3	81 $\pm$ 2	75 $\pm$ 4
HRV				
LF:HF ratio	0.24 $\pm$ 0.04	0.24 $\pm$ 0.04	0.21 $\pm$ 0.04	0.23 $\pm$ 0.04

NM, normal rat; DM, diabetic rat; LV ejection fraction, left ventricular ejection; LF:HF ratio, low-frequency:high-frequency ratio. * $P < 0.05$ vs baseline and [†] $P < 0.05$ vs NM at 10 days after vehicle injection.

Table 2 Metabolic parameters, echocardiography, and heart rate variability (HRV) at 10 days after vehicle/STZ injection. Values are expressed as mean $\pm$ s.d. of five to six rats in each group

Parameters	NMV	DMV	DMI	DML	DMH	DMHI
Metabolic parameters						
Body weight (g)	316 $\pm$ 9	261 $\pm$ 14*	270 $\pm$ 20*	256 $\pm$ 12*	250 $\pm$ 16*	273 $\pm$ 16*
Food intake (g/day)	22 $\pm$ 2	24 $\pm$ 2*	24 $\pm$ 1*	24 $\pm$ 2*	25 $\pm$ 2*	25 $\pm$ 2*
Fasting blood glucose (mg/dl)	100 $\pm$ 5	394 $\pm$ 33*	400 $\pm$ 40*	392 $\pm$ 26*	414 $\pm$ 48*	407 $\pm$ 36*
Plasma insulin (ng/ml)	1.51 $\pm$ 0.60	0.67 $\pm$ 0.29*	0.55 $\pm$ 0.13*	0.52 $\pm$ 0.31*	0.56 $\pm$ 0.27*	0.50 $\pm$ 0.30*
Plasma MDA (μ mol/ml)	1.04 $\pm$ 0.02	1.04 $\pm$ 0.02	1.03 $\pm$ 0.03	1.04 $\pm$ 0.02	1.03 $\pm$ 0.02	1.03 $\pm$ 0.02
Echocardiography						
Fractional shortening (%)	55 $\pm$ 2	53 $\pm$ 2	55 $\pm$ 4	52 $\pm$ 4	54 $\pm$ 5	52 $\pm$ 3
LV ejection fraction (%)	80 $\pm$ 2	77 $\pm$ 3	76 $\pm$ 4	75 $\pm$ 4	75 $\pm$ 4	74 $\pm$ 5
HRV						
LF:HF ratio	0.21 $\pm$ 0.04	0.22 $\pm$ 0.03	0.22 $\pm$ 0.05	0.23 $\pm$ 0.03	0.26 $\pm$ 0.05	0.23 $\pm$ 0.06

NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg; LV ejection fraction, left ventricular ejection fraction; LF:HF ratio, low-frequency:high-frequency ratio; * P < 0.05 vs NMV.

Effects of pharmacological interventions on plasma MDA in diabetic rats

At 4 weeks after the STZ injection, plasma MDA levels were significantly increased in DMV rats when compared with NMV rats (Tables 3, 4 and 5). All treatments in diabetic rats significantly decreased the plasma MDA level, compared with DMV rats.

Effects of pharmacological interventions on echocardiography in diabetic rats

During week 4 after the STZ injection, diabetic rats developed cardiac contractile dysfunction that was characterized by significantly decreased %FS and %LVEF when compared with NMV rats (Tables 3, 4 and 5). Moreover, beginning at the fourth week for the insulin group and combined drug group,

and in the eighth week after both doses of PCA treatment, a significant increase in the %FS and %LVEF occurred in diabetic rats when compared with DMV rats (Tables 3, 4 and 5). Interestingly, beginning at week 8 of PCA treatment at a dose of 100 mg/kg in diabetic rats, the echocardiography results did not differ from those for DMI rats. During week 12 of the combined drug treatment in diabetic rats, the echocardiography was restored to a normal status, whereas DMI alone could not achieve a similar benefit as %FS and %LVEF were still lower than those in the normal rats.

Effects of pharmacological interventions on HRV in diabetic rats

At 4 weeks after STZ injection, diabetic control rats had a significantly increased LF:HF ratio when compared with NMV rats, indicating a cardiac autonomic imbalance

Table 3 Effects of pharmacological interventions on metabolic parameters, echocardiography, and heart rate variability (HRV) at week 4 of treatment. Values are expressed as mean $\pm$ s.d. of five to six rats in each group

Parameters	NMV	DMV	DMI	DML	DMH	DMHI
Metabolic parameters						
Body weight (g)	397 $\pm$ 23	237 $\pm$ 40*	310 $\pm$ 18* [†]	234 $\pm$ 25* [‡]	241 $\pm$ 16* [‡]	296 $\pm$ 30* [†]
Food intake (g/day)	23 $\pm$ 2	34 $\pm$ 2*	29 $\pm$ 1* [†]	33 $\pm$ 2* [‡]	33 $\pm$ 2* [‡]	29 $\pm$ 2* [†]
Fasting blood glucose (mg/dl)	99 $\pm$ 5	431 $\pm$ 88*	166 $\pm$ 25* [†]	316 $\pm$ 42* ^{†,‡}	291 $\pm$ 58* ^{†,‡}	155 $\pm$ 42* [†]
Plasma insulin (ng/ml)	1.30 $\pm$ 0.29	0.32 $\pm$ 0.13*	1.50 $\pm$ 0.86 [†]	0.57 $\pm$ 0.20* [‡]	0.46 $\pm$ 0.09* [‡]	1.28 $\pm$ 0.10 [†]
Plasma MDA (μ mol/ml)	1.03 $\pm$ 0.08	1.14 $\pm$ 0.10*	1.05 $\pm$ 0.05 [†]	1.03 $\pm$ 0.04 [†]	1.00 $\pm$ 0.02 [†]	0.98 $\pm$ 0.06 [†]
Echocardiography						
Fractional shortening (%)	58 $\pm$ 4	43 $\pm$ 3*	55 $\pm$ 3 [†]	43 $\pm$ 2* [‡]	44 $\pm$ 2* [‡]	56 $\pm$ 2 [†]
LV ejection fraction (%)	82 $\pm$ 3	68 $\pm$ 4*	79 $\pm$ 3 [†]	68 $\pm$ 2* [‡]	69 $\pm$ 3* [‡]	80 $\pm$ 3 [†]
HRV						
LF:HF ratio	0.23 $\pm$ 0.08	0.45 $\pm$ 0.06*	0.269 $\pm$ 0.025 [†]	0.42 $\pm$ 0.04* [‡]	0.41 $\pm$ 0.03* [‡]	0.25 $\pm$ 0.06 [†]

NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg; LV ejection fraction, left ventricular ejection fraction; LF:HF ratio, low-frequency:high-frequency ratio. * P < 0.05 vs NMV, [†] P < 0.05 vs DMV, and [‡] P < 0.05 vs DMI.

Table 4 Effects of pharmacological interventions on metabolic parameters, echocardiography, and heart rate variability (HRV) at week 8 of treatment. Values are expressed as mean $\pm$ s.d. of five to six rats in each group

Parameters	NMV	DMV	DMI	DML	DMH	DMHI
Metabolic parameters						
Body weight (g)	411 $\pm$ 46	233 $\pm$ 23*	323 $\pm$ 32*, [†]	244 $\pm$ 23*, [‡]	243 $\pm$ 17*, [‡]	319 $\pm$ 24*, [†]
Food intake (g/day)	24 $\pm$ 1	36 $\pm$ 1*	30 $\pm$ 1*, [†]	34 $\pm$ 1*, [‡]	35 $\pm$ 2*, [‡]	31 $\pm$ 2*, [†]
Fasting blood glucose (mg/dl)	94 $\pm$ 6	441 $\pm$ 42*	164 $\pm$ 42*, [†]	331 $\pm$ 46*, ^{†,‡}	270 $\pm$ 53*, ^{†,‡}	156 $\pm$ 31*, [†]
Plasma insulin (ng/ml)	1.71 $\pm$ 1.14	0.43 $\pm$ 0.14*	1.49 $\pm$ 1.01 [†]	0.42 $\pm$ 0.08*, [‡]	0.46 $\pm$ 0.19*, [‡]	2.01 $\pm$ 0.91 [†]
Plasma MDA (μ mol/ml)	1.87 $\pm$ 0.03	2.09 $\pm$ 0.08*	1.88 $\pm$ 0.03 [†]	1.90 $\pm$ 0.04 [†]	1.89 $\pm$ 0.03 [†]	1.88 $\pm$ 0.02 [†]
Echocardiography						
Fractional shortening (%)	58 $\pm$ 4	33 $\pm$ 4*	55 $\pm$ 5 [†]	49 $\pm$ 1*, ^{†,‡}	51 $\pm$ 1*, [†]	55 $\pm$ 4 [†]
LV ejection fraction (%)	82 $\pm$ 3	55 $\pm$ 4*	79 $\pm$ 4 [†]	74 $\pm$ 1*, ^{†,‡}	75 $\pm$ 2*, [†]	80 $\pm$ 3 [†]
HRV						
LF:HF ratio	0.23 $\pm$ 0.06	0.55 $\pm$ 0.06*	0.31 $\pm$ 0.10 [†]	0.44 $\pm$ 0.06*, ^{†,‡}	0.37 $\pm$ 0.07*, [†]	0.30 $\pm$ 0.06 [†]

NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg; LV ejection fraction, left ventricular ejection fraction; LF:HF ratio, low-frequency/high-frequency ratio. * P < 0.05 vs NMV, [†] P < 0.05 vs DMV, and [‡] P < 0.05 vs DMI.

(Table 3). During the fourth week of insulin and of combined drug treatment, and week 8 of both dosages of PCA, there was a significant decrease in the LF:HF ratio in diabetic rats, compared with DMV rats (Tables 3, 4 and 5). Interestingly, beginning at week 8 of PCA treatment at a dose of 100 mg/kg in diabetic rats, the HRV did not differ from that for insulin treatment in diabetic rats.

Effects of pharmacological interventions on plasma HbA1c levels in diabetic rats

At week 12 of treatment, DMV had significantly elevated plasma HbA1c levels when compared with NMV rats. PCA (both doses), insulin, and combined drug treatments in diabetic rats decreased the plasma HbA1c levels, compared with DMV rats (Fig. 1A).

Effects of pharmacological interventions on cardiac MDA levels in diabetic rats

At the end of treatment, cardiac MDA levels were significantly increased in DMV rats, compared with NMV rats (Fig. 1B). PCA (both doses), insulin, and combined drug treatments in diabetic rats significantly decreased cardiac MDA levels, compared with DMV rats. Interestingly, in DMH, the cardiac MDA level decreased similarly to that of to DMI rats.

Effects of pharmacological interventions on cardiac mitochondrial function and morphology in diabetic rats

Cardiac mitochondrial ROS production at the end of treatment was significantly increased in DMV rats,

Table 5 Effects of pharmacological intervention on metabolic parameters, echocardiography, and heart rate variability (HRV) at week 12 of treatment. Values are expressed as mean $\pm$ s.d. of five to six rats in each group

Parameters	NMV	DMV	DMI	DML	DMH	DMHI
Metabolic parameters						
Body weight (g)	444 $\pm$ 23	225 $\pm$ 12*	353 $\pm$ 18*, [†]	243 $\pm$ 21*, [‡]	246 $\pm$ 19*, [‡]	339 $\pm$ 23*, [†]
Food intake (g/day)	24 $\pm$ 3	38 $\pm$ 3*	33 $\pm$ 2*, [†]	35 $\pm$ 2*, [‡]	34 $\pm$ 3*, [‡]	32 $\pm$ 1*, [†]
Fasting blood glucose (mg/dl)	102 $\pm$ 7	541 $\pm$ 64*	184 $\pm$ 26*, [†]	312 $\pm$ 90*, ^{†,‡}	270 $\pm$ 58*, ^{†,‡}	176 $\pm$ 27*, [†]
Plasma insulin (ng/ml)	1.23 $\pm$ 0.21	0.39 $\pm$ 0.07*	1.39 $\pm$ 0.81 [†]	0.56 $\pm$ 0.14*, [‡]	0.49 $\pm$ 0.22*, [‡]	1.23 $\pm$ 0.80 [†]
Plasma MDA (μ mol/ml)	2.14 $\pm$ 0.02	2.25 $\pm$ 0.05*	2.16 $\pm$ 0.03 [†]	2.19 $\pm$ 0.03*, [†]	2.19 $\pm$ 0.04*, [†]	2.13 $\pm$ 0.02 [†]
Echocardiography						
Fractional shortening (%)	57 $\pm$ 3	33 $\pm$ 3*	52 $\pm$ 3*, [†]	49 $\pm$ 3*, [†]	48 $\pm$ 5*, [†]	56 $\pm$ 3*, [†]
LV ejection fraction (%)	82 $\pm$ 2	54 $\pm$ 5*	75 $\pm$ 3*, [†]	74 $\pm$ 3*, [†]	73 $\pm$ 5*, [†]	80 $\pm$ 3*, [†]
HRV						
LF:HF ratio	0.24 $\pm$ 0.05	0.54 $\pm$ 0.13*	0.31 $\pm$ 0.04 [†]	0.41 $\pm$ 0.06*, [†]	0.34 $\pm$ 0.04*, [†]	0.29 $\pm$ 0.14 [†]

NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg; LV ejection fraction, left ventricular ejection fraction; LF:HF ratio, low-frequency/high-frequency ratio. * P < 0.05 vs NMV, [†] P < 0.05 vs DMV, and [‡] P < 0.05 vs DMI.

Figure 1

Effects of pharmacological interventions on plasma HbA1c and cardiac MDA levels. At the end of treatments, (A) plasma HbA1c level was increased in DMV rats, all treatments significantly decreased plasma HbA1c levels in diabetic rats. (B) Cardiac MDA level was elevated in DMV rats, and all treatments decreased cardiac MDA level in diabetic rats. Values are expressed as mean $\pm$ S.E.M. NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH+insulin, 4 U/kg. * P <0.05 vs NMV, † P <0.05 vs DMV, ‡ P <0.05 vs DMI.

compared with NMV rats (Fig. 2A). PCA (both doses), insulin, and combined drug treatments in diabetic rats significantly reduced the mitochondrial ROS levels, compared with DMV rats. For mitochondrial membrane potential change, the red:green fluorescence intensity ratio was significantly decreased in DMV rats, compared with NMV rats, indicating cardiac mitochondrial membrane depolarization. All treatments attenuated mitochondrial depolarization, compared with DMV rats (Fig. 2B). For assessment of cardiac mitochondrial swelling, the absorbance was significantly decreased in the DMV rats, compared with NMV rats, indicating mitochondrial swelling (Fig. 2C). Moreover, PCA (both doses), insulin, and combined drug treatments significantly decreased mitochondrial swelling in diabetic rats, compared with DMV rats (Fig. 2C). The TEM images of

cardiac mitochondria from each group are shown in Fig. 2D. Compared with the intact cardiac mitochondria of NMV rats, unfolded cristae were mostly found in DMV rats, indicating mitochondrial swelling. All treatments attenuated mitochondrial swelling, as indicated by decreased unfolded cristae in diabetic rats.

Effects of pharmacological interventions on BAX and BCL2 protein expression in the heart of diabetic rats

At week 12 after treatment, in the DMV rats, although BAX expression was not changed (Fig. 3A), BCL2 expression was significantly decreased compared with NMV rats (Fig. 3B). PCA (both doses), insulin, and combined drug treatments significantly increased BCL2 protein levels in diabetic rats, compared with DMV rats (Fig. 3B and C).

Discussion

This study demonstrated that STZ-induced type 1 diabetic rats were characterized by hyperglycemia, insulin deficiency, body weight loss, polyphagia, increased oxidative damage, depressed HRV, cardiac contractile dysfunction, cardiac mitochondrial dysfunction, and decreased cardiac anti-apoptotic BCL2 protein levels. The insulin and combined drug treatments increased the plasma insulin level, elevated body weight, and reduced food intake. In addition, all treatments in diabetic rats decreased the blood glucose level, reduced oxidative damage, improved HRV, attenuated cardiac dysfunction, prevented cardiac mitochondrial dysfunction, and increased BCL2 protein expression in diabetic rats. Interestingly, PCA at a dose of 100 mg/kg improved HRV and cardiac dysfunction similarly to insulin treatment in diabetic rats. Moreover, diabetic rats treated with combined drugs had echocardiography results restored to a normal status, whereas insulin alone could not achieve this. Insulin and combined drug treatments in diabetic rats also restored HRV.

A previous study demonstrated that PCA administration prevented the increase in plasma glucose and HbA1c levels and the decrease in plasma insulin levels in STZ-induced diabetic rats (Harini & Pugalendi 2010). PCA normalized the activities of gluconeogenic enzymes such as glucose 6-phosphatase and fructose 1,6-bisphosphatase, as well as that of the glycolytic enzyme glucokinase (Harini & Pugalendi 2010). In STZ-induced diabetic mice, dietary supplementation with PCA improved glycemic control and attenuated homeostatic disorders (Lin *et al.* 2009, 2011). Consistent with our findings, PCA treatment alone decreased hyperglycemia, whereas it could not

Figure 2

Effects of pharmacological interventions on cardiac mitochondrial function and morphology. (A) Cardiac mitochondrial ROS production was increased in DMV rats; all treatments reduced mitochondrial ROS production in diabetic rats. (B) Cardiac mitochondrial swelling was increased in DMV rats; all treatments prevented mitochondrial swelling in diabetic rats. (C) Cardiac mitochondrial membrane depolarization was increased in DMV rats; all treatments prevented mitochondrial membrane depolarization in

diabetic rats. (D) Representative TEM images of mitochondrial morphology. Values are expressed as mean $\pm$ s.e.m. NMV, normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg. * $P < 0.05$ vs NMV and $^{\dagger}P < 0.05$ vs DMV.

increase insulin levels in STZ-induced diabetic rats. As PCA has been shown to exert an insulin-like activity by increasing glucose uptake via enhancing GLUT4 translocation and adiponectin secretion caused by the increased PPAR γ activity in adipocytes (Scazzocchio *et al.* 2011), this could be the mechanism responsible for its glycemic effect observed in this study.

The increased ROS generation and impaired antioxidant defenses both contribute to oxidative stress in diabetes. Most of the ROS generated within cells are from mitochondria that were exposed to hyperglycemia (Brownlee 1995, Shen *et al.* 2006). The MDA is an index of oxidative damage. Results from previous studies indicated that plasma and tissue MDA levels are elevated in STZ-induced diabetic rats, and that PCA decreased plasma, renal, and cardiac MDA levels, as well as elevated antioxidant defense in STZ-induced diabetic mice (Lin *et al.* 2009, Naowaboot *et al.* 2009). Consistent with results from our study, diabetic rats treated with PCA had

decreased oxidative damage in both plasma and cardiac tissues. PCA also attenuated cardiac mitochondrial ROS production in diabetic rats, especially PCA at a dose of 100 mg/kg, which exhibited a strong potency similar to that of the insulin treatment. Increased ROS has been proposed to amplify hyperglycemia-induced MAPK isoform activation and increase advanced glycation end product (AGE) formation (Brownlee 1995, Koya & King 1998), which could promote cell injury that contributes to the development of cardiac dysfunction in diabetes. HbA1c is one of the AGEs, a form of hemoglobin that is measured primarily to identify the average plasma glucose concentration over prolonged periods of time. Monitoring HbA1c in type 1 diabetic patients may improve the outcomes (Larsen *et al.* 1990). Results from a previous study indicated that PCA decreased plasma HbA1c levels and tissue AGEs in STZ-induced diabetic mice (Lin *et al.* 2011). Consistent with our findings, PCA decreased plasma HbA1c levels in STZ-induced diabetic rats.

Figure 3

Effects of pharmacological interventions on cardiac BAX and BCL2 levels. (A) Cardiac BAX expression was not different among groups. (B) Cardiac BCL2 expression was decreased in DMV rats; all treatments increased cardiac BCL2 expression in diabetic rats. (C) Representative blot bands for BAX and BCL2 in each group. Values are expressed as mean $\pm$ s.e.m. NMV,

normal rats treated with vehicle; DMV, diabetic rats treated with vehicle; DMI, diabetic rats treated with insulin, 4 U/kg; DML, diabetic rats treated with PCA, 50 mg/kg; DMH, diabetic rats treated with PCA, 100 mg/kg; DMHI, DMH + insulin, 4 U/kg. * $P < 0.05$ vs NMV and $^{\dagger}P < 0.05$ vs DMV.

In human studies, progressive autonomic dysfunction and depressed HRV in diabetic patients indicated cardiac autonomic neuropathy (Malpas & Maling 1990, Kudat *et al.* 2006). The results from an animal study indicated that HRV was depressed in the STZ-induced diabetic rat (Fazan *et al.* 1997). As depressed HRV has been associated with increased oxidative damage, such as increased plasma MDA levels (Pavithran *et al.* 2008), the prevention of depressed HRV in diabetic rats by PCA could be due to its ability to decrease oxidative stress in this study.

Results from previous studies indicated that the cardiac function index including ejection fraction and stroke volume was depressed at 4 weeks following STZ-induced diabetes (Crespo *et al.* 2008, 2011). Consistent with our findings, echocardiography showed a reduction of %FS and %LVEF in diabetic rats in this study. Moreover, mitochondria are known to be the source of energy required for the heart to function properly. The effects of PCA in reducing oxidative stress as well as attenuating cardiac mitochondrial dysfunction could be responsible for improved cardiac function as observed in this study. Supporting this statement is the fact that beginning at 4 weeks for insulin or combined drug treatments and at 8 weeks for both doses of PCA treatment, improved cardiac function was observed in STZ-induced diabetic rats included in our study. It is important to note that despite the fact that PCA improved cardiac function, insulin could provide this improvement faster (by the fourth week of treatment) than PCA (by the eighth week).

Overproduction of ROS in diabetes can activate the mitochondrial pathway, resulting in loss of mitochondrial membrane integrity and release of cytochrome *c*, apoptotic protease-activating factor 1, and other pro-apoptotic factors in the cytoplasm. Maintenance of mitochondrial membrane potential depends on pro-apoptotic (Bax) and anti-apoptotic (Bcl2) members of Bcl2 family, causing or preventing cytochrome *c* release. PCA has been shown to prevent apoptosis by attenuating the change in the mitochondrial membrane permeability, decreasing oxidative damage, and increasing anti-apoptotic BCL2 expression in MPP⁺-induced mitochondrial dysfunction and apoptotic cell death in PC12 cells (Guan *et al.* 2006). These findings were consistent with our study demonstrating that PCA improved cardiac mitochondrial function and increased cardiac BCL2 level in diabetic rats.

In summary, in the hearts of STZ-induced diabetic rats, cardiac autonomic imbalance and cardiac dysfunction were developed. PCA, insulin, and combined drug treatments attenuated these adverse effects by attenuating

hyperglycemia, oxidative damage, cardiac autonomic imbalance, and cardiac mitochondrial dysfunction, and increasing anti-apoptotic protein. These findings indicated that PCA could exert beneficial effects on the heart in type 1 DM and may be used as a supplement to prevent cardiac complications in diabetic patients.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Faculty of Medicine Chiang Mai University Endowment Fund (N C), the Thailand Research Fund RTA5580006 (N C) and BRG 57 (S C C), TRG 57 (S K), Chiang Mai University Excellent Center Award (N C), and Udon Thani Rajabhat University Fund (Y S).

References

- An D & Rodrigues B 2006 Role of changes in cardiac metabolism in development of diabetic cardiomyopathy. *American Journal of Physiology. Heart and Circulatory Physiology* **291** H1489–H1506. (doi:10.1152/ajpheart.00278.2006)
- Apaijai N, Pintana H, Chattipakorn SC & Chattipakorn N 2012 Cardio-protective effects of metformin and vildagliptin in adult rats with insulin resistance induced by a high-fat diet. *Endocrinology* **153** 3878–3885. (doi:10.1210/en.2012-1262)
- Apaijai N, Pintana H, Chattipakorn SC & Chattipakorn N 2013 Effects of vildagliptin versus sitagliptin, on cardiac function, heart rate variability and mitochondrial function in obese insulin-resistant rats. *British Journal of Pharmacology* **169** 1048–1057. (doi:10.1111/bph.12176)
- Asbun J & Villarreal FJ 2006 The pathogenesis of myocardial fibrosis in the setting of diabetic cardiomyopathy. *Journal of the American College of Cardiology* **47** 693–700. (doi:10.1016/j.jacc.2005.09.050)
- Boudina S & Abel ED 2007 Diabetic cardiomyopathy revisited. *Circulation* **115** 3213–3223. (doi:10.1161/CIRCULATIONAHA.106.679597)
- Brownlee M 1995 Advanced protein glycosylation in diabetes and aging. *Annual Review of Medicine* **46** 223–234. (doi:10.1146/annurev.med.46.1.223)
- Chinda K, Palee S, Surinkaew S, Phornphutkul M, Chattipakorn S & Chattipakorn N 2013 Cardioprotective effect of dipeptidyl peptidase-4 inhibitor during ischemia–reperfusion injury. *International Journal of Cardiology* **167** 451–457. (doi:10.1016/j.ijcard.2012.01.011)
- Chou TH, Ding HY, Lin RJ, Liang JY & Liang CH 2010 Inhibition of melanogenesis and oxidation by protocatechuic acid from *Origanum vulgare* (oregano). *Journal of Natural Products* **73** 1767–1774. (doi:10.1021/np100281g)
- Crespo MJ, Zalacain J, Dunbar DC, Cruz N & Arocho L 2008 Cardiac oxidative stress is elevated at the onset of dilated cardiomyopathy in streptozotocin-diabetic rats. *Journal of Cardiovascular Pharmacology and Therapeutics* **13** 64–71. (doi:10.1177/1074248407307854)
- Crespo MJ, Marrero M, Cruz N, Quidgley J, Creagh O, Torres H & Rivera K 2011 Diabetes alters cardiovascular responses to anaesthetic induction agents in STZ-diabetic rats. *Diabetes & Vascular Disease Research* **8** 299–302. (doi:10.1177/1479164111421035)
- Dyntar D, Sergeev P, Klisic J, Ambuhl P, Schaub MC & Donath MY 2006 High glucose alters cardiomyocyte contacts and inhibits myofibrillar formation. *Journal of Clinical Endocrinology and Metabolism* **91** 1961–1967. (doi:10.1210/jc.2005-1904)

- Fazan RJr, Ballejo G, Salgado MC, Moraes MF & Salgado HC 1997 Heart rate variability and baroreceptor function in chronic diabetic rats. *Hypertension* **30** 632–635. (doi:10.1161/01.HYP.30.3.632)
- Guan S, Jiang B, Bao YM & An LJ 2006 Protocatechuic acid suppresses MPP⁺-induced mitochondrial dysfunction and apoptotic cell death in PC12 cells. *Food and Chemical Toxicology* **44** 1659–1666. (doi:10.1016/j.fct.2006.05.004)
- Harini R & Pugalendi KV 2010 Antihyperglycemic effect of protocatechuic acid on streptozotocin-diabetic rats. *Journal of Basic and Clinical Physiology and Pharmacology* **21** 79–91. (doi:10.1515/JBCPP.2010.21.1.79)
- Incharoen T, Thephinlap C, Srichairatanakool S, Chattipakorn S, Winichagoon P, Fucharoen S, Vadolas J & Chattipakorn N 2007 Heart rate variability in β -thalassemic mice. *International Journal of Cardiology* **121** 203–204. (doi:10.1016/j.ijcard.2006.08.076)
- Jiang H, Fang J, Wu B, Yin G, Sun L, Qu J, Barger SW & Wu S 2011 Overexpression of serine racemase in retina and overproduction of D-serine in eyes of streptozotocin-induced diabetic retinopathy. *Journal of Neuroinflammation* **8** 119. (doi:10.1186/1742-2094-8-119)
- Kay CD, Mazza GJ & Holub BJ 2005 Anthocyanins exist in the circulation primarily as metabolites in adult men. *Journal of Nutrition* **135** 2582–2588.
- Koya D & King GL 1998 Protein kinase C activation and the development of diabetic complications. *Diabetes* **47** 859–866. (doi:10.2337/diabetes.47.6.859)
- Krishna KM, Gopal GS, Chalam CR, Madan K, Kumar VK, Prakash GJ & Annapurna A 2005 The influence of sulindac on diabetic cardiomyopathy: a non-invasive evaluation by Doppler echocardiography in streptozotocin-induced diabetic rats. *Vascular Pharmacology* **43** 91–100. (doi:10.1016/j.vph.2005.02.012)
- Kudat H, Akkaya V, Sozen AB, Salman S, Demirel S, Ozcan M, Atilgan D, Yilmaz MT & Guven O 2006 Heart rate variability in diabetes patients. *Journal of International Medical Research* **34** 291–296. (doi:10.1177/147323000603400308)
- Kumar S & Sitasawad SL 2009 N-acetylcysteine prevents glucose/glucose oxidase-induced oxidative stress, mitochondrial damage and apoptosis in H9c2 cells. *Life Sciences* **84** 328–336. (doi:10.1016/j.lfs.2008.12.016)
- Kumfu S, Chattipakorn S, Chinda K, Fucharoen S & Chattipakorn N 2012 T-type calcium channel blockade improves survival and cardiovascular function in thalassemic mice. *European Journal of Haematology* **88** 535–548. (doi:10.1111/j.1600-0609.2012.01779.x)
- Larsen ML, Horder M & Mogensen EF 1990 Effect of long-term monitoring of glycosylated hemoglobin levels in insulin-dependent diabetes mellitus. *New England Journal of Medicine* **323** 1021–1025. (doi:10.1056/NEJM199010113231503)
- Lekawanvijit S, Kompa AR, Zhang Y, Wang BH, Kelly DJ & Krum H 2012 Myocardial infarction impairs renal function, induces renal interstitial fibrosis, and increases renal KIM-1 expression: implications for cardiorenal syndrome. *American Journal of Physiology. Heart and Circulatory Physiology* **302** H1884–H1893. (doi:10.1152/ajpheart.00967.2011)
- Lin HH, Chen JH, Huang CC & Wang CJ 2007 Apoptotic effect of 3,4-dihydroxybenzoic acid on human gastric carcinoma cells involving JNK/p38 MAPK signaling activation. *International Journal of Cancer* **120** 2306–2316. (doi:10.1002/ijc.22571)
- Lin CY, Huang CS, Huang CY & Yin MC 2009 Anticoagulatory, antiinflammatory, and antioxidative effects of protocatechuic acid in diabetic mice. *Journal of Agricultural and Food Chemistry* **57** 6661–6667. (doi:10.1021/jf9015202)
- Lin CY, Tsai SJ, Huang CS & Yin MC 2011 Antiglycative effects of protocatechuic acid in the kidneys of diabetic mice. *Journal of Agricultural and Food Chemistry* **59** 5117–5124. (doi:10.1021/jf200103f)
- Liu CL, Wang JM, Chu CY, Cheng MT & Tseng TH 2002 *In vivo* protective effect of protocatechuic acid on tert-butyl hydroperoxide-induced rat hepatotoxicity. *Food and Chemical Toxicology* **40** 635–641. (doi:10.1016/S0278-6915(02)00002-9)
- Malpas SC & Maling TJ 1990 Heart-rate variability and cardiac autonomic function in diabetes. *Diabetes* **39** 1177–1181. (doi:10.2337/diab.39.10.1177)
- Masella R, Vari R, D'Archivio M, Di Benedetto R, Matarrese P, Malorni W, Scazzocchio B & Giovannini C 2004 Extra virgin olive oil biophenols inhibit cell-mediated oxidation of LDL by increasing the mRNA transcription of glutathione-related enzymes. *Journal of Nutrition* **134** 785–791.
- Min SW, Ryu SN & Kim DH 2010 Anti-inflammatory effects of black rice, cyanidin-3-O- β -D-glycoside, and its metabolites, cyanidin and protocatechuic acid. *International Immunopharmacology* **10** 959–966. (doi:10.1016/j.intimp.2010.05.009)
- Naowaboot J, Pannangpetch P, Kukongviriyapan V, Kukongviriyapan U, Nakmareong S & Itharat A 2009 Mulberry leaf extract restores arterial pressure in streptozotocin-induced chronic diabetic rats. *Nutrition Research* **29** 602–608. (doi:10.1016/j.nutres.2009.06.002)
- Ohuchi H, Suzuki H, Yasuda K, Arakaki Y, Echigo S & Kamiya T 2000 Heart rate recovery after exercise and cardiac autonomic nervous activity in children. *Pediatric Research* **47** 329–335. (doi:10.1203/00006450-200003000-00008)
- Palee S, Weerateerangkul P, Chinda K, Chattipakorn SC & Chattipakorn N 2013 Mechanisms responsible for beneficial and adverse effects of rosiglitazone in a rat model of acute cardiac ischaemia-reperfusion. *Experimental Physiology* **98** 1028–1037. (doi:10.1113/expphysiol.2012.070433)
- Pavithran P, Nandeesha H, Sathiyapriya V, Bobby Z & Madanmohan T 2008 Short-term heart variability and oxidative stress in newly diagnosed essential hypertension. *Clinical and Experimental Hypertension* **30** 486–496. (doi:10.1080/10641960802251875)
- Prachayasakul W, Kerdphoo S, Petsophonakul P, Pongchaidecha A, Chattipakorn N & Chattipakorn SC 2011 Effects of high-fat diet on insulin receptor function in rat hippocampus and the level of neuronal corticosterone. *Life Sciences* **88** 619–627. (doi:10.1016/j.lfs.2011.02.003)
- Raher MJ, Thibault HB, Buys ES, Kuruppu D, Shimizu N, Brownell AL, Blake SL, Rieusset J, Kaneki M, Derumeaux G *et al.* 2008 A short duration of high-fat diet induces insulin resistance and predisposes to adverse left ventricular remodeling after pressure overload. *American Journal of Physiology. Heart and Circulatory Physiology* **295** H2495–H2502. (doi:10.1152/ajpheart.00139.2008)
- Scazzocchio B, Vari R, Filesi C, D'Archivio M, Santangelo C, Giovannini C, Iacovelli A, Silecchia G, Li Volti G, Galvano F *et al.* 2011 Cyanidin-3-O- β -glucoside and protocatechuic acid exert insulin-like effects by upregulating PPAR γ activity in human omental adipocytes. *Diabetes* **60** 2234–2244. (doi:10.2337/db10-1461)
- Shen X, Zheng S, Metreveli NS & Epstein PN 2006 Protection of cardiac mitochondria by overexpression of MnSOD reduces diabetic cardiomyopathy. *Diabetes* **55** 798–805. (doi:10.2337/diabetes.55.03.06.db05-1039)
- Shi GF, An LJ, Jiang B, Guan S & Bao YM 2006 Alpinia protocatechuic acid protects against oxidative damage *in vitro* and reduces oxidative stress *in vivo*. *Neuroscience Letters* **403** 206–210. (doi:10.1016/j.neulet.2006.02.057)
- Sies H 1991 Oxidative stress: from basic research to clinical application. *American Journal of Medicine* **91** 31S–38S. (doi:10.1016/0002-9343(91)90281-2)
- Sroka Z & Cisowski W 2003 Hydrogen peroxide scavenging, antioxidant and anti-radical activity of some phenolic acids. *Food and Chemical Toxicology* **41** 753–758. (doi:10.1016/S0278-6915(02)00329-0)
- Surinkaew S, Kumphune S, Chattipakorn S & Chattipakorn N 2013 Inhibition of p38 MAPK during ischemia, but not reperfusion, effectively attenuates fatal arrhythmia in ischemia/reperfusion heart. *Journal of Cardiovascular Pharmacology* **61** 133–141. (doi:10.1097/FJC.0b013e318279b7b1)
- Tarozzi A, Morroni F, Hrelia S, Angeloni C, Marchesi A, Cantelli-Forti G & Hrelia P 2007 Neuroprotective effects of anthocyanins and their *in vivo*

- metabolites in SH-SY5Y cells. *Neuroscience Letters* **424** 36–40. (doi:10.1016/j.neulet.2007.07.017)
- Thummasorn S, Kumfu S, Chattipakorn S & Chattipakorn N 2011 Granulocyte-colony stimulating factor attenuates mitochondrial dysfunction induced by oxidative stress in cardiac mitochondria. *Mitochondrion* **11** 457–466. (doi:10.1016/j.mito.2011.01.008)
- Trost S & LeWinter M 2001 Diabetic cardiomyopathy. *Current Treatment Options in Cardiovascular Medicine* **3** 481–492. (doi:10.1007/s11936-001-0022-9)
- Tseng TH, Kao TW, Chu CY, Chou FP, Lin WL & Wang CJ 2000 Induction of apoptosis by hibiscus protocatechuic acid in human leukemia cells via reduction of retinoblastoma (RB) phosphorylation and Bcl-2 expression. *Biochemical Pharmacology* **60** 307–315. (doi:10.1016/S0006-2952(00)00322-1)
- Vari R, D'Archivio M, Filesi C, Carotenuto S, Scazzocchio B, Santangelo C, Giovannini C & Masella R 2011 Protocatechuic acid induces anti-oxidant/detoxifying enzyme expression through JNK-mediated Nrf2 activation in murine macrophages. *Journal of Nutritional Biochemistry* **22** 409–417. (doi:10.1016/j.jnutbio.2010.03.008)
- Vitaglione P, Donnarumma G, Napolitano A, Galvano F, Gallo A, Scalfi L & Fogliano V 2007 Protocatechuic acid is the major human metabolite of cyanidin-glucosides. *Journal of Nutrition* **137** 2043–2048.
- Wang D, Wei X, Yan X, Jin T & Ling W 2010 Protocatechuic acid, a metabolite of anthocyanins, inhibits monocyte adhesion and reduces atherosclerosis in apolipoprotein E-deficient mice. *Journal of Agricultural and Food Chemistry* **58** 12722–12728. (doi:10.1021/jf103427j)
- Wang D, Zou T, Yang Y, Yan X & Ling W 2011 Cyanidin-3-O- β -glucoside with the aid of its metabolite protocatechuic acid, reduces monocyte infiltration in apolipoprotein E-deficient mice. *Biochemical Pharmacology* **82** 713–719. (doi:10.1016/j.bcp.2011.04.007)
- Yan JJ, Jung JS, Hong YJ, Moon YS, Suh HW, Kim YH, Yun-Choi HS & Song DK 2004 Protective effect of protocatechuic acid isopropyl ester against murine models of sepsis: inhibition of TNF- α and nitric oxide production and augmentation of IL-10. *Biological & Pharmaceutical Bulletin* **27** 2024–2027. (doi:10.1248/bpb.27.2024)
- Yin MC, Lin CC, Wu HC, Tsao SM & Hsu CK 2009 Apoptotic effects of protocatechuic acid in human breast, lung, liver, cervix, and prostate cancer cells: potential mechanisms of action. *Journal of Agricultural and Food Chemistry* **57** 6468–6473. (doi:10.1021/jf9004466)
- Yip EC, Chan AS, Pang H, Tam YK & Wong YH 2006 Protocatechuic acid induces cell death in HepG2 hepatocellular carcinoma cells through a c-Jun N-terminal kinase-dependent mechanism. *Cell Biology and Toxicology* **22** 293–302. (doi:10.1007/s10565-006-0082-4)
- Zhang X, Shi GF, Liu XZ, An LJ & Guan S 2011 Anti-ageing effects of protocatechuic acid from *Alpinia* on spleen and liver antioxidative system of senescent mice. *Cell Biochemistry and Function* **29** 342–347. (doi:10.1002/cbf.1757)
- Zhao SM, Wang YL, Guo CY, Chen JL & Wu YQ 2014 Progressive decay of Ca²⁺ homeostasis in the development of diabetic cardiomyopathy. *Cardiovascular Diabetology* **13** 75. (doi:10.1186/1475-2840-13-75)

Received in final form 17 July 2014

Accepted 28 July 2014

Accepted Preprint published online 29 July 2014