STAT4 contributes to adipose tissue inflammation and atherosclerosis

A D Dobrian¹, M A Hatcher¹, J J Brotman¹, E V Galkina², P Taghavi-Moghadam²,
H Pei⁴, B A Haynes¹ and J L Nadler³

Departments of ¹Physiological Sciences, ²Microbiology and Molecular Cell Biology, and ³Internal Medicine,
Eastern Virginia Medical School, 700W Olney Road, Norfolk, Virginia 23505, USA

⁴Division of Inflammation Biology, La Jolla Institute for Allergy and Immunology, San Diego, La Jolla,
California, USA

Correspondence
should be addressed
to A D Dobrian
Email
dobriaad@evms.edu

Abstract

Adipose tissue (AT) inflammation is an emerging factor contributing to cardiovascular disease. STAT4 is a transcription factor expressed in adipocytes and in immune cells and contributes to AT inflammation and insulin resistance in obesity. The objective of this study was to determine the effect of STAT4 deficiency on visceral and peri-aortic AT inflammation in a model of atherosclerosis without obesity. *Stat4*^{−/−} *Apoe*^{−/−} mice and *Apoe*^{−/−} controls were kept either on chow or Western diet for 12 weeks. Visceral and peri-aortic AT were collected and analyzed for immune composition by flow cytometry and for cytokine/chemokine expression by real-time PCR. *Stat4*^{−/−} *Apoe*^{−/−} and *Apoe*^{−/−} mice had similar body weight, plasma glucose, and lipids. Western diet significantly increased macrophage, CD4⁺, CD8⁺, and NK cells in peri-aortic and visceral fat in *Apoe*^{−/−} mice. In contrast, in *Stat4*^{−/−} *Apoe*^{−/−} mice, a Western diet failed to increase the percentage of immune cells infiltrating the AT. Also, IL12p40, TNFα, CCL5, CXCL10, and CX3CL1 were significantly reduced in the peri-aortic fat in *Stat4*^{−/−} *Apoe*^{−/−} mice. Importantly, *Stat4*^{−/−} *Apoe*^{−/−} mice on a Western diet had significantly reduced plaque burden vs *Apoe*^{−/−} controls. In conclusion, STAT4 deletion reduces inflammation in peri-vascular and visceral AT and this may contribute via direct or indirect effects to reduced atheroma formation.

Key Words

- ▶ peri-aortic fat
- ▶ T lymphocytes
- ▶ chemokines
- ▶ macrophages

Journal of Endocrinology
(2015) 227, 13–24

Introduction

Excessive accumulation of visceral adipose tissue (AT) is independently associated with cardiovascular disease (See *et al.* 2007). In addition, the fat around the conduit vessels such as the aorta as well as the pericardial fat were shown to play a role in the progression of atherosclerosis (Ohman *et al.* 2011) and were associated with coronary atherosclerotic plaque formation in humans (Konishi *et al.* 2010a,b). Therefore, both visceral and perivascular AT accumulation and inflammation emerge as key contributors to atherosclerosis. The *Apoe*^{−/−} mouse model of

atherosclerosis displays accelerated plaque formation on high cholesterol diet. However, this mouse model neither gains weight nor develops insulin resistance as a result of high fat diet feeding (Gao *et al.* 2007, Kawashima *et al.* 2009). This phenotype was attributed, at least in part, to the inability of visceral AT to accumulate excess lipids resulting in a more sensitive adipocyte phenotype and reduced inflammation (Huang *et al.* 2006, 2013, Hofmann *et al.* 2008). This raises the question whether various AT depots may contribute to the development of

atherosclerosis in this model. While the contribution of visceral fat in atherosclerosis development in this model was not reported, the peri-vascular fat was proven causal for the development of atherosclerosis in *Apoe*^{-/-} mice fed a Western diet (Ohman *et al.* 2011).

The mechanisms contributing to atherosclerosis by visceral and perivascular fat are incompletely understood. In atherosclerotic *Apoe*^{-/-} mice, peri-adventitial AT produces elevated levels of interleukin 6 (IL6), IL1A, and MIP1A (Lohmann *et al.* 2009) and in a model of obesity with angiotensin II infusion peri-aortic AT induces inflammation and enhances aneurism formation (Police *et al.* 2009). The TLR/JAK-STAT pathway is activated in human peri-vascular adipocytes from patients with atherosclerosis (Police *et al.* 2009). STAT4 is downstream of the Jak/Tyk kinases and, upon phosphorylation in response to IL12 or other cytokines, induces expression of genes involved in the proliferation and differentiation of various hematopoietic and non-hematopoietic cells (Darnell 1997, Horvath & Darnell 1997, Imada & Leonard 2000, Leonard & Lin 2000). STAT4 is expressed in T and NK cells and has a prominent role for IL12-induced Th1 cell differentiation and for NK cell activation (Watford *et al.* 2004, Kaplan 2005, Good *et al.* 2009). IL12 is also highly expressed in rodent and human atherosclerotic lesions and several studies have shown that approaches to reduce IL12 levels prevent atherosclerosis (Zhao *et al.* 2002, Davenport & Tipping 2003, Hauer *et al.* 2005, Zhang *et al.* 2006, Eid *et al.* 2009). Importantly, recent findings indicate that STAT4 has a determinant role for optimal human Th1 lineage development (Chang *et al.* 2009). Our group showed that STAT4 is markedly activated in the balloon injured carotid artery of the obese Zucker rat, and that an IL12-signaling inhibitor can reduce STAT4 activation and vascular injury responses (Pei *et al.* 2006). In addition, STAT4-deficient mice are protected from developing insulin resistance on a high fat diet, in part due to reduced immune cell trafficking in visceral AT and reduced pro-inflammatory cytokine production by adipocytes (Dobrian *et al.* 2013). Collectively, these results suggest that activation of STAT4 may participate in vascular inflammatory responses in part via modulation of AT inflammation. To directly address this hypothesis we examined the effect of STAT4 deficiency on visceral and peri-aortic AT inflammation in *Stat4*^{-/-} *Apoe*^{-/-} mice, a model of atherosclerosis lacking the confounding effects of insulin resistance and obesity. A key finding is the significant effect of STAT4 deficiency on immune composition as well as pro-inflammatory cytokine and chemokine production mainly in the peri-aortic fat. The anti-inflammatory effect of STAT4 deficiency was

significant in the mice fed a high cholesterol diet and was associated with the reduced atherosclerotic plaque burden, suggesting that the activation of this pathway in AT may be a contributor to accelerated diet-related atherosclerosis.

Materials and methods

Animals and diets

All procedures involving animals were approved by the IACUC of Eastern Virginia Medical School and University of Virginia at Charlottesville. Female *Stat4*^{-/-} *Apoe*^{-/-} or *Apoe*^{-/-} mice were bred in our colonies and at 8–10 weeks of age were either fed a Western diet (0.15% cholesterol, Harlan, Madison, WI, USA) or were maintained on regular rodent chow for 12 weeks (*n* = 6–10 mice/group). All of the mice were between 20 and 22 weeks of age at euthanasia. Mice were housed in a pathogen-free facility, and food and water were provided a regular supply of food and allowed to feed *ad libitum* throughout the experiment. Body weights were measured weekly.

Lipids, glucose, and insulin

Lipids, glucose, and insulin were measured in all of the groups in non-fasted terminal plasma. Triglycerides, total cholesterol, LDL, and HDL cholesterol were measured using a colorimetric kit from Wako (Richmond, VA, USA). Plasma glucose was measured according to manufacturer's instructions using a colorimetric method (BioVision, Inc., Milpitas, CA, USA). Insulin was measured by ELISA using a commercially available kit (Mercodia, Winston Salem, NC, USA).

Stromal vascular fraction preparation

Samples of peri-gonadal (visceral) and peri-aortic AT (0.1–0.3 g) were digested with collagenase as described before with minor modifications (Weisberg *et al.* 2003, Cole *et al.* 2010). The floating adipocytes were collected and washed and the infranatant was removed and centrifuged at 500 *g* for 5 min to pellet the stromal vascular fraction (SVF). SVF was used for flow cytometry to determine immune cell content.

Flow cytometry

Counted SVF cells were incubated for 30 min, at room temperature with one of the following combinations of fluorophore-conjugated primary antibodies: *Cocktail1* (for macrophage phenotyping): CD11b-Pacific Blue,

CD45-PerCP, Cd11c-PE, F4/80-Alexa 647, and CD206-FITC; *Cocktail2* (for T cell phenotyping): CD3-Pacific Orange; CD4-APC; CD8-FITC; CD45-PerCP, NK40.6-Pacific Blue. All of the antibodies were from BD Pharmingen (San Jose, CA, USA) or from BioLegend (San Diego, CA, USA). Cells were analyzed on a BD upgraded FACS Caliber Flow Cytometer (eight-colors) using FlowJo Software (Tree Star, Inc., Ashland, OR, USA).

Immunohistochemistry

ATs were fixed in 10% buffered formalin overnight then embedded in paraffin and following antigen retrieval were incubated overnight with polyclonal rabbit anti-mouse antibody for CD45 (Abcam, Cambridge, MA, USA; 1:100 dilution) or rat anti-mouse F4/80 antibody (Abcam; 1:100 dilution). Secondary antibody staining was performed using the Vectastain ABC Kit (Vector Laboratories, Burlingame, CA, USA) and detected with 3,3'-diaminobenzidine. Slides were counterstained using Mayer's hematoxylin. Sections incubated with non-immune IgG (Pierce, Rockford, IL, USA) instead of the primary antibody were used as method controls. All pictures were captured with an Olympus microscope using 200 $\times$ magnification. Quantification of the immunohistochemical data was done using a MetaMorph Software ver6.3 (Molecular Devices, Downingtown, PA, USA) with an established arbitrary threshold. Data was normalized to section area and expressed as arbitrary units.

Macrophage polarization

Peritoneal macrophages were isolated by lavage with sterile PBS and cultured in RPMI-1640 media supplemented with 10% FBS, 1% penicillin/streptomycin, 1% glutamax, 1% HEPES, 0.5% non-essential amino acids (NEAA), 0.5% sodium pyruvate, and 50 μ M beta-mercaptoethanol (BME). After 24 h, the adherent cells were treated with either a combination of IFN γ (150 U/ml) and LPS (100 ng/ml) or with IL4 (20 U/ml) and IL13 (15 U/ml) for additional 24 h to induce polarization towards 'classically' (M1) or 'alternatively' (M2) activated macrophage phenotypes respectively. Bone-marrow-derived macrophages (BMDM) were isolated from the femoral and tibial bones and were differentiated in culture for 7 days using RPMI-1640 media supplemented with 10% FBS, 1% penicillin/streptomycin, 1% glutamax, and 10 ng/ml M-CSF. Subsequently, cells were treated for 24 h to induce 'classical' M1 activation (as described above) or 'metabolic' activation (MMe) using a combination of glucose (30 mM), insulin (10 nM), and palmitate (0.4 mM), as described by Kratz *et al.* (2014).

Real-time PCR

RNA from total AT was extracted and reverse transcribed as previously described (Dobrian *et al.* 2010). Real-time PCR was performed using TaqMan probes from Applied Biosystems. β -actin was used to normalize the data. Results were expressed as fold change by the $2^{-\Delta\Delta C_t}$ method using *Apoe*^{-/-} mice as a control group.

En face staining

The aortic trees of mice were collected, fixed *in situ* by perfusion with PBS containing 4% paraformaldehyde, then additionally fixed for 24 h and stained with Sudan IV (Tangirala *et al.* 1995). *En face* stained aortic tissues were analyzed for the percent surface areas occupied by lesions using Photoshop Software.

In vitro chemokine production by aortic cells in response to STAT4-dependent AT secretome

Visceral AT from *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice was excised and minced into ~ 1 mm³ pieces and incubated for 24 h in DMEM supplemented with 1% penicillin/streptomycin. The following day, aortic suspensions were generated by collagenase digestion from *Apoe*^{-/-} mice. The aortic cells were incubated with the AT conditioned media (CM) at $\sim 10^6$ cells/well for 24 h and then cells were washed, collected by centrifugation and processed for RNA extraction and gene expression analysis by real-time PCR.

Statistical analysis

Statistical analysis was performed using GraphPad Prism Software (GraphPad Software, Inc., La Jolla, CA, USA). Student's *t*-test unpaired analysis was used for all data comparisons between the *Stat4*^{-/-}*Apoe*^{-/-} mice and *Apoe*^{-/-} controls. For comparisons of more than two groups, one-way ANOVA was used for data analysis. For experiments including two independent variables (diet and genotype), data was analyzed using two-way ANOVA. Data were expressed as mean $\pm$ S.E.M. and the null hypothesis was rejected for a *P* value <0.05.

Results

STAT4 deficiency does not change body weight, systemic lipids, glucose, nor insulin

Stat4^{-/-}*Apoe*^{-/-} or *Apoe*^{-/-} control mice were kept on either a chow or a Western diet for 12 weeks. Mice on both

chow (not shown) and Western diets (Table 1) had similar body weights. Also, randomly fed plasma glucose, insulin, triglycerides, total, HDL, and LDL cholesterol were similar between the two groups for both the chow-fed and Western-diet-fed mice. The lipid profile in the Western-diet-fed mice were in the expected range with significantly elevated triglycerides and LDL+VLDL plasma cholesterol levels (Table 1).

Effect of STAT4 deficiency on lymphocyte population in visceral and peri-aortic ATs

STAT4 is abundantly expressed in the T and NK cells of mice and humans, where it plays an important role in Th1 polarization and in the cytotoxic response of NK cells following cytokine challenge. Therefore, we compared the lymphocyte composition in the STAT4-deficient and -sufficient *Apoe*^{-/-} mice. Flow cytometry analysis of peri-aortic and visceral ATs of *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice on a Western diet showed significantly increased numbers of CD45+ cells in the peri-aortic AT of both groups of mice (Fig. 1A). While CD45+ cellularity was similar in the visceral fat of *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice, the numbers of CD45+ cells was significantly reduced in the peri-aortic fat of *Stat4*^{-/-}*Apoe*^{-/-} compared to *Apoe*^{-/-} controls (Fig. 1A). Similar results were obtained by AT immunohistochemistry using a CD45 antibody (Fig. 1B). To further determine the composition of the CD45+ lymphocytes, we next analyzed the CD3+ T and NK cell populations in the visceral and peri-aortic ATs of mice on chow or Western diets. In the *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} chow-fed mice,

the relative percentages of CD3+, CD3+CD4+, CD3+CD8+, and NK cells were similar for both fat depots (not shown). In the Western-diet-fed mice, the relative percentages of CD3+ cells in the CD45+ gate were also similar for both groups in both AT depots (Fig. 1C and D). However, significant differences of the relative percentages of CD4+, CD8+, and NK+ cells were found between the *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice in both the visceral and peri-aortic depots. In visceral and peri-aortic fat of *Stat4*^{-/-}*Apoe*^{-/-} mice, we found a predominant population of CD3+CD4+ cells (~20%) compared to the low relative percentages of CD3+CD8+ cells (8.2 and 2.1% in peri-aortic and visceral fat respectively) (Fig. 1C and D). In stark contrast, the *Apoe*^{-/-} mice showed high percentages of CD3+CD8+ cells in both depots (22.7 and 27.8%, in the peri-aortic and visceral fat respectively) compared to only 2.3% CD3+CD4+ cells in peri-aortic fat and 1.9% in visceral fat (Fig. 1C and D). In addition, STAT4 deficiency resulted in significantly lower percentages of NK cells in both the visceral and peri-aortic depots of *Stat4*^{-/-}*Apoe*^{-/-} compared to *Apoe*^{-/-} mice (Fig. 1C and D). Collectively, this data indicates that in response to Western diet feeding, in *Stat4*^{-/-}*Apoe*^{-/-} mice, the increase in the CD3+ cells in AT is biased towards a CD4+ vs a CD8+ population in a depot-independent fashion, while in the *Apoe*^{-/-} mice there is a predominant CD8+ population in both depots. Also, the relative percentage of visceral and peri-aortic AT NK40.6+ cells is dramatically reduced in *Stat4*^{-/-}*Apoe*^{-/-} compared to *Apoe*^{-/-} mice on a Western diet.

Effect of STAT4 deficiency on antigen-presenting cells and macrophage polarization in visceral and peri-aortic AT

The antigen-presenting cells are important contributors to AT inflammation. STAT4 is expressed in activated blood monocytes and dendritic cells, and therefore STAT4 deficiency may impact numbers and phenotype of resident AT APC directly or via indirect changes in adaptive immunity and inflammation. In contrast to C57Bl6 mice on a high fat diet (Lumeng *et al.* 2007a,b), visceral AT of Western-diet-fed *Apoe*^{-/-} mice has lower percentages of CD11b+F4/80+ cells (33.5%) in the CD45+ gate (Fig. 2B). Virtually all of these cells were CD11c+F4/80+ canonical macrophages, characterized by a pro-inflammatory phenotype, while only ~2.5% were CD206+F4/80+ alternatively activated (M2) macrophages. In peri-aortic fat of the same mice, the percentage of CD11b+F4/80+ cells in the CD45+ gate was ~1.86%. Most of the cells had an 'alternatively' activated M2

Table 1 Body weight and plasma glucose, insulin, and lipids in Western diet fed mice. Female *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice were kept on a Western diet for 12 weeks starting at 8–10 weeks of age. Body weights were measured at the time of euthanasia. Plasma glucose, insulin, triglycerides, total cholesterol, LDL cholesterol, and HDL cholesterol were measured in randomly fed mice at the time of euthanasia. Results are from 8 to 10 mice and were expressed as mean ± S.E.M.

	<i>Stat4</i> ^{-/-} <i>Apoe</i> ^{-/-}	<i>Apoe</i> ^{-/-}	P value
Body weight (g)	36.8 ± 2.81	33.1 ± 1.12	>0.05
Glucose (mg/dl)	141 ± 5.9	127.3 ± 9.2	>0.05
Insulin (µg/l)	3.5 ± 0.8	3.9 ± 0.3	>0.05
Triglycerides (mg/dl)	172.3 ± 14.5	178 ± 12.1	>0.05
Total cholesterol (mg/dl)	1468.3 ± 73.7	1346.1 ± 84.3	>0.05
HDL cholesterol (mg/dl)	29.3 ± 7.6	28.4 ± 8.2	>0.05
LDL cholesterol (mg/dl)	1346.2 ± 114	1312.4 ± 130.3	>0.05

Figure 1

Effect of STAT4 deficiency on lymphocyte numbers and relative composition in visceral and peri-aortic adipose tissue (AT). (A) CD45⁺ cells in peri-aortic and visceral ATs were determined by flow cytometry and normalized to tissue weight; *significant compared to visceral AT for the same mice group. Also shown is a representative gating strategy for the forward scattered and CD45⁺ gates in *Apoe*^{-/-} mice. (B) Immunohistochemistry for CD45 in the peri-aortic and visceral fat of *Stat4*^{-/-}*Apoe*^{-/-} and

Apoe^{-/-} mice. Relative percentages of CD3⁺, CD3+CD4⁺, CD3+CD8⁺, and NK40.6⁺ were determined in the CD45⁺ gate in peri-aortic (C) and visceral (D) ATs of mice on a Western diet. Representative FACS plots in visceral fat are also shown. Data is from *n*=5 to 7 mice/group. #, significant compared to visceral AT. A full colour version of this figure is available at <http://dx.doi.org/10.1530/JOE-15-0098>.

macrophage phenotype, while only ~0.6% of the cells were CD11c+F4/80⁺ canonical macrophages (Fig. 2A and B). Compared to *Apoe*^{-/-} mice on the Western diet, the *Stat4*^{-/-}*Apoe*^{-/-} mice had ~1/3 of the percentage of CD11b+F4/80⁺ cells and three- to 3.5-fold higher percentage of CD206+F4/80⁺ M2 macrophages in both the visceral and peri-aortic depots (Fig. 2B). Also, the relative abundance of the F4/80⁺ macrophages was 2.5-fold higher in visceral fat of *Apoe*^{-/-} mice compared to *Stat4*^{-/-}*Apoe*^{-/-} mice, as determined by immunohistochemistry (Fig. 2C). Also, *Stat4*^{-/-}*Apoe*^{-/-} mice had virtually no CD11c+F4/80⁺ canonical macrophages in either of the depots (Fig. 2A and B). Interestingly, *Apoe*^{-/-} mice on a high fat diet showed a significant increase (~6.5-fold) in the percentage of CD11c+F4/80⁻ cells (dendritic cells) compared to *Stat4*^{-/-}*Apoe*^{-/-} mice in the peri-aortic but not visceral fat. Also, *Apoe*^{-/-} and *Stat4*^{-/-}*Apoe*^{-/-} mice on a chow diet have very low percentages of CD11b+F4/80⁺ macrophages (~3.5%) and of

CD11c+F4/80⁻ (dendritic cells) (<1%) of the total CD45⁺ cells in visceral fat and even lower numbers in the peri-aortic fat and no significant differences were found between the groups (not shown). Collectively, this data show that in face of dietary challenge, STAT4 deficiency leads to reduced proportion of pro-inflammatory 'canonical' macrophages and increased percentage of 'alternatively activated' macrophages in AT depots. These changes may result in reduced overall cytokine production and inflammation in *Stat4*^{-/-}*Apoe*^{-/-} mice compared to *Apoe*^{-/-} controls.

To determine if the differences found in the phenotype of AT macrophages upon STAT4 deficiency are due to intrinsic differences in response to cytokines or metabolic cues, we isolated peritoneal and BMDM and induced *in vitro* polarization with various combinations of cytokines or metabolic activators. Peritoneal macrophages isolated from *Apoe*^{-/-} and *Stat4*^{-/-}*Apoe*^{-/-} mice were treated *in vitro* with either LPS+IFN γ or with IL4+IL13 to

Figure 2

Effect of STAT4 deficiency on macrophage abundance, polarization, and inflammatory profile. Relative composition of macrophages in peri-aortic fat (A) and visceral fat (B) of mice on a Western diet ($n=5-7$); representative FACS plots for CD11c+F4/80+ are also shown. (C) Representative micrographs and quantitation of F4/80 staining in visceral adipose tissue of *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice on a Western diet ($n=4$ /group); AU, arbitrary units. (D) M1 and M2 macrophage marker gene expression in peritoneal macrophages were measured by real-time PCR and expressed as fold change compared to unstimulated

macrophages (M0); $n=3$ independent experiments performed in duplicate; *significantly different compared to M1/M2 and M0; #significantly different compared to *Apoe*^{-/-} group. (E) M1 and MMe (metabolically activated) markers were measured in bone marrow-derived macrophages by real-time PCR; results represent mean $\pm$ S.E.M. from $n=4$ independent experiments performed in duplicate; *significant compared to *Apoe*^{-/-} group. Statistical analysis for (D) and (E) was performed using one-way ANOVA. A full colour version of this figure is available at <http://dx.doi.org/10.1530/JOE-15-0098>.

drive M1 or M2 differentiation (Fig. 2D). The unstimulated macrophages showed similar expression of iNOS (M1 marker) and CD206 (M2 marker). Also, macrophages from both groups responded robustly to cytokine challenge and showed significantly increased M1 and M2 markers upon respective stimulations (Fig. 2D). Interestingly, CD206 expression was significantly increased (~ 1.8 -fold) in macrophages of *Stat4*^{-/-}*Apoe*^{-/-} mice compared to *Apoe*^{-/-} controls following *in vitro* treatments (Fig. 2D) paralleling the phenotype of resident AT

macrophages found *in vivo*. BMDM from both mice groups were also generated by *in vitro* culture of bone marrow cells in presence of M-CSF. Recent evidence showed that *in vitro* metabolic challenge induces a distinct macrophage phenotype that resembles the M1 classically activated macrophages but also expresses anti-inflammatory markers such as PPARC. Similar to peritoneal macrophages, unstimulated BMDM from both mice groups had similar expression of all of the tested genes (not shown). Also, BMDM from both mice groups showed robust activation

of several M1 markers such as iNOS and arginase along with pro-inflammatory cytokines such as TNF α and IL1 β (Fig. 2E). In addition, M1 macrophages from *Stat4*^{-/-}*Apoe*^{-/-} mice showed decreased iNOS and IL1b expression compared to *Apoe*^{-/-} mice (Fig. 2E). Interestingly, expression of *Abca1* gene was also significantly increased in M1 macrophages from *Stat4*^{-/-}*Apoe*^{-/-} mice compared to controls, suggesting a potential ability of these macrophages to limit lipid loading by upregulating cellular cholesterol efflux (Fig. 2E). BMDM were also treated with a combination of insulin/glucose/palmitic acid to achieve 'metabolic activation' (MME; Kratz *et al.* 2014). MMEs from *Stat4*^{-/-}*Apoe*^{-/-} mice showed reduced activation of the pro-inflammatory markers iNOS and TNF α compared to *Apoe*^{-/-} controls along with increased expression of the anti-inflammatory genes PPAR γ and the *Abca1* transporter (Fig. 2E). No differences were found in expression of TNF α , arginase, and *Plin2* (Fig. 2E). These findings indicate that macrophages from *Stat4*^{-/-}*Apoe*^{-/-} mice assume a lower pro-inflammatory phenotype compared to *Apoe*^{-/-} mice in response to both M1 and MME polarization.

Role of STAT4 deficiency on AT cytokines and chemokines expression

We further investigated whether the reduced pro-inflammatory phenotype of macrophages from *Stat4*^{-/-}*Apoe*^{-/-} mice found both *in vivo* and *in vitro* is mirrored by a reduction in pro-inflammatory cytokine and chemokine gene expression in ATs. We found that in *Apoe*^{-/-} mice on a Western diet, gene expression of IL12p40 was twofold increased in visceral fat and approximately sixfold increased in the peri-aortic fat compared to *Stat4*^{-/-}*Apoe*^{-/-} mice (Fig. 3A and B). In the peri-aortic fat there was also approximately fourfold increased TNF α gene expression in *Apoe*^{-/-} mice compared to *Stat4*^{-/-}*Apoe*^{-/-} mice. Expression of PPAR γ was reduced approximately twofold in *Apoe*^{-/-} mice in both fat depots compared to *Stat4*^{-/-}*Apoe*^{-/-} mice (Fig. 3A and B). In addition, expression of CCL5, CXCL10, and CX3CL1 were all significantly increased in the peri-aortic fat of the *Apoe*^{-/-} mice compared to *Stat4*^{-/-}*Apoe*^{-/-} mice while expression of CCL2 and CXCL16 was similar between groups (Fig. 3C).

Pro-inflammatory response of aortic cells to STAT4-deficient visceral AT secretome

To determine whether STAT4 may effect aortic wall pro-inflammatory gene expression via the STAT4 dependent

Figure 3

Expression of pro-inflammatory cytokines and chemokines in *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice on a Western diet. Cytokine gene expression was measured in visceral (A) and peri-aortic (B) adipose tissue (AT) using real-time PCR; results are expressed as fold change in gene expression in *Apoe*^{-/-} mice compared to *Stat4*^{-/-}*Apoe*^{-/-} group and represent mean $\pm$ S.E.M. of *n* = 8 mice/group; chemokine expression in peri-aortic AT (C) was also measured by real-time PCR in *n* = 6 mice /group; *significantly different (*P* < 0.05) compared to *Stat4*^{-/-}*Apoe*^{-/-} group.

AT secretome we examined *in vitro* the response of aortic cells from *Apoe*^{-/-} mice to CM generated from visceral AT of *Stat4*^{-/-}*Apoe*^{-/-} or *Apoe*^{-/-} controls. Briefly, we excised and incubated AT from *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice in DMEM, and collected the 24-h CM. The CM was further incubated for 24 h with fresh preparations of aortic cell suspensions from *Apoe*^{-/-} mice (Fig. 4A). Following incubation, aortic cells were washed and collected by centrifugation and gene expression of various macrophage and T cell chemokines and activation markers were performed by real-time PCR (Fig. 4B). Gene expression for CX3CL1, CCL5, TNF α , and IL12p40 was not detectable in *Apoe*^{-/-} aortic cells incubated with either of the CM. However, in cells incubated with *Apoe*^{-/-} CM, mRNA expression of CCL2 was significantly higher compared to cells incubated with *Stat4*^{-/-}*Apoe*^{-/-} CM (Fig. 4B). This result shows that visceral fat secretome may induce chemokine expression in aortic cells, at least in part, via a STAT4-dependent mechanism. Interestingly, we found no significant difference in the expression of the macrophage activation marker *I-A^b* (*MHC-II*) between aortic cells incubated with AT CM from *Stat4*^{-/-}*Apoe*^{-/-} or *Apoe*^{-/-} mice (Fig. 4B). This result suggests that visceral AT secretome can directly induce increased expression of some of the pro-inflammatory genes in cells of the aortic wall and that STAT4 deficiency may limit inflammation.

Figure 4

Stat4^{-/-}*Apoe*^{-/-} visceral adipose tissue secretome decreases gene expression of CCL2 in *Apoe*^{-/-} aortic cells. (A) Experimental schematic demonstrating the approach. See 'Materials and methods' for more detailed description. (B) Real-time PCR analysis of CCL2 and I-A^b in aortic cell suspensions following incubation with visceral AT CM from either *Stat4*^{-/-}*Apoe*^{-/-} or *Apoe*^{-/-} mice. Data is expressed as 2^{-ΔΔCt} normalized to 18s as a housekeeping gene. Results represent mean ± S.E.M. from *n*=4 independent experiments performed in duplicate. A full colour version of this figure is available at <http://dx.doi.org/10.1530/JOE-15-0098>.

STAT4 deficiency reduces aortic plaque burden in mice on a Western diet

In this study we analyzed a small sample of mice (*n*=4 on chow and *n*=8 on a Western diet) to confirm previous data showing reduced atherosclerosis in *Stat4*^{-/-}*Apoe*^{-/-} mice compared to *Apoe*^{-/-} controls (manuscript under revision). Our data shows that compared to *Apoe*^{-/-} controls, the *Stat4*^{-/-}*Apoe*^{-/-} mice on a Western diet have a prominent and significant ~41% reduction in aortic plaque area by *en face* staining (Fig. 5). In age-matched *Stat4*^{-/-}*Apoe*^{-/-} mice on chow diet, the plaque burden is also slightly but not significantly reduced (Fig. 5). There is significant interaction between the diet and the genotype (*P*<0.05), indicating that STAT4 deficiency is particularly protective in the context of high dietary cholesterol challenge.

Discussion

The contribution of AT to systemic inflammation and cardiovascular disease is well documented. In this study we report for the first time the effect of STAT4 deficiency

on peri-vascular and visceral AT inflammation in atherosclerotic *Apoe*^{-/-} mice. STAT4 deficiency also reduces the development of atherosclerosis, suggesting that reduction of peri-vascular and visceral inflammation may have a contributing athero-protective role. We show that STAT4 deficiency reduces pro-inflammatory cytokines and chemokines as well as lymphocyte number and composition and macrophage polarization in peri-vascular fat of mice on a Western diet. In *Apoe*^{-/-} mice, the increased atherosclerotic burden on a Western diet compared to chow is largely due to a dramatic increase in circulating VLDL+LDL cholesterol (Plump *et al.* 1992). We show that protection by STAT4 deficiency is independent of circulating lipids as the STAT4-deficient mice have similarly increased levels of LDL+VLDL cholesterol to *Apoe*^{-/-} controls (Table 1).

We previously showed that STAT4 deficiency improves glucose tolerance and increases insulin sensitivity in a mouse model of obesity and insulin resistance (Dobrian *et al.* 2013). This effect was accompanied by reduced AT inflammation due to more insulin-sensitive and less hypertrophic adipocytes and reduced lymphocyte infiltration in the visceral AT. However, *Apoe*^{-/-} mice are lean and insulin sensitive even when fed a Western diet (Gao *et al.* 2007, Kawashima *et al.* 2009). Therefore, the anti-inflammatory effects of STAT4 in this model cannot be attributed to an improved metabolic profile or to reduced visceral adiposity. Indeed, we did not find any differences in glucose, insulin or body weight between the *Stat4*^{-/-}*Apoe*^{-/-} mice and the *Apoe*^{-/-} controls (Table 1).

Figure 5

STAT4 deficiency reduces atherosclerotic lesion burden in *Apoe*^{-/-} mice on a Western diet. Female *Stat4*^{-/-}*Apoe*^{-/-} and *Apoe*^{-/-} mice were kept on a Western diet or fed regular chow diet for 12 weeks starting at 8–10 weeks of age. (A) Representative *en face* analysis using Sudan IV staining of whole aortas from Western-diet-fed mice. (B) Lesion analysis from *n*=4 mice/group for the chow diet and from *n*=8 mice/group for the Western diet groups. Data represent % stained area of the total aortic tissue. Statistical analysis was performed using two-way ANOVA. A full colour version of this figure is available at <http://dx.doi.org/10.1530/JOE-15-0098>.

In this study we focused our investigation on two of the more relevant fat depots, visceral and peri-aortic, known to exert vasocrine effects in both mice and humans (Lakka *et al.* 2002, Berg & Scherer 2005, Yudkin *et al.* 2005, Lohmann *et al.* 2009, Ouwers *et al.* 2010). While visceral AT contributes to accelerated atherosclerosis in *Apoe*^{-/-} mice, the peri-vascular AT in particular plays a causative role in atherosclerotic plaque formation in this mouse model (Ohman *et al.* 2008, 2011). Our data shows significantly higher numbers of CD45+ lymphocytes in the peri-aortic but not in the visceral fat of *Apoe*^{-/-} compared to *Stat4*^{-/-}*Apoe*^{-/-} mice on a Western diet (Fig. 1). In particular, the number of CD3+CD8+ T lymphocytes was dramatically increased in the peri-aortic fat in *Apoe*^{-/-} mice on a Western diet compared to *Stat4*^{-/-}*Apoe*^{-/-} mice. Previous studies indicated that expansion of CD3+ subsets in response to IL12 stimulation is dependent on STAT4 (Yoo *et al.* 2002). We found that in visceral fat of *Stat4*^{-/-}*Apoe*^{-/-} mice expression of IL12p40 subunit is reduced compared to *Apoe*^{-/-} controls, suggesting lower levels of functional IL12. In this study we also found that expression of a number of potent chemokines for T lymphocytes or monocytes, such as CCL5, CXCL10, and CX3CL1, is reduced in the peri-aortic fat of *Stat4*^{-/-}*Apoe*^{-/-} mice compared to controls (Fig. 3). STAT4 is involved in the regulation of several chemokine receptors that are preferentially expressed in Th1 cells, such as CXCR3 and CCR5 (Kim *et al.* 2006). We have previously shown (Dobrian *et al.* 2013) that the expression of these and other chemokine receptors are selectively decreased in CD8+ but not CD4+ cells of STAT4-deficient mice, compared to WT controls. It is therefore possible that reduced chemokine/cytokine production in peri-aortic and visceral fat combined with reduced migratory capacity of CD8+ cells due to a reduction in the chemokine receptors results in a relative reduction of CD8+ cells recruited in the ATs of *Stat4*^{-/-}*Apoe*^{-/-} mice. Cytotoxic CD8(+) T lymphocytes represent up to 50% of the total numbers of leukocytes found in advanced human plaques and dominate early immune responses in mouse lesions (Gewaltig *et al.* 2008, Kolbus *et al.* 2010). A recent publication showed that depletion of this lymphocyte subset in *Apoe*^{-/-} mice fed a high fat diet ameliorates atherosclerosis (Kyaw *et al.* 2013). Therefore, peri-vascular and possibly visceral fat may constitute a reservoir of CD8+ cells for the aortic wall and reduction of this T cell subset may be part of the mechanism responsible for athero-protection reported in this paper and by our group in preliminary observations in *Stat4*^{-/-}*Apoe*^{-/-} mice (manuscript under revision). An in-depth analysis of the

atherosclerotic phenotype and related vascular mechanisms in *Stat4*^{-/-}*Apoe*^{-/-} mice are under publication elsewhere.

There was also a prominent reduction in the number of NK40.6+ cells in *Stat4*^{-/-}*Apoe*^{-/-} mice on a Western diet compared to *Apoe*^{-/-} controls, in both peri-aortic and visceral fat. Like CD8+ cells, NK cells appear to play an important role in the development of atherosclerosis (Major *et al.* 2004, Kolbus *et al.* 2010). STAT4 is abundantly expressed in NK cells and is key for IFN γ production and increased cytotoxicity. Therefore, reduction in the relative composition of NK cells in visceral and peri-aortic fat may also contribute to reduced vascular wall inflammation in STAT4-deficient mice.

Macrophages represent a major component of visceral AT in rodents (Lumeng *et al.* 2007a,b) but are less abundant in the peri-vascular fat. In *Apoe*^{-/-} mice, the number of macrophages is reportedly very low in various AT depots. We also found that only a minor proportion of up to ~2.2% of the total CD45+ cells in peri-aortic fat of *Apoe*^{-/-} mice is represented by CD11b+F4/80+ macrophages and is similar for *Stat4*^{-/-}*Apoe*^{-/-} mice (Fig. 2). Even in visceral fat, the proportion of CD11b+F4/80+ cells is ~30% in *Apoe*^{-/-} mice and only ~10% in *Stat4*^{-/-}*Apoe*^{-/-} mice. Therefore, in these mice the tropism of macrophages may be more prominent for other tissues, including the aorta. Interestingly, we found that STAT4 deficiency induces a bias towards CD206+ macrophages in both depots *in vivo* and that peritoneal macrophages of *Stat4*^{-/-}*Apoe*^{-/-} mice have a more robust response to 'alternative' activation *in vitro* (Fig. 2). It is known that the CD206+ (alternatively activated) macrophages are a source of the anti-inflammatory IL10 and IL4 cytokines as opposed to the M1 macrophages that are an abundant source of IL12, IL6, IL1 β , and other pro-inflammatory cytokines (Lee *et al.* 2012). We also found that in the peri-aortic fat, gene expression of IL12p40 and TNF α was decreased in *Stat4*^{-/-}*Apoe*^{-/-} mice. The data suggests that STAT4 deficiency drives M2 polarization of AT macrophages without affecting the number of resident cells. This conclusion is also supported by reduced *in vitro* propensity of the STAT4-deficient BMDM to differentiate into a M1 phenotype (Fig. 2) as well as increased PPARC expression in AT (Fig. 3) known to promote M2 polarization (Bouhleh *et al.* 2007, Odegaard *et al.* 2007). Also, virtually the entire population of F4/80+ macrophages was CD11c+ in both visceral and peri-aortic fat of *Apoe*^{-/-} mice, while in *Stat4*^{-/-}*Apoe*^{-/-} mice the CD11c+ macrophages were not detectable. Canonical CD11c+ macrophages are known to have a marked pro-inflammatory phenotype

in both mice and humans (Geissmann *et al.* 2010) and can therefore be a source of circulating cytokines and chemokines impacting on immune cell recruitment and inflammation in the vascular wall. Importantly, several recent studies have described a combined M1/M2 phenotype of macrophages in obesity in mice and humans, suggesting they can adapt a more complex phenotype (Zeyda *et al.* 2007, Riboldi *et al.* 2013). Although mice on the *Apoe*^{-/-} background are lean and normoglycemic, we tested the paradigm that metabolic activation of BMDM from *Stat4*^{-/-}*Apoe*^{-/-} mice may lead to phenotypically different macrophages compared to *Apoe*^{-/-} cells. Indeed, *in vitro* stimulation of BMDM with a combination of insulin/glucose/palmitate resulted in significantly reduced expression of IL1B and iNOS in cells from *Stat4*^{-/-}*Apoe*^{-/-} mice compared to *Apoe*^{-/-} controls. Also, metabolic activation has been shown to up-regulate anti-inflammatory genes such as PPARC and lead to more efficient metabolism due to an increase in *Abca1* and *Plin2* genes (Kratz *et al.* 2014). In *Stat4*^{-/-}*Apoe*^{-/-} macrophages PPARC gene expression was significantly elevated compared to *Apoe*^{-/-}, suggesting a more efficient anti-inflammatory response in the former in response to metabolic activation. This may be relevant especially upon Western diet feeding when circulating lipids are dramatically increased and may lead to a predominant pro-inflammatory macrophage phenotype. Finally, we found that *Abca1* gene expression was significantly increased in MME macrophages of *Stat4*^{-/-}*Apoe*^{-/-} mice compared to *Apoe*^{-/-} controls. This finding suggests that STAT4 deficiency promotes a more efficient cholesterol efflux from macrophages and therefore could contribute to the reduced atheroma formation in *Stat4*^{-/-}*Apoe*^{-/-} mice on a Western diet via a reduction of foam cells.

Many studies demonstrate a link between the inflammation within AT, insulin resistance and accelerated atherosclerosis (Mazzone *et al.* 2008). AT releases inflammatory cytokines and bioactive mediators that are potent inducers of structural and functional changes in the vasculature (Brown *et al.* 2014). Collectively referred to as the AT secretome, these factors may originate from adipocytes or other cells in AT, such as immune cells and vascular cells. We examined the effect of STAT4-sufficient and -deficient AT secretome on aortic gene expression of pro-inflammatory cytokines and chemokines in *Apoe*^{-/-} mice. Interestingly, we found that CCL2 expression in aortic cells of *Apoe*^{-/-} mice following incubation with AT CM of *Stat4*^{-/-}*Apoe*^{-/-} mice is significantly lower compared to CM from *Apoe*^{-/-} mice (Fig. 4). This finding indicates that visceral AT secretome can

support inflammation and atherosclerosis development at least through changes in the aortic expression of CCL2, one of the major chemokines necessary for MF recruitment into the aorta. Although we did not test the effect of peri-aortic fat secretome on aortic cells due to the limited tissue availability, the similarly immune cells composition and phenotype found between visceral and peri-aortic depots suggest comparable responses. Also, although the CM component or combination responsible for the response is unknown, the effect is STAT4-related and further studies are granted to pinpoint the mechanism.

Collectively our data show that STAT4 deficiency in *Apoe*^{-/-} mice reduces immune cell infiltration, pro-inflammatory cytokine and chemokine expression and macrophage polarization in peri-aortic and visceral AT. These effects may contribute to the athero-protective effect of STAT4 deficiency in this model, particularly in mice on a Western diet. Potential mechanisms include reduction of the reservoir of pro-inflammatory cells or pro-inflammatory cytokines and chemokines to the aortic wall or blunted expression of pro-inflammatory genes in cells of the aortic wall. Ultimately, STAT4 deficiency limits vascular inflammation potentially leading to reduced plaque burden in diet-induced accelerated atherosclerosis. Our study suggests that selective inhibitors of the Jak/Tyk/STAT4 pathway may have possible cardiovascular health benefits beyond their current indications as therapeutics for autoimmune and chronic inflammatory diseases.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the PO1 HL55798 to J L N and by the RO1 HL112605 to J L N and A D D.

Acknowledgements

The authors wish to thank Dr Mark Kaplan for providing the STAT4 null mice. The authors also acknowledge the excellent technical support of Lindsey Glenn BS, Jack Lindsey MS, Ashley James BS, and Mr Ciriaco Villafior.

References

- Berg AH & Scherer PE 2005 Adipose tissue, inflammation, and cardiovascular disease. *Circulation Research* **96** 939–949. (doi:10.1161/01.RES.0000163635.62927.34)

- Bouhrel MA, Derudas B, Rigamonti E, Dievart R, Brozek J, Haulon S, Zawadzki C, Jude B, Torpier G, Marx N *et al.* 2007 PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metabolism* **6** 137–143. (doi:10.1016/j.cmet.2007.06.010)
- Brown NK, Zhou Z, Zhang J, Zeng R, Wu J, Eitzman DT, Chen YE & Chang L 2014 Perivascular adipose tissue in vascular function and disease: a review of current research and animal models. *Arteriosclerosis, Thrombosis, and Vascular Biology* **34** 1621–1630. (doi:10.1161/ATVBAHA.114.303029)
- Chang HC, Han L, Goswami R, Nguyen ET, Pelloso D, Robertson MJ & Kaplan MH 2009 Impaired development of human Th1 cells in patients with deficient expression of STAT4. *Blood* **113** 5887–5890. (doi:10.1182/blood-2008-09-179820)
- Cole BK, Keller SR, Wu R, Carter JD, Nadler JL & Nunemaker CS 2010 Valsartan protects pancreatic islets and adipose tissue from the inflammatory and metabolic consequences of a high-fat diet in mice. *Hypertension* **55** 715–721. (doi:10.1161/HYPERTENSIONAHA.109.148049)
- Darnell JE Jr 1997 STATs and gene regulation. *Science* **277** 1630–1635. (doi:10.1126/science.277.5332.1630)
- Davenport P & Tipping PG 2003 The role of interleukin-4 and interleukin-12 in the progression of atherosclerosis in apolipoprotein E-deficient mice. *American Journal of Pathology* **163** 1117–1125. (doi:10.1016/S0002-9440(10)63471-2)
- Dobrian AD, Ma Q, Lindsay JW, Leone KA, Ma K, Coben J, Galkina EV & Nadler JL 2010 Dipeptidyl peptidase IV inhibitor sitagliptin reduces local inflammation in adipose tissue and in pancreatic islets of obese mice. *American Journal of Physiology. Endocrinology and Metabolism* **300** E410–E421. (doi:10.1152/ajpendo.00463.2010)
- Dobrian AD, Galkina EV, Ma Q, Hatcher M, Aye SM, Butcher MJ, Ma K, Haynes BA, Kaplan MH & Nadler JL 2013 STAT4 deficiency reduces obesity-induced insulin resistance and adipose tissue inflammation. *Diabetes* **62** 4109–4121. (doi:10.2337/db12-1275)
- Eid RE, Rao DA, Zhou J, Lo SF, Ranjbaran H, Gallo A, Sokol SI, Pfau S, Pober JS & Tellides G 2009 Interleukin-17 and interferon- γ are produced concomitantly by human coronary artery-infiltrating T cells and act synergistically on vascular smooth muscle cells. *Circulation* **119** 1424–1432. (doi:10.1161/CIRCULATIONAHA.108.827618)
- Gao J, Katagiri H, Ishigaki Y, Yamada T, Ogihara T, Imai J, Uno K, Hasegawa Y, Kanzaki M, Yamamoto TT *et al.* 2007 Involvement of apolipoprotein E in excess fat accumulation and insulin resistance. *Diabetes* **56** 24–33. (doi:10.2337/db06-0144)
- Geissmann F, Gordon S, Hume DA, Mowat AM & Randolph GJ 2010 Unravelling mononuclear phagocyte heterogeneity. *Nature Reviews. Immunology* **10** 453–460. (doi:10.1038/nri2784)
- Gewaltig J, Kummer M, Koella K, Cathomas G & Biedermann BC 2008 Requirements for CD8 T-cell migration into the human arterial wall. *Human Pathology* **39** 1756–1762. (doi:10.1016/j.humpath.2008.04.018)
- Good SR, Thieu VT, Mathur AN, Yu Q, Stritesky GL, Yeh N, O'Malley JT, Perumal NB & Kaplan MH 2009 Temporal induction pattern of STAT4 target genes defines potential for Th1 lineage-specific programming. *Journal of Immunology* **183** 3839–3847. (doi:10.4049/jimmunol.0901411)
- Hauer AD, Uyttenhove C, de Vos P, Stroobant V, Renauld JC, van Berkel TJ, van Snick J & Kuiper J 2005 Blockade of interleukin-12 function by protein vaccination attenuates atherosclerosis. *Circulation* **112** 1054–1062. (doi:10.1161/CIRCULATIONAHA.104.533463)
- Hofmann SM, Perez-Tilve D, Greer TM, Coburn BA, Grant E, Basford JE, Tschop MH & Hui DY 2008 Defective lipid delivery modulates glucose tolerance and metabolic response to diet in apolipoprotein E-deficient mice. *Diabetes* **57** 5–12. (doi:10.2337/db07-0403)
- Horvath CM & Darnell JE 1997 The state of the STATs: recent developments in the study of signal transduction to the nucleus. *Current Opinion in Cell Biology* **9** 233–239. (doi:10.1016/S0955-0674(97)80067-1)
- Huang ZH, Reardon CA & Mazzone T 2006 Endogenous ApoE expression modulates adipocyte triglyceride content and turnover. *Diabetes* **55** 3394–3402. (doi:10.2337/db06-0354)
- Huang ZH, Reardon CA, Subbiah PV, Getz GS & Mazzone T 2013 ApoE derived from adipose tissue does not suppress atherosclerosis or correct hyperlipidemia in apoE knockout mice. *Journal of Lipid Research* **54** 202–213. (doi:10.1194/jlr.M031906)
- Imada K & Leonard WJ 2000 The Jak–STAT pathway. *Molecular Immunology* **37** 1–11. (doi:10.1016/S0161-5890(00)00018-3)
- Kaplan MH 2005 STAT4: a critical regulator of inflammation *in vivo*. *Immunologic Research* **31** 231–242. (doi:10.1385/IR:31:3:231)
- Kawashima Y, Chen J, Sun H, Lann D, Hajjar RJ, Yakar S & Leroith D 2009 Apolipoprotein E deficiency abrogates insulin resistance in a mouse model of type 2 diabetes mellitus. *Diabetologia* **52** 1434–1441. (doi:10.1007/s00125-009-1378-8)
- Kim SH, Gunst KV & Sarvetnick N 2006 STAT4/6-dependent differential regulation of chemokine receptors. *Clinical Immunology* **118** 250–257. (doi:10.1016/j.clim.2003.10.002)
- Kolbus D, Ramos OH, Berg KE, Persson J, Wigren M, Bjorkbacka H, Fredrikson GN & Nilsson J 2010 CD8 $^{+}$ T cell activation predominate early immune responses to hypercholesterolemia in Apoe(–)/(–) mice. *BMC Immunology* **11** 58. (doi:10.1186/1471-2172-11-58)
- Konishi M, Sugiyama S, Sato Y, Oshima S, Sugamura K, Nozaki T, Ohba K, Matsubara J, Sumida H, Nagayoshi Y *et al.* 2010a Pericardial fat inflammation correlates with coronary artery disease. *Atherosclerosis* **213** 649–655. (doi:10.1016/j.atherosclerosis.2010.10.007)
- Konishi M, Sugiyama S, Sugamura K, Nozaki T, Ohba K, Matsubara J, Matsuzawa Y, Sumida H, Nagayoshi Y, Nakaura T *et al.* 2010b Association of pericardial fat accumulation rather than abdominal obesity with coronary atherosclerotic plaque formation in patients with suspected coronary artery disease. *Atherosclerosis* **209** 573–578. (doi:10.1016/j.atherosclerosis.2009.10.008)
- Kratz M, Coats BR, Hisert KB, Hagman D, Mutskov V, Peris E, Schoenfelt KQ, Kuzma JN, Larson I, Billing PS *et al.* 2014 Metabolic dysfunction drives a mechanistically distinct proinflammatory phenotype in adipose tissue macrophages. *Cell Metabolism* **20** 614–625. (doi:10.1016/j.cmet.2014.08.010)
- Kyaw T, Winship A, Tay C, Kanellakis P, Hosseini H, Cao A, Li P, Tipping P, Bobik A & Toh BH 2013 Cytotoxic and proinflammatory CD8 $^{+}$ T lymphocytes promote development of vulnerable atherosclerotic plaques in apoE-deficient mice. *Circulation* **127** 1028–1039. (doi:10.1161/CIRCULATIONAHA.112.001347)
- Lakka HM, Lakka TA, Tuomilehto J & Salonen JT 2002 Abdominal obesity is associated with increased risk of acute coronary events in men. *European Heart Journal* **23** 706–713. (doi:10.1053/euhj.2001.2889)
- Lee MJ, Wu Y & Fried SK 2012 Adipose tissue remodeling in pathophysiology of obesity. *Current Opinion in Clinical Nutrition and Metabolic Care* **13** 371–376. (doi:10.1097/MCO.0b013e32833aabef)
- Leonard WJ & Lin JX 2000 Cytokine receptor signaling pathways. *Journal of Allergy and Clinical Immunology* **105** 877–888. (doi:10.1067/mai.2000.106899)
- Lohmann C, Schafer N, von Lukowicz T, Sokrates Stein MA, Boren J, Rutti S, Wahli W, Donath MY, Luscher TF & Matter CM 2009 Atherosclerotic mice exhibit systemic inflammation in periaortic and visceral adipose tissue, liver, and pancreatic islets. *Atherosclerosis* **207** 360–367. (doi:10.1016/j.atherosclerosis.2009.05.004)
- Lumeng CN, Bodzin JL & Saltiel AR 2007a Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *Journal of Clinical Investigation* **117** 175–184. (doi:10.1172/JCI29881)
- Lumeng CN, Deyoung SM, Bodzin JL & Saltiel AR 2007b Increased inflammatory properties of adipose tissue macrophages recruited during diet-induced obesity. *Diabetes* **56** 16–23. (doi:10.2337/db06-1076)
- Major AS, Wilson MT, McCaleb JL, Ru Su Y, Stanic AK, Joyce S, Van Kaer L, Fazio S & Linton MF 2004 Quantitative and qualitative differences in proatherogenic NKT cells in apolipoprotein E-deficient mice.

- Arteriosclerosis, Thrombosis, and Vascular Biology* **24** 2351–2357. (doi:10.1161/01.ATV.0000147112.84168.87)
- Mazzone T, Chait A & Plutzky J 2008 Cardiovascular disease risk in type 2 diabetes mellitus: insights from mechanistic studies. *Lancet* **371** 1800–1809. (doi:10.1016/S0140-6736(08)60768-0)
- Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V, Mukundan L, Red Eagle A, Vats D, Brombacher F, Ferrante AW *et al.* 2007 Macrophage-specific PPAR γ controls alternative activation and improves insulin resistance. *Nature* **447** 1116–1120. (doi:10.1038/nature05894)
- Ohman MK, Shen Y, Obimba CI, Wright AP, Warnock M, Lawrence DA & Eitzman DT 2008 Visceral adipose tissue inflammation accelerates atherosclerosis in apolipoprotein E-deficient mice. *Circulation* **117** 798–805. (doi:10.1161/CIRCULATIONAHA.107.717595)
- Ohman MK, Luo W, Wang H, Guo C, Abdullah W, Russo HM & Eitzman DT 2011 Perivascular visceral adipose tissue induces atherosclerosis in apolipoprotein E deficient mice. *Atherosclerosis* **219** 33–39. (doi:10.1016/j.atherosclerosis.2011.07.012)
- Ouwens DM, Sell H, Greulich S & Eckel J 2010 The role of epicardial and perivascular adipose tissue in the pathophysiology of cardiovascular disease. *Journal of Cellular and Molecular Medicine* **14** 2223–2234. (doi:10.1111/j.1582-4934.2010.01141.x)
- Pei H, Gu J, Thimmalapura PR, Mison A & Nadler JL 2006 Activation of the 12-lipoxygenase and signal transducer and activator of transcription pathway during neointima formation in a model of the metabolic syndrome. *American Journal of Physiology. Endocrinology and Metabolism* **290** E92–E102. (doi:10.1152/ajpendo.00133.2005)
- Plump AS, Smith JD, Hayek T, Aalto-Setälä K, Walsh A, Verstuyft JG, Rubin EM & Breslow JL 1992 Severe hypercholesterolemia and atherosclerosis in apolipoprotein E-deficient mice created by homologous recombination in ES cells. *Cell* **71** 343–353. (doi:10.1016/0092-8674(92)90362-G)
- Police SB, Thatcher SE, Charnigo R, Daugherty A & Cassis LA 2009 Obesity promotes inflammation in periaortic adipose tissue and angiotensin II-induced abdominal aortic aneurysm formation. *Arteriosclerosis, Thrombosis, and Vascular Biology* **29** 1458–1464. (doi:10.1161/ATVBAHA.109.192658)
- Riboldi E, Porta C, Morlacchi S, Viola A, Mantovani A & Sica A 2013 Hypoxia-mediated regulation of macrophage functions in pathophysiology. *International Immunology* **25** 67–75. (doi:10.1093/intimm/dxs110)
- See R, Abdullah SM, McGuire DK, Khera A, Patel MJ, Lindsey JB, Grundy SM & de Lemos JA 2007 The association of differing measures of overweight and obesity with prevalent atherosclerosis: the Dallas Heart Study. *Journal of the American College of Cardiology* **50** 752–759. (doi:10.1016/j.jacc.2007.04.066)
- Tangirala RK, Rubin EM & Palinski W 1995 Quantitation of atherosclerosis in murine models: correlation between lesions in the aortic origin and in the entire aorta, and differences in the extent of lesions between sexes in LDL receptor-deficient and apolipoprotein E-deficient mice. *Journal of Lipid Research* **36** 2320–2328.
- Watford WT, Hissong BD, Bream JH, Kanno Y, Muul L & O'Shea JJ 2004 Signaling by IL-12 and IL-23 and the immunoregulatory roles of STAT4. *Immunological Reviews* **202** 139–156. (doi:10.1111/j.0105-2896.2004.00211.x)
- Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL & Ferrante AW Jr 2003 Obesity is associated with macrophage accumulation in adipose tissue. *Journal of Clinical Investigation* **112** 1796–1808. (doi:10.1172/JCI200319246)
- Yoo JK, Cho JH, Lee SW & Sung YC 2002 IL-12 provides proliferation and survival signals to murine CD4⁺ T cells through phosphatidylinositol 3-kinase/Akt signaling pathway. *Journal of Immunology* **169** 3637–3643. (doi:10.4049/jimmunol.169.7.3637)
- Yudkin JS, Eringa E & Stehouwer CD 2005 “Vasocrine” signalling from perivascular fat: a mechanism linking insulin resistance to vascular disease. *Lancet* **365** 1817–1820. (doi:10.1016/S0140-6736(05)66585-3)
- Zeyda M, Farmer D, Todoric J, Aszmann O, Speiser M, Gyori G, Zlabinger GJ & Stulnig TM 2007 Human adipose tissue macrophages are of an anti-inflammatory phenotype but capable of excessive pro-inflammatory mediator production. *International Journal of Obesity* **31** 1420–1428. (doi:10.1038/sj.ijo.0803632)
- Zhang X, Niessner A, Nakajima T, Ma-Krupa W, Kopecky SL, Frye RL, Goronzy JJ & Weyand CM 2006 Interleukin 12 induces T-cell recruitment into the atherosclerotic plaque. *Circulation Research* **98** 524–531. (doi:10.1161/01.RES.0000204452.46568.57)
- Zhao L, Cuff CA, Moss E, Wille U, Cyrus T, Klein EA, Pratico D, Rader DJ, Hunter CA, Pure E *et al.* 2002 Selective interleukin-12 synthesis defect in 12/15-lipoxygenase-deficient macrophages associated with reduced atherosclerosis in a mouse model of familial hypercholesterolemia. *Journal of Biological Chemistry* **277** 35350–35356. (doi:10.1074/jbc.M205738200)

Received in final form 25 July 2015

Accepted 5 August 2015