Depot- and sex-specific effects of maternal obesity in offspring's adipose tissue

Simon Lecoutre¹, Barbara Deracinois¹, Christine Laborie¹, Delphine Eberlé¹, Céline Guinez¹, Polina E Panchenko², Jean Lesage¹, Didier Vieau¹, Claudine Junien^{2,3}, Anne Gabory² and Christophe Breton¹

¹Univ. Lille, EA4489, Équipe Malnutrition Maternelle et Programmation des Maladies Métaboliques, F59000 Lille, France

²INRA, UMR1198 Biologie du Développement et Reproduction, F-78350 Jouy-en-Josas, France

³UVSQ, Université Versailles-Saint-Quentin-en-Yvelines, Versailles, France

Correspondence
should be addressed
to C Breton

Email
christophe.breton@univ-lille1.fr

Abstract

According to the Developmental Origin of Health and Disease (DOHaD) concept, alterations of nutrient supply in the fetus or neonate result in long-term programming of individual body weight (BW) setpoint. In particular, maternal obesity, excessive nutrition, and accelerated growth in neonates have been shown to sensitize offspring to obesity. The white adipose tissue may represent a prime target of metabolic programming induced by maternal obesity. In order to unravel the underlying mechanisms, we have developed a rat model of maternal obesity using a high-fat (HF) diet (containing 60% lipids) before and during gestation and lactation. At birth, newborns from obese dams (called HF) were normotrophs. However, HF neonates exhibited a rapid weight gain during lactation, a key period of adipose tissue development in rodents. In males, increased BW at weaning (+30%) persists until 3 months of age. Nine-month-old HF male offspring was normoglycemic but showed mild glucose intolerance, hyperinsulinemia, and hypercorticotesteronemia. Despite no difference in BW and energy intake, HF adult male offspring was predisposed to fat accumulation showing increased visceral (gonadal and perirenal) depot weights and hyperleptinemia. However, only perirenal adipose tissue depot exhibited marked adipocyte hypertrophy and hyperplasia with elevated lipogenic (i.e. sterol-regulated element binding protein 1 (*Srebp1*), fatty acid synthase (*Fas*), and leptin) and diminished adipogenic (i.e. peroxisome proliferator-activated receptor gamma (*Pparγ*), 11β-hydroxysteroid dehydrogenase type 1 (*11β-Hsd1*)) mRNA levels. By contrast, very few metabolic variations were observed in HF female offspring. Thus, maternal obesity and accelerated growth during lactation program offspring for higher adiposity via transcriptional alterations of visceral adipose tissue in a depot- and sex-specific manner.

Key Words

- ▶ high-fat diet
- ▶ visceral adiposity
- ▶ gene expression
- ▶ adipocyte size
- ▶ developmental origin of health and disease
- ▶ dimorphism

Journal of Endocrinology
(2016) **230**, 39–53

Introduction

The rising prevalence of obesity in the world is considered a global epidemic (Popkin *et al.* 2012). Obesity is characterized by accumulation and functional alterations

of white adipose tissue (WAT) predisposing the individuals to increased risk of metabolic pathologies (Sun *et al.* 2011). The expansion of WAT results from hyperplasia (increase

in adipocyte number) and/or hypertrophy (increase in adipocyte size) along with modifications of tissue sensitivity to circulating hormones (Björntorp & Sjöström 1971). Obesity is the result of a complex interaction between genetic and environmental factors (Bouchard 2009). According to the Developmental Origin of Health and Disease (DOHaD) concept also called 'developmental programming' or 'conditioning' (Barker 2004, Hanson & Gluckman 2014), alterations of nutrient supply in the fetus or neonate result in long-term programming of individual body weight (BW) setpoint. Epidemiological studies initially showed that maternal undernutrition leading to fetal growth restriction was associated with higher adiposity in adulthood (Ravelli *et al.* 1999). Clinical studies have also shown that maternal obesity, excessive nutrition, and accelerated growth in neonates sensitize offspring to obesity (Leddy *et al.* 2008).

Thus, WAT may represent a prime target of metabolic programming induced by maternal obesity. Perturbations to the perinatal nutrient supply may affect adipocyte development, leading to persistent alterations in their number and functional properties (Lukaszewski *et al.* 2013, Lecoutre & Breton 2014, 2015). Indeed, in fetuses and neonates, adipocyte stem cells are still plastic and potentially sensitive to maternal factors (Tang & Lane 2012). In humans, the number of adipocyte is set early in life and is a major determinant of fat mass in adulthood (Spalding *et al.* 2008). The timing of adipose tissue development, which differs between species, determines the window of vulnerability to potential adverse environment. In rodents, adipose tissue growth and adipogenesis mainly take place during the last week of gestation and accelerate throughout lactation, whereas in larger mammals, these processes occur before birth. However, there is now convincing evidence that adipogenesis occurs throughout the lifetime (Muhlhauser & Smith 2009).

Little is known about the programming mechanisms that may account for long-lasting perturbation of adipogenesis and WAT metabolism in offspring from obese dams. To unravel the underlying mechanisms, several animal models of maternal obesity have been developed using high-fat (HF) or cafeteria diet applied during the preconception, gestation, and/or lactation periods in dams (Williams *et al.* 2014). These studies confirmed that maternal obesity has common long-term metabolic consequences sensitizing the offspring to metabolic syndrome features. In particular, maternal obesity at conception programs enhanced adipogenesis and lipogenesis from the fetal period to adulthood resulting

in higher WAT mass and larger adipocytes (Muhlhauser & Smith 2009, Borengasser *et al.* 2013, Murabayashi *et al.* 2013). Overfeeding during lactation and/or postweaning periods leads to accelerated growth and consistently worsens adipogenesis and lipogenesis programming (Desai & Ross 2011, Guberman *et al.* 2013, Masuyama & Hiramatsu 2014). Programmed upregulation of the key adipogenic factor peroxisome proliferator-activated receptor gamma (PPAR γ) is one of the characteristic features of fat expansion in the offspring of obese dams (Samuelsson *et al.* 2008, Muhlhauser & Smith 2009, Sen & Simmons 2010, Dahlhoff *et al.* 2014, Desai *et al.* 2015). Obesity-prone offspring rats from obese mothers also exhibited modified fatty acid composition within WAT (Benkalfat *et al.* 2011). However, in rodents, few studies have examined depot- and sex-specific consequences of maternal obesity in offspring's WAT and there is little agreement among them (Sun *et al.* 2012, Ornellas *et al.* 2013, Dahlhoff *et al.* 2014, Masuyama & Hiramatsu 2014).

In this study, we examined whether maternal obesity differently programs adipocyte number and morphology using a model of maternal obesity in rats fed a HF diet before and during gestation and lactation. We also profiled gene expression in two visceral fat depots (gonadal and perirenal WAT) in adult offspring of both sexes. Here, we demonstrate the maternal obesity and accelerated growth during lactation program offspring for higher adiposity via transcriptional alterations of visceral adipose tissue in a depot- and sex-specific manner.

Materials and methods

Animals

Four-week-old virgin female Wistar rats (1 month) were purchased from Charles River Laboratories (L'Arbresle, France) and were housed in individual cages in a humidity-controlled room with a 12h light:12h darkness cycle. Food and water were available *ad libitum*. After 2 weeks of acclimatization on a control (C) diet (3.85 kcal/g with 10% of total calories as fat consisting of soybean oil (5.6%) and lard (4.4%), 70% as carbohydrate, and 20% as protein; D12450J, Research Diets, New Brunswick, NJ, USA), female rats were fed either a HF diet (5.24 kcal/g with 60% of total calories as fat consisting of soybean oil (5.6%) and lard (54.4%), 20% as carbohydrate, and 20% as protein; D12492, Research Diets, New Brunswick, NJ, USA) or a C diet for 16 weeks ($n=12$ per group). After 14 weeks of HF diet, 20-week-old female (5 months) rats were subjected to an oral glucose tolerance test (OGTT). Plasma levels

of insulin, leptin, and corticosterone were also measured after 16 h overnight fasting in both groups. After mating with a male rat fed a C diet, 22-week-old pregnant females were transferred into individual cages with free access to water and continued on their respective diets (C or HF diet) throughout gestation and lactation. Maternal BW was measured weekly until delivery. At parturition, pups were weighed and sexed. Litter size was adjusted to eight pups per dam (four males and four females). During lactation, BWs of dams and pups were assessed on postnatal days (PND) 1, 4, 7, 11, 14, 17, and 21. At weaning (PND21), dams were killed and glycemia as well as plasma levels of leptin insulin and corticosterone were determined after 16 h overnight fasting. To obviate any litter effects, animals used for each experiment were randomly chosen in different litters and only a limited number of animals (one to two males and females) were used from each litter. After weaning, male (M) and female (F) offspring from C or HF dams were housed individually with free access to water and C diet, divided into four groups (CM, CF, HFM, and HFF ($n=16$ per group) and weighed weekly until 9 months of age. Animal use authorization by the French Ministry of Agriculture (No. 04860) has been granted to our laboratory for experimentation with rats. Experiments were conducted in accordance with the principles of laboratory animal care (European Communities Council Directive of 1986, 86/609/EEC).

Food intake and metabolic parameters

Food consumption was recorded weekly from weaning to adulthood until killing the animal in the four groups. Food intake of rats was measured once a day at the beginning of the light phase (09:00 h) by subtracting the uneaten food from the initial amount. Weight-related energy intake is defined as the energy content of the food ingested (kcal) expressed relative to BW (g). Twenty-four-week-old (6 months) offspring was placed in metabolic cage (Bioseb, Vitrolles, France). After an acclimatization period, food intake was recorded for each 24 h period during 1 week to investigate light/dark phase food intake rhythm.

Oral glucose tolerance test

For OGTT, rats were fasted overnight. Basal blood glucose level defined as T0 was determined using a glucometer (Glucotrend 2, Roche Diagnostics) before oral glucose administration (2 g/kg BW). Tail vein blood glucose was then measured at 0, 30, 60, 90, and 120 min after administration.

Endocrine parameters

Plasma hormone levels were evaluated in 30-week-old (7 months) and 36-week-old offspring (9 months) while killing. Blood glucose was determined as described above. Plasma leptin and adiponectin concentrations were measured using murine ELISA kits (Diagnostic Systems Laboratories, Inc. Webster TX, USA and Adipogen Inc, Korea, respectively). Plasma corticosterone levels were determined by a competitive enzyme immunoassay (Immunodiagnostic Systems Ltd, Boldon, UK). Plasma insulin concentrations were measured by ELISA (DRG International, Inc, Springfield Township, NJ, USA). Plasma apelin content was determined by ELISA (Phoenix Pharmaceuticals, Burlingame, CA, USA). The assay sensitivity was 0.07 ng/mL (insulin), 0.04 ng/mL (leptin), 0.1 ng/mL (adiponectin), 0.55 ng/mL (corticosterone), and the intra- and inter-assay coefficients of variation were 4 and 9.1% (insulin), 5.4 and 7.3% (leptin), 4.4 and 6.1% (adiponectin), 4.9 and 7.8% (corticosterone), respectively. Assay kits were used to determine the contents of plasma triglycerides and total cholesterol (61238 Triglyceride Enzymatique PAP100, 61218 Cholesterol Liquide, BioMérieux, France) as well as free cholesterol and free fatty acid (FFA) (references 279-47106 and 999-75406, Wako Chemicals, Neuss, Germany). Each sample was measured in duplicate.

Plasma and tissue collections

Thirty-six-week-old rats (9 months) were rapidly weighed and killed by decapitation between 09:00 and 10:00 h after 16 h overnight fasting. Trunk blood samples were collected into prechilled tubes containing EDTA (20 μ L of a 5% solution) and centrifuged at 4000g for 10 min at 4°C. Plasma was stored at -20°C. Several tissues (brown adipose tissue, liver, heart, kidney, and adrenal gland) as well as gonadal (GWAT) and perirenal (PWAT) fat pads were weighed, frozen in liquid nitrogen and stored at -80°C. For histology experiments, animals were fixed by intracardiac perfusion using buffered 4% paraformaldehyde solution.

Gene expression analysis

GWAT and PWAT gene expression levels were determined in the four groups using RT-quantitative PCR as previously validated (Lukaszewski *et al.* 2011). Briefly, total RNA was extracted and purified using RNeasy lipid tissue kit (Qiagen) according to the manufacturer's

recommendations. The yield of total RNA was quantified on a Multiskan GO Microplate Spectrophotometer (Thermo Scientific). The quality of total RNA was assessed by determining the 260/280 and the 260/230 absorbance ratio and by agarose gel electrophoresis. First-strand cDNAs were synthesized using ThermoScript RT Kit (Invitrogen and Life Technologies). Relative expression levels of RNA per sample were quantified by SYBR Green assay on a Roche Light Cycler 480 sequence detection assay (Roche). Primer sequences are presented in Table 1. For each transcript, PCR was performed in duplicate with 10 µL final reaction volumes with 1 µL of cDNA, 8 µL of QuantiTect SYBR Green Master mix (Qiagen), and 0.5 µL of each primer set (Table 1). PCR was conducted using the following cycle parameters:

10 min at 95°C, and 40 three-step cycles of 15 s at 95°C, 20 s at 60°C, and 30 s at 72°C. A pool of cDNA from control tissues was used as a standard for quantitative correction. All cDNA samples were applied in dilution of 1:10 to obtain results within the range of the standard. Analysis of transcript level was carried out using the determination of the threshold cycle Ct for each reaction corrected by the efficiency. The level of gene expression was normalized to the reference gene transcript cyclophilin A RNA.

Morphometric analysis of adipose tissue

Fat pad mass as well as cell-size distributions were measured. GWAT and PWAT from the four groups

Table 1 Primers used for RT-quantitative PCR analysis.

Target	Sequence	NCBI detected transcript no.	Length of amplicon (bp)
Leptin			
Forward	5'-GTTCTGTGGCTTTGGTCCT-3'	NM_013076	99
Reverse	5'-CTGGTGACMTGGTCTTGATGA-3'		
Ob-Rb			
Forward	5'-GCATGCAGAATCAGTGATATTGG-3'	NM_012596	81
Reverse	5'-CAAGCTGTATCGACACTGATTTCTTC-3'		
Adiponectin			
Forward	5'-CAGGAGATGCTGGAATGACA-3'	NM_144744	114
Reverse	5'-CTACGCTGAATGCTGAGTGAT-3'		
Fas			
Forward	5'-GGACATGGTCACAGACGATGAC-3'	NM_017332	94
Reverse	5'-GTCGAACTTGGACAGATCCTTCA-3'		
C/Ebpa			
Forward	5'-AGTTGACCAGTGACAATGACCG-3'	NM_012524	94
Reverse	5'-TCAGGCAGCTGGCGGAAGAT-3'		
Pparγ			
Forward	5'-TTGGCCATATTTATAGCTGTCATTATT-3'	NM_001138838	76
Reverse	5'-TGCTCTCGATGGGCTTCA-3'		
Srebp1			
Forward	5'-CATCGACTACATCCGCTTCTTACA-3'	NC_005109	97
Reverse	5'-GTCTTTCAGTGATTGCTTTGTGA-3'		
Gr			
Forward	5'-GTGAAATGGGCAAAGGCGATAC-3'	NM_012576	98
Reverse	5'-GCAAATGCCATGAGAAACATCC-3'		
Mr			
Forward	5'-CTGTGCCCCGCTTTGATGGTAG-3'	NM_013131	100
Reverse	5'-GGGATGCTGGCTTCAGGGTAAT-3'		
11β-Hsd1			
Forward	5'-GGACTGGACATGCTCATTCTCA-3'	NM_017080	84
Reverse	5'-GCTTCTTCGCACAGAGTGGATA-3'		
11β-Hsd2			
Forward	5'-ATGTGAACCTCTGGGAGAAGCG-3'	NM_012524	100
Reverse	5'-CCATGCAAGTGCTCAATGTAGTC-3'		
Housekeeping gene Ppia (Cyclophilin A)			
Forward	5'-ATTCATGTGCCAGGGTGGT-3'	NM_017101	115
Reverse	5'-GATGCCAGGACCTGTATGCT-3'		

See text for definitions. Accession numbers, primers sequences, and size of PCR products are indicated. Accession numbers correspond to the mRNA sequences.

($n=6$ per group) were postfixed for 24h in 4% paraformaldehyde in phosphate buffer saline and embedded in paraffin. Fixed tissues were then cut into serial 10 μm sections, mounted on gelatin-coated slides and stained with Groat's hematoxylin and phloxin (2%), according to standard laboratory protocols. Sections were examined using light microscopy (Leica DM IRE2) and photomicrographs were captured at $\times 20$ magnification. The surface of adipocytes was evaluated in ten randomly selected fields of vision for a total of at least 250 adipocytes using ImageJ software (NIH). Total cell number is a direct measure reflecting hyperplasia. The number of cells was estimated using the formula as described previously (Lemonnier 1972).

Statistical analysis

All data are expressed as mean $\pm$ S.E.M. Statistical analysis was carried out using GraphPad Prism5 (GraphPad). A direct comparison between a pair of groups was made using an unpaired Student's *t*-test or a two-way ANOVA for repeated measures followed by a Bonferroni *post hoc* test, where appropriate. A *P* value of <0.05 was considered to be statistically significant.

Results

Effects of HF diet on maternal parameters

HF-fed female rats gained more weight than C females (ANOVA $P<0.0001$) (Fig. 1A). After 14 weeks of HF diet, females had about two-fold higher plasma leptin concentration (4.3 ± 0.6 vs 2.2 ± 0.4 ng/mL, $P<0.05$) compared with C females (Table 2). No difference was observed in basal glycemia, plasma insulin, and corticosterone levels. However, HF-fed females showed a more pronounced increase in glucose levels during OGTT with a higher area under the curve (AUC) (Fig. 1B), reflecting impaired glucose tolerance (ANOVA, $P<0.0001$). HF-fed dams displayed a 20% increase in BW at the end of the gestation (Fig. 1A). At weaning, HF-fed dams exhibited about three-fold higher plasma leptin (2.5 ± 0.4 vs 0.9 ± 0.2 ng/mL, $P<0.05$) and corticosterone levels (796.2 ± 187 vs 278.1 ± 86.8 ng/mL, $P<0.05$) (Table 2), while showing a marked increase in all fat pads weights (data not shown). No variation was observed in glycemia or plasma insulin levels.

Figure 1

Maternal phenotype. BW of dams fed either a C or a HF diet during preconception, gestation, and lactation (A). Concentration of blood glucose during OGTT after 14 weeks of diet and area under the curve (B) ($n=12$ per group). *Effect of maternal HF diet vs maternal C diet (** $P<0.01$; *** $P<0.001$).

Effects of maternal obesity on offspring growth during lactation

Maternal obesity did not impact the litter size (C: 10 ± 2 pups vs HF: 9 ± 3 pups) or the birth weight of offspring (C: 6.2 ± 0.1 g vs HF: 6.3 ± 0.1 g). However, both sexes of HF neonates exhibited rapid weight gain during lactation (ANOVA, $P<0.0001$) (Fig. 2). *Post hoc* analysis revealed difference in BW from PND10 in male offspring (Fig. 2A) and from PND17 in female offspring (Fig. 2B). At weaning, HF male offspring exhibited a 30% increase, whereas HF female offspring only showed a 10% increase in BW (Fig. 2A and B).

Effects of maternal obesity on offspring growth and energy intake from weaning to adulthood

Weaned offspring was fed a C diet until 9 months of age. The BW difference between C and HF male rats persisted until 12 weeks of age (Fig. 3A). Among females, BW equalized between C and HF rats as early as 1 week after weaning (Fig. 3B). In both sexes, HF rats exhibited similar weight-normalized energy intake during adulthood, suggesting that they were not hyperphagic

Table 2 Plasma parameters of dams fed either a control (C) or a high-fat (HF) diet.

Variable	Dams after 14 weeks of diet		Diet effect <i>P</i> -value	Dams at weaning		Diet effect <i>P</i> -value
	C	HF		C	HF	
Glucose (dg/mL)	95.1 ± 4.7	96 ± 2.2	NS	107 ± 8.7	114.3 ± 21	NS
Insulin (μU/mL)	8.6 ± 0.4	9.3 ± 1.3	NS	10.4 ± 1	12.4 ± 0.6	NS
Leptin (ng/mL)	2.2 ± 0.4	4.3 ± 0.6	<0.05	0.9 ± 0.2	2.5 ± 0.4	<0.05
Corticosterone (ng/mL)	756.4 ± 206.6	1143 ± 317.6	NS	278.1 ± 86.8	796.2 ± 187.7	<0.05

Values are means ± s.e.m. (*n* = 12 females per group).

(Fig. 3C and D). However, HF 4-week-old female offspring exhibited significantly increased weight-normalized energy intake as compared with C rats (Fig. 3D). This transient postweaning hyperphagia may reflect female-specific regulations of food intake from milk to solid diet. HF 6-month-old rats placed in metabolic cages exhibited a lower food intake during the dark phase and a higher food intake during the light phase, resulting in modified light/dark phase food intake rhythm (data not shown).

Effects of maternal obesity on adult offspring metabolic parameters

At 7 months of age, C and HF offspring had comparable fasting blood glucose concentrations. During OGTT, HF males displayed increased glucose levels at 30 min ($P < 0.01$) (Fig. 4A) with a trend toward higher AUC ($P = 0.09$) (Fig. 4B), reflecting mild glucose intolerance. No difference was observed in HF female rats (Fig. 4C and D). In HF male rats, serum corticosterone concentration was about two-fold higher than that in C animals after fasting (188.7 ± 27.94 vs 90.42 ± 14.18 ng/mL, $P < 0.01$) and feeding (57.4 ± 9.4 vs 23 ± 3.4 ng/mL, $P < 0.05$) conditions (Fig. 5A). No difference was observed in HF female rats (Fig. 5B).

Unlike females, HF 9-month-old male rats displayed increased PWAT (45.8 ± 2.4 vs 39.2 ± 2.6 mg/g BW, $P < 0.05$) and GWAT weights, when normalized to

BW (35 ± 1.6 vs 30 ± 1.7 mg/g BW, $P < 0.05$) compared with C rats (Table 3). These findings were consistent with an increase in plasma leptin levels (Table 4). HF females exhibited a decrease in interscapular brown fat pad weight, but this was not observed in HF males (1.54 ± 0.16 vs 1.97 ± 0.1 mg/g BW, $P < 0.05$) (Table 3).

At 9 months of age, HF male rats had about 1.5-fold higher plasma insulin (43.63 ± 3.79 vs 33.47 ± 3.23 μU/mL, $P < 0.05$) and leptin concentrations (15.06 ± 1.25 vs 10.15 ± 1.11 ng/mL, $P < 0.05$) compared with C rats, whereas no difference was observed in HF females (Table 4). The increased Homeostasis Model Assessment-insulin resistance (HOMA-IR) index (+38.5%, $P < 0.05$) suggests that HF male rats had decreased insulin sensitivity compared with C rats.

Effects of maternal obesity on adult offspring adipose tissue morphometric parameters

As shown in representative photographs (Fig. 6A), PWAT of HF male offspring exhibited an increase in average adipocyte area (Fig. 6B) and total cell number (Fig. 6C) compared with C rats. Adipocytes measuring 7500 μm² or less represented 65% of all adipocytes in C male offspring, whereas they represented only 40% of all adipocytes in HF male offspring. This indicates that maternal HF diet decreased the frequency of small-sized adipocytes (Fig. 6D). In particular, the proportion of adipocytes measuring 2500 μm² or less displayed a marked sixfold decrease in HF male offspring. In addition, these animals showed greater percentage of large-sized adipocytes (7500–40,000 μm²) (60% vs 35%) compared with C rats. Although no changes in average adipocyte area (Fig. 7A and B) and total cell number (Fig. 7C) were observed in GWAT of HF males, a marked five-fold reduction in frequency of 0–2500 size adipocytes similarly occurred (Fig. 7D). By contrast, both fat pads of HF female offspring (Figs 6 and 7) showed no major changes.

Figure 2

Rat offspring growth curves from birth to weaning. BW of male (A) and female (B) from C or HF dams (*n* = 16 per group). *Effect of maternal HF diet vs maternal C diet (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).

Figure 3

Rat offspring growth curves and food intake from weaning to adulthood. BWs of male (A) and female (B) from C or HF dams. Weekly weight-related energy intake of male (C) and female (D) from C or HF dams ($n=16$ per group). Weight-related energy intake is defined as the energy content of the food ingested (kcal) expressed relative to BW (g). *Effect of maternal HF diet vs maternal C diet ($*P<0.05$; $**P<0.01$; $***P<0.001$).

Effects of maternal obesity on adult offspring adipose tissue gene expression profile

Maternal obesity led to pronounced changes in PWAT gene expression in HF adult male offspring. RT-quantitative PCR data showed that leptin mRNA content was increased (+1.5-fold) in HF male rats compared with C animals (Fig. 8A). This is in agreement with the increased serum leptin levels (Table 4). In PWAT, maternal obesity resulted in increased mRNA levels for genes promoting *de novo* lipogenesis such as fatty acid synthase (*Fas*, +1.6-fold) and sterol regulatory element-binding protein-1 (*Srebp1*, +1.7-fold) in HF male offspring. In addition, mRNA

expression levels of genes involved in adipogenesis such as *Ppar γ* and 11 β -hydroxysteroid dehydrogenase type 1 (*11 β -Hsd1*) were decreased (–1.3- and –1.5-fold, respectively) in HF male offspring vs C rats (Fig. 8A). By contrast, maternal obesity did not affect GWAT mRNA expression profiles to the same extent in HF male rats, except for a downregulation of Ob-Rb (Fig. 9A). Compared with HF males, less change in gene expression profile was observed in HF female adult offspring in both fat pads. HF female offspring still showed increased CCAAT/enhancer binding protein α (*C/Ep α*) mRNA expression levels (+2-fold) in PWAT (Fig. 8B) and lower adiponectin

Figure 4

Oral glucose tolerance test in 7-month-old adult offspring. Time course of plasma glucose in male (A) and female (C) from C or HF dams ($n=16$ per group). The respective AUCs are shown (B, D). *Effect of maternal HF diet vs maternal C diet ($**P<0.01$).

Figure 5

Plasma corticosterone concentrations in fasted and fed conditions in 7-month-old adult offspring. Plasma levels in male (A) and female (B) from C or HF dams ($n=16$ per group). Effect of maternal HF diet vs maternal C diet (** $P<0.001$). #Effect of fasted condition vs fed condition (* $P<0.001$).

(-1.4 -fold) and glucocorticoid receptor (*Gr*) (-1.4 -fold) mRNA expression levels in GWAT (Fig. 9B). This was consistent with differences in fat depot weights and plasma leptin concentrations observed in HF male vs female offspring (Tables 3 and 4).

Discussion

The main finding of this study is that maternal obesity has long-lasting consequences on visceral WAT of adult rat offspring in a depot- and sex-specific manner. In particular, we showed that HF adult male offspring exhibit greater visceral fat pad weights with adipocyte hypertrophy and hyperplasia, despite no difference in BW and energy intake. Our findings disagree with other studies (Kirk *et al.* 2009, Nivoit *et al.* 2009, Desai *et al.* 2014), reporting that maternal obesity prior and throughout pregnancy and lactation programs hyperphagia and marked increased BW in adult rat offspring. The discrepancy between our results and those of others may reflect differences in fetal and/or postnatal programming. These differences may depend on the duration of maternal HF feeding, the dietary lipid content (percentage of lipids vs carbohydrates), the lipid composition (saturated vs unsaturated), and the palatability (presence of sweetened condensed milk) of the diet and therefore the severity of maternal obesity (i.e. weight gain, hormonal status, inflammation grade, and so on). It may also depend on the difference of genetic background of rat strain used (Wistar vs Sprague–Dawley) (Zambrano *et al.* 2010).

In our model, HF adult offspring might have developed a modification of WAT's sensitivity to circulating hormones. Indeed, perirenal WAT depot exhibited changes in lipogenic and adipogenic pathways that may favor triglyceride storage in mature adipocytes. Consistent with increased fat

Table 3 Body composition of 9-month-old offspring from control (C) or obese (HF) dams.

Variable	Male offspring		Female offspring		Diet effect P-value	Effect of sex P	Effect of maternal diet P	Interaction P
	C	HF	C	HF				
Body weight (g)	563.7 ± 10.4	581.3 ± 9.9	291.06 ± 6.38	282.89 ± 8.34	NS	<0.0001	0.65	0.22
Perirenal adipose tissue (mg/g BW)	39.2 ± 2.6	45.8 ± 2.4	38 ± 4.72	37.94 ± 2.04	NS	0.16	0.14	0.12
Gonadal adipose tissue (mg/g BW)	30 ± 1.7	35 ± 1.6	28.88 ± 3.43	26.99 ± 2.15	NS	0.03	0.17	0.03
Brown fat pad (mg/g BW)	1.2 ± 0.05	1.2 ± 0.08	1.97 ± 0.1	1.54 ± 0.16	NS	<0.0001	0.02	0.02
Liver (mg/g BW)	22.43 ± 0.4	23.37 ± 0.4	23.43 ± 0.64	23.11 ± 0.48	NS	0.57	0.66	0.28
Heart (mg/g BW)	2.38 ± 0.04	2.28 ± 0.04	2.75 ± 0.19	3.16 ± 0.11	NS	<0.0001	0.15	0.02
Kidney (mg/g BW)	4.8 ± 0.2	4.9 ± 0.09	6.16 ± 0.143	6.17 ± 0.13	NS	<0.0001	0.35	0.39
Adrenal (mg/g BW)	0.1 ± 0.01	0.1 ± 0.01	0.23 ± 0.013	0.19 ± 0.02	NS	<0.0001	0.03	0.1

Values are mean ± S.E.M. ($n=16$ rats per group).

Table 4 Plasma parameters of 9-month-old offspring from control (C) or obese (HF) dams.

Variable	Male offspring		Diet effect P-value	Female offspring		Diet effect P-value	Effect of sex P	Effect of maternal diet P	Interaction P
	C	HF		C	HF				
Glucose (dg/mL)	105.7 ± 2.76	104 ± 1.63	NS	99 ± 3.7	87.89 ± 2.84	NS	0.0003	0.04	0.12
NEFA (mM)	0.63 ± 0.03	0.62 ± 0.03	NS	0.72 ± 0.04	0.74 ± 0.09	NS	0.02	0.77	0.74
Triacylglycerol (mg/dL)	198.9 ± 16.94	216.7 ± 16.08	NS	91.77 ± 10.94	76.01 ± 18.41	NS	<0.0001	0.95	0.73
Total cholesterol (mM)	0.99 ± 0.04	1.01 ± 0.04	NS	0.66 ± 0.04	0.59 ± 0.04	NS	<0.0001	0.55	0.29
Insulin (µU/mL)	33.47 ± 3.2	43.63 ± 3.79	<0.05	10.79 ± 2.07	10.5 ± 1.86	NS	<0.0001	0.16	0.13
Leptin (ng/mL)	10.15 ± 1.11	15.06 ± 1.25	<0.01	3.27 ± 0.51	5.76 ± 1.35	NS	<0.0001	0.002	0.3
Adiponectin (µg/mL)	13.73 ± 1.1	14.5 ± 1.12	NS	22.12 ± 1.56	21.93 ± 3.93	NS	<0.0001	0.85	0.76
Apelin (ng/mL)	7.1 ± 0.86	6.21 ± 1.08	NS	3.99 ± 0.25	2.97 ± 0.2	NS	0.001	0.29	0.93

Values are mean ± s.e.m. (n = 16 rats per group).

deposition, HF adult male offspring displayed higher serum leptin concentration. Hyperleptinemia may be interpreted as a leptin-resistant state (Kirk *et al.* 2009, Sun *et al.* 2012). First, leptin is known to activate adipogenesis by promoting preadipocyte differentiation (Bol *et al.* 2008, Guo *et al.* 2009). However, despite hyperleptinemia, the number of small adipocytes was markedly decreased in both depots of HF adult male offspring. In agreement with these findings, we observed that *Pparγ* gene expression was downregulated. Our findings disagree with other studies that reported an upregulation of *Pparγ* contents in WAT along with enhanced adipogenesis in obese-prone offspring from malnourished dams (Samuelsson *et al.* 2008, Muhlhausler & Smith 2009, Sen & Simmons 2010, Desai *et al.* 2015). This discrepancy may be due to differences in the establishment of epigenetic marks during adipogenesis and/or hormonal environment, tissue sensitivity, as well as inflammatory status in adipose tissue of adult offspring (Breton 2013). The decrease in gene expression might be seen as an adaptive mechanism to limit fat accumulation (Lukaszewski *et al.* 2011). Indeed, an increase in the lipogenic capacity of adipose tissue is expected during the 'dynamic phase of obesity', when fat stores are rapidly expanding. However, during long-lasting and stable obesity, the decreased expression of lipogenic genes may prevent a further development of fat mass (Ortega *et al.* 2010). Further experiments on the kinetic of fat deposition and the transcriptional profile of lipogenic genes during the development of WAT are needed to address this question. In agreement with this hypothesis, several studies described a relationship between obesity and lower expression and/or activity of *Pparγ* in visceral WAT. These modifications appear to be strongly associated with the pathogenesis of metabolic syndrome (Zhang *et al.* 1996, Fujiki *et al.* 2009). Second, given the antilipogenic leptin action on mature adipocytes (Huan *et al.* 2003, Jiang *et al.* 2009), HF adult male offspring had a trend toward reduced leptin receptor contents and did not show any suppression of *Fas* gene expression, but instead a markedly increased expression. In accordance with these findings, we reported that hyperleptinemic rat offspring from undernourished dams exhibited impaired leptin sensitivity with reduced pSTAT3 in WAT (Lukaszewski *et al.* 2011).

Moreover, HF adult male offspring exhibited hyperinsulinemia with elevated *Srebp1* and *Fas* mRNA levels in PWAT, two genes that are known to be upregulated by insulin levels. This suggests that adipose tissue remains sensitive to insulin. This is in agreement with a greater insulin-induced AKT phosphorylation and the upregulation of lipogenic pathways observed in WAT of HF offspring (Borengasser *et al.* 2013). HF adult

Figure 6

Morphometric analysis of perirenal white adipose tissue in 9-month-old offspring. Representative photomicrographs of paraffin-embedded sections (scale bars=100 μm) (A), average area (B), total cell number (C), and percentage of adipocytes in a given size range (area in μm²) (D) in male and female offspring from C or HF dams ($n=6$ per group). *Effect of maternal HF diet vs maternal C diet ($*P<0.05$; $**P<0.01$). #Effect of male vs female ($#P<0.05$; $##P<0.01$).

male offspring also displayed hypercortisolemia. Several lines of evidence prompted us to explore whether sensitivity of WAT to glucocorticoid (GC) was modified in offspring. First, GC alone or in interaction with insulin regulates the differentiation of preadipocytes and

lipogenic genes (Campbell *et al.* 2011). Second, a close link between chronic excess of plasma GC levels and/or increased GC sensitivity within WAT (i.e. modifications of GR, MR, 11β-hydroxysteroid dehydrogenase type 1 (11β-HSD1), and 11β-hydroxysteroid dehydrogenase

Figure 7

Morphometric analysis of gonadal white adipose tissue in 9-month-old offspring. Representative photomicrographs of paraffin-embedded sections (scale bars=100 μm) (A), average area (B), total cell number (C), and percentage of adipocytes in a given size range (area in μm²) (D) in male and female offspring from C or HF dams ($n=6$ per group). *Effect of maternal HF diet vs maternal C diet ($*P<0.05$; $**P<0.01$). #Effect of male vs female ($#P<0.05$; $##P<0.01$).

Figure 8

mRNA expression levels of perirenal white adipose tissue in 9-month-old offspring. Gene expression was measured in male (A) and female (B) by quantitative real-time PCR, corrected for the mean expression of reference gene (cyclophilin A) ($n=10$ per group). Gene symbols are detailed in Table 1. *Effect of maternal HF diet vs maternal C diet ($*P<0.05$; $**P<0.01$).

type 2 (11β-HSD2) contents) and fat expansion has been observed in offspring from malnourished dams (Gnanalingham *et al.* 2005, Lukaszewski *et al.* 2011, Guo *et al.* 2013). Third, increased expression of GR and

11β-HSD1 in visceral adipose tissue has been associated with the development of obesity in rats overfed during lactation (Boullu-Ciocca *et al.* 2008). We observed a depot-specific downregulation of 11β-Hsd1 mRNA in

Figure 9

mRNA expression levels of gonadal white adipose tissue in 9-month-old offspring. Gene expression was measured in male (A) and female (B) by quantitative real-time PCR, corrected for the mean expression of reference gene (cyclophilin A) ($n=10$ per group). Gene symbols are detailed in Table 1. *Effect of maternal HF diet vs maternal C diet ($*P<0.05$).

PWAT in HF adult offspring as previously reported in 3-month-old offspring of obese mice (Samuelsson *et al.* 2008). We also showed that the ratio between *11 β -Hsd1* and *11 β -Hsd2* expression that controls local balance between active and inactive GC metabolites (Lee *et al.* 2014) was decreased. As described in obesity-prone progeny from undernourished dams, it may diminish intratissular GC responsiveness and represent an adaptive mechanism to counteract excess fat storage (Lukaszewski *et al.* 2011). We cannot exclude that HF offspring may have decreased energy expenditure. Indeed, additional programming mechanisms such as elevated FFA transport and/or lower lipolysis/ β -oxydation activities within WAT might account for increased triglyceride storage (Dahlhoff *et al.* 2014).

At birth, HF offspring had a normal birth weight and, then, exhibited a rapid weight gain during lactation, a key period of adipose tissue development. Adipocyte stem cells are also very sensitive to maternal factors during this developmental period (Tang & Lane 2012). Adipocyte number expansion that is set earlier in obese individual may be a major determinant for increased fat mass in adulthood (Spalding *et al.* 2008). Obesity may arise from increased lipid storage in mature adipocytes during the perinatal period. In line with these findings, we showed that maternal obesity predisposes adult offspring to adiposity by increasing the number of adipocytes and the average fat cell volume. The accelerated postnatal growth in offspring is frequently associated with persisting adiposity throughout life. Several models have shed light on the importance of energy intake and milk composition during the lactation period for adipose tissue programming. Indeed, pups from mothers exposed to HF diet only during lactation (Sun *et al.* 2012, Desai *et al.* 2014, White & Tchoukalova 2014) and neonates reared in small litters, representing a model of postnatal overfeeding (Boullu-Ciocca *et al.* 2008), also displayed persistent hypertrophic adipocytes with enhanced adipogenic and lipogenic mRNA expression levels. However, maternal obesity before conception and gestation is also able to program similar outcomes during the embryonic period. Indeed, despite normal fetal weight, fetus from mice fed a HF diet prior and throughout pregnancy displayed larger adipocytes (Murabayashi *et al.* 2013, Umekawa *et al.* 2015) and increased mRNA expression levels of *Zfp423*, a key transcriptional factor initiating adipogenic commitment (Yang *et al.* 2013). Adult mouse offspring also exhibited increased mRNAs levels of several genes involved in *de novo* lipogenesis and lipid droplet size in visceral

WAT (Dahlhoff *et al.* 2014). Similarly, obesity-prone rat offspring from obese dams that were induced intragastric HF diet feeding displayed an increase in adipogenic and lipogenic pathways (Shankar *et al.* 2008).

Finally, we showed that maternal obesity sensitizes adult rat offspring to increased visceral adiposity in a depot- and sex-specific manner. Indeed, among GWAT and PWAT, only the latter shows marked programming features in HF male offspring. By contrast, very few variations were observed in WAT of HF female offspring. In line with these findings, studies have previously demonstrated the heterogeneity of the adipose lineage. All adipogenic stem cells and adipocytes do not behave equally during adipogenesis. Indeed, each fat depot has a unique developmental gene expression signature (Yamamoto *et al.* 2010). Fat stem cells are influenced by the anatomic location of the depot and/or the hormonal microenvironment, as well as aging, gender, and metabolic health (Williams *et al.* 2014). Thus, intrinsic genetic depot-specific differences in adipose stem cells result in different adipogenic potential, gene expression profile, growth rate, and biological properties (i.e. hormone sensitivity) between visceral and subcutaneous fat pads, but also between each specific visceral fat pad. The fact that different adipocyte precursors might determine the development and the function of specific fat pads led to the notion that each WAT depot could be considered a separate mini-organ (Berry *et al.* 2013).

Among programming mechanisms, inappropriate hormone levels during the perinatal period, are a key factor leading to persistent deregulation of energy homeostasis in progeny. It may result in long-term fat expansion with permanent changes in plasma hormone levels in adult offspring (Breton 2013). Consistent with this notion, maternal obesity prolonged and amplified the plasma leptin surge in offspring in a sex-specific manner (Kirk *et al.* 2009, Masuyama & Hiramatsu 2014). Maternal HF diet during lactation was also associated with increased insulin and leptin levels in milk (Vogt *et al.* 2014). Leptin that displays differential morphogenesis effects on male and female adipocytes (Guo *et al.* 2009) might account for WAT's programming dimorphism. However, despite the marked lactation effect in HF male vs female neonates, gender-specific modifications of plasma hormone levels and/or adipose tissue hormonal sensitivity remain to be determined.

Maternal obesity may also affect epigenetic mechanisms during adipogenesis. These modifications might be persistent and have long-term effects on

the expression of adipogenic and lipogenic genes. We hypothesize that maternal obesity affects offspring's energy and hormonal status modifying activity of the enzymatic components of the epigenetic machinery. It may cause epigenetic modifications that reprogram offspring's adipose tissue. Differences in fat cell embryonic origin, development, genetic, and hormonal sensitivity may result in a depot-specific programming effects that may predispose offspring to higher adiposity (Öst & Pospisilik 2015). Indeed, maternal obesity in mice induces increased gene expression of *Zfp423* with lower promoter methylation levels in fetal offspring (Yang *et al.* 2013). Similarly, weanling rats from obese dams display increased *Zfp423* and *C/Ebpβ* mRNA expression levels with alterations in DNA methylation of CpG sites (Borengasser *et al.* 2013). Maternal HF diet during pregnancy also results in histone modifications within leptin and adiponectin promoter regions with gene expression modifications in mouse offspring (Masuyama & Hiramatsu 2012).

Few studies have reported that maternal obesity programs metabolic alterations and adiposity differently in a sex-dependent manner in progeny (Sun *et al.* 2012, Ornellas *et al.* 2013, Dahlhoff *et al.* 2014, Masuyama & Hiramatsu 2014). The basis of the sex-specific programming effects remains unclear, but could reflect direct interactions between nutritional signals and sex hormones in the tissues of the developing fetus (Aiken & Ozanne 2013). In human, numerous studies suggest that sex differences in fetal growth in response to adverse pregnancy conditions are likely to be mediated by sex-specific adaptation of the placenta (Clifton 2010). Similarly, sex-specific programming effects in rat offspring from obese dams might be due to sex-specific differences in placental response to maternal obesity (Reynolds *et al.* 2015). Epigenetic mechanisms may also contribute to placental programming in a dimorphic manner. Thus, the consumption of HF diet during pregnancy appears to differently affect placental methylation and placental gene expression patterns in male and female mice offsprings (Gallou-Kabani *et al.* 2010). Thus, sex-specific differences in terms of epigenetic modulations may be associated with developmentally programmed phenotypes. It is possible that postnatal hormonal milieu, which is different between male and female offspring, modify the programming of adipose tissue induced by maternal obesity. This may result in gender-specific outcomes in relation to different sex-steroids (Dunn *et al.* 2011). Thus, a better knowledge of the epigenome changes in response

to maternal obesity may provide a promising way forward to reverse adverse programming of adiposity.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was supported by grants from the French 'Heart and Arteries' Foundation and grants of the French Ministry of Education.

Acknowledgments

The authors thank Valérie Montel, Anne Dicks-Coopman, Charlene Pourpe, and Capucine Besengez for their excellent technical assistance.

References

- Aiken CE & Ozanne SE 2013 Sex differences in developmental programming models. *Reproduction* **145** R1–R13. (doi:10.1530/REP-11-0489)
- Barker DJP 2004 Developmental origins of adult health and disease. *Journal of Epidemiology and Community Health* **58** 114–115. (doi:10.1136/jech.58.2.114)
- Benkalfat NB, Merzouk H, Bouanane S, Merzouk S-A, Bellenger J, Gresti J, Tessier C & Narce M 2011 Altered adipose tissue metabolism in offspring of dietary obese rat dams. *Clinical Science* **121** 19–28. (doi:10.1042/CS20100534)
- Berry DC, Stenesen D, Zeve D & Graff JM 2013 The developmental origins of adipose tissue. *Development* **140** 3939–3949. (doi:10.1242/dev.080549)
- Björntorp P & Sjöström L 1971 Number and size of adipose tissue fat cells in relation to metabolism in human obesity. *Metabolism: Clinical and Experimental* **20** 703–713. (doi:10.1016/0026-0495(71)90084-9)
- Bol VV, Reusens BM & Remacle CA 2008 Postnatal catch-up growth after fetal protein restriction programs proliferation of rat preadipocytes. *Obesity* **16** 2760–2763. (doi:10.1038/oby.2008.417)
- Borengasser SJ, Zhong Y, Kang P, Lindsey F, Ronis MJJ, Badger TM, Gomez-Acevedo H & Shankar K 2013 Maternal obesity enhances white adipose tissue differentiation and alters genome-scale DNA methylation in male rat offspring. *Endocrinology* **154** 4113–4125. (doi:10.1210/en.2012-2255)
- Bouchard C 2009 Childhood obesity: are genetic differences involved? *American Journal of Clinical Nutrition* **89** 1494S–1501S. (doi:10.3945/ajcn.2009.27113C)
- Boullu-Ciocca S, Achard V, Tassistro V, Dutour A & Grino M 2008 Postnatal programming of glucocorticoid metabolism in rats modulates high-fat diet-induced regulation of visceral adipose tissue glucocorticoid exposure and sensitivity and adiponectin and proinflammatory adipokines gene expression in adulthood. *Diabetes* **57** 669–677. (doi:10.2337/db07-1316)
- Breton C 2013 The hypothalamus – adipose axis is a key target of developmental programming by maternal nutritional manipulation. *Journal of Endocrinology* **216** R19–R31 (doi:10.1530/JOE-12-0157)
- Campbell JE, Peckett AJ, D'souza AM, Hawke TJ & Riddell MC 2011 Adipogenic and lipolytic effects of chronic glucocorticoid exposure. *American Journal of Physiology: Cell Physiology* **300** C198–C209. (doi:10.1152/ajpcell.00045.2010)

- Clifton VL 2010 Review: sex and the human placenta: mediating differential strategies of fetal growth and survival. *Placenta* **31** (Supplement) S33–S39. (doi:10.1016/j.placenta.2009.11.010)
- Dahlhoff M, Pfister S, Blutke A, Rozman J, Klingenspor M, Deutsch MJ, Rathkolb B, Fink B, Gimpfl M, Hrabě de Angelis M, *et al.* 2014 Periconceptional obesogenic exposure induces sex-specific programming of disease susceptibilities in adult mouse offspring. *Biochimica et Biophysica Acta* **1842** 304–317. (doi:10.1016/j.bbadis.2013.11.021)
- Desai M & Ross MG 2011 Fetal programming of adipose tissue: effects of intrauterine growth restriction and maternal obesity/high-fat diet. *Seminars in Reproductive Medicine* **29** 237–245. (doi:10.1055/s-0031-1275517)
- Desai M, Jellyman JK, Han G, Beall M, Lane RH & Ross MG 2014 Maternal obesity and high-fat diet program offspring metabolic syndrome. *American Journal of Obstetrics and Gynecology* **211** 237.e1–e237.e13. (doi:10.1016/j.ajog.2014.03.025)
- Desai M, Jellyman JK, Han G, Lane RH & Ross MG 2015 Programmed regulation of rat offspring adipogenic transcription factor (PPAR γ) by maternal nutrition. *Journal of Developmental Origins of Health and Disease* **6** 530–538. (doi:10.1017/S2040174415001440)
- Dunn GA, Morgan CP & Bale TL 2011 Sex-specificity in transgenerational epigenetic programming. *Hormones and Behavior* **59** 290–295. (doi:10.1016/j.yhbeh.2010.05.004)
- Fujiki K, Kano F, Shiota K & Murata M 2009 Expression of the peroxisome proliferator activated receptor gamma gene is repressed by DNA methylation in visceral adipose tissue of mouse models of diabetes. *BMC Biology* **7** 38. (doi:10.1186/1741-7007-7-38)
- Gallou-Kabani C, Gabory A, Tost J, Karimi M, Mayeur S, Lesage J, Boudadi E, Gross M-S, Taurelle J, Vigé A, *et al.* 2010 Sex- and diet-specific changes of imprinted gene expression and DNA methylation in mouse placenta under a high-fat diet. *PLoS ONE* **5** e14398. (doi:10.1371/journal.pone.0014398)
- Gnanalingham MG, Mostyn A, Symonds ME & Stephenson T 2005 Ontogeny and nutritional programming of adiposity in sheep: potential role of glucocorticoid action and uncoupling protein-2. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **289** R1407–R1415. (doi:10.1152/ajpregu.00375.2005)
- Guberman C, Jellyman JK, Han G, Ross MG & Desai M 2013 Maternal high-fat diet programs rat offspring hypertension and activates the adipose renin-angiotensin system. *American Journal of Obstetrics and Gynecology* **209** 262.e1–e8. (doi:10.1016/j.ajog.2013.05.023)
- Guo K, Mogen J, Struzzi S & Zhang Y 2009 Preadipocyte transplantation: an in vivo study of direct leptin signaling on adipocyte morphogenesis and cell size. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **296** R1339–R1347. (doi:10.1152/ajpregu.90691.2008)
- Guo C, Li C, Myatt L, Nathanielsz PW & Sun K 2013 Sexually dimorphic effects of maternal nutrient reduction on expression of genes regulating cortisol metabolism in fetal baboon adipose and liver tissues. *Diabetes* **62** 1175–1185. (doi:10.2337/db12-0561)
- Hanson MA & Gluckman PD 2014 Early developmental conditioning of later health and disease: physiology or pathophysiology? *Physiological Reviews* **94** 1027–1076. (doi:10.1152/physrev.00029.2013)
- Huan J-N, Li J, Han Y, Chen K, Wu N & Zhao AZ 2003 Adipocyte-selective reduction of the leptin receptors induced by antisense RNA leads to increased adiposity, dyslipidemia, and insulin resistance. *Journal of Biological Chemistry* **278** 45638–45650. (doi:10.1074/jbc.M304165200)
- Jiang L, Wang Q, Yu Y, Zhao F, Huang P, Zeng R, Qi RZ, Li W & Liu Y 2009 Leptin contributes to the adaptive responses of mice to high-fat diet intake through suppressing the lipogenic pathway. *PLoS ONE* **4** e6884. (doi:10.1371/journal.pone.0006884)
- Kirk SL, Samuelsson A-M, Argenton M, Dhonye H, Kalamatianos T, Poston L, Taylor PD & Coen CW 2009 Maternal obesity induced by diet in rats permanently influences central processes regulating food intake in offspring. *PLoS ONE* **4** e5870. (doi:10.1371/journal.pone.0005870)
- Lecoutre S & Breton C 2014 The cellularity of offspring's adipose tissue is programmed by maternal nutritional manipulations. *Adipocyte* **3** 256–262. (doi:10.4161/adip.29806)
- Lecoutre S & Breton C 2015 Maternal nutritional manipulations program adipose tissue dysfunction in offspring. *Frontiers in Physiology* **6** 158. (doi:10.3389/fphys.2015.00158)
- Leddy MA, Power ML & Schulkin J 2008 The impact of maternal obesity on maternal and fetal health. *Reviews in Obstetrics & Gynecology* **1** 170–178.
- Lee M-J, Pramyothin P, Karastergiou K & Fried SK 2014 Deconstructing the roles of glucocorticoids in adipose tissue biology and the development of central obesity. *Biochimica et Biophysica Acta* **1842** 473–481. (doi:10.1016/j.bbadis.2013.05.029)
- Lemonnier D 1972 Effect of age, sex, and sites on the cellularity of the adipose tissue in mice and rats rendered obese by a high-fat diet. *Journal of Clinical Investigation* **51** 2907–2915. (doi:10.1172/JCI107115)
- Lukaszewski M-A, Mayeur S, Fajardy I, Delahaye F, Dutriez-Casteloot I, Montel V, Dickes-Coopman A, Laborie C, Lesage J, Vieau D, *et al.* 2011 Maternal prenatal undernutrition programs adipose tissue gene expression in adult male rat offspring under high-fat diet. *American Journal of Physiology: Endocrinology and Metabolism* **301** E548–E559. (doi:10.1152/ajpendo.00011.2011)
- Lukaszewski M-A, Eberlé D, Vieau D & Breton C 2013 Nutritional manipulations in the perinatal period program adipose tissue in offspring. *American Journal of Physiology: Endocrinology and Metabolism* **305** E1195–E1207. (doi:10.1152/ajpendo.00231.2013)
- Masuyama H & Hiramatsu Y 2012 Effects of a high-fat diet exposure in utero on the metabolic syndrome-like phenomenon in mouse offspring through epigenetic changes in adipocytokine gene expression. *Endocrinology* **153** 2823–2830. (doi:10.1210/en.2011-2161)
- Masuyama H & Hiramatsu Y 2014 Additive effects of maternal high fat diet during lactation on mouse offspring. *PLoS ONE* **9** e92805. (doi:10.1371/journal.pone.0092805)
- Muhlhausler B & Smith SR 2009 Early-life origins of metabolic dysfunction: role of the adipocyte. *Trends in Endocrinology and Metabolism* **20** 51–57. (doi:10.1016/j.tem.2008.10.006)
- Murabayashi N, Sugiyama T, Zhang L, Kamimoto Y, Umekawa T, Ma N & Sagawa N 2013 Maternal high-fat diets cause insulin resistance through inflammatory changes in fetal adipose tissue. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* **169** 39–44. (doi:10.1016/j.ejogrb.2013.02.003)
- Nivoit P, Morens C, Van Assche FA, Jansen E, Poston L, Remacle C & Reusens B 2009 Established diet-induced obesity in female rats leads to offspring hyperphagia, adiposity and insulin resistance. *Diabetologia* **52** 1133–1142. (doi:10.1007/s00125-009-1316-9)
- Ornellas F, Mello VS, Mandarim-de-Lacerda CA & Aguilá MB 2013 Sexual dimorphism in fat distribution and metabolic profile in mice offspring from diet-induced obese mothers. *Life Sciences* **93** 454–463. (doi:10.1016/j.lfs.2013.08.005)
- Ortega FJ, Mayas D, Moreno-Navarrete JM, Catalán V, Gómez-Ambrosi J, Esteve E, Rodríguez-Hermosa JJ, Ruiz B, Ricart W, Peral B, *et al.* 2010 The gene expression of the main lipogenic enzymes is downregulated in visceral adipose tissue of obese subjects. *Obesity* **18** 13–20. (doi:10.1038/oby.2009.202)
- Öst A & Pospisilik JA 2015 Epigenetic modulation of metabolic decisions. *Current Opinion in Cell Biology* **33** 88–94. (doi:10.1016/j.ceb.2014.12.005)
- Popkin BM, Adair LS & Ng SW 2012 Global nutrition transition and the pandemic of obesity in developing countries. *Nutrition Reviews* **70** 3–21. (doi:10.1111/j.1753-4887.2011.00456.x)
- Ravelli AC, van Der Meulen JH, Osmond C, Barker DJ & Bleker OP 1999 Obesity at the age of 50 y in men and women exposed to famine prenatally. *American Journal of Clinical Nutrition* **70** 811–816.
- Reynolds CM, Vickers MH, Harrison CJ, Segovia SA & Gray C 2015 Maternal high fat and/or salt consumption induces sex-specific

- inflammatory and nutrient transport in the rat placenta. *Physiological Reports* **3** e12399. (doi:10.14814/phy2.12399)
- Samuelsson A-M, Matthews PA, Argenton M, Christie MR, McConnell JM, Jansen EHJM, Piersma AH, Ozanne SE, Twinn DF, Remacle C, *et al.* 2008 Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. *Hypertension* **51** 383–392. (doi:10.1161/HYPERTENSIONAHA.107.101477)
- Sen S & Simmons RA 2010 Maternal antioxidant supplementation prevents adiposity in the offspring of Western diet-fed rats. *Diabetes* **59** 3058–3065. (doi:10.2337/db10-0301)
- Shankar K, Harrell A, Liu X, Gilchrist JM, Ronis MJJ & Badger TM 2008 Maternal obesity at conception programs obesity in the offspring. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **294** R528–R538. (doi:10.1152/ajpregu.00316.2007)
- Spalding KL, Arner E, Westermark PO, Bernard S, Buchholz BA, Bergmann O, Blomqvist L, Hoffstedt J, Näslund E, Britton T, *et al.* 2008 Dynamics of fat cell turnover in humans. *Nature* **453** 783–787. (doi:10.1038/nature06902)
- Sun K, Kusminski CM & Scherer PE 2011 Adipose tissue remodeling and obesity. *Journal of Clinical Investigation* **121** 2094–2101. (doi:10.1172/JCI45887.2094)
- Sun B, Purcell RH, Terrillion CE, Yan J, Moran TH & Tamashiro KLK 2012 Maternal high-fat diet during gestation or suckling differentially affects offspring leptin sensitivity and obesity. *Diabetes* **61** 2833–2841. (doi:10.2337/db11-0957)
- Tang QQ & Lane MD 2012 Adipogenesis: from stem cell to adipocyte. *Annual Review of Biochemistry* **81** 715–736. (doi:10.1146/annurev-biochem-052110-115718)
- Umekawa T, Sugiyama T, Du Q, Murabayashi N, Zhang L, Kamimoto Y, Yoshida T, Sagawa N & Ikeda T 2015 A maternal mouse diet with moderately high-fat levels does not lead to maternal obesity but causes mesenteric adipose tissue dysfunction in male offspring. *Journal of Nutritional Biochemistry* **26** 259–266. (doi:10.1016/j.jnutbio.2014.10.012)
- Vogt MC, Paeger L, Hess S, Steculorum SM, Awazawa M, Hampel B, Neupert S, Nicholls HT, Mauer J, Hausen AC, *et al.* 2014 Neonatal insulin action impairs hypothalamic neurocircuit formation in response to maternal high-fat feeding. *Cell* **156** 495–509. (doi:10.1016/j.cell.2014.01.008)
- White UA & Tchoukalova YD 2014 Sex dimorphism and depot differences in adipose tissue function. *Biochimica et Biophysica Acta: Molecular Basis of Disease* **1842** 377–392. (doi:10.1016/j.bbadis.2013.05.006)
- Williams L, Seki Y, Vuguin PM & Charron MJ 2014 Animal models of in utero exposure to a high fat diet: a review. *Biochimica et Biophysica Acta* **1842** 507–519. (doi:10.1016/j.bbadis.2013.07.006)
- Yamamoto Y, Gesta S, Lee KY, Tran TT, Saadatirad P & Kahn CR 2010 Adipose depots possess unique developmental gene signatures. *Obesity* **18** 872–878. (doi:10.1038/oby.2009.512)
- Yang Q-Y, Liang J-F, Rogers CJ, Zhao J-X, Zhu M-J & Du M 2013 Maternal obesity induces epigenetic modifications to facilitate Zfp423 expression and enhance adipogenic differentiation in fetal mice. *Diabetes* **62** 3727–3735. (doi:10.2337/db13-0433)
- Zambrano E, Martínez-Samayoa PM, Rodríguez-González GL & Nathanielsz PW 2010 Dietary intervention prior to pregnancy reverses metabolic programming in male offspring of obese rats. *Journal of Physiology* **588** 1791–1799. (doi:10.1113/jphysiol.2010.190033)
- Zhang B, Berger J, Hu E, Szalkowski D, White-Carrington S, Spiegelman BM & Moller DE 1996 Negative regulation of peroxisome proliferator-activated receptor-gamma gene expression contributes to the antiadipogenic effects of tumor necrosis factor-alpha. *Molecular Endocrinology* **10** 1457–1466. (doi:10.1210/mend.10.11.8923470)

Received in final form 7 April 2016

Accepted 27 April 2016

Accepted Preprint published online 27 April 2016