Collective timekeeping among cells of the master circadian clock

Jennifer A Evans

Department of Biomedical Sciences, Marquette University, Milwaukee, WI, USA

Correspondence
should be addressed
to J A Evans

Email
jennifer.evans@marquette.edu

Abstract

The suprachiasmatic nucleus (SCN) of the anterior hypothalamus is the master circadian clock that coordinates daily rhythms in behavior and physiology in mammals. Like other hypothalamic nuclei, the SCN displays an impressive array of distinct cell types characterized by differences in neurotransmitter and neuropeptide expression. Individual SCN neurons and glia are able to display self-sustained circadian rhythms in cellular function that are regulated at the molecular level by a 24 h transcriptional–translational feedback loop. Remarkably, SCN cells are able to harmonize with one another to sustain coherent rhythms at the tissue level. Mechanisms of cellular communication in the SCN network are not completely understood, but recent progress has provided insight into the functional roles of several SCN signaling factors. This review discusses SCN organization, how intercellular communication is critical for maintaining network function, and the signaling mechanisms that play a role in this process. Despite recent progress, our understanding of SCN circuitry and coupling is far from complete. Further work is needed to map SCN circuitry fully and define the signaling mechanisms that allow for collective timekeeping in the SCN network.

Journal of Endocrinology
(2016) **230**, R27–R49

Introduction

Daily rhythms in metabolic and endocrine function serve to anticipate predictable changes in the environment. These rhythms are not driven externally, but instead arise from an intrinsic cellular process that tracks the hours of the day. In other words, cells are daily clocks. Because cellular clocks tick with a period close to but not exactly 24 h, they are referred to as ‘circadian’ (i.e. about a day). Biological timekeeping at the circadian timescale ensures that physiological and behavioral processes occur at the appropriate time of day. In this respect, it is an essential and ubiquitous feature of life on this planet. For example, glucocorticoid release surges just before waking to proactively marshal important energy resources (Fig. 1, Takahashi *et al.* 1968). Likewise, the secretion of growth hormone is highest during slow-wave sleep when its

ability to repair and strengthen bones is most effective (Fig. 1, Takahashi *et al.* 1968). In both cases, the timing of these rhythms is programmed by a biological clock and influenced by sleep (Czeisler & Klerman 1999). Over the last few decades, we have gained a deep understanding of the mechanisms that produce circadian timekeeping at the molecular and cellular levels. A key remaining question is how the numerous clock cells of our bodies form a coordinated system.

In mammals, the circadian system is a hierarchical collection of tissue clocks located throughout the brain and body (Mohawk *et al.* 2012). Many of these clocks are endocrine tissues that regulate hormone synthesis and release across the day (Fig. 1). Examples include the pituitary gland, the pineal gland, the adrenal gland,

Figure 1

Daily hormone rhythms are regulated by the circadian timekeeping system. (A) Hormone release fluctuates over the circadian cycle, with peak times occurring at specific times of day. Figure based on data from (Takahashi *et al.* 1968, Gavrilu *et al.* 2003, Koutkia *et al.* 2005, Natalucci *et al.* 2005, Russell *et al.* 2008). (B) The circadian timekeeping system is a hierarchical collection of tissue clocks located throughout the brain and body, many of which are endocrine tissues that regulate hormone synthesis and release across the day.

adipose tissue, and immune cells. These myriad body clocks are coordinated by the suprachiasmatic nucleus (SCN) of the anterior hypothalamus (Mohawk & Takahashi 2011). The SCN serves as a master clock that receives cues from the environment (e.g., light) and relays them to downstream tissues through a variety of outputs (e.g. synaptic connections, humoral cues, behavioral and

physiological control of sleep and body temperature). In this manner, the SCN harmonizes the various body clocks with one another and with the local time zone. Classic work demonstrates that the SCN is necessary for maintaining circadian rhythms in numerous processes, including sleep, feeding, drinking, melatonin production, and reproductive function (Klein *et al.* 1991, Weaver 1998). Given its extensive influence on endocrine function, understanding SCN circuitry is of prime importance.

The SCN is a neuronal network of cellular clocks that coordinate with one another to form a functional and cohesive population (Welsh *et al.* 2010). Like other hypothalamic nuclei, the SCN produces a diverse range of neurotransmitters and neuropeptides (van den Pol & Tsujimoto 1985, Lee *et al.* 2013), which have been used to define distinct neuronal subclasses (Antle & Silver 2005). In addition to relaying information to downstream clocks, SCN signaling molecules serve as local coupling factors that are important for maintaining network function. In particular, when SCN neurons are embedded in the network, they produce rhythms that are more precise, higher amplitude, and more robust than those displayed by isolated SCN neurons. Thus, intercellular communication within the SCN network is critically important for ensuring the fidelity of outputs to downstream tissues. Moreover, SCN neurons coordinate as a population to encode environmental conditions that are critical to survival (e.g. seasonal changes in day length). Despite its importance, the process by which SCN neurons influence one another is not completely understood. For instance, it is not known precisely how different SCN factors contribute to network function, how these signals are transmitted, and how they act to influence cellular rhythms. This review will describe recent progress that has advanced the understanding of SCN circuitry and highlight several issues that remain outstanding.

Circadian timekeeping at the cellular level

As a tissue, the SCN displays daily rhythms in numerous cellular processes, including metabolism, electrical activity, gene/protein expression, peptide release, and response to photic stimulation (Klein *et al.* 1991, Weaver 1998). Most of these rhythms persist when the SCN is studied in isolation from the environment and the rest of the brain. Thus, autonomous rhythmicity is an intrinsic property of the SCN itself. A key question posed almost immediately after this discovery was whether SCN rhythms are a network- or cellular-driven phenomenon.

The simple answer: SCN neurons are intrinsic clock cells. Pioneering work demonstrated that individual SCN neurons are capable of sustaining cellular rhythms even when dispersed at low density (Welsh *et al.* 1995, Herzog *et al.* 1998, Honma *et al.* 1998). Based on these results, it is commonly stated that SCN neurons are autonomous, self-sustained clocks. But the reality of the situation is slightly more complex because SCN neurons communicate with one another in ways that strengthen cellular rhythms (see more in next section).

But how does a cellular clock keep circadian time? Research conducted over the last few decades has revealed an elegant molecular mechanism that operates in nearly every cell of the body (Buhr & Takahashi 2013). Briefly, cellular rhythms are generated by interlocking feedback loops controlling the daily transcription of 'clock genes' and 'clock-controlled genes' (Fig. 2). At its core, the molecular circadian clock is a delayed negative-feedback loop, with positive elements that drive transcription and negative elements that repress transcription on a daily basis. The positive elements are the transcription factors CLOCK and

BMAL1, whereas the negative elements are PERIOD and CRYPTOCHROME proteins. Daily transcription is initiated when CLOCK and BMAL1 form a dimer that activates the expression of a family of *Period* (*Per1*, *Per2*, *Per3*) and *Cryptochrome* genes (*Cry1*, *Cry2*). This increases the levels of PER and CRY proteins, which dimerize, translocate into the nucleus, and inhibit their own transcription by repressing CLOCK-BMAL1 function (Fig. 2). This transcriptional repression, along with ubiquitination and degradation, causes PER and CRY levels to decline. Falling levels of PER and CRY then allow for derepression of CLOCK-BMAL1, which reactivates the transcription of *Per* and *Cry*. And so, the cycle renews the following day. This type of negative-feedback mechanism forms the basis of circadian timekeeping in a wide range of organisms (bacteria, fungi, plants, insects, and mammals) although the genes involved differ across species (Mackey 2007).

It remains a key area of research to understand fully the molecular control of cellular rhythms (Zhang & Kay 2010). With each passing day, our understanding of the circadian network at this level grows. We now appreciate that there are additional gene networks that regulate the core loop described above. For example, one interconnected loop regulates *Bmal1* and *Clock* transcription via opposing actions of ROR and REV-ERB (Fig. 2). This ancillary loop regulates the precision, robustness, and amplitude of core clock gene expression. Also, there are numerous 'clock-controlled genes' that are influenced by the molecular clock either directly or indirectly. For example, the transcription factor D-binding protein (DBP) is directly controlled by CLOCK-BMAL1 (Fig. 2). By regulating the timing of DBP expression, the core molecular clock can influence myriad downstream targets. To further demonstrate the interlocked nature of these feedback loops, DBP can influence *Per* transcription by binding to an upstream promoter sequence. Moreover, clock genes may contain additional regulatory elements (e.g., CRE) through which their expression is regulated by cellular and environmental signals. Collectively, the circadian molecular clock operates in nearly every cell of the body to regulate at least half the genome (Zhang *et al.* 2014).

In SCN neurons, the molecular oscillator controls myriad cellular processes (Kuhlman 2007). Notably, SCN neurons typically display daily rhythms in membrane potential with high firing during the day and low firing during the night (Brown & Piggins 2007). The temporal patterning of SCN electrical activity has important implications for network signaling and outputs to downstream tissues, so this is a critical area of research. Yet this approach is often limited in the number of SCN

Figure 2

The molecular circadian clock comprises interlocking transcriptional-translational feedback loops. CLOCK and BMAL1 are transcription factors that bind to E-box elements within the promoter sequences of a variety of clock genes. The protein products of *Period* (*Per*) and *Cryptochrome* (*Cry*) genes form repressors that inhibit their own transcription. Additional feedback loops involve other clock genes that interact with the elements of the core loop to amplify and stabilize molecular clock function. For example, the protein products of *Rev-Erb* and *Ror* genes compete for binding at ROR elements to influence *Bmal1* and *Clock* transcription. Further, the protein products of clock-controlled genes, such as *Dbp*, can influence both downstream molecular targets and core clock genes by binding at D-box elements.

neurons that can be recorded at the same time. Recent technological advances enable a more comprehensive view of SCN network function using real-time imaging of molecular activity. Over the years, researchers have engineered an impressive array of mouse models with genetically encoded optical reporters (e.g. *Per1*-GFP, *Per1*-luciferase, *PER2::LUCIFERASE*, *Bmal1*-luciferase, and *Cry1*-luciferase) that allow one to track molecular clock function in real time (Kuhlman *et al.* 2000, Yamaguchi *et al.* 2003, Yoo *et al.* 2004, Noguchi *et al.* 2010, Maywood *et al.* 2013). In addition, genetic mouse models and viral tools have been developed to monitor daily rhythms in Ca^{+2} and other intracellular signals (Enoki *et al.* 2012, Brancaccio *et al.* 2013, Irwin & Allen 2013). Provided there is good penetrance and faithful recapitulation of native molecular function, real-time imaging approaches provide an unprecedented view of the dynamic clock because they can be used to monitor hundreds of SCN neurons simultaneously. Relative to standard electrophysiological approaches, they also allow for cellular recordings to be performed in a relatively noninvasive manner for many days in culture. Together with more traditional approaches, real-time imaging techniques provide excellent tools to investigate cellular clock function and network properties.

The SCN network: greater than the sum of its parts

It is now well established that the SCN contains multiple cellular clocks that interact with one another to form a functional network. As described above, individual SCN cells can express self-sustained circadian rhythms (Welsh *et al.* 1995, Herzog *et al.* 1998, Honma *et al.* 1998). Nevertheless, it is clear that SCN neurons are weaker clocks when isolated from one another. For instance, ~60% of SCN neurons are competent oscillators when dispersed in low-density culture, but only ~30% of neurons are rhythmic under conditions of complete isolation (Webb *et al.* 2009). Strikingly, this is much lower than that observed when SCN neurons are connected in the network, with >90% of cells sustaining rhythms in a slice preparation. Further, coupled SCN neurons display rhythms that are more precise, higher amplitude, and more robust to perturbation and stochastic noise (Herzog *et al.* 2004, Abraham *et al.* 2010, Buhr *et al.* 2010). Remarkably, network interactions can even preserve cellular clock function when the molecular oscillator is crippled by genetic defects (Nakamura *et al.* 2002, Liu *et al.* 2007, Ko *et al.* 2010, Evans *et al.* 2012). This series

of observations indicates that SCN neurons can keep time by themselves, but are more capable clocks when they receive signals from other cells in the network. Of note, this is not true for all types of clock cells. Fibroblasts are strong circadian oscillators that appear to depend very little on one another for sustaining cellular rhythms (Nagoshi *et al.* 2004, Leise *et al.* 2012). While this may be viewed as evidence that SCN neurons are 'weaker' clocks than fibroblasts, this is not necessarily the case. Fibroblast clocks may oscillate well as individual cells, but they lack network-level properties that can buffer them from clock gene mutations (Liu *et al.* 2007). By contrast, SCN neurons have strength in numbers because they influence and bolster one another.

Another key property of SCN neurons is that they are able to synchronize with one another. SCN neurons embedded within the network display well-coordinated rhythms both *in vivo* and *in vitro*. By contrast, SCN neurons that are dissociated 'run at different speeds' due to the expression of different period lengths (Welsh *et al.* 1995, Herzog *et al.* 1998, Honma *et al.* 1998). As a consequence, dissociated SCN neurons gradually lose synchrony with one another over time in culture. But SCN neurons remain synchronized when they are able to interact fully with one another. These results indicate that SCN neurons adopt a common period due to intercellular communication, which prevents loss of synchrony among SCN neurons. Studies of mutant SCN neurons likewise suggest that they interact with one another to determine overall period (Liu *et al.* 1997, Herzog *et al.* 1998, Low-Zeddis & Takahashi 2001). Due to their ability to synchronize, SCN neurons remain coordinated as a population, maintain tissue-level rhythms, and send strong outputs to downstream tissues. Importantly, the coherence of SCN rhythms is a key determinant of behavioral and physiological rhythmicity (Ciarleglio *et al.* 2009). This is a property that appears to be unique to the SCN because most other types of cells appear unable to synchronize with one another (Yamazaki *et al.* 2000, Nagoshi *et al.* 2004, Welsh *et al.* 2004). Cellular clocks that lack communication would be expected to rely on SCN-controlled signals to maintain tissue coherence (Farnell *et al.* 2011). For example, fibroblasts are strong cellular oscillators, but they desynchronize and lose population-level rhythms (Welsh *et al.* 2004). Interestingly, it may be their strongly autonomous cellular clock that prevents fibroblasts from synchronizing with one another (Locke *et al.* 2008).

Lastly, network interactions influence the relative timing of neuronal activity to regulate the waveform of their collective rhythm. When embedded in the network,

SCN neurons 'prefer' to adopt specific phase relationships with electrical and molecular rhythms that are slightly dispersed in time rather than occurring all at the same time (Quintero *et al.* 2003, Saeb-Parsy & Dyball 2003, Yamaguchi *et al.* 2003, Hamada *et al.* 2004, Rohling *et al.* 2006, Yan *et al.* 2007, Evans *et al.* 2011, Myung *et al.* 2012, Brancaccio *et al.* 2013). This is not a random event; rather, there are clear regional phase differences that are reproducible and stereotyped across animals. Phase mapping analyses of PERIOD2::LUCIFERASE (PER2::LUC) expression in the murine SCN reveal several consistent patterns (Evans *et al.* 2011). First, the caudal SCN typically assumes an earlier phase than the rostral SCN. Also, there are fairly complex gradients of expression, with the dorsal SCN phase leading more central and ventral regions. These spatiotemporal arrangements are intrinsically regulated by the network itself (Quintero *et al.* 2003, Yamaguchi *et al.* 2003, Evans *et al.* 2011), and yet can be modulated markedly by environmental lighting conditions (reviewed in Meijer *et al.* 2012, Evans & Gorman 2016).

In terms of environmental modulation of SCN temporal organization, one of the most studied contexts is seasonal changes in day length. Day length regulates the phase coherence of SCN neurons, with more clustered phases under short winter-like days than long summer-like days (Jagota *et al.* 2000, Hazlerigg *et al.* 2005, Inagaki *et al.* 2007, Naito *et al.* 2008, Evans *et al.* 2013, Myung *et al.* 2015). Temporal coherence of phase among SCN neurons influences the overall waveform of rhythms produced by the network, with longer duration of clock gene expression and electrical activity under long days. This photoperiodic encoding is determined largely by changes in SCN phase relationships rather than changes in cellular rhythms (Rohling *et al.* 2006, Brown & Piggins 2009). Effectively, photoperiodic encoding by the SCN network alters the patterning of outputs transmitted under summer vs winter conditions. Seasonal changes in SCN outputs provide time of year information to downstream tissues to produce seasonal changes in behavior and physiology (Schaap *et al.* 2003, Inagaki *et al.* 2007). Thus, the SCN network is both a daily clock and an annual calendar.

In addition to its ecological significance, plasticity in SCN phase relationships can be exploited to test mechanisms of intercellular communication (Evans *et al.* 2013, 2015). In this approach, SCN neurons are desynchronized by light *in vivo* and then allowed to resynchronize *in vitro*, so that the process of network coupling can be tracked in real time. Specifically, very long day lengths reorganize the SCN network, so that it adopts a highly polarized state

with two groups of SCN neurons cycling in antiphase. After release from these lighting conditions, SCN neurons within these two groups interact with one another and gradually resynchronize over the course of a week. Leveraging this form of network plasticity, the process of SCN coupling can be tracked in real time using an *ex vivo* slice preparation. By capturing the dynamic process of SCN communication, this analytical assay may provide a novel discovery tool for further defining the circuitry of the SCN network (Evans *et al.* 2013).

SCN network organization: functional differences among neuronal subclasses

The SCN contains local projection neurons that communicate with one another and with other hypothalamic structures (Abrahamson & Moore 2001, Moore *et al.* 2002). The axons of many SCN neurons terminate within the nucleus itself, thus forming local circuit connections and/or collaterals from longer range projections. Nearly all SCN neurons produce β -aminobutyric acid (GABA), yet they can be categorized into distinct subgroups based on co-expression of different neuropeptides (Moore & Speh 1993, Abrahamson & Moore 2001). The SCN is typically subdivided into two spatially segregated compartments: the shell and the core (Moore & Silver 1998). These two compartments contain distinct subclasses of neurons that differ neurochemically (Fig. 3). The SCN shell contains a dense population of neurons that express arginine vasopressin (AVP), as well as other types of neurons. The SCN core contains a variety of subclasses, including neurons that express vasoactive intestinal polypeptide (VIP) or gastrin-releasing peptide (GRP). These subgroups have distinct developmental patterns (Antle *et al.* 2005b, Bedont & Blackshaw 2015) and are thought to represent distinct subclasses with minimal overlap in neuropeptide expression. However, as the index of SCN peptides grows, it is becoming apparent that SCN neurons can co-express different peptides that may contribute to their functional activity (e.g. Geoghegan & Carter 2008, Atkins *et al.* 2010, Drouyer *et al.* 2010, Hundahl *et al.* 2012, Lee *et al.* 2015). Although the shell–core scheme of the SCN network continues to be a convenient construct, it will likely morph to become more sophisticated as understanding of SCN circuitry increases (Morin 2007, 2012).

The expression and location of AVP and VIP neurons are consistent across mammalian species, but regional anatomy and chemoarchitecture can vary (Morin 2007, Cassone *et al.* 1988). For example, the SCN of rodents

Figure 3

The SCN network is a heterogeneous population of cellular clocks. (A) Organization of the mouse SCN, illustrating the compartmentalization of cell types and inputs in the coronal plane. AVP, arginine vasopressin; PK2, prokineticin 2; CCK, cholecystokinin; CALB, calbindin; CALR, calretinin; SP, substance P; GRP, gastrin-releasing peptide; VIP, vasoactive intestinal polypeptide; 5-HT, serotonin (from dorsal raphe); NPY, neuropeptide Y (from intergeniculate leaflet of the thalamus, which also releases GABA, neurotensin, and enkephalin); Glut, glutamate (from retina); PACAP, pituitary adenylate cyclase-activating polypeptide (from retina); 3V, third ventricle; OC, optic chiasm. (B) The shell–core model of the SCN network, representing how each compartment is thought to contribute to pacemaker function.

contains neurons that express calcium-binding proteins, but the location and specific proteins expressed differ among rodent species. In the Syrian hamster, calbindin-expressing SCN neurons form a spatially segregated subgroup densely packed into the central region of the core compartment. The mouse SCN contains both calbindin and calretinin neurons, with greatest expression within the core compartment. However, neither population forms a dense subnucleus. By contrast, calretinin neurons in the rat appear in the SCN shell rather than the SCN core. Another example of SCN variability across species is neuropeptide Y (NPY) expression. The rodent SCN does not produce NPY, but the human SCN does (Moore 1991). The consequence of species variation in SCN chemoarchitecture warrants further study given its potential influence on SCN function.

In addition to differences in neuropeptide expression, SCN neurons can be distinguished based on their response to afferent input. For instance, light exposure during the night typically increases SCN electrical activity and gene expression (i.e. *C-fos*, *Period*), but does so only in a subset of neurons (Meijer *et al.* 1986, Jiao *et al.* 1999, Aggelopoulos & Meissl 2000, Schwartz *et al.* 2000, Hamada *et al.* 2001, Nakamura *et al.* 2004, Brown *et al.* 2011). Of those SCN neurons affected directly by retinal stimulation, some respond with sustained increases in electrical activity, some respond with phasic responses, and others are inhibited. Further, photoreponsive SCN neurons may be distinguished based on the source of retinal input they receive (Brown *et al.* 2011). Tract tracing studies suggest that there is a denser retinal projection to the SCN core than the SCN shell (Abrahamson & Moore 2001, Lokshin *et al.* 2015), which maps onto functional differences in photic response. After light exposure, the SCN core displays changes in gene expression that precede those in the SCN shell (Silver *et al.* 1996, Dardente *et al.* 2002, Yan & Okamura 2002, Yan & Silver 2002, 2004, Kuhlman *et al.* 2003, Nagano *et al.* 2003). Regional differences in photic responsiveness have functional consequences that can influence how the network adjusts following changes in environmental lighting conditions. For instance, after simulated travel across time zones, the SCN core re-entrains faster than the SCN shell (Nagano *et al.* 2003, Albus *et al.* 2005, Nakamura *et al.* 2005, Davidson *et al.* 2009, Rohling *et al.* 2011, Sellix *et al.* 2012), and this form of light-driven desynchrony is thought to contribute to symptoms of jetlag. Moreover, regional patterning of afferent projections to the SCN is not unique to the retina, with inputs from the thalamus, pretectum, and median raphe also being densest in the SCN core (Fig. 3, for review, see Morin & Allen 2006). Furthermore, the SCN shell may receive input from a distinct set of structures, at least in the rat (Moga & Moore 1997, Leak & Moore 2001). Future studies should test whether this is consistent across mammalian species given the potential implications.

SCN neurons can also be distinguished based on efferent projections. The SCN communicates with downstream targets using both synaptic and humoral signals (Mohawk & Takahashi 2011). In the case of synaptic connections, the SCN innervates structures in the hypothalamus, thalamus, and forebrain (Watts 1991, Kalsbeek & Buijs 2002), and these projections may originate from specific SCN regions. With regard to outputs from SCN neuronal subclasses, those originating from AVP and VIP neurons have been most extensively studied. AVP and VIP neurons often project to the same

target, but the relative patterning of innervation can be target-specific (Watts & Swanson 1987, Abrahamson & Moore 2001, Kalsbeek & Buijs 2002). For instance, both AVP and VIP neurons in the murine SCN project to the paraventricular nucleus and the dorsomedial nucleus of the hypothalamus, but each target contains more AVP fibers than VIP fibers (Abrahamson & Moore 2001). The functional roles of projections from distinct SCN regions are far from fully understood. Thus far, research indicates that both SCN compartments influence rhythms in downstream targets, yet there are differences in the role of outputs from different SCN regions (Yan *et al.* 2005, Zhou & Cheng 2005, Lee *et al.* 2009, Schwartz *et al.* 2009, Kalsbeek *et al.* 2010, Butler *et al.* 2012, Smarr *et al.* 2012, Wotus *et al.* 2013, Evans *et al.* 2015). One emerging theme is that the SCN shell appears to set the phase of downstream tissues; however, both SCN shell and core neurons provide signals to downstream tissues that can influence their rhythms. The efferent projection patterns of distinct SCN subclasses should be revisited in future work given recent advances in cell-type-specific tract tracing (Callaway & Luo 2015). A deeper understanding of this issue may provide insight into the functional role of outputs from different SCN subclasses.

Lastly, there is a rich body of work describing regional differences in SCN function. As mentioned above, SCN neurons display regional differences in the phase of electrical and molecular activities, which can be detected even within a chemically defined subclass (Hamada *et al.* 2004, Kalsbeek *et al.* 2006, Evans *et al.* 2011, Yoshikawa *et al.* 2015). Although the majority of SCN neurons tend to exhibit clustered phases of electrical and molecular activities, antiphase rhythms are expressed by subgroups of SCN neurons whose chemical identity remains ill defined (Nakamura *et al.* 2001, King *et al.* 2003, Lee *et al.* 2003). In addition, SCN neurons from distinct regions or peptidergic subclasses can exhibit different period length (Shinohara *et al.* 1995, Nakamura *et al.* 2001, Noguchi & Watanabe 2008, Myung *et al.* 2012). Further, daily patterns of electrical activity can differ markedly among neuronal subgroups (Shibata *et al.* 1984, Jiao *et al.* 1999, Jobst & Allen 2002, Saeb-Parsy & Dyball 2003, Belle *et al.* 2009). This suggests that SCN neurons can oscillate with different properties, but the presence of cellular rhythms may not be universal among SCN neurons. Many studies examining the SCN network have found that photoresponsive neurons within the SCN core display a less rhythmic or arrhythmic phenotype (Jiao *et al.* 1999, Hamada *et al.* 2001, Jobst & Allen 2002). However, recent work indicates that cellular rhythmicity is stochastic, not

restricted to neurons of a particular peptidergic subgroup, and sustained by network interactions (Webb *et al.* 2009). This provides new insight, in that it suggests that there is no single subclass of 'pacemaker' neuron in the SCN network, and that cellular rhythmicity is strongly influenced by network connectivity rather than chemical phenotype.

In summary, SCN neurons in different regions of the network can be distinguished in many ways. The most common model of SCN organization highlights functional distinctions between its shell and core compartments. In this scheme, the SCN core contains first-order neurons that receive and process photic input. SCN core neurons display strong rhythms in light-induced responses, but electrical and molecular rhythms of these neurons may be low amplitude or nonexistent. Nevertheless, input transmitted from SCN core neurons is important for coordinating neurons in the rest of the network. In contrast, neurons in the SCN shell are strongly rhythmic and play a key role in transmitting daily outputs to downstream structures. Overall, there is utility in this linear model, but some key aspects of SCN circuitry remain unexplained. For example, if SCN core neurons are not different from SCN shell neurons when studied in isolation, what accounts for their differences when they are embedded in the network? If specific subclasses of SCN core neurons do indeed lack a strong molecular oscillator, how do they maintain circadian rhythms in photic responsiveness? Presumably, these observations reflect that molecular and photoresponsive rhythms of SCN core neurons are influenced by intercellular communication, but the relevant signals remain undefined. Additional work investigating SCN signaling is expected to expand the understanding of its circuitry.

SCN coupling mechanisms

SCN neurons communicate through multiple mechanisms and signaling factors (van den Pol & Dudek 1993, Michel & Colwell 2001, Aton & Herzog 2005). An important role of synaptic signaling is based on observations that SCN neurons desynchronize when cultured with tetrodotoxin (TTX) to block Na⁺-dependent action potentials (Yamaguchi *et al.* 2003). However, there is also evidence that the SCN network can use other forms of coupling that do not depend on synaptic communication. For instance, SCN timekeeping can be maintained in the absence of Na⁺-dependent action potentials and Ca²⁺-dependent synaptic transmission (Schwartz *et al.* 1987, Earnest *et al.* 1991,

Schwartz 1991, Bouskila & Dudek 1993, Dudek *et al.* 1993, Shibata & Moore 1993). Further, circadian rhythms are maintained in some species *in vivo* under environmental conditions that severely compromise neuronal activity (Menaker 1961, Grahn *et al.* 1994). Further, the SCN displays metabolic rhythms at an embryonic age (Shibata & Moore 1987, Reppert 1992) that precedes the completion of synaptogenesis (Bedont & Blackshaw 2015). Collectively, this suggests that synaptic transmission is not the exclusive means by which SCN neurons can communicate with one another. Interestingly, functional studies indicate that SCN neurons can communicate through paracrine signaling (Maywood *et al.* 2011). In this work, co-cultured SCN slices were able to influence the rhythmic properties of one another even though they were unable to establish cross-slice synaptic connections. One potential mechanism driving this effect may be the nonsynaptic release of SCN neuropeptides from axons, dendrites, and somata (Castel *et al.* 1996). While it remains unclear which specific SCN neuropeptides are released at these sites, this form of communication may influence the function of neuronal networks (Ludwig & Leng 2006, van den Pol 2012). In the section below, recent insight into the roles of different SCN coupling factors has been reviewed. In particular, much has been learned about the ways in which VIP, GABA, and AVP signaling influence SCN function. Nevertheless, it remains a challenge to fully map the neurochemical, temporal, and spatial properties of SCN circuits.

Vasoactive intestinal polypeptide (VIP)

Over the past decade, clear evidence has emerged that indicates VIP is important for synchronizing SCN neurons (Vosko *et al.* 2007, Harmar 2003). As described above, VIP is produced by a subset of SCN neurons located within the ventral SCN core, with functional evidence suggesting that there are at least two subclasses of VIP neurons in the rat (Kawamoto *et al.* 2003). VIP is expected to have pervasive effects because VIP+ fibers innervate nearly all SCN regions (Card *et al.* 1981, Card & Moore 1984), and most SCN neurons express the VIP receptor, VPAC2 (Kalamatianos *et al.* 2004b, Kallo *et al.* 2004b, An *et al.* 2012). The influence of VIP signaling is expected to fluctuate daily since VIP and its receptor are expressed rhythmically in the SCN *in vivo* and *in vitro* (Takahashi *et al.* 1989, Glazer & Gozes 1994, Duncan *et al.* 1995, Shinohara *et al.* 1995, Cagampang *et al.* 1998, Dardente *et al.* 2004). The VPAC2 receptor is a $G\alpha_s$ -coupled receptor that activates adenylyl cyclase, cAMP, PKA, and

CRE-dependent transcription (Fig. 4A, Harmar *et al.* 1998, Couvineau & Laburthe 2011). Given that *Period* genes are activated by CREB, this suggests that VIP will influence cellular rhythms in SCN neurons. As expected, VIP signaling alters SCN electrical and molecular activities *in vivo* and *in vitro* (Piggins *et al.* 1995, Nielsen *et al.* 2002, Cutler *et al.* 2003, Itri & Colwell 2003, Pakhotin *et al.* 2006, Irwin & Allen 2010, Kudo *et al.* 2013). Moreover, it can phase-shift other SCN neuropeptide rhythms, such as AVP release (Watanabe *et al.* 2000). Consistent with VPAC2 being a $G\alpha_s$ -coupled receptor, the effects of VIP on SCN electrical and molecular rhythms depend on PKA signaling (Nielsen *et al.* 2002, Meyer-Spasche & Piggins 2004, An *et al.* 2011). However, VIP-induced effects also require activation of other intracellular cascades (i.e., mitogen-activated protein kinase, phospholipase C), which indicates that other signaling mechanisms may be involved (Fig. 4A).

Interest in the role of VIP signaling was especially piqued when it was discovered that the loss of VIP signaling is able to compromise circadian rhythmicity. When studied under constant dark conditions, the majority of mice lacking the gene for VIP or its receptor display either arrhythmic locomotor patterns or low-amplitude rhythms with altered period (Harmar *et al.* 2002, Colwell *et al.* 2003, Aton *et al.* 2005, Ciarleglio *et al.* 2009). Arrhythmic *Vip*-deficient mice also display loss of rhythms in SCN electrical activity and gene expression due to the lack of network synchrony and a decrease in the number of SCN neurons able to maintain viable cellular rhythms (Aton *et al.* 2005, Maywood *et al.* 2006, Brown *et al.* 2007, Hughes *et al.* 2008, Ciarleglio *et al.* 2009). Importantly, SCN cellular rhythms and synchrony can be rescued in *Vip* knockout slices by daily application of a VPAC2 agonist (Aton *et al.* 2005). Loss of cellular rhythms and synchrony likewise occurs in wildtype SCN slices exposed to a VPAC2 antagonist (Brown *et al.* 2007, Evans *et al.* 2013), which suggests that the effects of *VIP/VPAC2* deletion are not due to developmental abnormalities. Further, VIP signaling has important consequences for spatiotemporal arrangements of the SCN network, which involves intercellular communication triggered by $G\alpha_q$ signaling in VIP neurons (Brancaccio *et al.* 2013). Collectively, this work indicates that VIP is a local coupling factor that amplifies and entrains SCN neurons that are weak intrinsic oscillators. These effects are very important for maintaining cellular rhythms and synchronized network activity within the SCN network.

In addition to serving as a local coupling factor, VIP signaling also influences other processes. In particular, VIP

Figure 4

Inter-cellular signaling mechanisms that alter cellular activity and clock gene expression within SCN neurons. (A) SCN neurons express the VPAC2 receptor for VIP, which is a G-protein-coupled receptor with seven transmembrane domains. Upon VIP binding to VPAC2, $G\alpha_s$ activates adenyl cyclase, cAMP, PKA, and CRE-dependent transcription. (B) Within the SCN, both $GABA_A$ receptors and $GABA_B$ receptors are expressed. GABA binding to the ionotropic $GABA_A$ receptor allows for influx or efflux of Cl^- , depending on $[Cl^-]_i$. By contrast, the metabotropic $GABA_B$ receptor is coupled to $G\alpha_{i/o}$, which inhibits adenyl cyclase. (C) The receptors for both AVP and GRP are $G\alpha_q$ -coupled receptors that stimulate phospholipase C ($PLC\beta$), which in turn activates phosphatidylinositol 4,5-bisphosphate (PIP2) to cause diacyl glycerol (DAG)-mediated activation of protein kinase C (PKC) and inositol triphosphate (IP3)-induced release of intracellular calcium stores. Intracellular release of calcium activates calmodulin (CaM), which stimulates Ca^{2+} /calmodulin-dependent protein kinase (CaMK). Due to the expression of CRE-elements in *Period* genes, changes in the intracellular signaling cascades illustrated in A–C can alter the function of the molecular clock.

is involved in photic signaling (Shen *et al.* 2000, Colwell *et al.* 2003, Kuhlman *et al.* 2003, Hughes *et al.* 2004, Dragich *et al.* 2010, Lucassen *et al.* 2012, An *et al.* 2013), and light influences VIP expression in the SCN (Duncan *et al.* 1995, Shinohara & Inouye 1995, Isobe & Nishino 1998, Smith & Canal 2009, Francl *et al.* 2010). Deficits in VIP signaling cause abnormal responses to light, with one result being an unusual responsiveness to light during the day. Furthermore, loss of VIP causes disruptions in a wide variety of physiological processes, including cardiovascular function, metabolism, and reproduction (Shimizu *et al.* 1996, Bechtold *et al.* 2008, Sheward *et al.* 2010, Hannibal *et al.* 2011, Schroeder *et al.* 2011, Loh *et al.* 2014). This indicates that VIP signaling is important for maintaining endocrine function, which could be due to its role in the SCN network itself and/or its role as an output signal to downstream tissues (van der Beek *et al.* 1993, Kalsbeek *et al.* 1993, Gerhold *et al.* 2001, Egli *et al.* 2004, Kallo *et al.* 2004b, Gerhold & Wise 2006, Loh *et al.* 2011, Fahrenkrug *et al.* 2012). Also, age- and sex-related differences in VIP expression may influence changes in the strength and robustness of circadian function (Zhou *et al.* 1995, Kawakami *et al.* 1997, Krajnak *et al.* 1998a,b, Kallo *et al.* 2004a, Mahoney *et al.* 2009). Thus,

VIP signaling is an important modulator whose absence has widespread consequences for behavior and physiology. There continues to be a need to define the properties and mechanisms of VIP signaling to better understand how it regulates clock function at the cellular, network, and systems levels. For instance, recent work demonstrates that VIP can desynchronize the phase of SCN neurons if given at the wrong time or at high doses (An *et al.* 2013, Ananthasubramaniam *et al.* 2014). This suggests that VIP expression is tightly regulated within the SCN network. Of interest, 4E-BP1 has been identified recently as a molecular repressor of *Vip* mRNA translation (Cao *et al.* 2013). Future studies are expected to provide additional insight into the temporal and spatial features of VIP signaling, as well as modulatory mechanisms.

γ -aminobutyric acid (GABA)

Since it was first reported that nearly all SCN neurons express GABA, it has been suspected that this neurotransmitter regulates SCN communication. The SCN displays a high density of GABA terminals and cells (van den Pol & Tsujimoto 1985, Okamura *et al.* 1989, Decavel & Van den Pol 1990, Moore & Speh 1993,

Huhman *et al.* 1996, Castel & Morris 2000, Abrahamson & Moore 2001). Receptors for GABA include both ionotropic GABA_A receptors and metabotropic GABA_B receptors coupled to Gα_{i/o} (Fig. 4B, Bormann 2000), with both being expressed in the SCN (Gao *et al.* 1995, O'Hara *et al.* 1995, Naum *et al.* 2001, Belenky *et al.* 2003, 2008). As with other SCN factors, daily rhythms in GABA signaling are evident in the SCN, and these rhythms can be modulated by light exposure (Aguilar-Roblero *et al.* 1993, Huhman *et al.* 1996, Naum *et al.* 2001, Itri *et al.* 2004). Consistent with this anatomical work, SCN neurons are sensitive to GABA signaling (Liou *et al.* 1990, Mason *et al.* 1991, Gillespie *et al.* 1997, 1999, Jiang *et al.* 1997b, Strecker *et al.* 1997, Biggs & Prosser 1998, Cardinali & Golombek 1998, Mintz *et al.* 2002, Ehlen *et al.* 2008). Collectively, these studies indicate that GABA regulates SCN neuronal activity, modulates photic signaling, and serves as an output signal to downstream tissues (Wang *et al.* 2003). For modulation of photic signaling, an important role has been demonstrated for both GABA_A and GABA_B receptors (Gillespie *et al.* 1997, Biggs & Prosser 1998, Belenky *et al.* 2003, Novak *et al.* 2004, Ehlen & Paul 2009). By contrast, GABA_A receptor signaling is thought to be the strongest local regulator of SCN network activity (Kim & Dudek 1992, Shimura *et al.* 1996, Strecker *et al.* 1997, Albus *et al.* 2005, Fan *et al.* 2015). Over the last two decades, work has revealed surprising ways in which GABA can modulate SCN function.

One of the first surprises discovered about GABA signaling in the SCN concerns the polarity of responses to this neurotransmitter. GABA is typically defined as an inhibitory neurotransmitter; however, GABA can depolarize and increase intracellular calcium concentration in SCN neurons (Wagner *et al.* 1997, Choi *et al.* 2008, Irwin & Allen 2009). Although excitatory GABAergic responses are not always observed (Bos & Mirmiran 1993, Dudek *et al.* 1993, Gribkoff *et al.* 1999), this may stem at least in part from spatial or temporal differences in recordings performed in different laboratories (De Jeu & Pennartz 2002, Albus *et al.* 2005, Choi *et al.* 2008, Ikeda *et al.* 2013, Alamilla *et al.* 2014, DeWoskin *et al.* 2015). Indeed, the SCN displays regional differences in the expression of the chloride co-transporters (Belenky *et al.* 2008, 2010, Choi *et al.* 2008) that determine chloride reversal potential (Vogt 2015). Adding to this complexity, it has been demonstrated recently that the polarity of GABA responses in the SCN is influenced by the duration of daily light exposure (Farajnia *et al.* 2014, Myung *et al.* 2015). This may have great relevance for the ability of the

SCN to serve as an annual calendar, and firmly places the SCN on the growing list of mature neural networks that exhibit plasticity in the polarity of GABA responses (Marty & Llano 2005).

Given that the nature of SCN responses to GABA remain unclear, it should come as little surprise that the precise role of GABA in SCN coupling has proven difficult to define. Early work demonstrated that dissociated SCN neurons exposed to GABA will synchronize their electrical rhythms via GABA_A signaling (Liu & Reppert 2000, Shirakawa *et al.* 2000). Additional studies further indicated that GABA_A signaling is involved in the transfer of resetting information from the SCN core to the SCN shell (Albus *et al.* 2005, Han *et al.* 2012). However, GABA receptor antagonists do not desynchronize SCN neurons *in vitro* (Aton *et al.* 2006). These results came as a surprise because they run counter to the hypothesis that GABA acts as a coupling factor. Recent work has provided some insight into these conflicting findings. GABA_A signaling does influence SCN coupling, but surprisingly it destabilizes phase relationships of SCN neurons (Evans *et al.* 2013, Freeman *et al.* 2013, DeWoskin *et al.* 2015, Myung *et al.* 2015). Normally, this effect of GABA_A signaling is hard to detect because VIP signaling is a potent synchronizing agent. But in the absence of VIP, SCN neurons desynchronize due to GABA_A signaling (Evans *et al.* 2013, Freeman *et al.* 2013). In fact, antagonism of GABA_A signaling will 'fix' a *Vip* KO slice and prevent desynchrony from occurring. This indicates that the loss of synchrony that emerges during VIP deficiency is not due to 'passive' desynchrony caused by cellular period differences but rather 'active' desynchronizing responses elicited by GABA signaling. This provides us with a new view into SCN circuitry because it indicates that some SCN signaling mechanisms promote synchronization, while others may cause desynchronization. This also reveals that SCN coupling factors may be arranged into pairs that oppose one another's effects. Building on this theme of interactive coupling mechanisms, GABA_A signaling can interact with VIP signaling in either an antagonistic or cooperative manner depending on the state of the network (Evans *et al.* 2013), which may account for earlier work demonstrating synchronizing effects (Liu & Reppert 2000, Shirakawa *et al.* 2000, Albus *et al.* 2005, Han *et al.* 2012). Together with previous work, this strongly suggests that the functional role of a coupling mechanism can change based on the experience, history, or age of the animal (Evans *et al.* 2013, Bedont *et al.* 2014, Wang *et al.* 2014).

Arginine vasopressin (AVP)

Although AVP has been traditionally viewed more as an SCN output signal, a more direct role in network synchronization has been suggested recently. AVP is rhythmically expressed both *in vivo* and *in vitro* (Cassone *et al.* 1988, Sukhov *et al.* 1993, Shinohara *et al.* 1998, Dardente *et al.* 2004, Van der Veen *et al.* 2005, Miller *et al.* 2006, Noguchi & Watanabe 2008, Mahoney *et al.* 2009, Yoshikawa *et al.* 2015). Of the three known AVP receptors, V_{1a} (V1) and V_{1b} (V3) are expressed within the SCN (Kalamatianos *et al.* 2004a, Li *et al.* 2009), which are both $G\alpha_q$ -coupled receptors that stimulate phospholipase C to cause DAG-mediated activation of PKC and IP3-induced mobilization of intracellular calcium (Fig. 4C, Maybauer *et al.* 2008). Although AVP was one of the first SCN neuropeptides to be discovered, it was deemed not necessary for circadian rhythms based on work in rats with a spontaneous loss-of-function mutation (Grobowski *et al.* 1981, Boer *et al.* 1998). Further work, however, revealed that these rats display lower amplitude rhythms of sleep, melatonin release, and corticosterone (Schroder *et al.* 1988, Brown & Nunez 1989, Wideman *et al.* 2000), and this led to the suggestion that AVP was mostly an important output signal (Jin *et al.* 1999, Tousson & Meissl 2004). In support of a network role, however, AVP neurons project locally within the SCN itself (Castel *et al.* 1990, Romijn *et al.* 1997) and exogenous AVP is able to regulate the cellular activity of SCN neurons (Liou & Albers 1989, Mihai *et al.* 1994, Ingram *et al.* 1998). However, it was unclear whether these responses actually produced functional consequences because AVP application *in vitro* or *in vivo* does not phase-shift SCN rhythms. Nevertheless, recent work has revealed that AVP is able to influence the function of the SCN network. V_{1a} and V_{1b} receptor knockout mice display a pronounced circadian phenotype, in that they are resistant to jetlag and re-entrain almost instantly following a shift in the light:dark cycle (Yamaguchi *et al.* 2013). Wildtype mice receiving AVP receptor antagonists directed to the SCN likewise shift quickly, thus discounting a developmental basis of this effect. Rapid recovery following simulated jetlag is also observed in mice lacking *Bmal1* in AVP neurons, which can be reversed by SCN-specific rescue of molecular clock function (Mieda *et al.* 2015). Collectively, this work suggests that AVP signaling is involved in setting the pace of re-entrainment. It has been proposed that the loss of AVP signaling causes a change in SCN coupling that allows the entire network to shift rapidly to the new time zone. Consistent with this, application of AVP can

synchronize SCN neurons collected from mice deficient in VIP signaling (Maywood *et al.* 2011). Further work testing the specific role of AVP signaling in SCN coupling is warranted.

Gastrin-releasing peptide (GRP)

The potential role of GRP in SCN coupling remains to be determined, although there is clear evidence that GRP signaling is involved in photic processing. GRP is rhythmically expressed in the SCN and modulated by light (Shinohara *et al.* 1993, Dardente *et al.* 2004, Lee *et al.* 2013). The GRP receptor, also known as the bombesin 2 (BB2) receptor, is likewise rhythmically expressed within the SCN in a manner influenced by light exposure (Karatsoreos *et al.* 2006). Like the AVP receptor, BB2 is a $G\alpha_q$ -coupled receptor (Fig. 4C, Jensen *et al.* 2008). GRP neurons within the SCN receive retinal input that induces cellular responses (Dardente *et al.* 2002, Lesauter *et al.* 2011). Further, GRP can phase-shift SCN rhythms *in vitro* and locomotor rhythms *in vivo* in a pattern similar to those induced by light and VIP (Piggins *et al.* 1995, Gillespie *et al.* 1997, McArthur *et al.* 2000, Aida *et al.* 2002, Piggins *et al.* 2005, Antle *et al.* 2005a, Gamble *et al.* 2007, Biello 2009). These effects of GRP are dependent on BB2 receptor signaling, CREB-dependent transcription, clock gene activation, and changes in fast delayed rectifier potassium currents (Aida *et al.* 2002, Piggins *et al.* 2005, Gamble *et al.* 2007, 2011). However, the role of GRP may not be limited to photic signaling because GRP can enhance cellular rhythms in SCN slices collected from mice deficient in VIP signaling (Brown *et al.* 2005, Maywood *et al.* 2006, 2011). Further, cellular rhythmicity is attenuated by a BB2 receptor antagonist, but only when applied in the absence of VIP signaling (Brown *et al.* 2005). Together with the work on GABA described above, this suggests that VIP is a very strong modulator of SCN cellular rhythms that may mask the effects of other signaling mechanisms.

Other neuroactive substances

In addition to these well-studied subgroups, there are likely novel subclasses of SCN neurons that play a role in network coupling. The SCN produces dozens of signaling factors, and the number of recognized SCN peptides continues to increase (van den Pol & Tsujimoto 1985, Lee *et al.* 2013, 2015). A recent study demonstrated that SCN network function is regulated by neuromedin S, which is an SCN peptide produced by both AVP and VIP neurons

(Lee *et al.* 2015). Also, a forward peptidomics screen identified little SAS as a novel peptide produced in the SCN core that relays photic signals independent of VIP- or GRP-dependent signaling (Atkins *et al.* 2010). Another interesting development is that intra-SCN glutamatergic signaling may play a role in coupling the left and right SCN (Michel *et al.* 2013), and yet glutamate is not expressed by SCN neurons (Strecker *et al.* 1997). Communication between the left and right SCN likely differs from that coupling neurons within each SCN (Bouskila & Dudek 1993), and additional studies may shed new light on mechanisms mediating this coupling. Overall, this work highlights that the SCN remains a complex structure, and it is likely that additional coupling signals will be identified in the future.

Nonsynaptic interactions

One early hypothesis posited a role of local electric field effects produced by changes in the membrane potential of adjacent cells. Ephaptic interactions produced by electrical field potentials in dendrites can synchronize electrical activity in the mammalian neocortex, but it is unlikely that this mechanism couples oscillators within the SCN given its nonlaminar organization (Van den Pol 1980, van Esseveldt *et al.* 2000). Nevertheless, functional and anatomical evidence suggests that SCN neurons can communicate through nonsynaptic release of neuropeptides (Castel *et al.* 1996, Maywood *et al.* 2011). Moreover, electrotonic communication through low-resistance gap junctions may influence SCN coupling (Jiang *et al.* 1997a, Colwell 2000, Jobst *et al.* 2004, Long *et al.* 2005). Gap junctions are channels that allow the exchange of small molecules between neurons and/or glia in close apposition (Bennett *et al.* 1991, Rash *et al.* 2000, 2001, Connors & Long 2004). Gap junctions can be found in both SCN neurons and glia, with the diffusion of labeled molecules (i.e. dye coupling) occurring mostly between homotypic cells located in the same SCN compartment (Jiang *et al.* 1997a, Colwell 2000). Studies using paired intracellular recordings and dye coupling suggest that electrical coupling varies as a function of circadian phase (Colwell 2000, Long *et al.* 2005), which suggests that communication through gap junctions is not a passive process but one that is actively regulated in a dynamic fashion. Further evidence that gap junctions likely influence SCN function stems from pharmacological experiments demonstrating that SCN electrical rhythms become broader, arrhythmic, or bimodal after octanol

or halothane application (Prosser *et al.* 1994, Shinohara *et al.* 2000a,b, Shirakawa *et al.* 2001). These specific changes in waveform are thought to reflect changes in SCN phase relationships caused by altered coupling, although this has yet to be tested directly. Further, the cellular location of these gap junctions remains unclear since these pharmacological inhibitors would be expected to affect both neurons and glia, as well as producing off-target effects. A specific role for neuronal gap junctions is indicated by work demonstrating that SCN electrical coupling is dependent on the expression of the neuron-specific gap junction protein connexin36 (Long *et al.* 2005). Signaling through these gap junctions is thought to influence circadian behavior because mice lacking connexin36 display reduced amplitude of locomotor rhythms under constant darkness (Long *et al.* 2005). Another factor that may modulate circadian behavior via gap junction signaling is the expression of the polysialylated form of neural cell adhesion molecule (Lee *et al.* 1995, Glass *et al.* 2003), although this protein is also involved in synaptic transmission. This work suggests that gap junctions influence SCN function, but additional studies are needed to delineate their precise role.

Astrocytes

While most circadian research has focused on the functional properties of SCN neurons, the SCN also contains glial cells (Guldner 1983). SCN astrocytes can be detected by labeling glial fibrillary acidic protein (GFAP), which displays a daily rhythm that is modulated by light (Lavialle & Serviere 1993, Moriya *et al.* 2000, Gerics *et al.* 2006, Becquet *et al.* 2008, Lindley *et al.* 2008, Canal *et al.* 2009). Like other cell types, astrocytes are intrinsic oscillators that display daily rhythms in metabolic function (Schwartz & Gainer 1977, van den Pol *et al.* 1992, Lavialle & Serviere 1993, Womac *et al.* 2009, Burkeen *et al.* 2011) and clock gene/protein expression (van den Pol *et al.* 1992, Prolo *et al.* 2005, Cheng *et al.* 2009, Yagita *et al.* 2010, Duhart *et al.* 2013). The available evidence points to several ways in which SCN glia and neurons interact (Jackson 2011); however, the precise role of glia in the SCN network is not well characterized. In general, astrocytes provide physical support, release pro-survival factors, influence synaptic clearance, and produce gliotransmitters that interact with presynaptic and postsynaptic receptors (Faissner *et al.* 2010). Anatomical and functional evidence suggests that SCN astrocytes may influence light-induced resetting by

regulating glutamate release from retinal terminals (van den Pol *et al.* 1992, Lavaille & Serviere 1995, Tamada *et al.* 1998, Moriya *et al.* 2000, Lavialle *et al.* 2001, Girardet *et al.* 2010). Interestingly, VIP and AVP neurons in the SCN core and shell compartments display differences in daily rhythms of glial coverage of dendrites, with higher coverage of VIP and AVP neurons during the night and day, respectively (Becquet *et al.* 2008). This suggests that SCN glia may differentially regulate signaling to these two subpopulations across the circadian cycle. Consistent with a glial role in SCN photic responses and/or interneuronal coupling, mice that have a mutation in GFAP have altered locomotor activity rhythms in LL (Moriya *et al.* 2000). More direct evidence for glial involvement in SCN coupling is provided by work demonstrating that SCN electrical rhythms become bimodal in the presence of the glial metabolism antagonist, fluorocitrate (Prosser *et al.* 1994). These changes in the waveform of SCN electrical rhythms are thought to be due to altered communication among SCN neuronal subpopulations (Wang *et al.* 2014), which could be tested further with real-time imaging techniques. Communication between SCN neurons and astrocytes is likely bidirectional since damped rhythms in astrocytes are enhanced by co-culture with SCN explants, but not cortical explants (Prolo *et al.* 2005). Further, VIP influences the phase and amplitude of astrocyte rhythms in a dose-dependent manner (Marpegan *et al.* 2009). Collectively, this work suggests that SCN astrocytes and neurons interact with one another, and future studies are expected to clarify the functional consequences of this relationship.

Conclusions

In many ways, mapping the circuitry of the SCN remains a challenge. Defining the specific role of any given signaling mechanism can be complicated given that multiple processes interact and modulate one another (Shinohara *et al.* 2000b, Itri *et al.* 2004, Haas *et al.* 2011, Wang *et al.* 2014). In addition, it remains a challenge to separate the role of a given neuromodulator in mediating intra-network coupling from its role in processing input and transmitting output. Despite the difficulty in addressing these issues, it remains critical to address how SCN cells integrate the various signals provided by other cell types in the network. The combinatorial effects of SCN coupling factors have yet to be systemically investigated, although some work has addressed this issue in the context of photic signaling (Albers *et al.* 1991, Piggins *et al.* 1995). It is likely

that future technological advances will prove critical for achieving a deeper understanding of SCN circuitry. Going forward, it will be important to define the precise cellular location of receptors for putative coupling factors. Further, whether these receptors are co-expressed spatially and temporally on specific subclasses of SCN neurons should be addressed in future work. Given the potential for rhythms in each component of SCN signaling (e.g. signal release, receptor expression, physical connectivity, and astrocyte function), advanced techniques for imaging cellular connections and responses will continue to be essential for making progress in this area.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

Funding support from the NIH (R01NS091234) and the Whitehall Foundation (2014-12-65) is acknowledged.

References

- Abraham U, Granada AE, Westermark PO, Heine M, Kramer A & Herzel H 2010 Coupling governs entrainment range of circadian clocks. *Molecular Systems Biology* **6** 438. (doi:10.1038/msb.2010.92)
- Abrahamson EE & Moore RY 2001 Suprachiasmatic nucleus in the mouse: retinal innervation, intrinsic organization and efferent projections. *Brain Research* **916** 172–191. (doi:10.1016/S0006-8993(01)02890-6)
- Aggelopoulos NC & Meissl H 2000 Responses of neurones of the rat suprachiasmatic nucleus to retinal illumination under photopic and scotopic conditions. *Journal of Physiology* **523** 211–222. (doi:10.1111/j.1469-7793.2000.t01-1-00211.x)
- Aguilar-Roblero R, Verduzco-Carbajal L, Rodriguez C, Mendez-Franco J, Moran J & de la Mora MP 1993 Circadian rhythmicity in the GABAergic system in the suprachiasmatic nuclei of the rat. *Neuroscience Letters* **157** 199–202. (doi:10.1016/0304-3940(93)90736-5)
- Aida R, Moriya T, Araki M, Akiyama M, Wada K, Wada E & Shibata S 2002 Gastrin-releasing peptide mediates photic entrainable signals to dorsal subsets of suprachiasmatic nucleus via induction of *Period* gene in mice. *Molecular Pharmacology* **61** 26–34. (doi:10.1124/mol.61.1.26)
- Alamilla J, Perez-Burgos A, Quinto D & Aguilar-Roblero R 2014 Circadian modulation of the Cl(–) equilibrium potential in the rat suprachiasmatic nuclei. *BioMed Research International* **2014** 424982. (doi:10.1155/2014/424982)
- Albers HE, Liou SY, Stopa EG & Zoeller RT 1991 Interaction of colocalized neuropeptides: functional significance in the circadian timing system. *Journal of Neuroscience* **11** 846–851.
- Albus H, Vansteensel MJ, Michel S, Block GD & Meijer JH 2005 A GABAergic mechanism is necessary for coupling dissociable ventral and dorsal regional oscillators within the circadian clock. *Current Biology* **15** 886–893. (doi:10.1016/j.cub.2005.03.051)
- An S, Irwin RP, Allen CN, Tsai CA & Herzog ED 2011 Vasoactive intestinal polypeptide requires parallel changes in adenylate cyclase and phospholipase C to entrain circadian rhythms to a predictable

- phase. *Journal of Neurophysiology* **105** 2289–2296. (doi:10.1152/jn.00966.2010)
- An S, Tsai C, Ronecker J, Bayly A & Herzog ED 2012 Spatiotemporal distribution of vasoactive intestinal polypeptide receptor 2 in mouse suprachiasmatic nucleus. *Journal of Comparative Neurology* **520** 2730–2741. (doi:10.1002/cne.23078)
- An S, Harang R, Meeker K, Granados-Fuentes D, Tsai CA, Mazuski C, Kim J, Doyle FJ III, Petzold LR & Herzog ED 2013 A neuropeptide speeds circadian entrainment by reducing intercellular synchrony. *PNAS* **110** E4355–E4361. (doi:10.1073/pnas.1307088110)
- Ananthasubramanian B, Herzog ED & Herzl H 2014 Timing of neuropeptide coupling determines synchrony and entrainment in the mammalian circadian clock. *PLoS Computational Biology* **10** e1003565. (doi:10.1371/journal.pcbi.1003565)
- Antle MC & Silver R 2005 Orchestrating time: arrangements of the brain circadian clock. *Trends in Neurosciences* **28** 145–151. (doi:10.1016/j.tins.2005.01.003)
- Antle MC, Kriegsfeld LJ & Silver R 2005a Signaling within the master clock of the brain: localized activation of mitogen-activated protein kinase by gastrin-releasing peptide. *Journal of Neuroscience* **25** 2447–2454.
- Antle MC, LeSauter J & Silver R 2005b Neurogenesis and ontogeny of specific cell phenotypes within the hamster suprachiasmatic nucleus. *Developmental Brain Research* **157** 8–18. (doi:10.1016/j.devbrainres.2005.02.017)
- Atkins N Jr, Mitchell JW, Romanova EV, Morgan DJ, Cominski TP, Ecker JL, Pinter JE, Sweedler JV & Gillette MU 2010 Circadian integration of glutamatergic signals by little SAAs in novel suprachiasmatic circuits. *PLoS ONE* **5** e12612. (doi:10.1371/journal.pone.0012612)
- Aton SJ & Herzog ED 2005 Come together, right...now: synchronization of rhythms in a mammalian circadian clock. *Neuron* **48** 531–534.
- Aton SJ, Colwell CS, Harnar AJ, Waschek J & Herzog ED 2005 Vasoactive intestinal polypeptide mediates circadian rhythmicity and synchrony in mammalian clock neurons. *Nature Neuroscience* **8** 476–483.
- Aton SJ, Huettner JE, Straume M & Herzog ED 2006 GABA and Gi/o differentially control circadian rhythms and synchrony in clock neurons. *PNAS* **103** 19188–19193. (doi:10.1073/pnas.0607466103)
- Bechtold DA, Brown TM, Luckman SM & Piggins HD 2008 Metabolic rhythm abnormalities in mice lacking VIP-VPAC2 signaling. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **294** R344–R351.
- Becquet D, Girardet C, Guillaumond F, Francois-Bellan AM & Bosler O 2008 Ultrastructural plasticity in the rat suprachiasmatic nucleus. Possible involvement in clock entrainment. *Glia* **56** 294–305. (doi:10.1002/glia.20613)
- Bedont JL & Blackshaw S 2015 Constructing the suprachiasmatic nucleus: a watchmaker's perspective on the central clockworks. *Frontiers in Systems Neuroscience* **9** 74. (doi:10.3389/fnsys.2015.00074)
- Bedont JL, LeGates TA, Slat EA, Byerly MS, Wang H, Hu J, Rupp AC, Qian J, Wong GW, Herzog ED, et al. 2014 *Lhx1* controls terminal differentiation and circadian function of the suprachiasmatic nucleus. *Cell Reports* **7** 609–622. (doi:10.1016/j.celrep.2014.03.060)
- Belenky MA, Sagiv N, Fritschy JM & Yarom Y 2003 Presynaptic and postsynaptic GABA_A receptors in rat suprachiasmatic nucleus. *Neuroscience* **118** 909–923. (doi:10.1016/S0306-4522(03)00062-9)
- Belenky MA, Yarom Y & Pickard GE 2008 Heterogeneous expression of gamma-aminobutyric acid and gamma-aminobutyric acid-associated receptors and transporters in the rat suprachiasmatic nucleus. *Journal of Comparative Neurology* **506** 708–732. (doi:10.1002/cne.21553)
- Belenky MA, Sollars PJ, Mount DB, Alper SL, Yarom Y & Pickard GE 2010 Cell-type specific distribution of chloride transporters in the rat suprachiasmatic nucleus. *Neuroscience* **165** 1519–1537. (doi:10.1016/j.neuroscience.2009.11.040)
- Belle MD, Diekmann CO, Forger DB & Piggins HD 2009 Daily electrical silencing in the mammalian circadian clock. *Science* **326** 281–284. (doi:10.1126/science.1169657)
- Bennett MV, Barrio LC, Bargiello TA, Spray DC, Hertzberg E & Saez JC 1991 Gap junctions: new tools, new answers, new questions. *Neuron* **6** 305–320. (doi:10.1016/0896-6273(91)90241-Q)
- Biello SM 2009 Circadian clock resetting in the mouse changes with age. *Age* **31** 293–303. (doi:10.1007/s11357-009-9102-7)
- Biggs KR & Prosser RA 1998 GABA_B receptor stimulation phase-shifts the mammalian circadian clock in vitro. *Brain Research* **807** 250–254. (doi:10.1016/S0006-8993(98)00820-8)
- Boer GJ, van Esseveldt LE & Rietveld WJ 1998 Cellular requirements of suprachiasmatic nucleus transplants for restoration of circadian rhythm. *Chronobiology International* **15** 551–566. (doi:10.3109/07420529808998707)
- Bormann J 2000 The 'ABC' of GABA receptors. *Trends in Pharmacological Sciences* **21** 16–19. (doi:10.1016/S0165-6147(99)01413-3)
- Bos NP & Mirmiran M 1993 Effects of excitatory and inhibitory amino acids on neuronal discharges in the cultured suprachiasmatic nucleus. *Brain Research Bulletin* **31** 67–72. (doi:10.1016/0361-9230(93)90012-Z)
- Bouskila Y & Dudek FE 1993 Neuronal synchronization without calcium-dependent synaptic transmission in the hypothalamus. *PNAS* **90** 3207–3210. (doi:10.1073/pnas.90.8.3207)
- Brancaccio M, Maywood ES, Chesham JE, Loudon AS & Hastings MH 2013 A G_q-Ca²⁺ axis controls circuit-level encoding of circadian time in the suprachiasmatic nucleus. *Neuron* **78** 714–728. (doi:10.1016/j.neuron.2013.03.011)
- Brown MH & Nunez AA 1989 Vasopressin-deficient rats show a reduced amplitude of the circadian sleep rhythm. *Physiology & Behavior* **46** 759–762.
- Brown TM & Piggins HD 2007 Electrophysiology of the suprachiasmatic circadian clock. *Progress in Neurobiology* **82** 229–255. (doi:10.1016/j.pneurobio.2007.05.002)
- Brown TM & Piggins HD 2009 Spatiotemporal heterogeneity in the electrical activity of suprachiasmatic nuclei neurons and their response to photoperiod. *Journal of Biological Rhythms* **24** 44–54. (doi:10.1177/0748730408327918)
- Brown TM, Hughes AT & Piggins HD 2005 Gastrin-releasing peptide promotes suprachiasmatic nuclei cellular rhythmicity in the absence of vasoactive intestinal polypeptide-VPAC2 receptor signaling. *Journal of Neuroscience* **25** 11155–11164. (doi:10.1523/JNEUROSCI.3821-05.2005)
- Brown TM, Colwell CS, Waschek JA & Piggins HD 2007 Disrupted neuronal activity rhythms in the suprachiasmatic nuclei of vasoactive intestinal polypeptide-deficient mice. *Journal of Neurophysiology* **97** 2553–2558. (doi:10.1152/jn.01206.2006)
- Brown TM, Wynne J, Piggins HD & Lucas RJ 2011 Multiple hypothalamic cell populations encoding distinct visual information. *Journal of Physiology* **589** 1173–1194. (doi:10.1111/jphysiol.2010.199877)
- Buhr ED & Takahashi JS 2013 Molecular components of the mammalian circadian clock. *Handbook of Experimental Pharmacology* **217** 3–27. (doi:10.1007/978-3-642-25950-0)
- Buhr ED, Yoo SH & Takahashi JS 2010 Temperature as a universal resetting cue for mammalian circadian oscillators. *Science* **330** 379–385. (doi:10.1126/science.1195262)
- Burke JF, Womac AD, Earnest DJ & Zoran MJ 2011 Mitochondrial calcium signaling mediates rhythmic extracellular ATP accumulation in suprachiasmatic nucleus astrocytes. *Journal of Neuroscience* **31** 8432–8440. (doi:10.1523/JNEUROSCI.6576-10.2011)
- Butler MP, Rainbow MN, Rodriguez E, Lyon SM & Silver R 2012 Twelve-hour days in the brain and behavior of split hamsters. *European Journal of Neuroscience* **36** 2556–2566. (doi:10.1111/j.1460-9568.2012.08166.x)
- Cagampang FRA, Sheward WJ, Harnar AJ, Piggins HD & Coen CW 1998 Circadian changes in the expression of vasoactive intestinal peptide

- 2 receptor mRNA in the rat suprachiasmatic nuclei. *Molecular Brain Research* **54** 108–112. (doi:10.1016/S0169-328X(97)00327-6)
- Callaway EM & Luo L 2015 Monosynaptic circuit tracing with glycoprotein-deleted rabies viruses. *Journal of Neuroscience* **35** 8979–8985. (doi:10.1523/JNEUROSCI.0409-15.2015)
- Canal MM, Mohammed NM & Rodriguez JJ 2009 Early programming of astrocyte organization in the mouse suprachiasmatic nuclei by light. *Chronobiology International* **26** 1545–1558. (doi:10.3109/07420520903398542)
- Cao R, Robinson B, Xu H, Gkogkas C, Khoutorsky A, Alain T, Yanagiya A, Nevarko T, Liu AC, Amir S, *et al.* 2013 Translational control of entrainment and synchrony of the suprachiasmatic circadian clock by mTOR/4E-BP1 signaling. *Neuron* **79** 712–724. (doi:10.1016/j.neuron.2013.06.026)
- Card JP & Moore RY 1984 The suprachiasmatic nucleus of the golden hamster: immunohistochemical analysis of cell and fiber distribution. *Neuroscience* **13** 415–431. (doi:10.1016/0306-4522(84)90240-9)
- Card JP, Brecha N, Karten HJ & Moore RY 1981 Immunocytochemical localization of vasoactive intestinal polypeptide-containing cells and processes in the suprachiasmatic nucleus of the rat: light and electron microscopic analysis. *Journal of Neuroscience* **1** 1289–1303.
- Cardinali DP & Golombek DA 1998 The rhythmic GABAergic system. *Neurochemical Research* **23** 607–614. (doi:10.1023/A:1022426519297)
- Cassone VM, Speh JC, Card JP & Moore RY 1988 Comparative anatomy of the mammalian hypothalamic suprachiasmatic nucleus. *Journal of Biological Rhythms* **3** 71–91. (doi:10.1177/074873048800300106)
- Castel M & Morris JF 2000 Morphological heterogeneity of the GABAergic network in the suprachiasmatic nucleus, the brain's circadian pacemaker. *Journal of Anatomy* **196** 1–13. (doi:10.1046/j.1469-7580.2000.19610001.x)
- Castel M, Feinstein N, Cohen S & Harari N 1990 Vasopressinergic innervation of the mouse suprachiasmatic nucleus: an immunoelectron microscopic analysis. *Journal of Comparative Neurology* **298** 172–187. (doi:10.1002/cne.902980204)
- Castel M, Morris J & Belenky M 1996 Non-synaptic and dendritic exocytosis from dense-cored vesicles in the suprachiasmatic nucleus. *Neuroreport* **7** 543–547. (doi:10.1097/00001756-199601310-00040)
- Cheng HY, Alvarez-Saavedra M, Dziema H, Choi YS, Li A & Obrietan K 2009 Segregation of expression of *mPeriod* gene homologs in neurons and glia: possible divergent roles of *mPeriod1* and *mPeriod2* in the brain. *Human Molecular Genetics* **18** 3110–3124. (doi:10.1093/hmg/ddp252)
- Choi HJ, Lee CJ, Schroeder A, Kim YS, Jung SH, Kim JS, Kim do Y, Son EJ, Han HC, Hong SK, *et al.* 2008 Excitatory actions of GABA in the suprachiasmatic nucleus. *Journal of Neuroscience* **28** 5450–5459. (doi:10.1523/JNEUROSCI.5750-07.2008)
- Ciarleglio CM, Gamble KL, Axley JC, Strauss BR, Cohen JY, Colwell CS & McMahon DG 2009 Population encoding by circadian clock neurons organizes circadian behavior. *Journal of Neuroscience* **29** 1670–1676. (doi:10.1523/JNEUROSCI.3801-08.2009)
- Colwell CS 2000 Rhythmic coupling among cells in the suprachiasmatic nucleus. *Journal of Neurobiology* **43** 379–388. (doi:10.1002/1097-4695(20000615)43:4<379::aid-neu6>3.0.co;2-0)
- Colwell CS, Michel S, Itri J, Rodriguez W, Tam J, Lelievre V, Hu Z, Liu X & Waschek JA 2003 Disrupted circadian rhythms in VIP- and PHI-deficient mice. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **285** R939–R949.
- Connors BW & Long MA 2004 Electrical synapses in the mammalian brain. *Annual Review of Neuroscience* **27** 393–418. (doi:10.1146/annurev.neuro.26.041002.131128)
- Couvineau A & Laburthe M 2011 VPAC receptors: structure, molecular pharmacology and interaction with accessory proteins. *British Journal of Pharmacology* **166** 42–50. (doi:10.1111/j.1476-5381.2011.01676.x)
- Cutler DJ, Haraura M, Reed HE, Shen S, Sheward WJ, Morrison CF, Marston HM, Harmar AJ & Piggins HD 2003 The mouse VPAC2 receptor confers suprachiasmatic nuclei cellular rhythmicity and responsiveness to vasoactive intestinal polypeptide *in vitro*. *European Journal of Neuroscience* **17** 197–204. (doi:10.1046/j.1460-9568.2003.02425.x)
- Czeisler CA & Klerman EB 1999 Circadian and sleep-dependent regulation of hormone release in humans. *Recent Progress in Hormone Research* **54** 97–130.
- Dardente H, Poirel VJ, Klosen P, Pevet P & Masson-Pevet M 2002 *Per* and neuropeptide expression in the rat suprachiasmatic nuclei: compartmentalization and differential cellular induction by light. *Brain Research* **958** 261–271. (doi:10.1016/S0006-8993(02)03563-1)
- Dardente H, Menet JS, Challet E, Tournier BB, Pevet P & Masson-Pevet M 2004 Daily and circadian expression of neuropeptides in the suprachiasmatic nuclei of nocturnal and diurnal rodents. *Brain Research. Molecular Brain Research* **124** 143–151. (doi:10.1016/j.molbrainres.2004.01.010)
- Davidson AJ, Castanon-Cervantes O, Leise TL, Molyneux PC & Harrington ME 2009 Visualizing jet lag in the mouse suprachiasmatic nucleus and peripheral circadian timing system. *European Journal of Neuroscience* **29** 171–180. (doi:10.1111/j.1460-9568.2008.06534.x)
- De Jeu M & Pennartz C 2002 Circadian modulation of GABA function in the rat suprachiasmatic nucleus: excitatory effects during the night phase. *Journal of Neurophysiology* **87** 834–844.
- Decavel C & Van den Pol AN 1990 GABA: a dominant neurotransmitter in the hypothalamus. *Journal of Comparative Neurology* **302** 1019–1037. (doi:10.1002/cne.903020423)
- DeWoskin D, Myung J, Belle MD, Piggins HD, Takumi T & Forger DB 2015 Distinct roles for GABA across multiple timescales in mammalian circadian timekeeping. *PNAS* **112** E3911–E3919. (doi:10.1073/pnas.1420753112)
- Dragich JM, Loh DH, Wang LM, Vosko AM, Kudo T, Nakamura TJ, Odom IH, Tateyama S, Hagopian A, Waschek JA, *et al.* 2010 The role of the neuropeptides PACAP and VIP in the photic regulation of gene expression in the suprachiasmatic nucleus. *European Journal of Neuroscience* **31** 864–875. (doi:10.1111/j.1460-9568.2010.07119.x)
- Drouyer E, LeSauter J, Hernandez AL & Silver R 2010 Specializations of gastrin-releasing peptide cells of the mouse suprachiasmatic nucleus. *Journal of Comparative Neurology* **518** 1249–1263. (doi:10.1002/cne.22272)
- Dudek FE, Kim YI & Bouskila Y 1993 Electrophysiology of the suprachiasmatic nucleus: synaptic transmission, membrane properties, and neuronal synchronization. *Journal of Biological Rhythms* **8** (Supplement) S33–S37.
- Duhart JM, Leone MJ, Paladino N, Evans JA, Castanon-Cervantes O, Davidson AJ & Golombek DA 2013 Suprachiasmatic astrocytes modulate the circadian clock in response to TNF- α . *Journal of Immunology* **191** 4656–4664. (doi:10.4049/jimmunol.1300450)
- Duncan MJ, Cheng X & Heller KS 1995 Photoperiodic exposure and time of day modulate the expression of *arginine vasopressin* mRNA and *vasoactive intestinal peptide* mRNA in the suprachiasmatic nuclei of Siberian hamsters. *Brain Research. Molecular Brain Research* **32** 181–186. (doi:10.1016/0169-328X(95)00072-Z)
- Earnest DJ, Digiorgio SM & Sladek CD 1991 Effects of tetrodotoxin on the circadian pacemaker mechanism in suprachiasmatic explants *in vitro*. *Brain Research Bulletin* **26** 677–682. (doi:10.1016/0361-9230(91)90160-L)
- Egli M, Bertram R, Sellix MT & Freeman ME 2004 Rhythmic secretion of prolactin in rats: action of oxytocin coordinated by vasoactive intestinal polypeptide of suprachiasmatic nucleus origin. *Endocrinology* **145** 3386–3394. (doi:10.1210/en.2003-1710)
- Ehlen JC & Paul KN 2009 Regulation of light's action in the mammalian circadian clock: role of the extrasynaptic GABA $_A$ receptor. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **296** R1606–R1612. (doi:10.1152/ajpregu.90878.2008)
- Ehlen JC, Novak CM, Karom MC, Gamble KL & Albers HE 2008 Interactions of GABA $_A$ receptor activation and light on *Period* mRNA

- expression in the suprachiasmatic nucleus. *Journal of Biological Rhythms* **23** 16–25. (doi:10.1177/0748730407310785)
- Enoki R, Kuroda S, Ono D, Hasan MT, Ueda T, Honma S & Honma K 2012 Topological specificity and hierarchical network of the circadian calcium rhythm in the suprachiasmatic nucleus. *PNAS* **109** 21498–21503. (doi:10.1073/pnas.1214415110)
- Evans JA & Gorman MR 2016 In synch but not in step: Circadian clock circuits regulating plasticity in daily rhythms. *Neuroscience* **320** 259–290. (doi:10.1016/j.neuroscience.2016.01.072)
- Evans JA, Leise TL, Castanon-Cervantes O & Davidson AJ 2011 Intrinsic regulation of spatiotemporal organization within the suprachiasmatic nucleus. *PLoS ONE* **6** e15869. (doi:10.1371/journal.pone.0015869)
- Evans JA, Pan H, Liu AC & Welsh DK 2012 *Cry1^{-/-}* circadian rhythmicity depends on SCN intercellular coupling. *Journal of Biological Rhythms* **27** 443–452. (doi:10.1177/0748730412461246)
- Evans JA, Leise TL, Castanon-Cervantes O & Davidson AJ 2013 Dynamic interactions mediated by nonredundant signaling mechanisms couple circadian clock neurons. *Neuron* **80** 973–983. (doi:10.1016/j.neuron.2013.08.022)
- Evans JA, Suen TC, Callif BL, Mitchell AS, Castanon-Cervantes O, Baker KM, Kloehn I, Baba K, Teubner BJ, Ehlen JC, *et al.* 2015 Shell neurons of the master circadian clock coordinate the phase of tissue clocks throughout the brain and body. *BMC Biology* **13** 43. (doi:10.1186/s12915-015-0157-x)
- Fahrenkrug J, Georg B, Hannibal J & Jorgensen HL 2012 Altered rhythm of adrenal clock genes, *Star* and serum corticosterone in VIP receptor 2-deficient mice. *Journal of Molecular Neuroscience* **48** 584–596.
- Faissner A, Pyka M, Geissler M, Sobik T, Frischknecht R, Gundelfinger ED & Seidenbecher C 2010 Contributions of astrocytes to synapse formation and maturation – potential functions of the perisynaptic extracellular matrix. *Brain Research Reviews* **63** 26–38. (doi:10.1016/j.brainresrev.2010.01.001)
- Fan J, Zeng H, Olson DP, Huber KM, Gibson JR & Takahashi JS 2015 Vasoactive intestinal polypeptide (VIP)-expressing neurons in the suprachiasmatic nucleus provide sparse GABAergic outputs to local neurons with circadian regulation occurring distal to the opening of postsynaptic GABA_A ionotropic receptors. *Journal of Neuroscience* **35** 1905–1920. (doi:10.1523/JNEUROSCI.2661-14.2015)
- Farajnia S, van Westering TL, Meijer JH & Michel S 2014 Seasonal induction of GABAergic excitation in the central mammalian clock. *PNAS* **111** 9627–9632. (doi:10.1073/pnas.1319820111)
- Farnell YF, Shende VR, Neuendorff N, Allen GC & Earnest DJ 2011 Immortalized cell lines for real-time analysis of circadian pacemaker and peripheral oscillator properties. *European Journal of Neuroscience* **33** 1533–1540. (doi:10.1111/j.1460-9568.2011.07629.x)
- Francl JM, Kaur G & Glass JD 2010 Regulation of vasoactive intestinal polypeptide release in the suprachiasmatic nucleus circadian clock. *Neuroreport* **21** 1055–1059. (doi:10.1097/WNR.0b013e32833fcb4)
- Freeman GM Jr, Krock RM, Aton SJ, Thaben P & Herzog ED 2013 GABA networks destabilize genetic oscillations in the circadian pacemaker. *Neuron* **78** 799–806. (doi:10.1016/j.neuron.2013.04.003)
- Gamble KL, Allen GC, Zhou T & McMahon DG 2007 Gastrin-releasing peptide mediates light-like resetting of the suprachiasmatic nucleus circadian pacemaker through cAMP response element-binding protein and *Per1* activation. *Journal of Neuroscience* **27** 12078–12087. (doi:10.1523/JNEUROSCI.1109-07.2007)
- Gamble KL, Kudo T, Colwell CS & McMahon DG 2011 Gastrin-releasing peptide modulates fast delayed rectifier potassium current in *Per1*-expressing SCN neurons. *Journal of Biological Rhythms* **26** 99–106. (doi:10.1177/0748730410396678)
- Gao B, Fritschy JM & Moore RY 1995 GABA_A-receptor subunit composition in the circadian timing system. *Brain Research* **700** 142–156. (doi:10.1016/0006-8993(95)00944-L)
- Gavrila A, Peng CK, Chan JL, Mietus JE, Goldberger AL & Mantzoros CS 2003 Diurnal and ultradian dynamics of serum adiponectin in healthy men: comparison with leptin, circulating soluble leptin receptor, and cortisol patterns. *Journal of Clinical Endocrinology & Metabolism* **88** 2838–2843.
- Geoghegan D & Carter DA 2008 A novel site of adult doublecortin expression: neuropeptide neurons within the suprachiasmatic nucleus circadian clock. *BMC Neuroscience* **9** 2. (doi:10.1186/1471-2202-9-2)
- Gerhold LM & Wise PM 2006 Vasoactive intestinal polypeptide regulates dynamic changes in astrocyte morphometry: impact on gonadotropin-releasing hormone neurons. *Endocrinology* **147** 2197–2202. (doi:10.1210/en.2005-1262)
- Gerhold LM, Horvath TL & Freeman ME 2001 Vasoactive intestinal peptide fibers innervate neuroendocrine dopaminergic neurons. *Brain Research* **919** 48–56. (doi:10.1016/S0006-8993(01)02993-6)
- Gerics B, Szalay F & Hajos F 2006 Glial fibrillary acidic protein immunoreactivity in the rat suprachiasmatic nucleus: circadian changes and their seasonal dependence. *Journal of Anatomy* **209** 231–237. (doi:10.1111/j.1469-7580.2006.00593.x)
- Gillespie CF, Mintz EM, Marvel CL, Huhman KL & Albers HE 1997 GABA(A) and GABA(B) agonists and antagonists alter the phase-shifting effects of light when microinjected into the suprachiasmatic region. *Brain Research* **759** 181–189. (doi:10.1016/S0006-8993(97)00235-7)
- Gillespie CF, Van Der Beek EM, Mintz EM, Mickley NC, Jasnow AM, Huhman KL & Albers HE 1999 GABAergic regulation of light-induced c-Fos immunoreactivity within the suprachiasmatic nucleus. *Journal of Comparative Neurology* **411** 683–692. (doi:10.1002/(sici)1096-9861(19990906)411:4<683::aid-cne12>3.0.co;2-j)
- Girardet C, Becquet D, Blanchard MP, Francois-Bellan AM & Bosler O 2010 Neuroglial and synaptic rearrangements associated with photic entrainment of the circadian clock in the suprachiasmatic nucleus. *European Journal of Neuroscience* **32** 2133–2142. (doi:10.1111/j.1460-9568.2011.07712.x)
- Glass JD, Watanabe M, Fedorkova L, Shen H, Ungers G & Rutishauser U 2003 Dynamic regulation of polysialylated neural cell adhesion molecule in the suprachiasmatic nucleus. *Neuroscience* **117** 203–211. (doi:10.1016/S0306-4522(02)00817-5)
- Glazer R & Gozes I 1994 Diurnal oscillation in vasoactive intestinal peptide gene expression independent of environmental light entraining. *Brain Research* **644** 164–167. (doi:10.1016/0006-8993(94)90360-3)
- Grahn DA, Miller JD, Hough VS & Heller HC 1994 Persistence of circadian rhythmicity in hibernating ground squirrels. *American Journal of Physiology* **266** R1251–R1258.
- Gribkoff VK, Pieschl RL, Wisialowski TA, Park WK, Strecker GJ, de Jeu MT, Pennartz CM & Dudek FE 1999 A reexamination of the role of GABA in the mammalian suprachiasmatic nucleus. *Journal of Biological Rhythms* **14** 126–130. (doi:10.1177/074873099129000515)
- Groblewski TA, Nunez AA & Gold RM 1981 Circadian rhythms in vasopressin deficient rats. *Brain Research Bulletin* **6** 125–130. (doi:10.1016/S0361-9230(81)80036-6)
- Guldner FH 1983 Numbers of neurons and astroglial cells in the suprachiasmatic nucleus of male and female rats. *Experimental Brain Research* **50** 373–376.
- Haas JS, Zavala B & Landisman CE 2011 Activity-dependent long-term depression of electrical synapses. *Science* **334** 389–393. (doi:10.1126/science.1207502)
- Hamada T, LeSauter J, Venuti JM & Silver R 2001 Expression of *Period* genes: rhythmic and nonrhythmic compartments of the suprachiasmatic nucleus pacemaker. *Journal of Neuroscience* **21** 7742–7750.
- Hamada T, Antle MC & Silver R 2004 Temporal and spatial expression patterns of canonical clock genes and clock-controlled genes in the suprachiasmatic nucleus. *European Journal of Neuroscience* **19** 1741–1748. (doi:10.1111/j.1460-9568.2004.03275.x)
- Han S, Yu FH, Schwartz MD, Linton JD, Bosma MM, Hurley JB, Catterall WA & de la Iglesia HO 2012 Na(V)1.1 channels are critical for intercellular communication in the suprachiasmatic nucleus and

- for normal circadian rhythms. *PNAS* **109** E368–E377. (doi:10.1073/pnas.1115729109)
- Hannibal J, Hsiung HM & Fahrenkrug J 2011 Temporal phasing of locomotor activity, heart rate rhythmicity, and core body temperature is disrupted in VIP receptor 2-deficient mice. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* **300** R519–R530.
- Harmar AJ 2003 An essential role for peptidergic signalling in the control of circadian rhythms in the suprachiasmatic nuclei. *Journal of Neuroendocrinology* **15** 335–338. (doi:10.1046/j.1365-2826.2003.01005.x)
- Harmar AJ, Arimura A, Gozes I, Journot L, Laburthe M, Pisegna JR, Rawlings SR, Robberecht P, Said SI, Sreedharan SP, *et al.* 1998 International union of pharmacology. XVIII. Nomenclature of receptors for vasoactive intestinal peptide and pituitary adenylate cyclase-activating polypeptide. *Pharmacological Reviews* **50** 265–270.
- Harmar AJ, Marston HM, Shen S, Spratt C, West KM, Sheward WJ, Morrison CF, Dorin JR, Piggins HD, Reubi JC, *et al.* 2002 The VPAC(2) receptor is essential for circadian function in the mouse suprachiasmatic nuclei. *Cell* **109** 497–508. (doi:10.1016/S0092-8674(02)00736-5)
- Hazlerigg DG, Ebling FJ & Johnston JD 2005 Photoperiod differentially regulates gene expression rhythms in the rostral and caudal SCN. *Current Biology* **15** R449–R450.
- Herzog ED, Takahashi JS & Block GD 1998 Clock controls circadian period in isolated suprachiasmatic nucleus neurons. *Nature Neuroscience* **1** 708–713. (doi:10.1038/3708)
- Herzog ED, Aton SJ, Numano R, Sakaki Y & Tei H 2004 Temporal precision in the mammalian circadian system: a reliable clock from less reliable neurons. *Journal of Biological Rhythms* **19** 35–46. (doi:10.1177/0748730403260776)
- Honma S, Shirakawa T, Katsuno Y, Namihira M & Honma K 1998 Circadian periods of single suprachiasmatic neurons in rats. *Neuroscience Letters* **250** 157–160. (doi:10.1016/S0304-3940(98)00464-9)
- Hughes AT, Fahey B, Cutler DJ, Coogan AN & Piggins HD 2004 Aberrant gating of photic input to the suprachiasmatic pacemaker of mice lacking the VPAC2 receptor. *Journal of Neuroscience* **24** 3522–3526. (doi:10.1523/JNEUROSCI.5345-03.2004)
- Hughes AT, Guilding C, Lennox L, Samuels RE, McMahon DG & Piggins HD 2008 Live imaging of altered *Period1* expression in the suprachiasmatic nuclei of *Vipr2^{-/-}* mice. *Journal of Neurochemistry* **106** 1646–1657. (doi:10.1111/j.1471-4159.2008.05520.x)
- Huhman KL, Hennessey AC & Albers HE 1996 Rhythms of *glutamic acid decarboxylase* mRNA in the suprachiasmatic nucleus. *Journal of Biological Rhythms* **11** 311–316. (doi:10.1177/074873049601100404)
- Hundahl CA, Fahrenkrug J, Hay-Schmidt A, Georg B, Faltoft B & Hannibal J 2012 Circadian behaviour in neuroglobin deficient mice. *PLoS ONE* **7** e34462. (doi:10.1371/journal.pone.0034462)
- Ikeda Y, Kumagai H, Skach A, Sato M & Yanagisawa M 2013 Modulation of circadian glucocorticoid oscillation via adrenal opioid-CXCR7 signaling alters emotional behavior. *Cell* **155** 1323–1336. (doi:10.1016/j.cell.2013.10.052)
- Inagaki N, Honma S, Ono D, Tanahashi Y & Honma K 2007 Separate oscillating cell groups in mouse suprachiasmatic nucleus couple photoperiodically to the onset and end of daily activity. *PNAS* **104** 7664–7669. (doi:10.1073/pnas.0607713104)
- Ingram CD, Ciobanu R, Coculescu IL, Tanasescu R, Coculescu M & Mihai R 1998 Vasopressin neurotransmission and the control of circadian rhythms in the suprachiasmatic nucleus. *Progress in Brain Research* **119** 351–364.
- Irwin RP & Allen CN 2009 GABAergic signaling induces divergent neuronal Ca²⁺ responses in the suprachiasmatic nucleus network. *European Journal of Neuroscience* **30** 1462–1475. (doi:10.1111/j.1460-9568.2009.06944.x)
- Irwin RP & Allen CN 2010 Neuropeptide-mediated calcium signaling in the suprachiasmatic nucleus network. *European Journal of Neuroscience* **32** 1497–1506. (doi:10.1111/j.1460-9568.2010.07411.x)
- Irwin RP & Allen CN 2013 Simultaneous electrophysiological recording and calcium imaging of suprachiasmatic nucleus neurons. *Journal of Visualized Experiments* **82** 50794. (doi:10.3791/50794)
- Isobe Y & Nishino H 1998 AVP rhythm in the suprachiasmatic nucleus in relation to locomotor activity under constant light. *Peptides* **19** 827–832. (doi:10.1016/S0196-9781(98)00021-7)
- Itri J & Colwell CS 2003 Regulation of inhibitory synaptic transmission by vasoactive intestinal peptide (VIP) in the mouse suprachiasmatic nucleus. *Journal of Neurophysiology* **90** 1589–1597. (doi:10.1152/jn.00332.2003)
- Itri J, Michel S, Waschek JA & Colwell CS 2004 Circadian rhythm in inhibitory synaptic transmission in the mouse suprachiasmatic nucleus. *Journal of Neurophysiology* **92** 311–319. (doi:10.1152/jn.01078.2003)
- Jackson FR 2011 Glial cell modulation of circadian rhythms. *Glia* **59** 1341–1350. (doi:10.1002/glia.21097)
- Jagota A, de la Iglesia HO & Schwartz WJ 2000 Morning and evening circadian oscillations in the suprachiasmatic nucleus *in vitro*. *Nature Neuroscience* **3** 372–376. (doi:10.1038/73943)
- Jensen RT, Battey JF, Spindel ER & Benya RV 2008 International union of pharmacology. LXVIII. Mammalian bombesin receptors: nomenclature, distribution, pharmacology, signaling, and functions in normal and disease states. *Pharmacological Reviews* **60** 1–42. (doi:10.1124/pr.107.07108)
- Jiang Z-G, Yang Y-Q & Allen CN 1997a Tracer and electrical coupling of rat suprachiasmatic nucleus neurons. *Neuroscience* **77** 1059–1066.
- Jiang ZG, Yang Y, Liu ZP & Allen CN 1997b Membrane properties and synaptic inputs of suprachiasmatic nucleus neurons in rat brain slices. *Journal of Physiology* **499** 141–159.
- Jiao YY, Lee TM & Rusak B 1999 Photic responses of suprachiasmatic area neurons in diurnal degus (*Octodon degus*) and nocturnal rats (*Rattus norvegicus*). *Brain Research* **817** 93–103. (doi:10.1016/S0006-8993(98)01218-9)
- Jin X, Shearman LP, Weaver DR, Zylka MJ, de Vries GJ & Reppert SM 1999 A molecular mechanism regulating rhythmic output from the suprachiasmatic circadian clock. *Cell* **96** 57–68. (doi:10.1016/S0092-8674(00)80959-9)
- Jobst EE & Allen CN 2002 Calbindin neurons in the hamster suprachiasmatic nucleus do not exhibit a circadian variation in spontaneous firing rate. *European Journal of Neuroscience* **16** 2469–2474. (doi:10.1046/j.1460-9568.2002.02309.x)
- Jobst EE, Robinson DW & Allen CN 2004 Potential pathways for intercellular communication within the calbindin subnucleus of the hamster suprachiasmatic nucleus. *Neuroscience* **123** 87–99. (doi:10.1016/j.neuroscience.2003.08.059)
- Kalamatianos T, Kallo I & Coen CW 2004a Ageing and the diurnal expression of the mRNAs for vasopressin and for the V1a and V1b vasopressin receptors in the suprachiasmatic nucleus of male rats. *Journal of Neuroendocrinology* **16** 493–501.
- Kalamatianos T, Kallo I, Piggins HD & Coen CW 2004b Expression of VIP and/or PACAP receptor mRNA in peptide synthesizing cells within the suprachiasmatic nucleus of the rat and in its efferent target sites. *Journal of Comparative Neurology* **475** 19–35.
- Kallo I, Kalamatianos T, Piggins HD & Coen CW 2004a Ageing and the diurnal expression of mRNAs for vasoactive intestinal peptide and for the VPAC2 and PAC1 receptors in the suprachiasmatic nucleus of male rats. *Journal of Neuroendocrinology* **16** 758–766.
- Kallo I, Kalamatianos T, Wiltshire N, Shen S, Sheward WJ, Harmar AJ & Coen CW 2004b Transgenic approach reveals expression of the VPAC2 receptor in phenotypically defined neurons in the mouse suprachiasmatic nucleus and in its efferent target sites. *European Journal of Neuroscience* **19** 2201–2211.

- Kalsbeek A & Buijs RM 2002 Output pathways of the mammalian suprachiasmatic nucleus: coding circadian time by transmitter selection and specific targeting. *Cell and Tissue Research* **309** 109–118. (doi:10.1007/s00441-002-0577-0)
- Kalsbeek A, Teclemariam-Mesbah R & Pevet P 1993 Efferent projections of the suprachiasmatic nucleus in the golden hamster (*Mesocricetus auratus*). *Journal of Comparative Neurology* **332** 293–314. (doi:10.1002/cne.903320304)
- Kalsbeek A, Perreau-Lenz S & Buijs RM 2006 A network of (autonomic) clock outputs. *Chronobiology International* **23** 521–535. (doi:10.1080/07420520600651073)
- Kalsbeek A, Fliers E, Hofman MA, Swaab DF & Buijs RM 2010 Vasopressin and the output of the hypothalamic biological clock. *Journal of Neuroendocrinology* **22** 362–372. (doi:10.1111/j.1365-2826.2010.01956.x)
- Karatsoreos IN, Romeo RD, McEwen BS & Silver R 2006 Diurnal regulation of the gastrin-releasing peptide receptor in the mouse circadian clock. *European Journal of Neuroscience* **23** 1047–1053. (doi:10.1111/j.1460-9568.2006.04633.x)
- Kawakami F, Okamura H, Tamada Y, Maebayashi Y, Fukui K & Ibata Y 1997 Loss of day-night differences in *VIP* mRNA levels in the suprachiasmatic nucleus of aged rats. *Neuroscience Letters* **222** 99–102. (doi:10.1016/S0304-3940(97)13355-9)
- Kawamoto K, Nagano M, Kanda F, Chihara K, Shigeyoshi Y & Okamura H 2003 Two types of *VIP* neuronal components in rat suprachiasmatic nucleus. *Journal of Neuroscience Research* **74** 852–857. (doi:10.1002/jnr.10751)
- Kim YI & Dudek FE 1992 Intracellular electrophysiological study of suprachiasmatic nucleus neurons in rodents: inhibitory synaptic mechanisms. *Journal of Physiology* **458** 247–260. (doi:10.1113/jphysiol.1992.sp019416)
- King VM, Chahad-Ehlers S, Shen S, Harmar AJ, Maywood ES & Hastings MH 2003 A hVIPR transgene as a novel tool for the analysis of circadian function in the mouse suprachiasmatic nucleus. *European Journal of Neuroscience* **17** 822–832. (doi:10.1046/j.1460-9568.2003.02487.x)
- Klein DC, Moore RY & Reppert SM Eds 1991 *Suprachiasmatic Nucleus: The Mind's Clock*. New York, NY, USA: University Oxford Press.
- Ko CH, Yamada YR, Welsh DK, Buhr ED, Liu AC, Zhang EE, Ralph MR, Kay SA, Forger DB & Takahashi JS 2010 Emergence of noise-induced oscillations in the central circadian pacemaker. *PLoS Biology* **8** e1000513. (doi:10.1371/journal.pbio.1000513)
- Koutkia P, Schurgin S, Berry J, Breu J, Lee BS, Klibanski A & Grinspoon S 2005 Reciprocal changes in endogenous ghrelin and growth hormone during fasting in healthy women. *American Journal of Physiology: Endocrinology and Metabolism* **289** E814–E822.
- Krajnak K, Kashon ML, Rosewell KL & Wise PM 1998a Aging alters the rhythmic expression of vasoactive intestinal polypeptide mRNA but not arginine vasopressin mRNA in the suprachiasmatic nuclei of female rats. *Journal of Neuroscience* **18** 4767–4774.
- Krajnak K, Kashon ML, Rosewell KL & Wise PM 1998b Sex differences in the daily rhythm of vasoactive intestinal polypeptide but not arginine vasopressin messenger ribonucleic acid in the suprachiasmatic nuclei. *Endocrinology* **139** 4189–4196.
- Kudo T, Tahara Y, Gamble KL, McMahon DG, Block GD & Colwell CS 2013 Vasoactive intestinal peptide produces long-lasting changes in neural activity in the suprachiasmatic nucleus. *Journal of Neurophysiology* **110** 1097–1106. (doi:10.1152/jn.00114.2013)
- Kuhlman SJ 2007 Biological rhythms workshop IB: neurophysiology of SCN pacemaker function. *Cold Spring Harbor Symposia on Quantitative Biology* **72** 21–33. (doi:10.1101/sqb.2007.72.061)
- Kuhlman SJ, Quintero JE & McMahon DG 2000 GFP fluorescence reports *Period 1* circadian gene regulation in the mammalian biological clock. *Neuroreport* **11** 1479–1482. (doi:10.1097/00001756-200005150-00023)
- Kuhlman SJ, Silver R, Le Sauter J, Bult-Ito A & McMahon DG 2003 Phase resetting light pulses induce *Per1* and persistent spike activity in a subpopulation of biological clock neurons. *Journal of Neuroscience* **23** 1441–1450.
- Lavaille M & Serviere J 1993 Circadian fluctuations in GFAP distribution in the Syrian hamster suprachiasmatic nucleus. *Neuroreport* **4** 1243–1246. (doi:10.1097/00001756-199309000-00008)
- Lavaille M & Serviere J 1995 Developmental study in the circadian clock of the golden hamster: a putative role of astrocytes. *Developmental Brain Research* **86** 275–282. (doi:10.1016/0165-3806(95)00039-G)
- Lavaille M, Begue A, Papillon C & Vilaplana J 2001 Modifications of retinal afferent activity induce changes in astroglial plasticity in the hamster circadian clock. *Glia* **34** 88–100. (doi:10.1002/glia.1044)
- Leak RK & Moore RY 2001 Topographic organization of suprachiasmatic nucleus projection neurons. *Journal of Comparative Neurology* **433** 312–334. (doi:10.1002/cne.1142)
- Lee W, Watanabe M & Glass JD 1995 Photoperiod affects the expression of neural cell adhesion molecule and polysialic acid in the hypothalamus of the Siberian hamster. *Brain Research* **690** 64–72. (doi:10.1016/0006-8993(95)00588-H)
- Lee HS, Nelms JL, Nguyen M, Silver R & Lehman MN 2003 The eye is necessary for a circadian rhythm in the suprachiasmatic nucleus. *Nature Neuroscience* **6** 111–112. (doi:10.1038/nn1006)
- Lee ML, Swanson BE & de la Iglesia HO 2009 Circadian timing of REM sleep is coupled to an oscillator within the dorsomedial suprachiasmatic nucleus. *Current Biology* **19** 848–852. (doi:10.1016/j.cub.2009.03.051)
- Lee JE, Zamborg L, Southey BR, Atkins N Jr, Mitchell JW, Li M, Gillette MU, Kelleher NL & Sweedler JV 2013 Quantitative peptidomics for discovery of circadian-related peptides from the rat suprachiasmatic nucleus. *Journal of Proteome Research* **12** 585–593. (doi:10.1021/pr300605p)
- Lee IT, Chang AS, Manandhar M, Shan Y, Fan J, Izumo M, Ikeda Y, Motoike T, Dixon S, Seinfeld JE, et al. 2015 Neuromedin s-producing neurons act as essential pacemakers in the suprachiasmatic nucleus to couple clock neurons and dictate circadian rhythms. *Neuron* **85** 1086–1102. (doi:10.1016/j.neuron.2015.02.006)
- Leise TL, Wang CW, Gitis PJ & Welsh DK 2012 Persistent cell-autonomous circadian oscillations in fibroblasts revealed by six-week single-cell imaging of PER2::LUC bioluminescence. *PLoS ONE* **7** e33334. (doi:10.1371/journal.pone.0033334)
- Lesauter J, Silver R, Cloues R & Witkovsky P 2011 Light exposure induces short- and long-term changes in the excitability of retinorecipient neurons in suprachiasmatic nucleus. *Journal of Neurophysiology* **106** 576–588. (doi:10.1152/jn.00060.2011)
- Li JD, Burton KJ, Zhang C, Hu SB & Zhou QY 2009 Vasopressin receptor V1a regulates circadian rhythms of locomotor activity and expression of clock-controlled genes in the suprachiasmatic nuclei. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **296** R824–R830. (doi:10.1152/ajpregu.90463.2008)
- Lindley J, Deurveilher S, Rusak B & Semba K 2008 Transforming growth factor- α and glial fibrillary acidic protein in the hamster circadian system: daily profile and cellular localization. *Brain Research* **1197** 94–105. (doi:10.1016/j.brainres.2007.12.053)
- Liou SY & Albers HE 1989 Single unit response of suprachiasmatic neurons to arginine vasopressin (AVP) is mediated by a V1-like receptor in the hamster. *Brain Research* **477** 336–343. (doi:10.1016/0006-8993(89)91424-8)
- Liou SY, Shibata S, Albers HE & Ueki S 1990 Effects of GABA and anxiolytics on the single unit discharge of suprachiasmatic neurons in rat hypothalamic slices. *Brain Research Bulletin* **25** 103–107. (doi:10.1016/0361-9230(90)90259-3)
- Liu C & Reppert SM 2000 GABA synchronizes clock cells within the suprachiasmatic circadian clock. *Neuron* **25** 123–128. (doi:10.1016/S0896-6273(00)80876-4)
- Liu C, Weaver DR, Strogatz SH & Reppert SM 1997 Cellular construction of a circadian clock: period determination in the suprachiasmatic nuclei. *Cell* **91** 855–860. (doi:10.1016/S0092-8674(00)80473-0)

- Liu AC, Welsh DK, Ko CH, Tran HG, Zhang EE, Priest AA, Buhr ED, Singer O, Meeker K, Verma IM, *et al.* 2007 Intercellular coupling confers robustness against mutations in the SCN circadian clock network. *Cell* **129** 605–616. (doi:10.1016/j.cell.2007.02.047)
- Locke JC, Westermark PO, Kramer A & Herzel H 2008 Global parameter search reveals design principles of the mammalian circadian clock. *BMC Systems Biology* **2** 22. (doi:10.1186/1752-0509-2-22)
- Loh DH, Dragich JM, Kudo T, Schroeder AM, Nakamura TJ, Waschek JA, Block GD & Colwell CS 2011 Effects of vasoactive intestinal peptide genotype on circadian gene expression in the suprachiasmatic nucleus and peripheral organs. *Journal of Biological Rhythms* **26** 200–209. (doi:10.1177/0748730411401740)
- Loh DH, Kuljis DA, Azuma L, Wu Y, Truong D, Wang HB & Colwell CS 2014 Disrupted reproduction, estrous cycle, and circadian rhythms in female mice deficient in vasoactive intestinal peptide. *Journal of Biological Rhythms* **29** 355–369. (doi:10.1177/0748730414549767)
- Lokshin M, LeSauter J & Silver R 2015 Selective distribution of retinal input to mouse SCN revealed in analysis of sagittal sections. *Journal of Biological Rhythms* **30** 251–257. (doi:10.1177/0748730415584058)
- Long MA, Jutras MJ, Connors BW & Burwell RD 2005 Electrical synapses coordinate activity in the suprachiasmatic nucleus. *Nature Neuroscience* **8** 61–66. (doi:10.1038/nn1361)
- Low-Zeddies SS & Takahashi JS 2001 Chimera analysis of the *Clock* mutation in mice shows that complex cellular integration determines circadian behavior. *Cell* **105** 25–42. (doi:10.1016/S0092-8674(01)00294-X)
- Lucassen EA, van Diepen HC, Houben T, Michel S, Colwell CS & Meijer JH 2012 Role of vasoactive intestinal peptide in seasonal encoding by the suprachiasmatic nucleus clock. *European Journal of Neuroscience* **35** 1466–1474. (doi:10.1111/j.1460-9568.2012.08054.x)
- Ludwig M & Leng G 2006 Dendritic peptide release and peptide-dependent behaviours. *Nature Reviews Neuroscience* **7** 126–136. (doi:10.1038/nrn1845)
- Mackey SR 2007 Biological rhythms workshop IA: molecular basis of rhythms generation. *Cold Spring Harbor Symposia on Quantitative Biology* **72** 7–19. (doi:10.1101/sqb.2007.72.060)
- Mahoney MM, Ramanathan C, Hagenauer MH, Thompson RC, Smale L & Lee T 2009 Daily rhythms and sex differences in vasoactive intestinal polypeptide, VIPR2 receptor and arginine vasopressin mRNA in the suprachiasmatic nucleus of a diurnal rodent, *Arvicanthis niloticus*. *European Journal of Neuroscience* **30** 1537–1543. (doi:10.1111/j.1460-9568.2009.06936.x)
- Marpegan L, Krall TJ & Herzog ED 2009 Vasoactive intestinal polypeptide entrains circadian rhythms in astrocytes. *Journal of Biological Rhythms* **24** 135–143. (doi:10.1177/0748730409332042)
- Marty A & Llano I 2005 Excitatory effects of GABA in established brain networks. *Trends in Neurosciences* **28** 284–289. (doi:10.1016/j.tins.2005.04.003)
- Mason R, Biello SM & Harrington ME 1991 The effects of GABA and benzodiazepines on neurones in the suprachiasmatic nucleus (SCN) of Syrian hamsters. *Brain Research* **552** 53–57. (doi:10.1016/0006-8993(91)90659-J)
- Maybauer MO, Maybauer DM, Enkhbaatar P & Traber DL 2008 Physiology of the vasopressin receptors. *Best Practice & Research Clinical Anaesthesiology* **22** 253–263.
- Maywood ES, Reddy AB, Wong GK, O'Neill JS, O'Brien JA, McMahon DG, Harmar AJ, Okamura H & Hastings MH 2006 Synchronization and maintenance of timekeeping in suprachiasmatic circadian clock cells by neuropeptidergic signaling. *Current Biology* **16** 599–605. (doi:10.1016/j.cub.2006.02.023)
- Maywood ES, Chesham JE, O'Brien JA & Hastings MH 2011 A diversity of paracrine signals sustains molecular circadian cycling in suprachiasmatic nucleus circuits. *PNAS* **108** 14306–14311. (doi:10.1073/pnas.1101767108)
- Maywood ES, Drynan L, Chesham JE, Edwards MD, Dardente H, Fustin JM, Hazlerigg DG, O'Neill JS, Codner GF, Smyllie NJ, *et al.* 2013 Analysis of core circadian feedback loop in suprachiasmatic nucleus of *mCry1-luc* transgenic reporter mouse. *PNAS* **110** 9547–9552. (doi:10.1073/pnas.1220894110)
- McArthur AJ, Coogan AN, Ajpru S, Sugden D, Biello SM & Piggins HD 2000 Gastrin-releasing peptide phase-shifts suprachiasmatic nuclei neuronal rhythms *in vitro*. *Journal of Neuroscience* **20** 5496–5502.
- Meijer JH, Groos GA & Rusak B 1986 Luminance coding in a circadian pacemaker: the suprachiasmatic nucleus of the rat and the hamster. *Brain Research* **382** 109–118. (doi:10.1016/0006-8993(86)90117-4)
- Meijer JH, Colwell CS, Rohling JH, Houben T & Michel S 2012 Dynamic neuronal network organization of the circadian clock and possible deterioration in disease. *Progress in Brain Research* **199** 143–162. (doi:10.1016/B978-0-444-59427-3.00009-5)
- Menaker M 1961 The free running period of the bat clock; seasonal variations at low body temperature. *Journal of Cellular and Comparative Physiology* **57** 81–86. (doi:10.1002/jcp.1030570204)
- Meyer-Spasche A & Piggins HD 2004 Vasoactive intestinal polypeptide phase-advances the rat suprachiasmatic nuclei circadian pacemaker *in vitro* via protein kinase A and mitogen-activated protein kinase. *Neuroscience Letters* **358** 91–94. (doi:10.1016/j.neulet.2003.12.114)
- Michel S & Colwell CS 2001 Cellular communication and coupling within the suprachiasmatic nucleus. *Chronobiology International* **18** 579–600. (doi:10.1081/CBI-100106074)
- Michel S, Marek R, Vanderleest HT, Vansteensel MJ, Schwartz WJ, Colwell CS & Meijer JH 2013 Mechanism of bilateral communication in the suprachiasmatic nucleus. *European Journal of Neuroscience* **37** 964–971. (doi:10.1111/ejn.12109)
- Mieda M, Ono D, Hasegawa E, Okamoto H, Honma K, Honma S & Sakurai T 2015 Cellular clocks in AVP neurons of the SCN are critical for interneuronal coupling regulating circadian behavior rhythm. *Neuron* **85** 1103–1116. (doi:10.1016/j.neuron.2015.02.005)
- Mihai R, Juss TS & Ingram CD 1994 Suppression of suprachiasmatic nucleus neurone activity with a vasopressin receptor antagonist: possible role for endogenous vasopressin in circadian activity cycles *in vitro*. *Neuroscience Letters* **179** 95–99. (doi:10.1016/0304-3940(94)90943-1)
- Miller BH, Olson SL, Levine JE, Turek FW, Horton TH & Takahashi JS 2006 Vasopressin regulation of the proestrous luteinizing hormone surge in wild-type and Clock mutant mice. *Biology of Reproduction* **75** 778–784. (doi:10.1095/biolreprod.106.052845)
- Mintz EM, Jasnow AM, Gillespie CF, Huhman KL & Albers HE 2002 GABA interacts with photic signaling in the suprachiasmatic nucleus to regulate circadian phase shifts. *Neuroscience* **109** 773–778. (doi:10.1016/S0306-4522(01)00519-X)
- Moga MM & Moore RY 1997 Organization of neural inputs to the suprachiasmatic nucleus in the rat. *Journal of Comparative Neurology* **389** 508–534. (doi:10.1002/(SICI)1096-9861(19971222)389:3<508::AID-CNE11>3.0.CO;2-H)
- Mohawk JA & Takahashi JS 2011 Cell autonomy and synchrony of suprachiasmatic nucleus circadian oscillators. *Trends in Neurosciences* **34** 349–358. (doi:10.1016/j.tins.2011.05.003)
- Mohawk JA, Green CB & Takahashi JS 2012 Central and peripheral circadian clocks in mammals. *Annual Review of Neuroscience* **35** 445–462. (doi:10.1146/annurev-neuro-060909-153128)
- Moore RY 1991 Disorders of circadian function and the human circadian timing system. In *Suprachiasmatic Nucleus: The Mind's Clock*. Eds DC Klein, RY Moore & SM Reppert. New York, NY, USA: University Oxford Press.
- Moore RY & Silver R 1998 Suprachiasmatic nucleus organization. *Chronobiology International* **15** 475–487. (doi:10.3109/07420529808998703)
- Moore RY & Speh JC 1993 GABA is the principal neurotransmitter of the circadian system. *Neuroscience Letters* **150** 112–116. (doi:10.1016/0304-3940(93)90120-A)
- Moore RY, Speh JC & Leak RK 2002 Suprachiasmatic nucleus organization. *Cell and Tissue Research* **309** 89–98. (doi:10.1007/s00441-002-0575-2)

- Morin LP 2007 SCN organization reconsidered. *Journal of Biological Rhythms* **22** 3–13. (doi:10.1177/0748730406296749)
- Morin LP 2012 Neuroanatomy of the extended circadian rhythm system. *Experimental Neurology* **243** 4–20. (doi:10.1016/j.expneurol.2012.06.026)
- Morin LP & Allen CN 2006 The circadian visual system, 2005. *Brain Research. Brain Research Reviews* **51** 1–60. (doi:10.1016/j.brainresrev.2005.08.003)
- Moriya T, Yoshinobu Y, Kouzu Y, Katoh A, Gomi H, Ikeda M, Yoshioka T, Itohara S & Shibata S 2000 Involvement of glial fibrillary acidic protein (GFAP) expressed in astroglial cells in circadian rhythm under constant lighting conditions in mice. *Journal of Neuroscience Research* **60** 212–218. (doi:10.1002/(sici)1097-4547(20000415)60:2<212::aid-jnr10>3.0.co;2-p)
- Myung J, Hong S, Hatanaka F, Nakajima Y, De Schutter E & Takumi T 2012 Period coding of *Bmal1* oscillators in the suprachiasmatic nucleus. *Journal of Neuroscience* **32** 8900–8918. (doi:10.1523/JNEUROSCI.5586-11.2012)
- Myung J, Hong S, DeWoskin D, De Schutter E, Forger DB & Takumi T 2015 GABA-mediated repulsive coupling between circadian clock neurons in the SCN encodes seasonal time. *PNAS* **112** E3920–E3929. (doi:10.1073/pnas.1421200112)
- Nagano M, Adachi A, Nakahama K, Nakamura T, Tamada M, Meyer-Bernstein E, Sehgal A & Shigeyoshi Y 2003 An abrupt shift in the day/night cycle causes desynchrony in the mammalian circadian center. *Journal of Neuroscience* **23** 6141–6151.
- Nagoshi E, Saini C, Bauer C, Laroche T, Naef F & Schibler U 2004 Circadian gene expression in individual fibroblasts: cell-autonomous and self-sustained oscillators pass time to daughter cells. *Cell* **119** 693–705. (doi:10.1016/j.cell.2004.11.015)
- Naito E, Watanabe T, Tei H, Yoshimura T & Ebihara S 2008 Reorganization of the suprachiasmatic nucleus coding for day length. *Journal of Biological Rhythms* **23** 140–149. (doi:10.1177/0748730408314572)
- Nakamura W, Honma S, Shirakawa T & Honma K 2001 Regional pacemakers composed of multiple oscillator neurons in the rat suprachiasmatic nucleus. *European Journal of Neuroscience* **14** 666–674. (doi:10.1046/j.0953-816x.2001.01684.x)
- Nakamura W, Honma S, Shirakawa T & Honma K 2002 Clock mutation lengthens the circadian period without damping rhythms in individual SCN neurons. *Nature Neuroscience* **5** 399–400. (doi:10.1080/1028415021000055943)
- Nakamura TJ, Fujimura K, Ebihara S & Shinohara K 2004 Light response of the neuronal firing activity in the suprachiasmatic nucleus of mice. *Neuroscience Letters* **371** 244–248. (doi:10.1016/j.neulet.2004.09.005)
- Nakamura W, Yamazaki S, Takasu NN, Mishima K & Block GD 2005 Differential response of *Period1* expression within the suprachiasmatic nucleus. *Journal of Neuroscience* **25** 5481–5487. (doi:10.1523/JNEUROSCI.0889-05.2005)
- Natalucci G, Riedl S, Gleiss A, Zidek T & Frisch H 2005 Spontaneous 24-h ghrelin secretion pattern in fasting subjects: maintenance of a meal-related pattern. *European Journal of Endocrinology* **152** 845–850. (doi:10.1530/eje.1.01919)
- Naum OG, Fernanda Rubio M & Golombek DA 2001 Rhythmic variation in gamma-aminobutyric acid(A)-receptor subunit composition in the circadian system and median eminence of Syrian hamsters. *Neuroscience Letters* **310** 178–182. (doi:10.1016/S0304-3940(01)02129-2)
- Nielsen HS, Hannibal J & Fahrenkrug J 2002 Vasoactive intestinal polypeptide induces *per1* and *per2* gene expression in the rat suprachiasmatic nucleus late at night. *European Journal of Neuroscience* **15** 570–574. (doi:10.1046/j.0953-816x.2001.01882.x)
- Noguchi T & Watanabe K 2008 Regional differences in circadian period within the suprachiasmatic nucleus. *Brain Research* **1239** 119–126. (doi:10.1016/j.brainres.2008.08.082)
- Noguchi T, Michihata T, Nakamura W, Takumi T, Shimizu R, Yamamoto M, Ikeda M, Ohmiya Y & Nakajima Y 2010 Dual-color luciferase mouse directly demonstrates coupled expression of two clock genes. *Biochemistry* **49** 8053–8061. (doi:10.1021/bi100545h)
- Novak CM, Ehlen JC, Huhman KL & Albers HE 2004 GABA(B) receptor activation in the suprachiasmatic nucleus of diurnal and nocturnal rodents. *Brain Research Bulletin* **63** 531–535. (doi:10.1016/j.brainresbull.2004.05.001)
- O'Hara BF, Andretic R, Heller HC, Carter DB & Kilduff TS 1995 GABAA, GABAC, and NMDA receptor subunit expression in the suprachiasmatic nucleus and other brain regions. *Brain Research. Molecular Brain Research* **28** 239–250.
- Okamura H, Berod A, Julien JF, Geffard M, Kitahama K, Mallet J & Bobillier P 1989 Demonstration of GABAergic cell bodies in the suprachiasmatic nucleus: in situ hybridization of *glutamic acid decarboxylase (GAD)* mRNA and immunocytochemistry of GAD and GABA. *Neuroscience Letters* **102** 131–136. (doi:10.1016/0304-3940(89)90067-0)
- Pakhotin P, Harmar AJ, Verkhatsky A & Piggins H 2006 VIP receptors control excitability of suprachiasmatic nuclei neurones. *Pflügers Archiv* **452** 7–15. (doi:10.1007/s00424-005-0003-z)
- Piggins HD, Antle MC & Rusak B 1995 Neuropeptides phase shift the mammalian circadian pacemaker. *Journal of Neuroscience* **15** 5612–5622.
- Piggins HD, Goguen D & Rusak B 2005 Gastrin-releasing peptide induces c-Fos in the hamster suprachiasmatic nucleus. *Neuroscience Letters* **384** 205–210. (doi:10.1016/j.neulet.2005.03.072)
- Prolo LM, Takahashi JS & Herzog ED 2005 Circadian rhythm generation and entrainment in astrocytes. *Journal of Neuroscience* **25** 404–408. (doi:10.1523/JNEUROSCI.4133-04.2005)
- Prosser RA, Edgar DM, Heller HC & Miller JD 1994 A possible glial role in the mammalian circadian clock. *Brain Research* **643** 296–301. (doi:10.1016/0006-8993(94)90036-1)
- Quintero JE, Kuhlman SJ & McMahon DG 2003 The biological clock nucleus: a multiphasic oscillator network regulated by light. *Journal of Neuroscience* **23** 8070–8076.
- Rash JE, Staines WA, Yasumura T, Patel D, Furman CS, Stelmack GL & Nagy JI 2000 Immunogold evidence that neuronal gap junctions in adult rat brain and spinal cord contain connexin-36 but not connexin-32 or connexin-43. *PNAS* **97** 7573–7578. (doi:10.1073/pnas.97.13.7573)
- Rash JE, Yasumura T, Dudek FE & Nagy JI 2001 Cell-specific expression of connexins and evidence of restricted gap junctional coupling between glial cells and between neurons. *Journal of Neuroscience* **21** 1983–2000.
- Reppert S 1992 Pre-natal development of a hypothalamic biological clock. *Progress in Brain Research* **93** 119–132.
- Rohling J, Meijer JH, VanderLeest HT & Admiraal J 2006 Phase differences between SCN neurons and their role in photoperiodic encoding; a simulation of ensemble patterns using recorded single unit electrical activity patterns. *Journal of Physiology* **100** 261–270.
- Rohling JH, Vanderleest HT, Michel S, Vansteensel MJ & Meijer JH 2011 Phase resetting of the mammalian circadian clock relies on a rapid shift of a small population of pacemaker neurons. *PLoS ONE* **6** e25437. (doi:10.1371/journal.pone.0025437)
- Romijn HJ, Sluiter AA, Pool CW, Wortel J & Buijs RM 1997 Evidence from confocal fluorescence microscopy for a dense, reciprocal innervation between AVP-, somatostatin-, VIP/PHI-, GRP-, and VIP/PHI/GRP-immunoreactive neurons in the rat suprachiasmatic nucleus. *European Journal of Neuroscience* **9** 2613–2623. (doi:10.1111/j.1460-9568.1997.tb01691.x)
- Russell W, Harrison RF, Smith N, Darzy K, Shalet S, Weetman AP & Ross RJ 2008 Free triiodothyronine has a distinct circadian rhythm that is delayed but parallels thyrotropin levels. *Journal of Clinical Endocrinology & Metabolism* **93** 2300–2306.

- Saeb-Parsy K & Dyball RE 2003 Defined cell groups in the rat suprachiasmatic nucleus have different day/night rhythms of single-unit activity *in vivo*. *Journal of Biological Rhythms* **18** 26–42. (doi:10.1177/0748730402239674)
- Schaap J, Albus H, VanderLeest HT, Eilers PH, Detari L & Meijer JH 2003 Heterogeneity of rhythmic suprachiasmatic nucleus neurons: implications for circadian waveform and photoperiodic encoding. *PNAS* **100** 15994–15999. (doi:10.1073/pnas.2436298100)
- Schroder H, Stehle J & Henschel M 1988 Twenty-four-hour pineal melatonin synthesis in the vasopressin-deficient Brattleboro rat. *Brain Research* **459** 328–332. (doi:10.1016/0006-8993(88)90648-8)
- Schroeder A, Loh DH, Jordan MC, Roos KP & Colwell CS 2011 Circadian regulation of cardiovascular function: a role for vasoactive intestinal peptide. *American Journal of Physiology: Heart and Circulatory Physiology* **300** H241–H250. (doi:10.1152/ajpheart.00190.2010)
- Schwartz WJ 1991 Further evaluation of the tetrodotoxin-resistant circadian pacemaker in the suprachiasmatic nuclei. *Journal of Biological Rhythms* **6** 149–158. (doi:10.1177/074873049100600205)
- Schwartz WJ & Gainer H 1977 Suprachiasmatic nucleus: use of ¹⁴C-labeled deoxyglucose uptake as a functional marker. *Science* **197** 1089–1091. (doi:10.1126/science.887940)
- Schwartz WJ, Gross RA & Morton MT 1987 The suprachiasmatic nuclei contain a tetrodotoxin-resistant circadian pacemaker. *PNAS* **84** 1694–1698. (doi:10.1073/pnas.84.6.1694)
- Schwartz WJ, Carpino A Jr, de la Iglesia HO, Baler R, Klein DC, Nakabeppu Y & Aronin N 2000 Differential regulation of *fos* family genes in the ventrolateral and dorsomedial subdivisions of the rat suprachiasmatic nucleus. *Neuroscience* **98** 535–547. (doi:10.1016/S0306-4522(00)00140-8)
- Schwartz MD, Wotus C, Liu T, Friesen WO, Borjigin J, Oda GA & de la Iglesia HO 2009 Dissociation of circadian and light inhibition of melatonin release through forced desynchronization in the rat. *PNAS* **106** 17540–17545. (doi:10.1073/pnas.0906382106)
- Sellix MT, Evans JA, Leise TL, Castanon-Cervantes O, Hill DD, DeLisser P, Block GD, Menaker M & Davidson AJ 2012 Aging differentially affects the re-entrainment response of central and peripheral circadian oscillators. *Journal of Neuroscience* **32** 16193–16202. (doi:10.1523/JNEUROSCI.3559-12.2012)
- Shen S, Spratt C, Sheward WJ, Kallo I, West K, Morrison CF, Coen CW, Marston HM & Harmar AJ 2000 Overexpression of the human VPAC2 receptor in the suprachiasmatic nucleus alters the circadian phenotype of mice. *PNAS* **97** 11575–11580. (doi:10.1073/pnas.97.21.11575)
- Sheward WJ, Naylor E, Knowles-Barley S, Armstrong JD, Brooker GA, Seckl JR, Turek FW, Holmes MC, Zee PC & Harmar AJ 2010 Circadian control of mouse heart rate and blood pressure by the suprachiasmatic nuclei: behavioral effects are more significant than direct outputs. *PLoS ONE* **5** e9783. (doi:10.1371/journal.pone.0009783)
- Shibata S & Moore RY 1987 Development of neuronal activity in the rat suprachiasmatic nucleus. *Brain Research* **431** 311–315. (doi:10.1016/0165-3806(87)90220-3)
- Shibata S & Moore RY 1993 Tetrodotoxin does not affect circadian rhythms in neuronal activity and metabolism in rodent suprachiasmatic nucleus *in vitro*. *Brain Research* **606** 259–266. (doi:10.1016/0006-8993(93)90993-W)
- Shibata S, Liou S, Ueki S & Oomura Y 1984 Influence of environmental light-dark cycle and enucleation on activity of suprachiasmatic neurons in slice preparations. *Brain Research* **302** 75–81. (doi:10.1016/0006-8993(84)91286-1)
- Shimizu K, Nagai K & Nakagawa H 1996 An immunotoxin, anti-VIP antibody-ricin A chain conjugate eliminates neurons in the hypothalamic suprachiasmatic nucleus selectively and abolishes the circadian rhythm of water intake. *Brain Research Bulletin* **41** 369–378. (doi:10.1016/S0361-9230(96)00070-6)
- Shimura M, Harata N, Tamai M & Akaike N 1996 Allosteric modulation of GABAA receptors in acutely dissociated neurons of the suprachiasmatic nucleus. *American Journal of Physiology* **270** C1726–C1734.
- Shinohara K & Inouye ST 1995 Photic information coded by vasoactive intestinal polypeptide and neuropeptide Y. *Neuroscience & Biobehavioral Reviews* **19** 349–352.
- Shinohara K, Tominaga K, Isobe Y & Inouye ST 1993 Photic regulation of peptides located in the ventrolateral subdivision of the suprachiasmatic nucleus of the rat: daily variations of vasoactive intestinal polypeptide, gastrin-releasing peptide, and neuropeptide Y. *Journal of Neuroscience* **13** 793–800.
- Shinohara K, Honma S, Katsuno Y, Abe H & Honma K 1995 Two distinct oscillators in the rat suprachiasmatic nucleus *in vitro*. *PNAS* **92** 7396–7400. (doi:10.1073/pnas.92.16.7396)
- Shinohara K, Honma S, Katsuno Y, Abe H & Honma K 1998 Circadian release of amino acids in the suprachiasmatic nucleus *in vitro*. *Neuroreport* **9** 137–140. (doi:10.1097/00001756-199801050-00027)
- Shinohara K, Funabashi T, Mitushima D & Kimura F 2000a Effects of gap junction blocker on vasopressin and vasoactive intestinal polypeptide rhythms in the rat suprachiasmatic nucleus *in vitro*. *Neuroscience Research* **38** 43–47.
- Shinohara K, Hiruma H, Funabashi T & Kimura F 2000b GABAergic modulation of gap junction communication in slice cultures of the rat suprachiasmatic nucleus. *Neuroscience* **96** 591–596.
- Shirakawa T, Honma S, Katsuno Y, Oguchi H & Honma KI 2000 Synchronization of circadian firing rhythms in cultured rat suprachiasmatic neurons. *European Journal of Neuroscience* **12** 2833–2838. (doi:10.1046/j.1460-9568.2000.00170.x)
- Shirakawa T, Honma S & Honma K 2001 Multiple oscillators in the suprachiasmatic nucleus. *Chronobiology International* **18** 371–387. (doi:10.1081/CBI-100103962)
- Silver R, Romero MT, Besmer HR, Leak R, Nunez JM & LeSauter J 1996 Calbindin-D28K cells in the hamster SCN express light-induced Fos. *Neuroreport* **7** 1224–1228. (doi:10.1097/00001756-199604260-00026)
- Smarr BL, Morris E & de la Iglesia HO 2012 The dorsomedial suprachiasmatic nucleus times circadian expression of *Kiss1* and the luteinizing hormone surge. *Endocrinology* **153** 2839–2850. (doi:10.1210/en.2011-1857)
- Smith L & Canal MM 2009 Expression of circadian neuropeptides in the hypothalamus of adult mice is affected by postnatal light experience. *Journal of Neuroendocrinology* **21** 946–953. (doi:10.1111/j.1365-2826.2009.01914.x)
- Strecker GJ, Wuarin JP & Dudek FE 1997 GABAA-mediated local synaptic pathways connect neurons in the rat suprachiasmatic nucleus. *Journal of Neurophysiology* **78** 2217–2220.
- Sukhov RR, Walker LC, Rance NE, Price DL & Young WS 3rd 1993 Vasopressin and oxytocin gene expression in the human hypothalamus. *Journal of Comparative Neurology* **337** 295–306. (doi:10.1002/cne.903370210)
- Takahashi Y, Kipnis DM & Daughaday WH 1968 Growth hormone secretion during sleep. *Journal of Clinical Investigation* **47** 2079–2090. (doi:10.1172/JCI105893)
- Takahashi Y, Okamura H, Yanaihara N, Hamada S, Fujita S & Ibata Y 1989 Vasoactive intestinal peptide immunoreactive neurons in the rat suprachiasmatic nucleus demonstrate diurnal variation. *Brain Research* **497** 374–377. (doi:10.1016/0006-8993(89)90283-7)
- Tamada Y, Tanaka M, Munekawa K, Hayashi S, Okamura H, Kubo T, Hisa Y & Ibata Y 1998 Neuron-glia interaction in the suprachiasmatic nucleus: a double labeling light and electron microscopic immunocytochemical study in the rat. *Brain Research Bulletin* **45** 281–287. (doi:10.1016/S0361-9230(97)00403-6)
- Toussou E & Meissl H 2004 Suprachiasmatic nuclei grafts restore the circadian rhythm in the paraventricular nucleus of the hypothalamus. *Journal of Neuroscience* **24** 2983–2988. (doi:10.1523/JNEUROSCI.5044-03.2004)

- Van den Pol AN 1980 The hypothalamic suprachiasmatic nucleus of rat: intrinsic anatomy. *Journal of Comparative Neurology* **191** 661–702. (doi:10.1002/cne.901910410)
- van den Pol AN 2012 Neuropeptide transmission in brain circuits. *Neuron* **76** 98–115. (doi:10.1016/j.neuron.2012.09.014)
- van den Pol AN & Dudek FE 1993 Cellular communication in the circadian clock, the suprachiasmatic nucleus. *Neuroscience* **56** 793–811. (doi:10.1016/0306-4522(93)90128-3)
- van den Pol AN & Tsujimoto KL 1985 Neurotransmitters of the hypothalamic suprachiasmatic nucleus: immunocytochemical analysis of 25 neuronal antigens. *Neuroscience* **15** 1049–1086. (doi:10.1016/0306-4522(85)90254-4)
- van den Pol AN, Finkbeiner SM & Cornell-Bell AH 1992 Calcium excitability and oscillations in suprachiasmatic nucleus neurons and glia *in vitro*. *Journal of Neuroscience* **12** 2648–2664.
- van der Beek EM, Wiegant VM, van der Donk HA, van den Hurk R & Buijs RM 1993 Lesions of the suprachiasmatic nucleus indicate the presence of a direct vasoactive intestinal polypeptide-containing projection to gonadotrophin-releasing hormone neurons in the female rat. *Journal of Neuroendocrinology* **5** 137–144. (doi:10.1111/j.1365-2826.1993.tb00373.x)
- Van der Veen DR, Castillo MR, Van der Zee EA, Jansen K, Gerkema MP & Bult-Itto A 2005 Circadian dynamics of vasopressin in mouse selection lines: translation and release in the SCN. *Brain Research* **1060** 16–25. (doi:10.1016/j.brainres.2005.07.068)
- van Esseveldt KE, Lehman MN & Boer GJ 2000 The suprachiasmatic nucleus and the circadian time-keeping system revisited. *Brain Research. Brain Research Reviews* **33** 34–77. (doi:10.1016/S0165-0173(00)00025-4)
- Vogt K 2015 Diversity in GABAergic signaling. *Advances in Pharmacology* **73** 203–222. (doi:10.1016/bs.apha.2014.11.009)
- Vosko AM, Schroeder A, Loh DH & Colwell CS 2007 Vasoactive intestinal peptide and the mammalian circadian system. *General and Comparative Endocrinology* **152** 165–175. (doi:10.1016/j.ygcen.2007.04.018)
- Wagner S, Castel M, Gainer H & Yarom Y 1997 GABA in the mammalian suprachiasmatic nucleus and its role in diurnal rhythmicity. *Nature* **387** 598–603. (doi:10.1038/42468)
- Wang D, Cui LN & Renaud LP 2003 Pre- and postsynaptic GABA(B) receptors modulate rapid neurotransmission from suprachiasmatic nucleus to parvocellular hypothalamic paraventricular nucleus neurons. *Neuroscience* **118** 49–58. (doi:10.1016/S0306-4522(02)00906-5)
- Wang MH, Chen N & Wang JH 2014 The coupling features of electrical synapses modulate neuronal synchrony in hypothalamic suprachiasmatic nucleus. *Brain Research* **1550** 9–17. (doi:10.1016/j.brainres.2014.01.007)
- Watanabe K, Vanecsek J & Yamaoka S 2000 *In vitro* entrainment of the circadian rhythm of vasopressin-releasing cells in suprachiasmatic nucleus by vasoactive intestinal polypeptide. *Brain Research* **877** 361–366. (doi:10.1016/S0006-8993(00)02724-4)
- Watts AG 1991 The efferent projections of the suprachiasmatic nucleus: anatomical insights into the control of circadian rhythms. In *Suprachiasmatic Nucleus: The Mind's Clock*, pp 77–106. Eds DC Klein, RY Moore & SM Reppert. New York, NY, USA: University of Oxford Press.
- Watts AG & Swanson LW 1987 Efferent projections of the suprachiasmatic nucleus: II. Studies using retrograde transport of fluorescent dyes and simultaneous peptide immunohistochemistry in the rat. *Journal of Comparative Neurology* **258** 230–252. (doi:10.1002/cne.902580205)
- Weaver DR 1998 The suprachiasmatic nucleus: a 25-year retrospective. *Journal of Biological Rhythms* **13** 100–112. (doi:10.1177/074873098128999952)
- Webb AB, Angelo N, Huettner JE & Herzog ED 2009 Intrinsic, nondeterministic circadian rhythm generation in identified mammalian neurons. *PNAS* **106** 16493–16498. (doi:10.1073/pnas.0902768106)
- Welsh DK, Logothetis DE, Meister M & Reppert SM 1995 Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron* **14** 697–706. (doi:10.1016/0896-6273(95)90214-7)
- Welsh DK, Yoo SH, Liu AC, Takahashi JS & Kay SA 2004 Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Current Biology* **14** 2289–2295. (doi:10.1016/j.cub.2004.11.057)
- Welsh DK, Takahashi JS & Kay SA 2010 Suprachiasmatic nucleus: cell autonomy and network properties. *Annual Review of Physiology* **72** 551–577. (doi:10.1146/annurev-physiol-021909-135919)
- Wideman CH, Murphy HM & Nadzam GR 2000 Vasopressin deficiency provides evidence for separate circadian oscillators of activity and temperature. *Peptides* **21** 811–816. (doi:10.1016/S0196-9781(00)00213-8)
- Womac AD, Burkeen JF, Neuendorff N, Earnest DJ & Zoran MJ 2009 Circadian rhythms of extracellular ATP accumulation in suprachiasmatic nucleus cells and cultured astrocytes. *European Journal of Neuroscience* **30** 869–876. (doi:10.1111/j.1460-9568.2009.06874.x)
- Wotus C, Lilley TR, Neal AS, Suleiman NL, Schmuck SC, Smarr BL, Fischer BJ & de la Iglesia HO 2013 Forced desynchrony reveals independent contributions of suprachiasmatic oscillators to the daily plasma corticosterone rhythm in male rats. *PLoS ONE* **8** e68793.
- Yagita K, Yamanaka I, Emoto N, Kawakami K & Shimada S 2010 Real-time monitoring of circadian clock oscillations in primary cultures of mammalian cells using Tol2 transposon-mediated gene transfer strategy. *BMC Biotechnology* **10** 3. (doi:10.1186/1472-6750-10-3)
- Yamaguchi S, Isejima H, Matsuo T, Okura R, Yagita K, Kobayashi M & Okamura H 2003 Synchronization of cellular clocks in the suprachiasmatic nucleus. *Science* **302** 1408–1412. (doi:10.1126/science.1089287)
- Yamaguchi Y, Suzuki T, Mizoro Y, Kori H, Okada K, Chen Y, Fustin JM, Yamazaki F, Mizuguchi N, Zhang J, *et al.* 2013 Mice genetically deficient in vasopressin V1a and V1b receptors are resistant to jet lag. *Science* **342** 85–90. (doi:10.1126/science.1238599)
- Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M & Tei H 2000 Resetting central and peripheral circadian oscillators in transgenic rats. *Science* **288** 682–685. (doi:10.1126/science.288.5466.682)
- Yan L & Okamura H 2002 Gradients in the circadian expression of *Per1* and *Per2* genes in the rat suprachiasmatic nucleus. *European Journal of Neuroscience* **15** 1153–1162. (doi:10.1046/j.1460-9568.2002.01955.x)
- Yan L & Silver R 2002 Differential induction and localization of *mPer1* and *mPer2* during advancing and delaying phase shifts. *European Journal of Neuroscience* **16** 1531–1540. (doi:10.1046/j.1460-9568.2002.02224.x)
- Yan L & Silver R 2004 Resetting the brain clock: time course and localization of mPER1 and mPER2 protein expression in suprachiasmatic nuclei during phase shifts. *European Journal of Neuroscience* **19** 1105–1109.
- Yan L, Foley NC, Bobula JM, Kriegsfeld LJ & Silver R 2005 Two antiphase oscillations occur in each suprachiasmatic nucleus of behaviorally split hamsters. *Journal of Neuroscience* **25** 9017–9026.
- Yan L, Karatsoreos I, Lesauter J, Welsh DK, Kay S, Foley D & Silver R 2007 Exploring spatiotemporal organization of SCN circuits. *Cold Spring Harbor Symposia on Quantitative Biology* **72** 527–541. (doi:10.1101/sqb.2007.72.037)
- Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepka SM, Hong HK, Oh WJ, Yoo OJ, *et al.* 2004 PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *PNAS* **101** 5339–5346.

- Yoshikawa T, Nakajima Y, Yamada Y, Enoki R, Watanabe K, Yamazaki M, Sakimura K, Honma S & Honma K 2015 Spatiotemporal profiles of arginine vasopressin transcription in cultured suprachiasmatic nucleus. *European Journal of Neuroscience* **42** 2678–2689. (doi:10.1111/ejn.13061)
- Zhang EE & Kay SA 2010 Clocks not winding unravelling circadian networks. *Nature Reviews Molecular Cell Biology* **11** 764–776. (doi:10.1038/nrm2995)

- Zhang R, Lahens NF, Ballance HI, Hughes ME & Hogenesch JB 2014 A circadian gene expression atlas in implications for biology and medicine. *PNAS* **111** 16219–16224. (doi:10.1073/pnas.1408886111)
- Zhou QY & Cheng MY 2005 Prokineticin 2 and circadian clock output. *FEBS Journal* **272** 5703–5709.
- Zhou JN, Hofman MA & Swaab DF 1995 VIP neurons in the human SCN in relation to sex, age, and Alzheimer's disease. *Neurobiology of Aging* **16** 571–576.

Received in final form 11 April 2016

Accepted 6 May 2016

Accepted Preprint published online 6 May 2016