Selye's general adaptation syndrome: stress-induced gastro-duodenal ulceration and inflammatory bowel disease

George Fink

Florey Institute of Neuroscience and Mental Health, University of Melbourne, Parkville, Victoria, Australia

Correspondence should be addressed to G Fink
Email
george.fink@florey.edu.au

Abstract

Hans Selye in a note to *Nature* in 1936 initiated the field of stress research by showing that rats exposed to noxious stimuli responded by way of a 'general adaptation syndrome' (GAS). One of the main features of the GAS was the 'formation of acute erosions in the digestive tract, particularly in the stomach, small intestine and appendix'. This provided experimental evidence for the view based on clinical data that gastro-duodenal (peptic) ulcers could be caused by stress. This hypothesis was challenged by Marshall and Warren's Nobel Prize (2005)-winning discovery of a causal association between *Helicobacter pylori* and peptic ulcers. However, clinical and experimental studies suggest that stress can cause peptic ulceration in the absence of *H. pylori*. Predictably, the etiological pendulum of gastric and duodenal ulceration has swung from 'all stress' to 'all bacteria' followed by a sober realization that both factors play a role, separately as well as together. This raises the question as to whether stress and *H. pylori* interact, and if so, how? Stress has also been implicated in inflammatory bowel disease (IBD) and related disorders; however, there is no proof yet that stress is the primary etiological trigger for IBD. Central dopamine mechanisms seem to be involved in the stress induction of peptic ulceration, whereas activation of the sympathetic nervous system and central and peripheral corticotrophin-releasing factor appears to mediate stress-induced IBD.

Key Words

- ▶ peptic ulcers
- ▶ stress
- ▶ *Helicobacter pylori*
- ▶ inflammatory bowel disease
- ▶ brain-gut axis

Journal of Endocrinology
 (2017) **232**, F1–F5

Introduction

Eighty years ago, *Nature* published a note from Hans Selye entitled 'A Syndrome produced by Diverse Noxious Agents' (Selye 1936). Briefly, Selye reported that if experimental rats were 'damaged by acute non-specific noxious agents such as exposure to cold, surgical injury, excessive muscular exercise, or intoxications with sub lethal doses of diverse drugs (adrenaline, atropine, morphine, formaldehyde, etc.), a typical

syndrome appears'. Selye termed this the general adaptation syndrome (GAS) 'the symptoms of which are independent of the nature of the damaging agent or the pharmacological type of the drug employed and represent rather a response to damage as such.' The main features of the syndrome were a fall of body temperature and decrease in muscle tone; decrease in size of the thymus, spleen, lymph nodes and liver; acute erosions of the

digestive tract; enlargement of the adrenal glands and histological changes in the adrenal medullary chromaffin cells; increased secretion of pituitary 'thyrotropic and adrenotropic principles'. Selye's observations translate in modern terms to stress-induced immunosuppression, peptic ulceration and inflammatory bowel disease (IBD) and activation of the two main vertebrate stress response systems: the hypothalamic–pituitary–adrenal axis (HPA) and the sympathomedullary system.

Stress research has made huge advances since Selye's publication especially with respect to the fundamental neurobiology and neuroendocrinology of the stress response (McEwen 2007, Joels & Baram 2010, Henckens *et al.* 2016). However, uncertainties, irreproducible results and unresolved controversies remain (Fink 2011). These include, for example, the nature, if any, of genetic and epigenetic mechanisms that determine the susceptibility or resilience to stress; perinatal effects that might determine stress vulnerability in adults; the relative importance of the autonomic nervous versus the brain–pituitary–adrenocortical system in mental illness such as post-traumatic stress disorder, major depression and the psychoses; and whether stress and anxiety damage the human brain and if so whether the damage is reversible.

Here, attention will focus on the controversial issue of the role of stress versus *Helicobacter pylori* in the causation of gastro-duodenal ulceration. Brief mention of inflammatory bowel disease will also be made since this disorder, perhaps predicted by the GAS, might also be affected by stress.

Gastro-duodenal ulceration: stress vs *Helicobacter pylori*

The fact that emotions such as fear and anger affect acid secretions of the stomach was first documented in 1833 by US Army surgeon William Beaumont in his studies on Alexis St Martin a French Canadian fur trapper who accidentally sustained a massive gunshot wound in the stomach (Beaumont 1833). The wound resulted in a chronic gastric fistula that enabled Beaumont to make direct observations of the mucosa of the stomach and gastric secretions under different conditions. Beaumont's findings provided the basis for the subsequent experiments of Ivan Petrovich Pavlov carried out in dogs by means of a gastric pouch, which lead to the discovery of the conditioned reflex and the concept of the 'cephalic phase' of gastric and pancreatic secretion (Wood 2004). However, robust evidence that stress could induce gastro-duodenal

('peptic') ulcers was first adduced by Selye's GAS findings that stress induced the 'formation of acute erosions in the digestive tract, particularly in the stomach, small intestine and appendix' (Selye 1936).

The observations of Beaumont, Pavlov and Selye and the fact that massive perforation of the stomach and other parts of the gastrointestinal tract in the human were associated with brain lesions or 'neurogenic' factors (Oppen & Zimmerman 1938, Boles & Riggs 1941, Rosenthal & Tobias 1943) led to the view that stress was the main cause of gastro-duodenal (peptic) ulceration. The first description of neurogenic ulcers was by Harvey Cushing who, perhaps ironically, in later years with failing health developed an ulcer himself (Wijdsicks 2011). Psychological studies using life events as a measure showed that chronic stress in the human is associated with peptic ulceration when the stressor comprises personal threat or goal frustration (Piper & Tennant 1993).

The assumption that peptic ulceration was caused by stress seemed to be overturned when Barry Marshall and Robin Warren in the 1980s and 90s made their 2005 Nobel Prize-winning discovery that gastric and duodenal ulcers appeared to be caused by *H. pylori* (Marshall & Warren 1984, Marshall & Windsor 2005, Cover & Blaser 2009). This suggested that antibiotics together with proton pump inhibitors could be the optimal treatment for gastro-duodenal ulcers and dyspepsia in general, a simple and apparently benign approach that is only sometimes confounded by antibiotic resistance (Tay *et al.* 2012).

There is no doubt that a substantial number of peptic ulcers are associated with the presence of *H. pylori*. However, notwithstanding the importance of *H. pylori* for peptic ulceration (and gastric cancer), several studies in the human and in rodents show that stress plays a significant role in peptic ulceration (Matsushima *et al.* 1999, Kim *et al.* 2002, Levenstein 2009, McColl 2009). The role of stress in the causation of peptic ulceration in the human is exemplified by the fact that the Hanshin-Awaji earthquake, which occurred on 17 January 1995, was followed by a significant increase in the number of people with peptic ulceration (McColl 2009, Levenstein *et al.* 2015). Among the physically injured, stomach ulcers developed independently of *H. pylori* infection. Extensive burns also lead to stress ulcers that are likely to be independent of *H. pylori* status (McColl 2009). Furthermore, most people living with *H. pylori* never develop ulcers, about 30% of patients with ulcer do not harbor *H. pylori* infection and some patients in whom *H. pylori* colonization has been cleared by antibiotics subsequently develop new ulcers (Levenstein 2009).

The above points are supported by several control-balanced clinical studies. Thus, for example, a Finnish nationwide twin cohort study of lifestyle, stress and genes in peptic ulcer disease led to the conclusions that, first, familial aggregation of the peptic ulcer disease is modest and attributable almost solely to genetic factors, and, second, that the data did not support the concept that risk factors such as *H. pylori* infection could explain familial peptic ulcer disease (Räihä *et al.* 1998). A comprehensive study in Thailand of 70 patients with perforated peptic ulcer compared with a balanced number of control subjects led to the conclusion that stress, but not *H. pylori*, is associated with peptic ulcer disease in this Thai population (Wachirawat *et al.* 2003). Finally, a recent population-based Danish cohort ($n=2410$) study showed that psychological stress increased the incidence of peptic ulcer in subjects with or without *H. pylori* infection (Levenstein *et al.* 2015).

The case for the importance of psychosocial stress factors in the causation of peptic ulcer disease has been summarized as follows (Levenstein 2000, Jones 2006, Realo *et al.* 2015). First, despite the decreasing prevalence of *H. pylori* infection in Western societies, the burden of peptic ulcer disease remains high. Second, most *H. pylori*-infected individuals never develop peptic ulcers. Third, a proportion of patients develop peptic ulceration in the absence of *H. pylori*.

The precise mechanism by which stress can cause gastro-duodenal ulcers is yet to be determined. However, in addition to the possible effects of neuroendocrine (hypothalamic–pituitary–adrenal)-induced elevation of glucocorticoid levels on the gastro-duodenal mucosa (Levenstein *et al.* 2015), central and peripheral dopaminergic mechanisms remain contenders (Szabo 1979, Glavin & Szabo 1990, Ozdemir *et al.* 2007, Rasheed & Algsham 2012). The clue to a possible role of central dopamine came from clinical observations and hypotheses regarding the etiology of Parkinson's disease and schizophrenia. Thus, as early as 1965, Strang noted an association between Parkinson's disease, characterized by central dopamine deficiency, and gastro-duodenal ulceration (Strang 1965). This was especially the case in young Parkinsonian patients compared with control subjects. *Pari-passu* duodenal ulcers are rare in patients with schizophrenia, a condition in which dopaminergic activity is thought to be increased (Glavin & Szabo 1990). Experimental evidence for a central role of dopamine deficiency as a cause of duodenal ulceration comes from the effects of N-methyl-D-aspartate lesions of the substantia nigra (Landeira-Fernandez & Grijalva 2004) and the

systemic administration of the dopaminergic neurotoxin, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP). The latter induces nigrostriatal dopamine depletion the magnitude of which correlates with the severity of the duodenal ulcers (Glavin & Szabo 1990, Sikiric *et al.* 1999). Further evidence for the role of dopamine deficiency or excess in causing or protecting against gastro-duodenal ulceration, respectively, is derived from studies of dopamine agonists and antagonists (Szabo 1979, Glavin & Szabo 1990, Ozdemir *et al.* 2007, Rasheed & Algsham 2012).

Inflammatory bowel disease

Stress, in addition to its role in peptic ulceration, is thought to play a role in IBD and related disorders such as irritable bowel syndrome, adverse reactions to food antigens and gastroesophageal reflux disease (Konturek *et al.* 2011). This effect of stress is probably mediated by the sympathomedullary system and corticotrophin-releasing factor (CRF) acting both centrally and within the gut (Bunnett 2005, Tache *et al.* 2008, Lukewich *et al.* 2014). Notwithstanding the possible role of stress in bowel disorders and the fact that CRF and its receptors are regarded by some as potentially relevant therapeutic targets (Tache *et al.* 2008), it must be stressed that the signal factors that trigger IBD and related disorders remain to be determined. The story is far from complete.

Conclusion

Hans Selye's experimental rodent observations and inferences that stress can cause the 'formation of acute erosions in the digestive tract, particularly in the stomach, small intestine and appendix' also apply to the human. Thus, stress may play a role in IBD and related disorders, but proof that stress is a main let alone sole trigger awaits further rigorous research.

With respect to the stomach and duodenum, stress and *H. pylori* are both clearly risk factors for gastro-duodenal ulceration. Whether, and if so, how stress and *H. pylori* might interact, if at all, remains to be determined. Overall, the etiology of gastric and duodenal ulceration and the causal switch in dogma from 'all stress to all bacteria' followed by a sober realization that both factors might play a role have heuristic value for our understanding of disease pathogenesis.

Declaration of interest

The author declares that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work did not receive any specific grant from any funding agency in the public, commercial, or not-for-profit sector.

Acknowledgments

I am grateful to the Florey Institute of Neuroscience and Mental Health, the University of Melbourne, Victoria, Australia, and the Department of Neurobiology, the Weizmann Institute of Science, Rehovot, Israel for generously providing facilities that have enabled me to complete this work.

References

- Beaumont W 1833 Experiments and Observations on the Gastric Juice and the Physiology of Digestion, pp 280. Dover Edition, 1959: unabridged. New York, USA: Dover Publications Inc.
- Boles RS & Riggs HE 1941 Neurogenic factors in the production of acute gastric ulcer. *Journal of Nervous & Mental Disease* **94** 215–216. (doi:10.1097/00005053-194108000-00012)
- Bunnett NW 2005 The stressed gut: contributions of intestinal stress peptides to inflammation and motility. *PNAS* **102** 7409–7410. (doi:10.1073/pnas.0503092102)
- Cover TL & Blaser MJ 2009 Helicobacter pylori in health and disease. *Gastroenterology* **136** 1863–1873. (doi:10.1053/j.gastro.2009.01.073)
- Fink G 2011 Stress controversies: post-traumatic stress disorder, hippocampal volume, gastroduodenal ulceration. *Journal of Neuroendocrinology* **23** 107–117. (doi:10.1111/j.1365-2826.2010.02089.x)
- Glavin GB & Szabo S 1990 Dopamine in gastrointestinal disease. *Digestive Diseases and Sciences* **35** 1153–1161. (doi:10.1007/BF01537589)
- Henckens MJ, Deussing JM & Chen A 2016 Region-specific roles of the corticotropin-releasing factor-urocortin system in stress. *Nature Reviews Neuroscience* **17** 636–651. (doi:10.1038/nrn.2016.94)
- Joels M & Baram TZ 2010 The neurosymphony of stress. *Nature Reviews Neuroscience* **10** 459–466. (doi:10.1038/nrn2632)
- Jones MP 2006 The role of psychosocial factors in peptic ulcer disease: beyond Helicobacter pylori and NSAIDs. *Journal of Psychosomatic Research* **60** 407–412. (doi:10.1016/j.jpsychores.2005.08.009)
- Kim Y-H, Lee JH, Lee SS, Cho EY, Oh YL, Son HJ, Rhee P-L, Kim JJ, Koh KC, Paik SW, et al. 2002 Long-term stress and Helicobacter pylori infection independently induce gastric mucosal lesions in C57BL/6 mice. *Scandinavian Journal of Gastroenterology* **37** 1259–1264. (doi:10.1080/003655202761020515)
- Konturek PC, Brzozowski T & Konturek SJ 2011 Stress and the gut: pathophysiology, clinical consequences, diagnostic approach and treatment options. *Journal of Physiology and Pharmacology* **62** 591–599.
- Landeira-Fernandez J & Grijalva CV 2004 Participation of the substantia nigra dopaminergic neurons in the occurrence of gastric mucosal erosions. *Physiology & Behavior* **81** 91–99. (doi:10.1016/j.physbeh.2004.01.004)
- Levenstein S 2000 The very model of a modern etiology: a biopsychosocial view of peptic ulcer. *Psychosomatic Medicine* **62** 176–185. (doi:10.1097/00006842-200003000-00003)

- Levenstein S 2009 Helicobacter pylori and ulcers. Against reductionism. *British Medical Journal* **339** b3855. (doi:10.1136/bmj.b3855)
- Levenstein S, Rosenstock S, Jacobsen RK & Jorgensen T 2015 Psychological stress increases risk for peptic ulcer, regardless of Helicobacter pylori infection or use of nonsteroidal anti-inflammatory drugs. *Clinical Gastroenterology and Hepatology* **13** 498–506 e1. (doi:10.1016/j.cgh.2014.07.052)
- Lukewich MK, Rogers RC & Lomax AE 2014 Divergent neuroendocrine responses to localised and systemic inflammation. *Seminars in Immunology* **26** 402–408. (doi:10.106/j.smim.2014.01.004)
- Marshall BJ & Warren JR 1984 Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* **1** 1311–1315. (doi:10.1016/S0140-6736(84)91816-6)
- Marshall BJ & Windsor HM 2005 The relation of Helicobacter pylori to gastric adenocarcinoma and lymphoma: pathophysiology, epidemiology, screening, clinical presentation, treatment, and prevention. *Medical Clinics of North America* **89** 313–344. (doi:10.1016/j.mcna.2004.09.001)
- Matsushima Y, Aoyama N, Fukuda H, Kinoshita Y, Todo A, Himeno S, Fujimoto S, Kasuga M, Nakase H & Chiba T 1999 Gastric ulcer formation after the Hanshin-Awaji earthquake: a case study of Helicobacter pylori infection and stress-induced gastric ulcers. *Helicobacter* **4** 94–99. (doi:10.1046/j.1523-5378.1999.98290.x)
- McColl KE 2009 Helicobacter pylori-negative nonsteroidal anti-inflammatory drug-negative ulcer. *Gastroenterological Clinics of North America* **38** 353–361. (doi:10.1016/j.gtc.2009.03.004)
- McEwen BS 2007 Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiological Reviews* **87** 873–904. (doi:10.1152/physrev.00041.2006)
- Opper L & Zimmerman HM 1938 Ulcers of the digestive tract in association with cerebral lesions. *Yale Journal of Biology and Medicine* **11** 49–84.
- Ozdemir V, Jamal MM, Osapay K, Jadus MR, Sandor Z, Hashemzadeh M & Szabo S 2007 Cosegregation of gastrointestinal ulcers and schizophrenia in a large national inpatient discharge database: revisiting the 'brain-gut axis' hypothesis in ulcer pathogenesis. *Journal of Investigative Medicine* **55** 315–320. (doi:10.2310/6650.2007.00014)
- Piper DW & Tennant C 1993 Stress and personality in patients with chronic peptic ulcer. *Journal of Clinical Gastroenterology* **16** 211–214. (doi:10.1097/00004836-199304000-00009)
- Räihä I, Kempainen H, Kaprio J, Koskenvuo M & Sourander L 1998 Lifestyle, stress, and genes in peptic ulcer disease: a nationwide twin cohort study. *Archives of Internal Medicine* **158** 698–704. (doi:10.1001/archinte.158.7.698)
- Rasheed N & Alghasham A 2012 Central dopaminergic system and its implications in stress-mediated neurological disorders and gastric ulcers: short review. *Advances in Pharmacological Sciences* **2012** 1–11. (doi:10.1155/2012/182671)
- Realo A, Teras A, Kõõts-Ausmees L, Esko T, Metspalu A & Allik J 2015 The relationship between the Five-Factor Model personality traits and peptic ulcer disease in a large population-based adult sample. *Scandinavian Journal of Psychology* **56** 693–699. (doi:10.1111/sjop.12248)
- Rosenthal J & Tobias MJ 1943 Case report: acute phlegmonous gastritis with multiple perforations. Case report. *American Journal of Surgery* **59** 117–122. (doi:10.1016/S0002-9610(43)90515-X)
- Selye H 1936 A syndrome produced by diverse nocuous agents. *Nature* **138** 32. (doi:10.1038/138032a0)
- Sikiric P, Marovic A, Matoz W, Anic T, Buljat G, Mikus D, Stancic-Rokotov D, Separovic J, Seiwerth S, Grabarevic Z, et al. 1999 A behavioural study of the effect of pentadecapeptide BPC 157 in Parkinson's disease models in mice and gastric lesions induced by 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *Journal of Physiology* **93** 505–512. (doi:10.1016/s0928-4257(99)00119-9)
- Strang R 1965 The association of gastro-duodenal ulceration and Parkinson's disease. *Medical Journal of Australia* **1** 842–843.

- Szabo S 1979 Dopamine disorder in duodenal ulceration. *Lancet* **2** 880–882. (doi:10.1016/S0140-6736(79)92690-4)
- Tache Y & Brunnhuber S 2008 From Hans Selye's discovery of biological stress to the identification of corticotropin releasing factor signalling pathways: implication in stress-related functional bowel diseases. *Annals of the New York Academy of Science* **1148** 29–41. (doi:10.1196/annals.1410.007)
- Tay CY, Windsor HM, Thirriot F, Lu W, Conway C, Perkins TT & Marshall BJ 2012 *Helicobacter pylori* eradication in Western Australia using novel quadruple therapy combinations. *Alimentary Pharmacology and Therapeutics* **11–12** 1076–1083. (doi:10.1111/apt.12089)
- Wachirawat W, Hanucharurnkul S, Suriyawongpaisal P, Boonyapisit S, Levenstein S, Jearanaisilavong J, Atisook K, Boontong T & Theerabutr C 2003 Stress, but not *Helicobacter pylori*, is associated with peptic ulcer disease in a Thai population. *Journal of the Medical Association of Thailand* **86** 672–685.
- Wijdicks EF 2011 Cushing's ulcer: the eponym and his own. *Neurosurgery* **68** 1695–1698. (doi:10.1227/NEU.0b013e318212babf)
- Wood JD 2004 The first Nobel prize for integrated systems physiology: Ivan Petrovich Pavlov. *Physiology* **19** 326–330. (doi:10.1152/physiol.00034.2004)

Received in final form 31 October 2016

Accepted 20 December 2016

Accepted Preprint published online 20 December 2016