REVIEW

Physiological and pathological implications of retinoid action in the endometrium

Yanwen Jiang¹, Lu Chen¹, Robert N Taylor², Chunjin Li^{1,*} and Xu Zhou^{1,*}¹College of Animal Sciences, Jilin University, Changchun, Jilin, China²Departments of Obstetrics and Gynecology and Molecular Medicine and Translational Sciences, Wake Forest School of Medicine, Winston-Salem, North Carolina, USACorrespondence should be addressed to X Zhou or C Li: xzhou65@vip.sina.com or llcj158@163.com

*(C Li and X Zhou contributed equally to this work)

Abstract

Retinol (vitamin A) and its derivatives, collectively known as retinoids, are required for maintaining vision, immunity, barrier function, reproduction, embryogenesis and cell proliferation and differentiation. Despite the fact that most events in the endometrium are predominantly regulated by steroid hormones (estrogens and progesterone), accumulating evidence shows that retinoid signaling is also involved in the development and maintenance of the endometrium, stromal decidualization and blastocyst implantation. Moreover, aberrant retinoid metabolism seems to be a critical factor in the development of endometriosis, a common gynecological disease, which affects up to 10% of reproductive age women and is characterized by the ectopic localization of endometrial-like tissue in the pelvic cavity. This review summarizes recent advances in research on the mechanisms and molecular actions of retinoids in normal endometrial development and physiological function. The potential roles of abnormal retinoid signaling in endometriosis are also discussed. The objectives are to identify limitations in current knowledge regarding the molecular actions of retinoids in endometrial biology and to stimulate new investigations toward the development potential therapeutics to ameliorate or prevent endometriosis symptoms.

Key Words

- ▶ retinoids
- ▶ endometrial stroma
- ▶ implantation
- ▶ endometriosis

Journal of Endocrinology
(2018) **236**, R169–R188

Introduction

In many species, such as human, nonhuman primates and rodents, the endometrium consists of epithelial glands supported by a connective tissue stroma that undergoes cycles of proliferation and secretory activity. The uterine mucosa proliferates under the influence of estrogen. However, after ovulation, luteal progesterone changes the proliferative pattern to a secretory pattern that includes decidualization of endometrial stromal cells, which provides the nutritive and immune-privileged matrix for embryo implantation (Gellersen & Brosens 2014) and prevents the malignant transformation of endometrial

epithelium under unopposed estrogenic action (Cheng *et al.* 2008). Though progesterone is critical for the initiation and maintenance of decidualization in the estrogen-primed endometrium, other factors, including prostaglandins, prolactin, growth factors and extracellular matrix proteins, are important.

Retinoic acid (RA), the physiological active metabolite of vitamin A (retinol), controls multiple biological processes, including differentiation, apoptosis and cell survival, via its nuclear receptors RARs or nonclassical RA receptor peroxisome proliferator-activated

receptor β/δ (PPAR β/δ). The pathway of RA is also considered to be involved in the proliferation of epithelia and the transformation of endometrial stromal cells into specialized decidual cells. Numerous studies have demonstrated that the distribution of RA (Zheng *et al.* 2000), as well as the expression of RA receptors (Fukunaka *et al.* 2001, Ozaki *et al.* 2017), cellular retinol and RA-binding proteins (CRBP1, CRABP1 and CRABP2; Zheng & Ong 1998, Zheng *et al.* 2000), RA-synthesizing enzymes ALDH1A1 and ALDH1A2 (Napoli 1999, Duester 2000), and the RA-catabolizing enzyme CYP26A1 (Vermot *et al.* 2000) are highly and differentially regulated in the differentiating endometrium during the ovarian cycle and during the phase of blastocyst implantation (Vermot *et al.* 2000, Zheng 2000, Ozaki *et al.* 2017). The expression of these same genes can be modulated by ovarian steroid hormones (i.e., estradiol and progesterone) treatment (Vermot *et al.* 2000, Li & Ong 2003, Rühl *et al.* 2006, Fritzsche *et al.* 2007). The retinoid pathway is known to play vital roles in endometrial development and differentiation (Tanmahasamut & Sidell 2005, Wu *et al.* 2013, Nakajima *et al.* 2016), as well as endometrial neovascularization (Sidell *et al.* 2010) and blastocyst implantation (Han *et al.* 2010, Xia *et al.* 2010, Ma *et al.* 2012).

In addition, accumulating evidence shows that an aberrant RA metabolism can be a critical factor in the development of endometriosis (Pavone *et al.* 2011, Wieser *et al.* 2012, Pierzchalski *et al.* 2014, Yamagata *et al.* 2015), a common gynecological disease that affects up to 10% of reproductive-age women. This disease is characterized by the ectopic localization of endometrial-like tissues in the pelvic cavity, and its pathogenesis involves uncontrollable cell proliferation and is associated with local invasion and distant metastasis. Among the numerous aspects of endometrial behavior regulated by RA are matrix metalloproteinase secretion, gap junctional intracellular communication and the production of various cytokines involved in stromal cell growth, adhesion and differentiation (Wu *et al.* 2013). Interleukin 6 (IL-6), monocyte chemotactic protein 1 (MCP-1), tumor necrosis factor α (TNF- α), vascular endothelial growth factor (VEGF) and connexin43 are all aberrantly expressed in endometriotic lesions (Sawatsri *et al.* 2000, Sharpe-Timms 2001, Nozaki *et al.* 2006, Sidell *et al.* 2010, Wu *et al.* 2013). Thus, many seemingly discordant features of endometriosis, including decreased cell death, increased growth and migration, inflammation and enhanced invasive properties of intraperitoneally seeded endometrial cells, might be accounted for by the dysregulation of RA

signaling. This contention was recently supported in a mouse model of endometriosis where treatment with RA suppressed IL-6 and the establishment, growth and vascularity of peritoneal implants, promoted macrophage differentiation (Wieser *et al.* 2012).

RA plays fundamental roles in the normal maintenance of endometrial physiology (Zheng *et al.* 2000, Sidell *et al.* 2002, Ding *et al.* 2003, Kuroda *et al.* 2013), and aberrant RA metabolism might be a predisposing factor for the development of endometriosis (Pavone *et al.* 2011, Wieser *et al.* 2012, Pierzchalski *et al.* 2014, Yamagata *et al.* 2015). However, a comprehensive and systematic analysis and understanding of RA pathway in endometrial physiology and pathology remain lacking. In this review, the latest advances with regard to understanding the retinoid pathway are first discussed, followed by an assessment of the functional roles of this signaling network in endometrial development and physiological function. This review also summarizes evidence that supports fundamental defects in retinoid metabolism and action among women with endometriosis. The objectives are to identify limitations in our current knowledge regarding molecular retinoid actions in endometrial biology and to propose future investigations to develop therapeutic or preventative agents for clinical endometriosis management.

Retinoid metabolism and signaling

Retinol (vitamin A) is a lipid-soluble vitamin that cannot be synthesized *de novo* by animals and is obtained from the diet as either preformed retinoids or carotenoids within the intestine. In the enterocytes, proretinoid carotenoids, such as β -carotene, can be either cleaved and converted to retinoids or incorporated intact and unmodified along with dietary fat and cholesterol into chylomicrons. Dietary retinoids that are newly absorbed by the enterocytes are esterified to retinyl ester and packaged with dietary fat and cholesterol into chylomicrons, which are later secreted into the lymphatic system. Approximately 66%–75% of newly acquired retinoids are absorbed and stored in the liver as lipid droplet retinyl esters (Harrison 2005). The remainder is absorbed by extrahepatic tissues (Quadro 2004). In cases of dietary vitamin A deficiency, the stored retinoid is hydrolyzed and mobilized back to retinol, which can be bound by RBP4 and enter the bloodstream for transport to peripheral tissues. In blood, holo-RBP4 is bound to transthyretin (TTR), a carrier protein for thyroid hormones. In a fasting circulation, retinol-RBP4

is the preponderant retinoid form comprising >95% of retinoids, whereas, after a retinoid-rich meal, chylomicron retinyl ester concentrations exceed those of retinol-RBP4. In extrahepatic tissues, retinoids are acquired from the circulation in the form of carotenoids and retinyl esters from chylomicrons and retinol from retinol-RBP4-TTR ternary complexes. The uptake of retinol from retinol-RBP4-TTR into cells is mediated by the transmembrane-spanning protein stimulated by retinoic acid 6 (STRA6), which acts as a high-affinity cell surface receptor that binds extracellular holo-RBP, facilitates the dissociation of retinol from its carrier and transports it into cells (Kawaguchi *et al.* 2007, 2015).

Within target cells, carotenoids and retinyl esters can be both converted into retinol, which binds cellular retinol-binding proteins (CRBP1 and CRBP2).

Cellular retinol can be stored in the form of retinyl ester catalyzed by lecithin: retinol acyltransferase (LRAT) or converted into transcriptionally active RA metabolites via two enzymatic reactions. First, retinol is converted into retinaldehyde by alcohol dehydrogenases (ADHs) or retinol dehydrogenases (RDHs) (Fig. 1). Aldehyde dehydrogenases (ALDHs) then catalyze the conversion of retinaldehyde into RA (Conaway *et al.* 2013). In the retinal pigment epithelium of the eye, retinol can also be metabolized to 11-*cis*-retinal, which serves to regenerate the visual pigment rhodopsin.

Most of the physiologic actions of retinoids are accounted for by the transcriptional regulatory activity of physiological active metabolites, i.e., all-*trans*-retinoic acid (atRA or RA) and 9-*cis* retinoic acid (9cRA). The action of RA is mediated by nuclear receptors, RA receptors

Figure 1

Schematic representation of retinoid uptake, metabolism and signaling in endometrial stromal cells (ESC) and endometrial epithelial cells (EEC). ESC can take up retinoids from the circulation in the forms of retinyl esters in chylomicrons (following a retinoid-rich meal), RBP4-bound retinol (ROH) (during fasting) or albumin-bound RA (very low levels). The transport of retinol from retinol-RBP4 into ESC is mediated by STRA6; however, the cellular processes that mediate retinyl ester and RA uptake are not yet established. Carotenoids in chylomicrons also represent a source of retinoids and can be converted to retinal (Rald) by β -carotene-15, 15'-monooxygenase (BCMO1). Cellular retinol can be oxidized to retinal and RA by dehydrogenases (ADHs, RDHs and ALDHs) or be converted to retinyl ester, a retinoid storage form, catalyzed by LRAT. Retinyl esters can also be hydrolyzed to retinol catalyzed by retinyl ester hydrolase (REH). RA exerts autocrine and paracrine transcriptional regulatory effects through RARs, RXRs and PPAR β/δ or alternatively degraded by CYP26 enzymes in ESC and neighboring EEC. A full colour version of this figure is available at <https://doi.org/10.1530/JOE-17-0544>.

(RARs): RAR α , RAR β and RAR γ or peroxisome proliferator-activated receptor β/δ (PPAR β/δ). In conjunction with transcriptional coactivators and corepressors, RARs and PPARs heterodimerize with retinoid X receptors (RXRs: RXR α , RXR β and RXR γ) and associate with RA response elements (RAREs) or peroxisome proliferator response elements (PPREs) in the regulatory regions of specific target genes to regulate target gene transcription (Chambon 2005, Schug *et al.* 2008, Al Tanoury *et al.* 2013, Rochette-Egly 2014). RA partitioning between the two receptor signaling cascades has opposing effects on cell growth and apoptosis and the alternative pathways are regulated by the ratio of intracellular lipid-binding proteins, cellular RA-binding protein (CRABP2) and fatty acid-binding protein (FABP5; Schug *et al.* 2007). CRABP2 and FABP5 can bind and transport RA to RARs and PPAR β/δ , respectively (Delva *et al.* 1999, Dong *et al.* 1999). Furthermore, besides classical nuclear receptor signaling, RA stimulates rapid nongenomic signaling events through the stimulation of kinase phosphorylation by extra-nuclear RARs, which also can affect gene transcription (Al Tanoury *et al.* 2013, Rochette-Egly 2014). The primary biological ligand for RARs is atRA, whereas 9cRA can bind to RARs and RXRs.

The tissue distribution and levels of RA are regulated through its synthesis by ALDHs (Lin *et al.* 2003) and its catabolism into inactive polar compounds by CYP26 enzymes (Chithalen *et al.* 2002, Conaway *et al.* 2013). CRABPs and FABP5 also participate in the regulation of RA concentration (Dong *et al.* 1999, Noy 2000) and metabolism (Delva *et al.* 1999, Dong *et al.* 1999, Napoli 2016). In RA-sensitive tissues, CRABP1 can protect these vulnerable areas against the toxic effects of excessive amounts of RA (Ruberte *et al.* 1992). Furthermore, many of the components of RA biosynthesis are controlled by atRA via negative feedback. ATRA induces its own catabolism by inducing CYP26 transcription, allowing the degradation of excess atRA. Conversely, atRA can also regulate its biosynthesis by positive feedback, upregulating the expression of several RA biosynthesis components, such as RBP4, CRBP1 and STRA6 (Wu & Ross 2010).

Retinoid action in endometrial development and physiological function

The mammalian uterus develops from fused paramesonephric (Müllerian) ducts during embryonic development. The adult mammalian endometrium is divided into two steroid-responsive compartments, namely, an epithelial compartment consisting of luminal

and glandular epithelia and a stromal compartment filled with fibroblasts, microvascular cells, macrophages and perivascular and large granular lymphocytes that infiltrate the stroma during the luteal phase of the menstrual cycle (Punyadeera *et al.* 2003). Luminal epithelium is the first point of contact between the uterus and blastocyst during implantation, while glandular epithelia synthesize and secrete bioactive substances that contribute to uterine receptivity and stromal cell decidualization. The endometrial functional layer undergoes cyclic regeneration regulated by estradiol and progesterone during menstrual or estrous cycles. It initially proliferates under the influence of estrogen. However, after ovulation, in addition to estradiol, the ovary starts to produce progesterone, which changes the proliferative pattern to a secretory one, providing a suitable environment for blastocyst implantation. Following fertilization, the embryo implants into the uterine wall and trophoblast-derived gonadotropins support the pregnancy by maintaining corpus luteum production of progesterone and estradiol. In this case, the endometrial functional layer remains as decidua. If fertilization or implantation fails, the endometrium will be either remodeled (estrous cycle) or shed (menstrual cycle). Numerous factors are involved in the regulation of endometrial development and function and RA plays important roles in these processes.

Hormonal control of endometrial retinoid pathway

Numerous studies revealed that the retinoid pathway and distribution of RA are distinctly and highly regulated in human and rodent endometrial epithelium and stroma (Vermot *et al.* 2000, Rühl *et al.* 2006, Fritzsche *et al.* 2007). The biosynthesis of endogenous RA is catalyzed directly by ALDHs. Among them, ALDH1A1 (also known as RALDH1) and ALDH1A2 (also known as RALDH2) exhibit specific temporal and special expression patterns in human and rodent endometrium during the ovarian cycle and early pregnancy. The abundance of *Aldh1a1* mRNA increases during murine diestrus and proestrus, whereas *Aldh1a2* is highly induced in metestrus (Vermot *et al.* 2000). The tissue localization of ALDH1A1 and ALDH1A2 showed that *ALDH1A1* and *ALDH1A2* are expressed in glandular epithelial and stromal cells, respectively (Vermot *et al.* 2000). The highest *Aldh1a2* expression in mouse stromal cells is observed during the estrogenic phase of the estrous cycle, strongly suggesting that its endogenous expression is estrogen dependent. This speculation was confirmed by subsequent studies in the endometrium of mice, rats and

Table 1 Upregulated and downregulated genes involved in retinoid metabolism and signaling by estrogen and/or progesterone in the endometrium.

Genes	E ₂	P ₄	E ₂ + P ₄	Cell type	Species	References
<i>ALDH1A1</i>	↓ ^a	—		Epithelium	Mouse	Vermot <i>et al.</i> (2000), Rühl <i>et al.</i> (2006)
<i>ALDH1A2</i>	↑ ^b	—	↑	Stroma	Human, mouse, rat	Vermot <i>et al.</i> (2000), Deng <i>et al.</i> (2003), Li <i>et al.</i> (2004), Rühl <i>et al.</i> (2006)
<i>CYP26A1</i>	— ^c	↑	↑	Epithelium	Mouse	Fritzsche <i>et al.</i> (2007)
<i>CRABP2</i>	↑	^d		Epithelium	Mouse, rat	Wardlaw <i>et al.</i> (1997), Li & Ong (2003)
<i>RARα</i>	↑			Epithelium	Rat	Boehm <i>et al.</i> (1997)
<i>RARγ</i>	↑			Epithelium	Rat	Boehm <i>et al.</i> (1997)
<i>RXRα</i>	↑			Epithelium	Rat	Boehm <i>et al.</i> (1997)

^aDownregulation; ^bupregulation; ^cno effect; ^d(blank) remains to be established.E₂, estradiol; P₄, progesterone.

humans (Deng *et al.* 2003, Li *et al.* 2004, Rühl *et al.* 2006). Using ovariectomized mouse models, Rühl *et al.* (2006) demonstrated that *ALDH1A2* expression is rapidly (within 1–4 h) induced in stromal cells by estradiol but not by progesterone. By contrast, estradiol, but not progesterone, treatment decreases (within 4–24 h) *ALDH1A1* expression levels in glandular epithelium (Rühl *et al.* 2006). In a study by Deng *et al.* (2003), *ALDH1A2* expression was induced within human endometrium by estrogen replacement therapy. Thus, estradiol stimulates *ALDH1A2* expression in endometrial stroma but suppresses *ALDH1A1* expression in the endometrial glandular epithelium. By contrast, progesterone neither induces nor reduces the expression levels of the two genes (Table 1). The selective effects of estradiol are due to the differential expression of estrogen nuclear receptor isotypes ERα and ERβ (Katzenellenbogen & Korach 1997). In rat uteri, ERα is principally expressed in luminal and glandular epithelia, whereas ERβ is predominantly expressed in the stroma (Mowa & Iwanaga 2000a,b). As *ALDH1A2* expression might be selectively regulated via ERβ, further study should focus on using ERα/ERβ-selective ligands to dissect these estrogen-regulated pathways in the regulation of *ALDH* expression. Meanwhile, a potential estrogen response element was observed in the promoter of the mouse *Aldh1a2* gene (Wang *et al.* 2001). However, whether it plays a functional role remains unknown.

In addition to RA-synthesizing enzymes, RA levels are also regulated by RA-catabolizing enzymes, particularly the CYP26 family. These enzymes catalyze the conversion of RA into less-active polar metabolites (Conaway *et al.* 2013). CYP26A1, one of the most important members of the CYP26 family, is expressed in female reproductive tract (Vermot *et al.* 2000). In endometrial tissue obtained from premenopausal women, the *CYP26A1* mRNA levels were approximately 20 times higher during the late secretory phase than in the proliferative phase (Deng *et al.* 2003).

In mouse endometrium, although *CYP26A1* expression is undetectable during the normal ovarian cycle, it is strongly induced in luminal and glandular epithelia, 4 h after progesterone administration (Fritzsche *et al.* 2007) or 24 h after human chorionic gonadotrophin (hCG) administration (Vermot *et al.* 2000). Also, in mouse endometrial epithelium, *Cyp26a1* expression is strongly induced between 3.5 and 4.5 gestational days, i.e., when the developing blastocysts implant into the endometrium. Endometrial *Cyp26a1* expression can be mainly regulated by progesterone because not only *Cyp26a1* mRNA levels are strongly increased in uterine luminal and glandular epithelial cells by the administration of progesterone or the combination of progesterone and estradiol but also the simultaneous administration of an antiprogesterone inhibits the gene expression (Fritzsche *et al.* 2007) (Table 1). This regulation of *Cyp26a1* expression by progesterone may occur through direct activation of the progesterone receptor. RU486 (also known as mifepristone) is a synthetic steroid that exerts antiprogesterone action by competing with progesterone for receptor binding (Chan *et al.* 2003). Treatment with either RU486 or the combination of RU486 and progesterone showed an inhibition of *Cyp26a1* expression (Fritzsche *et al.* 2007). The mouse *Cyp26a1* promoter and its regulatory regions were partially characterized by Loudig and coworkers (2000, 2005). Their study showed the presence of two RA response elements on the *Cyp26a1* promoter, but no consensus binding sites for progesterone receptors. Thus, further studies are necessary to identify the possible regulatory elements responsible for the progesterone control of *Cyp26a1* expression.

Several binding proteins involved in the retinoid pathway are also distinctive in rodent and human endometria during the ovarian cycle (Loughney *et al.* 1995, Wardlaw *et al.* 1997). *Crbp1* expression peaked during diestrus, whereas *Crabp2* expression peaked

sharply during estrus (proliferative phase). In addition, *Crabp2* expression has been shown to be induced directly by estradiol administration in the uteri of ovariectomized rats (Li & Ong 2003) (Table 1). Immunohistochemical studies showed that CRABP2 is localized to the luminal epithelium, whereas CRBP1 in the stroma (Wardlaw *et al.* 1997). CRABP2 expression is associated with local RA levels (Bucco *et al.* 1997, Zheng *et al.* 2000). From these data, it can be concluded that the endometrial stroma upregulates ALDH1A2 and therefore RA production upon estrogen stimulation. RA then diffuses into the adjacent epithelium, where it is catabolized by CYP26A1 induced by progesterone during the late secretory phase or implantation.

Numerous studies demonstrated the expression of RARs and RXRs in human endometrial epithelial and stromal cells (Prentice *et al.* 1992, Loughney *et al.* 1995, Kumarendran *et al.* 1996, Fukunaka *et al.* 2001). Using Northern blot, Kumarendran *et al.* (1996) demonstrated that *RARβ* mRNA expression is 1.7-fold higher in epithelial samples from the proliferative phase than from the secretory phase. Meanwhile, no significant difference was observed between the expression levels of *RARα*, *RARγ* and *RXRα* mRNA in epithelial tissues in the proliferative phase and those in the secretory phases. Similarly, the expression of *RARα*, *RARβ*, *RARγ* and *RXRα* mRNA in human endometrial stroma did not change in the proliferative and secretory phases of the menstrual cycle (Kumarendran *et al.* 1996). Meanwhile, using immunohistochemistry and Western blot analysis, Fukunaka *et al.* (2001) investigated changes in expression and subcellular localization of RARs and RXRs in human endometrial epithelium during the menstrual cycle. Their results indicated that, in the nuclei of endometrial epithelia, RARs and RXRs are expressed strongly in the proliferative phase and reduced during the secretory phase, correlated with serum estradiol concentration and ER expression (Fukunaka *et al.* 2001). Although RXRs decrease during the secretory phase,

they are still expressed in the nuclei in the late secretory phase; by contrast, RARs are infrequently detected in the midsecretory and late secretory phases (Fukunaka *et al.* 2001). In stromal cells, the intensity of staining is strong in the proliferative phase but decreases in the secretory phase. Meanwhile, immunoreactivity to RXRs is strong throughout the cycle, similar to that in the nuclei of the epithelial cells (Fukunaka *et al.* 2001). In addition, ERα shows more intense staining in the endometrium of the proliferative phase, and PR shows positive staining in the endometrium of the mid-proliferative and early secretory phases (Fukunaka *et al.* 2001). The expression of retinoid receptors is associated with serum estradiol concentration and ER expression. Consistently, data from Boehm *et al.* (1997) showed that *RARα*, *RARγ* and *RXRα* expression levels are increased by estrogen in rat endometrial epithelium (Table 1). Thus, the expression of RARs is strongly affected by serum estrogen via ER in the endometrium. However, whether progesterone or PR expression inhibits the expression of retinoid receptors in endometrium remains to be determined.

Therefore, the endometrial retinoid pathway is mainly under the influence of ovarian steroid hormones, and RA and its receptors have regulatory effects on the endometrium during the ovarian cycle (Fig. 2).

Retinoid action in endometrial differentiation and development

Fate determination of endometrium in Müllerian duct

During development, the Müllerian ducts persist in females and differentiate into morphologically and functionally distinct oviducts, uterus and vagina. The oviductal epithelia are composed of ciliated cells and secretory cells (Yamanouchi *et al.* 2010); the uterine epithelia consist of simple columnar luminal and glandular epithelia (Kurita *et al.* 2000) and the vaginal epithelium

Figure 2

Hormonal control of the retinoid pathway in the endometrium. Estradiol (E_2) or E_2 + progesterone (P_4) can stimulate the expression of ALDH1A2, which can promote RA production, in stromal cells. RA produced by ESC can diffuse to EEC, where E_2 can promote the expression of RARs, PPAR β/δ and CRABP2. Thus, E_2 can increase RA signaling in the endometrium. However, P_4 can inhibit RA signaling through inducing CYP26A1 expression in EEC. A full colour version of this figure is available at <https://doi.org/10.1530/JOE-17-0544>.

develops into stratified cuboidal epithelium (Nakajima *et al.* 2011a). Nakajima *et al.* (2011a) performed tissue recombination experiments with epithelium and stroma to investigate the mechanisms underlying differentiation of the Müllerian ducts. The data showed that stromal factors determine the fate, differentiation and growth of Müllerian duct epithelia into the oviducts, uteri and vaginas of mice. DNA microarray analysis showed that the expression of several genes involved in retinoid metabolism is greater in the uteri than in the vaginas of neonatal mice (Suzuki *et al.* 2006, 2007). When pregnant rats are fed with a vitamin A-deficient diet, their fetuses exhibit incomplete Müllerian development (Wilson & Warkany 1948). Furthermore, Müllerian ducts of RAR or RXR KO mice are absent at embryonic day 12.5 (E12.5) (Mendelsohn *et al.* 1994, Kastner *et al.* 1997). These data suggest that retinoid signaling through RAR is essential for Müllerian duct development.

The important role of the retinoid pathway in fetal uterine stroma development was identified in the mouse. In fetal mice, *Rdh10* and *Aldh1a2* are expressed in the stroma of proximal Müllerian ducts, with expression levels decreasing toward the caudal ends (Nakajima *et al.* 2016). RDH10 is the primary enzyme responsible for catalyzing the conversion of retinol to retinaldehyde, the first step of RA synthesis, in embryonic mice (Farjo *et al.* 2011). Uterine *Cyp26a1* expression is essential for blastocyst implantation in pregnant mice (Han *et al.* 2010). However, *Cyp26a1* expression levels are low in Müllerian ducts, suggesting RA is not degraded there and more RA accumulates. Thus, RA levels might be increased in proximal Müllerian ducts. This hypothesis was confirmed in a reporter mouse expressing lacZ regulated by a strong RARE in the promoter. The data demonstrated that the highest endogenous RA levels are in the proximal Müllerian duct stroma, and the levels gradually decrease from the proximal to caudal regions (Nakajima *et al.* 2016). Thus, RA is produced and acts in the stroma of proximal and middle Müllerian ducts. *Rdh10* and *Aldh1a2* expression are significantly higher in the uteri of postnatal mice than those in their vaginas from postnatal day 2 (P2) to P90 (Nakajima *et al.* 2016). Moreover, mice fed from birth with vitamin A-deficient diets for 10 or 14 weeks exhibit squamous metaplasia in their uterine epithelia (Darwiche *et al.* 1993, Jetten *et al.* 1996). These data suggest a critical role of RA signaling in maintaining uterine epithelium.

In organ-cultured Müllerian ducts of RARE-lacZ mice at E14.5 and E17.5, retinal or RA treatment stimulated RA signaling in the stroma and induced uterine epithelial

differentiation, which was defined as a layer of columnar epithelial cells negative for oviductal and vaginal epithelial markers. By contrast, inhibition of RAR signaling with the pan-RAR antagonist AGN193109 (AGN; Nakajima *et al.* 2016) induces vaginal epithelial differentiation. Nakajima *et al.* (2016) cultured Müllerian duct explants at E14.5 or E17.5 with RA or RAR antagonists and grafted these under the renal capsules of host mice. At day 30 post grafting, control E14.5 Müllerian ducts developed oviductal and uterine epithelia, while AGN treatment irreversibly induced vaginal epithelia. Meanwhile, grafted E17.5 middle Müllerian ducts had only uterine epithelia, and AGN treatment irreversibly induced vaginal epithelia. In grafted E17.5 caudal Müllerian ducts, vaginal epithelia were observed, and RA treatment permanently induced uterine epithelia but did not induce oviductal epithelia. These data indicate that RA in stroma at developing stages induces uterine epithelia in Müllerian ducts, whereas the inhibition of RAR signaling induces vaginal epithelia. The fate of oviductal stroma is determined at E14.5 (Nakajima *et al.* 2016). In female mouse embryos, activin A induced by inhibin β A is essential for the differentiation of vaginal epithelia (Nakajima *et al.* 2011b), whereas RA inhibits inhibin β A expression in the urogenital sinus (Bryant *et al.* 2014). *Hoxa10* is expressed at the boundary between future uterine and oviductal stroma in the Müllerian ducts at E16.5 (Ma *et al.* 1998) and is connected with the determination of the borderline between oviduct and uterus (Benson *et al.* 1996). RA stimulates *Hoxa10* and *Hoxa11* expression in Müllerian ducts; however, AGN abolishes these. Although the *Hoxa10* promoter contains a RARE (Delacroix *et al.* 2010), RAR cannot bind to the predicted RARE in the Müllerian duct (Nakajima *et al.* 2016). Thus, RA seems to induce *Hoxa10* expression via indirect RAR signaling, and other factors might mediate the action of RA on the determination of the border between the uterine and vaginal stroma. These data support a model in which RA-RAR signaling can play a crucial role in the determination of the fate of epithelia and stroma in the female reproductive tract (Fig. 3).

Endometrial gland development

Uterine glands are essential for pregnancy on account of secreting or transporting bioactive substances that regulate uterine receptivity and blastocyst implantation. Uterine gland development or adenogenesis is a postnatal event in rodents, domestic animals and humans (Hu *et al.* 2004). At birth, murine uteri lack uterine glands and consist of simple luminal epithelium supported by undifferentiated mesenchyme. Between P0 and P9, glandular epithelial

Figure 3

Schematic representation of retinoid action in endometrial development and physiological function. RA signaling in the Müllerian duct determines the differentiation of epithelium to form the future uterus and vagina involving *Hoxa10*, inhibin β A and activin A. Besides, RA signaling stimulates endometrial GJIC formation through inducing Cx43 expression and neovascularization through inducing VEGF expression. RA prevents stromal cell decidualization through inhibiting PRL, IGFBP1 and HSD11B1 production and promotes maintenance of fibronectin, an indicator of the undifferentiated stromal phenotype. In addition, RA reduces factors like LIF and HB-EGF1, which are required for endometrial receptivity, and successful implantation during late secretory phase requires lower RA levels in the luminal epithelia. A full colour version of this figure is available at <https://doi.org/10.1530/JOE-17-0544>.

cells differentiate and bud from luminal epithelia. By P15, the uterine histoarchitecture resembles that of the adult (Spencer *et al.* 2012). Gland morphogenesis also occurs during endometrial regeneration following menses or parturition (Garry *et al.* 2010, Huang *et al.* 2012). Based on data from the available literature, the luminal and glandular epithelia of mouse uteri have common but distinctly expressed genes (Cha *et al.* 2012, Filant & Spencer 2013). The components of the retinoid pathway, including *Aldh1a1*, *Aldh1a3*, *Rdh1* and *Rdh10*, are enriched in the glandular epithelia of neonatal uteri. Through microarray analysis, the glandular epithelium-enriched genes in P10 uteri were found to be associated with branching morphogenesis, growth and RA biosynthesis. *Aldh1a1* is enriched in developing neonatal and adult endometrial glands (Filant & Spencer 2013). Furthermore, the expression levels of *Aldh1a1* and *Aldh1a3* on P10 are much lower in progesterone-induced uterine gland-knockout (PUGKO) mice than in control mice (Filant & Spencer 2013). In addition, retinyl palmitate increases glandular epithelial areas in the uteri of neonatal pigs treated for 14 days starting from birth (Vallet *et al.* 1995). These data suggest that the retinoid pathway is involved in uterine adenogenesis. Further studies are still needed to elucidate the functional role of the retinoid pathway and the mechanisms governing uterine adenogenesis in neonates. The involvement of RA signaling in postnatal mouse endometrial adenogenesis will require conditional postnatal deletion models, such as the PR-Cre system.

Endometrial proliferation

Ovarian estrogen induces the proliferation of epithelial cells and stromal fibroblasts during the pre-ovulatory

phase (Punyadeera *et al.* 2003). Studies in rodents suggested that RA plays an important role in modulating the effects of estrogen on the endometrium. As discussed earlier, endometria upregulate *ALDH1A2* upon estrogen stimulation during the proliferative phase (Vermot *et al.* 2000, Deng *et al.* 2003, Rühl *et al.* 2006). Furthermore, the expression levels of RA receptors (Fukunaka *et al.* 2001) and CRABP2 (Loughney *et al.* 1995, Wardlaw *et al.* 1997, Li & Ong 2003), which bind and deliver RA to nuclear RAR, are stimulated by estradiol in the proliferative phase in epithelial and stromal cells. These data suggest that estrogen coordinately upregulates RA production, transport and signaling in the proliferative phase of the ovarian cycle. This coordinated mechanism may play a role in antiproliferative effects that counterbalance estrogen-induced endometrial proliferation. Studies on vitamin A-deficient rats demonstrated that physiologic levels of RA suppress endometrial hyperplasia and metaplasia associated with chronic estrogen administration (Bo & Smith 1966). In addition, pharmacological doses of RA in immature ovariectomized rat suppress estrogen-induced endometrial cell proliferation (Boettger-Tong & Stancel 1995, Loughney & Redfern 1995). This might be a beneficial suppressive effect on estrogen-induced endometrial proliferation. RA can also effectively suppress the estrogen-stimulated proliferation of breast cancer cells and in several endometrial carcinoma lines (Carter *et al.* 1996, Toma *et al.* 1997). Targeting physiological RA accumulation in the endometrium by estradiol is a potential suppressor mechanism to prevent malignant transformation of hyperplastic endometrial cells during the ovarian cycle and gestation. In the case of vitamin A deficiency, uterine epithelium forms regions of

keratinized squamous metaplasia, indicating that RA is needed to maintain simple columnar uterine epithelium (Ponnamperuma *et al.* 1999).

Gap junction intercellular communications

Decidualized stromal cells not only acquire epithelioid structure and function, accumulating glycogen, lipids and subcellular organelles (Lawn *et al.* 1971) but also express interdigitating lamellar processes within clustered microdomains called gap junctions. These gap junctions act as membranous channels for the direct exchange of small molecules (e.g., second messengers) among adjacent cells or between these cells and their extracellular environments (Yamasaki *et al.* 1999). Gap junction intercellular communications (GJIC) are critical to the success of decidualization because their blockade suppresses the proliferation and differentiation of uterine stromal cells (Yu *et al.* 2011, Diao *et al.* 2013, Yu *et al.* 2014a, Winterhager & Kidder 2015). Connexin (CX) proteins are the major components of gap junctions, and CX43 (also known as GJA1) is the predominant CX protein in endometrial stromal cells (Jahn *et al.* 1995, Winterhager *et al.* 2009) and undergoes a variety of changes during the ovarian cycle (Mantena *et al.* 2006). The attenuation of CX43 expression disrupts GJIC between neighboring cells and impedes the differentiation of stromal cells in human and mouse endometrium (Laws *et al.* 2008, Yu *et al.* 2011). Endometrial CX43 may be regulated by hormonal changes, and the progesterone and estrogen levels in serum are reported to determine the expression levels of CX43 in the endometrium (Yu *et al.* 2014b, Winterhager & Kidder 2015). RA and other retinoids enhance GJIC and CX43 expression levels in numerous cell types, certain human cancer cell types, mouse fibroblasts and rat liver cells, through transcriptional and translational mechanisms (Stahl & Sies 1998, Carystinos *et al.* 2001). When treated with RA, CX43

expression in human endometrial stromal cells shows a dose-dependent increase at the mRNA and protein levels (Tanmahasamut & Sidell 2005) (Table 2). Concomitant with the increase in CX43 expression, stromal cells treated with RA exhibit a 2.5-fold enhancement in their GJIC as assessed by dye transfer experiments (Tanmahasamut & Sidell 2005). The ability of endometrial stromal cells to serve as an RA-responsive target is supported by the presence of RA nuclear receptors (i.e., RARs and RXRs) in these cells (Prentice 1992, Comptour *et al.* 2016). In addition to its expression level, the phosphorylation status of CX43 influences GJIC, including gap junction assembly and channel gating (Solan & Lampe 2009). In endometrial tissue, phosphorylation of CX43 mostly impairs GJIC (Tanmahasamut & Sidell 2005, Wu *et al.* 2013). Phosphorylation of CX43 is mediated by two serine/threonine protein kinase families, protein kinase C (PKC) (Oh *et al.* 1991) and MAPK (Warn-Cramer *et al.* 1996). RA induced the CX43 de-phosphorylation at serine 262 in human endometrial stromal cells (Tanmahasamut & Sidell 2005, Wu *et al.* 2013). RA antagonized the effects of 12-O-tetradecanoylphorbol-13-acetate (a PKC activator) and inhibited PKC activity in numerous cell systems (Cope 1986, Verma 1988, Nakagawa *et al.* 2003), although the activation of PKC activity by RA has also been reported (Kambhampati *et al.* 2003). Inhibition of MAPK activity by RA was previously reported (Nakagawa *et al.* 2003). The signaling pathway(s) involved in the CX43-dephosphorylating effects of RA remains to be elucidated, but protein phosphatase 2 has been implicated (Wu *et al.* 2013). Overall, these data support that retinoid signaling can enhance GJIC among endometrial stromal cells (Fig. 3).

Endometrial neovascularization

Increased vascular permeability and angiogenesis are vital to the successful stromal decidualization, embryo

Table 2 Effect of RA on the production of factors in the endometrial cells.

Factors	Regulation	Cell type	Species	References
CX43	↑	Stromal cells	Human	Tanmahasamut & Sidell (2005), Wu <i>et al.</i> (2013)
VEGF	↑	Stromal cells	Human, mouse	Matsumoto & Sato (2006), Sidell <i>et al.</i> (2010)
PRL	↓	Stromal cells	Human	Brar <i>et al.</i> (1996), Ozaki <i>et al.</i> (2017)
IGFBP1	↓	Stromal cells	Human	Brar <i>et al.</i> (1996), Ozaki <i>et al.</i> (2017)
HSD11B1	↓	Stromal cells	Human	Ozaki <i>et al.</i> (2017)
Fibronectin	↑	Stromal cells	Human	Brar <i>et al.</i> (1996)
LIF	↓	Epithelial cells	Mouse	Ma <i>et al.</i> (2012)
HB-EGF1	↓	Epithelial cells	Mouse	Ma <i>et al.</i> (2012)
HSD17B2	↑	Epithelial cells	Human	Cheng <i>et al.</i> (2008), Yamagata <i>et al.</i> (2015), Pavone <i>et al.</i> (2017)

↑, Upregulation; ↓, Downregulation.

implantation and placentation (Dey *et al.* 2004, Matsumoto & Sato 2006, Wang & Dey 2006, Matsumoto *et al.* 2007). VEGF, originally reported as a vascular permeability factor, is also a potent mitogen for endothelial cells and a key regulatory growth factor for vasculogenesis or angiogenesis (Ferrara *et al.* 1997). In human endometrial stromal cells, RA can combine with transcriptional activators of VEGF to stimulate the expression and secretion of VEGF through a translational mechanism mediated by reactive oxygen species (ROS; Sidell *et al.* 2010) (Table 2). In addition, GJIC plays a key role in endometrial neovascularization. The conditional deletion of *CX43* genes in stromal cells and the consequent disruption of GJIC result in the reduction of VEGF production and striking impairment in the development of new blood vessels within stromal compartments (Laws *et al.* 2008, Yu *et al.* 2011). Notably, *CX43* overexpression promotes VEGF secretion (Yu *et al.* 2016). As discussed earlier, retinoid signaling can enhance GJIC among endometrial stromal cells by stimulating *CX43* expression and dephosphorylation. Thus, retinoid signaling might promote neovascularization by enhancing stromal *CX43* expression and GJIC. Evidence showed that COX-2-derived prostaglandins participate in uterine angiogenesis during implantation and decidualization (Matsumoto *et al.* 2002). In *Cox-2* KO mice, failure of implantation and decidualization are primarily due to defects in VEGF signaling. cPGI (an analog of PGI₂) together with 9cRA improves poor implantation in *Cox-2*($-/-$) mice. Moreover, the administration of cPGI and 9cRA restore the expression of VEGF and angiogenesis (Matsumoto & Sato 2006). In addition, Saito *et al.* (2007) examined RA effects on *in vitro* capillary-like tube formation using human umbilical vein endothelial cells and demonstrated that RA as well as RAR agonist Am80 significantly induced capillary-like tube formation. The RA-induced tube formation was inhibited by RAR antagonist. Meanwhile, the RA-induced tube formation was completely abolished by coinubation with VEGF antibody or with VEGF receptor (VEGFR)-2 antibody, but not VEGFR-1 antibody (Saito *et al.* 2007). These data provide evidence that although ovarian steroid hormones primarily influence uterine vascular permeability and angiogenesis during the preimplantation period, RA-RAR signaling participates by regulating VEGF production (Fig. 3).

Stromal decidualization

Stromal decidualization denotes the transformation of mesenchymal stromal cells into specialized decidual cells, which acquire epithelioid structure and function, accumulating glycogen, lipids and subcellular

organelles (Lawn *et al.* 1971) in part through a process of mesenchymal-epithelial transformation (Yu *et al.* 2016). Decidual cells provide the nutritive and immune-privileged matrix that is required for blastocyst implantation and placental formation in mouse and human uteri (Zhang *et al.* 2013, Gellersen & Brosens 2014). Perturbed or inadequate decidualization leads to embryo miscarriage and early pregnancy failure (Kommagani *et al.* 2013). Accompanied by typical morphological changes, differentiating stromal cells express biochemical decidual markers, such as prolactin (PRL), insulin-like growth factor-binding protein-1 (IGFBP1) (Irwin *et al.* 1994, Brar *et al.* 1997) and 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) (Brosens *et al.* 1999, Kuroda *et al.* 2013). Data from Ozaki *et al.* (2017) showed that RARs, as well as CRABP2 and FABP5, which are responsible for binding and delivering RA to nuclear receptors RARs and PPAR β/δ respectively, decreased in decidualizing human endometrial stromal cells despite an increase in PPAR β/δ . Besides, decidualization was also associated with increased expression of CYP25A1 (Table 3). In addition, exposure of differentiating stromal cells to 10nM RA or retinal prevents the inhibition of the RAR pathway and perturbs expression of PRL, IGFBP1 and 11 β HSD1 (Ozaki *et al.* 2017). Brar *et al.* (1996) consistently demonstrated that RA treatment suppressed PRL and IGFBP1 production and promoted maintenance of fibronectin, an indicator of the undifferentiated stromal phenotype (Yu *et al.* 2016) (Table 2). Though PPAR β/δ and RXRs are highly expressed at implantation sites in human and rat endometrium (Fukunaka *et al.* 2001, Ding *et al.* 2003) and endometrial stromal cells can synthesize RA from retinol during decidualization and blastocyst implantation (Ulven *et al.* 2000, Deng *et al.* 2003), no evidence exists that RA, even at low concentrations, promotes stromal decidualization, at least not when assessed by the induction of responsive

Table 3 Upregulated and downregulated genes involved in retinoid metabolism and signaling in decidualizing endometrial stromal cells.

Genes	Upregulation or downregulation	Species	References
<i>RBP4</i>	↑	Human	Ozaki <i>et al.</i> (2017), Pavone <i>et al.</i> (2017)
<i>CYP26A1</i>	↑	Human	Ozaki <i>et al.</i> (2017)
<i>CRABP2</i>	↓	Human	Ozaki <i>et al.</i> (2017), Pavone <i>et al.</i> (2017)
<i>FABP5</i>	↓	Human	Ozaki <i>et al.</i> (2017)
<i>RARα</i>	↓	Human	Ozaki <i>et al.</i> (2017)
<i>PPARβ/δ</i>	↑	Human	Ozaki <i>et al.</i> (2017)

↑, Upregulation; ↓, Downregulation; E₂, estradiol; P₄, progesterone.

decidual markers (PRL, IGFBP1 and HSD11B1). At high concentrations, RA decreases cell viability and exhibits toxic effects (Ozaki *et al.* 2017). These data suggested that RA signaling is attenuated upon decidualization, and RA treatment suppresses the decidualization and contributes to the maintenance of stromal cells in an undecidualized state. Furthermore, RA was suggested to prevent the increase in intracellular cAMP accumulation in stromal cells that occurs during the induction of decidualization *in vitro* with progesterone and estradiol (Brar *et al.* 1996). Meanwhile, the activation of the cAMP signal pathway is essential and sufficient to induce PRL production in endometrial stromal cells (Tang *et al.* 1993, Gellersen *et al.* 1994), and inhibition of PRL production by RA cannot be restored by cAMP treatment (Brar *et al.* 1996). The inhibition of intracellular cAMP production by RA is a plausible mechanism by which RA suppresses stromal decidualization. On the other hand, endogenous RA biosynthesis from retinol is increased 1.4-fold in endometrial stromal cells subjected to *in vitro* decidualization compared to control cells (Sidell *et al.* 2010). Thus, more precise dose-response relationships should be established to fully understand the possible differential effects of RA signaling via the opposing CRABP2-RAR vs FABP5-PPAR β pathways.

Retinoid signaling in blastocyst implantation

In humans and rodents, implantation occurs via successive blastocyst apposition, attachment, and adhesion to the receptive luminal epithelium, followed by the penetration and invasion of trophoblast into the decidualized stroma (Dey *et al.* 2004). Successful implantation requires not only a receptive endometrium but also a preimplantation blastocyst temporally competent to engage in precise crosstalk with the maternal endometrial signals. Recent evidence suggested that two separate uterine signals during blastocyst implantation. One signal primes the trophoblast for attachment to the luminal epithelium, and the other results in the uptake of amino acids by the blastocyst, the motility of which is initiated for invasion (Gonzalez 2012). Signaling of endometrial receptivity to blastocyst attachment involves the actions of ovarian steroid hormones on the luminal epithelia and paracrine factors expressed by different endometrial cell types (Cha *et al.* 2012, Zhang *et al.* 2013).

Rodent studies demonstrated that the tight regulation of retinoid pathways play crucial roles in maintaining uterine receptivity and blastocyst implantation (Osteen *et al.* 2003, Han *et al.* 2010). In mice, the uterus becomes

receptive to blastocyst implantation by the afternoon of gestational day 4 but is refractory by the afternoon of day 5 (Zhang *et al.* 2013). No *Cyp26a1* mRNA was found in preimplantation endometrium, while its expression was specifically induced between 3.5 and 4.5 gestational days, i.e., the implantation period, was localized in luminal epithelia (Vermot *et al.* 2000, Han *et al.* 2010). The specific spatiotemporal expression pattern of *Cyp26a1* in preimplantation endometrium suggests that it is involved in blastocyst attachment and/or invasion to the endometrium but not in the initial preparations of the endometrium. This hypothesis was supported by data in which the number of implantation sites was significantly reduced when *Cyp26a1*-specific antisense oligos or anti-CYP26A1 antibodies were injected into the uterus on day 3 of pregnancy (Han *et al.* 2010). As described previously, the RA pathway is inhibited during the late secretory phase by endogenous expression of *Cyp26a1*. RA can inhibit matrix metalloproteases (Bruner-Tran *et al.* 2002, Osteen *et al.* 2003), thus, CYP26A1-mediated degradation of RA may facilitate blastocyst implantation (Osteen *et al.* 2003).

Pharmacological concentrations of RA are embryotoxic at the early postimplantation stage of development (Huang *et al.* 2001). This observation, along with the well-characterized functions played by CXs in embryonic implantation and development, suggests a possible link between known teratogenic effects of RA (Ross *et al.* 2000) and the possibility that RA at such concentrations may reduce the effective trophoblast invasive capacity through increased GJIC because CX43 expression and GJIC are dramatically reduced during the implantation window (Granot *et al.* 2000) potentially facilitating trophoblast invasion into the endometrial stroma.

Other studies established the importance of endometrial gland secretions, such as leukemia inhibitory factor (LIF), for endometrial receptivity and blastocyst implantation. LIF binds to its receptor LIFR present in luminal epithelia and activates downstream signaling via STAT3 (Niwa *et al.* 1998). Consequently, LIF induces the expression of luminal epithelia-enriched genes, including *HB-EGF1*, *IGFBP3* and *IRG*, which are associated with uterine receptivity (Kimber 2005). *Lif* KO mice are infertile because of the failure of blastocyst attachment to the uterine luminal epithelium. Excess RA (10 μ M) can significantly inhibit the expression of the *Lif*, *Hb-egf* and *Csf1* in endometrial epithelial cells isolated on day 4.5 of pregnancy (Ma *et al.* 2012) (Table 2).

These data suggest that successful implantation during late secretory phase requires lower RA levels in the

luminal epithelia than what occurs in the proliferative phase, and excess RA might negatively affect stromal decidualization, endometrial receptivity and blastocyst implantation. CYP26A1-mediated degradation of RA is a hallmark of the implantation phase.

Collectively, the endometrial retinoid pathway is distinctly regulated by synthesis catalyzed by ALDHs and catabolism catalyzed by CYP26A1 at different phases and in different compartments of the endometrium, which are mainly controlled by estradiol and progesterone. The retinoid pathway plays crucial roles in endometrial epithelial differentiation and proliferation, neovascularization and GJIC and negatively regulates stromal decidualization and blastocyst implantation (Fig. 3).

Retinoid pathway and endometriosis

Endometriosis is a nonmalignant, but potentially metastatic, gynecological condition, which is estrogen dependent and defined by the presence of hormonally responsive, ectopic implants of endometrial mucosa dispersed in extrauterine locations (Giudice & Kao 2004, Kennedy *et al.* 2005). Approximately 10% of all women and up to 30%–50% of symptomatic premenopausal women are affected and commonly suffer pelvic pain and/or infertility (Nnoaham *et al.* 2011, Stilley *et al.* 2012). Classical and neoclassical concepts of endometriosis etiology were reviewed comprehensively elsewhere (Bulun 2009, Taylor 2010, Burney & Giudice 2012, Bulun *et al.* 2015, Taylor *et al.* 2015) and will not be reiterated thoroughly in this review. Although the theories of endometriosis histogenesis remain controversial, recent findings suggest that defective epigenetic landscape possibly associated with deficient differentiation of endometrial tissue stem cells is a central mechanism responsible for the cellular origins of endometriosis (Bulun *et al.* 2015). Meanwhile, retinoid pathway is fundamentally flawed in endometriotic tissues and even systemically in women with endometriosis (Pavone *et al.* 2010, 2011, 2017, Pierzchalski *et al.* 2014, Taylor *et al.* 2015).

Studies from the group of Bulun showed altered expression of several genes involved in retinol uptake, metabolism and action in cells from patients with endometriosis (Pavone *et al.* 2010, 2011, 2017). Endometriotic stromal cells overall had decreased *CRABP2* and *CRBP1* expression and increased *CYP26B1* expression (Table 4), the latter resulting in the elimination of RA from cells. These results are consistent with decreased retinoid

uptake, metabolism and action within endometriotic lesions (Pavone *et al.* 2011). *ALDH1A2* expression levels were significantly reduced in endometriotic tissue and stromal cells (Table 4), resulting in reduced RA levels. Pierzchalski *et al.* (2014) directly quantified RA levels and biosynthesis from retinol in endometrial stromal cells, which were derived from corresponding eutopic and ectopic biopsies; retinol uptake and RA production predominantly occur in these stromal cells (Vermot *et al.* 2000, Pavone *et al.* 2011, Yamagata *et al.* 2014) (Fig. 1). The studies confirmed impaired RA biosynthesis in endometriotic implants. A major defect was the reduced expression of *CRBP1*, a retinol carrier protein serving as the preferred substrate for retinol dehydrogenase enzymes and the rate-limiting factor in RA biosynthesis (Napoli 2012). Moreover, *STRA6*, which mediates retinol uptake, showed an abnormally low expression and high levels of DNA methylation in endometriotic stromal cells (Pavone *et al.* 2011, Yamagata *et al.* 2015) (Table 4). Thus, reduced *STRA6*, *CRBP1* and *ALDH1A2* expression levels result in a significantly less efficient conversion of retinol to RA in stromal cells. In addition, *RARα* expression levels are strikingly low in tissues and stromal cells of endometriosis (Pavone *et al.* 2011) (Table 4). Transcriptional activation via the RA-*CRABP2*-*RAR* pathway can trigger cell cycle arrest (Donato *et al.* 2007) and apoptosis (Altucci *et al.* 2001, Kitareewan *et al.* 2002, Donato *et al.* 2005) and frequently leads to the inhibition of cell proliferation. Thus, reduced RA-*CRABP2*-*RAR* signaling can cause endometriotic cells to escape apoptosis and contribute to the survival of ectopic cells (Nasu *et al.* 2009, Pavone *et al.* 2010). These studies suggested that flaws in RA production and degradation might play a role in the pathogenesis of endometriosis.

Table 4 Changes in genes involved in retinoid uptake, metabolism and signaling in endometriosis.

Genes	Changes	Cell type	References
<i>CRBP1</i>	↓	Stromal cells	Pavone <i>et al.</i> (2011, 2017)
<i>CRABP2</i>	↓	Stromal cells	Pavone <i>et al.</i> (2010, 2011, 2017)
<i>ALDH1A2</i>	↓	Stromal cells	Pavone <i>et al.</i> (2011)
<i>STRA6</i>	↓	Stromal cells	Pavone <i>et al.</i> (2010, 2011), Yamagata <i>et al.</i> (2015)
<i>RARα</i>	↓	Stromal cells	Pavone <i>et al.</i> (2011)
<i>RXRα</i>	↓	Stromal cells	Pavone <i>et al.</i> (2011)
<i>CYP26B1</i>	↑	Stromal cells	Pavone <i>et al.</i> (2011, 2017)

↓, reduced; ↑, increased.

Estrogen plays a critical role in the establishment and maintenance of endometriosis (Osteen *et al.* 2005, Bulun 2009, Bulun *et al.* 2015). 17 β -hydroxysteroid dehydrogenase type 2 (HSD17B2) catalyzes the conversion of estradiol to estrone, a much less biologically potent estrogen, and plays a crucial role in local estradiol inactivation in the endometrium (Cheng *et al.* 2007). HSD17B2 is present in normal endometrial glandular cells but is lacking in the endometrium of women with endometriosis (Stewart 1994, Giudice *et al.* 2002). Pavone *et al.* (2017) incubated epithelial cells with human serum RBP4 for 48 h and found that HSD17B2 expression was significantly increased in a dose-dependent manner. In serum, retinol is bound by RBP4 in the form of retinol-RBP4 complex (holo-RBP4),

which then binds to transthyretin (TTR) and forms a ternary complex (D'Ambrosio *et al.* 2011). Binding of holo-RBP4 to TTR prevents the loss of the RBP4 from the circulation by filtration via the renal glomeruli, as apo-RBP4 cannot be bound by TTR and is lost in the urine. Since the human serum RBP4 used in the study of Pavone *et al.* (2017) may contain significant amounts of holo-RBP4, the increased *HSD17B2* expression might be caused by retinol from holo-RBP4. This hypothesis was supported by another study which demonstrated that RA stimulates the assembly of a multimeric complex composed of RAR α /RXR α tethered to transcription factors SP1 and SP3 on the *HSD17B2* promoter where it induces HSD17B2. The RA antagonist ANG abolishes RA-induced *HSD17B2* expression in endometrial cells (Cheng *et al.* 2008, Yamagata *et al.* 2015). Thus, RA appears to be one of the critical paracrine factors that stimulate the production of HSD17B2 in neighboring epithelial cells and an important mechanism for local estradiol inactivation in the endometrium (Table 2). As previously mentioned, as the RA pathway is fundamentally flawed in endometriotic tissues (Pavone *et al.* 2010), this provides a plausible explanation for the aberrant *HSD17B2* expression and the high local estradiol concentrations in endometriosis (Fig. 4).

As previously mentioned, CX43 is predominantly expressed in endometrial stromal cells (Jahn *et al.* 1995, Winterhager *et al.* 2009) and necessary for GJIC that allow the stroma to serve as an organized tissue. Regidor *et al.* (1997) showed that CX43 levels were reduced in human endometriosis lesions and this finding was confirmed by Yu *et al.* (2014b). They reported nearly exclusive CX43 immunostaining in the stromal compartment of normal biopsies, with reduced immunostaining and redistribution to scattered epithelia in the eutopic endometria of women with endometriosis. Moreover, stromal cells isolated from cases of endometriosis exhibited a reduction of ~45% in GJIC, confirmed at the levels of CX43 mRNA and protein expression and also functionally by lower Lucifer Yellow diffusion (Yu *et al.* 2014b). As previously mentioned, RA can stimulate CX43 expression and GJIC in stromal cells (Tanmahasamut & Sidell 2005, Wu *et al.* 2013). Thus, flawed retinoid action might cause decreased CX43 expression and GJIC, reducing decidualization capacity of stromal cells in endometriosis, which could contribute to the development or progression of endometriosis lesions and to the subfertile uterine phenotype associated with the syndrome.

TGF- β 1, a member of transforming growth factor- β (TGF- β) superfamily, has been supposed to play an extensive role in the onset and development of endometriosis, such

Figure 4

Schematic representation of aberrant retinoid signaling in endometriosis. In stromal cells of endometriosis, reduced STRA6, CRBP1 and ALDH1A2 expression levels result in a significantly less efficient conversion of retinol to RA. Meanwhile, increased CYP26A1 expression promotes RA degradation. Thus, decreased RA levels along with reduced RAR α and CRABP2 expression result in a reduced RA-RAR signaling, which results in reduced CX43 expression and GJIC in stromal cells. Meanwhile, reduced RA levels result in lower HSD17B2 production and high local concentrations of estradiol (E₂) in epithelial cells of endometriosis. The reduced RA signaling and increased local E₂ activity could enhance cell proliferation, invasiveness and impede apoptosis in endometriosis. The full colour version of this figure is available at <https://doi.org/10.1530/JOE-17-0544>.

Figure 5

Potential therapeutic mechanism of RA for endometriosis. Activation of the RA-RAR pathway can promote HSD17B2 production, which can stimulate the catabolism of E_2 and inhibit TGF- β 1 production. These actions of the RA-RAR pathway imply its potential to suppress the development of endometriosis. A full colour version of this figure is available at <https://doi.org/10.1530/JOE-17-0544>.

as preventing apoptosis during transit to the peritoneal cavity, promoting adhesion to peritoneum and invasion of endometriosis cells (Omwandho *et al.* 2010). With the exception of one study (Hao *et al.* 2000), two other studies reported that subjects with endometriosis exhibit higher levels of TGF- β 1 in peritoneal fluid (Oosterlynck *et al.* 1994, Ku'pker *et al.* 1998). Interestingly, it was reported that RA could inhibit the TGF- β 1 signaling pathway in the endometrial cells and many other cell types (Frenz & Liu 2000, Delgadillo *et al.* 2014, Hu *et al.* 2016). Although further studies are needed, it can be speculated that RA might suppress the development of endometriosis implants partly through repressing the TGF- β 1 signaling pathway (Fig. 5).

These data suggest two possible scenarios to consider the theories of the histogenesis of endometriosis associated with the aberrant retinoid pathway in ectopically growing endometrial cells. On the one hand, defects in RA signaling can result in high local concentrations of estradiol in endometriosis lesions due to deficient oxidation and inactivation of estradiol. On the other hand, reduced RA signaling, along with increased estradiol effects, could enhance cell proliferation and invasiveness and impede apoptosis in endometriosis (Fig. 4). With respect to eutopic endometrial function in endometriosis, impaired RA signaling can be attributed to dysfunctional stromal differentiation, leading to the reduced capacity for decidualization reported in these cases (Klemmt *et al.* 2006, Aghajanova *et al.* 2009, Lessey *et al.* 2013).

In conclusion, the precisely controlled retinoid signaling may play a critical role in a number of critical endometrial physiological events. In addition, the altered RA pathway may be a leading cause for the histogenesis of endometriosis.

Perspectives for future studies

RBP4, acting as the primary systemic and intercellular transporter of retinol, plays a key role in cellular retinol influx, efflux and exchange (Kawaguchi *et al.* 2015).

Pavone *et al.* (2017) observed that RBP4 secretion increases during the decidualization of human endometrial stromal cells. Immunoreactivity for RBP4 is consistently higher near the embryo implantation site compared with non-implantation sites in the baboon endometrium (Fazleabas *et al.* 1994). These data suggest a potential role of RBP4 production and secretion in decidualization and/or implantation, but still require further investigation. Of great interest is that complete deletion of *Aldh* genes in mice results in prenatal lethality, precluding investigation of their roles in postnatal organogenesis. The use of tissue-selective and conditional gene targeting approaches is necessary to better understand the role of these enzymes during the ovarian cycle and gestation. As previously mentioned, excess RA concentration is harmful to decidualization and implantation. Thus, actual retinoid concentrations in the endometrium, as have been performed in some limited studies (Pierzchalski *et al.* 2014), should be measured in future studies.

Accumulating evidence supports that treatment modalities that target the retinoid pathway may have therapeutic utility (Sokalska *et al.* 2013). RA treatment reduced cytokine concentrations and the number and size of lesions (Wieser *et al.* 2012). Thus, RA has the potential to suppress the development of endometriotic implants and agents that target the RA-shuttling system may be useful therapeutic targets in the future, bypassing the toxicity and teratogenicity encountered with RA analogs (Fig. 5). We propose that adjuvant or alternative medical therapies should be developed on the basis of these concepts. Their pharmacological actions and anticipated low side-effect profiles are predicted to be able to provide women with endometriosis with more treatment options for the long-term management of chronic and debilitating gynecologic diseases.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was supported by the National Natural Science Foundation of China (31372308 to X Z, 31772596 to X Z, 31672417 to C L and 31301969 to L C) and by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (USA) as part of the Cooperative Research Partnerships to Promote Workforce Diversity in the Reproductive Sciences (U01 HD66439 to R N T).

References

- Aghajanova L, Hamilton A, Kwintkiewicz J, Vo KC & Giudice LC 2009 Steroidogenic enzyme and key decidualization marker dysregulation in endometrial stromal cells from women with versus without endometriosis. *Biology of Reproduction* **80** 105–114. (<https://doi.org/10.1095/biolreprod.108.070300>)
- Al Tanoury Z, Piskunov A & Rochette-Egly C 2013 Vitamin A and retinoid signaling: genomic and nongenomic effects. *Journal of Lipid Research* **54** 1761–1775. (<https://doi.org/10.1194/jlr.R030833>)
- Altucci L, Rossin A, Raffelsberger W, Reitmair A, Chomienne C & Gronemeyer H 2001 Retinoic acid-induced apoptosis in leukemia cells is mediated by paracrine action of tumor-selective death ligand TRAIL. *Nature Medicine* **7** 680–686. (<https://doi.org/10.1038/89050>)
- Benson GV, Lim H, Paria BC, Satokata I, Dey SK & Maas RL 1996 Mechanisms of reduced fertility in Hoxa-10 mutant mice: uterine homeosis and loss of maternal Hoxa-10 expression. *Development* **122** 2687–2696.
- Bo WJ & Smith MS 1966 The effect of retinol and retinoic acid on the morphology of the rat uterus. *Anatomical Record* **156** 5–9. (<https://doi.org/10.1002/ar.1091560103>)
- Boehm N, Chateau D & Rochette-Egly C 1997 Retinoid receptors in rat vaginal and uterine epithelia: changes with ovarian steroids. *Molecular and Cellular Endocrinology* **132** 101–108. ([https://doi.org/10.1016/S0303-7207\(97\)00126-3](https://doi.org/10.1016/S0303-7207(97)00126-3))
- Boettger-Tong HL & Stancel GM 1995 Retinoic acid inhibits estrogen-induced uterine stromal and myometrial cell proliferation. *Endocrinology* **136** 2975–2983. (<https://doi.org/10.1210/endo.136.7.7789323>)
- Brar AK, Kessler CA, Meyer AJ, Cedars MI & Jikihara H 1996 Retinoic acid suppresses in-vitro decidualization of human endometrial stromal cells. *Molecular Human Reproduction* **2** 185–193. (<https://doi.org/10.1093/molehr/2.3.185>)
- Brar AK, Frank GR, Kessler CA, Cedars MI & Handwerger S 1997 Progesterone-dependent decidualization of the human endometrium is mediated by cAMP. *Endocrine* **6** 301–307. (<https://doi.org/10.1007/BF02820507>)
- Brosens JJ, Hayashi N & White JO 1999 Progesterone receptor regulates decidual prolactin expression in differentiating human endometrial stromal cells. *Endocrinology* **140** 4809–4820. (<https://doi.org/10.1210/endo.140.10.7070>)
- Bruner-Tran KL, Eisenberg E, Yeaman GR, Anderson TA, McBean J & Osteen KG 2002 Steroid and cytokine regulation of matrix metalloproteinase expression in endometriosis and the establishment of experimental endometriosis in nude mice. *Journal of Clinical Endocrinology and Metabolism* **87** 4782–4791. (<https://doi.org/10.1210/jc.2002-020418>)
- Bryant SL, Francis JC, Lokody IB, Wang H, Risbridger GP, Loveland KL & Swain A 2014 Sex specific retinoic acid signaling is required for the initiation of urogenital sinus bud development. *Developmental Biology* **395** 209–217. (<https://doi.org/10.1016/j.ydbio.2014.09.016>)
- Bucco RA, Zheng WL, Davis JT, Sierra-Rivera E, Osteen KG, Chaudhary AK & Ong DE 1997 Cellular retinoic acid-binding protein(II) presence in rat uterine epithelial cells correlates with their synthesis of retinoic acid. *Biochemistry* **36** 4009–4014. (<https://doi.org/10.1021/bi962094o>)
- Bulun SE 2009 Endometriosis. *New England Journal of Medicine* **360** 268–279. (<https://doi.org/10.1056/NEJMra0804690>)
- Bulun SE, Monsivais D, Kakinuma T, Furukawa Y, Bernardi L, Pavone ME & Dyson M 2015 Molecular biology of endometriosis: from aromatase to genomic abnormalities. *Seminars in Reproductive Medicine* **33** 220–224. (<https://doi.org/10.1055/s-0035-1554053>)
- Burney RO & Giudice LC 2012 Pathogenesis and pathophysiology of endometriosis. *Fertility and Sterility* **98** S11–S19. (<https://doi.org/10.1016/j.fertnstert.2012.06.029>)
- Carter CA, Pogribny M, Davidson A, Jackson CD, McGarrity LJ & Morris SM 1996 Effects of retinoic acid on cell differentiation and reversion toward normal in human endometrial adenocarcinoma (RL95–2) cells. *Anticancer Research* **16** 17–24.
- Carystinos GD, Bier A & Batist G 2001 The role of connexin-mediated cell-cell communication in breast cancer metastasis. *Journal of Mammary Gland Biology and Neoplasia* **6** 431–440. (<https://doi.org/10.1023/A:1014787014851>)
- Cha J, Sun X & Dey SK 2012 Mechanisms of implantation: strategies for successful pregnancy. *Nature Medicine* **18** 1754–1767. (<https://doi.org/10.1038/nm.3012>)
- Chambon P 2005 The nuclear receptor superfamily: a personal retrospect on the first two decades. *Molecular Endocrinology* **19** 1418–1428. (<https://doi.org/10.1210/me.2005-0125>)
- Chan CCW, Lao TT, Ho PC, Sung EOP & Cheung ANY 2003 The effect of mifepristone on the expression of steroid hormone receptors in human decidua and placenta: a randomized placebo-controlled double-blind study. *Journal of Clinical Endocrinology and Metabolism* **88** 5846–5850. (<https://doi.org/10.1210/jc.2003-030958>)
- Cheng YH, Imir A, Fenkci V, Yilmaz MB & Bulun SE 2007 Stromal cells of endometriosis fail to produce paracrine factors that induce epithelial 17 β -hydroxysteroid dehydrogenase type 2 gene and its transcriptional regulator Sp1: a mechanism for defective estradiol metabolism. *American Journal of Obstetrics and Gynecology* **196** 391.e1–e7. (<https://doi.org/10.1016/j.ajog.2006.12.014>)
- Cheng YH, Yin P, Xue Q, Yilmaz B, Dawson MI & Bulun SE 2008 Retinoic acid (RA) regulates 17 β -hydroxysteroid dehydrogenase type 2 expression in endometrium: interaction of RA receptors with specificity protein (SP) 1/SP3 for estradiol metabolism. *Journal of Clinical Endocrinology and Metabolism* **93** 1915–1923. (<https://doi.org/10.1210/jc.2007-1536>)
- Chithalen JV, Luu L, Petkovich M & Jones G 2002 HPLC-MS/MS analysis of the products generated from all-trans-retinoic acid using recombinant human CYP26A. *Journal of Lipid Research* **43** 1133–1142. (<https://doi.org/10.1194/jlr.M100343-JLR200>)
- Comptour A, Rouzaire M, Belville C, Bouvier D, Gallot D, Blanchon L & Sapin V 2016 Nuclear retinoid receptors and pregnancy: placental transfer, functions, and pharmacological aspects. *Cellular and Molecular Life Sciences* **73** 3823–3837. (<https://doi.org/10.1007/s00018-016-2332-9>)
- Conaway HH, Henning P & Lerner UH 2013 Vitamin A metabolism, action, and role in skeletal homeostasis. *Endocrine Reviews* **34** 766–797. (<https://doi.org/10.1210/er.2012-1071>)
- Cope FO 1986 The in vivo inhibition of mouse brain protein kinase-C by retinoic acid. *Cancer Letters* **30** 275–288. ([https://doi.org/10.1016/0304-3835\(86\)90052-2](https://doi.org/10.1016/0304-3835(86)90052-2))
- D'Ambrosio DN, Clugston RD & Blaner WS 2011 Vitamin A metabolism: an update. *Nutrients* **3** 63–103. (<https://doi.org/10.3390/nu3010063>)
- Darwiche N, Celli G, Sly L, Lancillotti F & De Luca LM 1993 Retinoid status controls the appearance of reserve cells and keratin expression in mouse cervical epithelium. *Cancer Research* **53** 2287–2299.
- Delacroix L, Moutier E, Altobelli G, Legras S, Poch O, Choukrallah MA, Bertin I, Jost B & Davidson I 2010 Cell-specific interaction of retinoic acid receptors with target genes in mouse embryonic fibroblasts and embryonic stem cells. *Cell and Molecular Biology* **30** 231–244. (<https://doi.org/10.1128/MCB.00756-09>)

- Delgadillo D, Barbier O, Sierra G & Reyes JL 2014 Retinoic acid improves recovery after nephrectomy and decreases renal TGF- β 1 expression. Gender-related effects. *Fundamental and Clinical Pharmacology* **28** 170–179. (<https://doi.org/10.1111/fcp.12013>)
- Delva L, Bastie J, Rochette-Egly C, Kraiba R, Balitrand N, Despouy G, Chambon P & Chomienne C 1999 Physical and functional interactions between cellular retinoic acid binding protein II and the retinoic acid-dependent nuclear complex. *Molecular and Cellular Biology* **19** 7158–7167. (<https://doi.org/10.1128/MCB.19.10.7158>)
- Deng L, Shipley GL, Loose-Mitchell DS, Stancel GM, Broadus R, Pickar JH & Davies PJ 2003 Coordinate regulation of the production and signaling of retinoic acid by estrogen in the human endometrium. *Journal of Clinical Endocrinology and Metabolism* **88** 2157–2163. (<https://doi.org/10.1210/jc.2002-021844>)
- Dey SK, Lim H, Das SK, Reese J, Paria BC, Daikoku T & Wang H 2004 Molecular cues to implantation. *Endocrine Reviews* **25** 341–373. (<https://doi.org/10.1210/er.2003-0020>)
- Diao H, Xiao S, Howerth EW, Zhao F, Li R, Ard MB & Ye X 2013 Broad gap junction blocker carbenoxolone disrupts uterine preparation for embryo implantation in mice. *Biology of Reproduction* **89** 31. (<https://doi.org/10.1095/biolreprod.113.110106>)
- Ding NZ, Ma XH, Diao HL, Xu LB & Yang ZM 2003 Differential expression of peroxisome proliferator-activated receptor delta at implantation sites and in decidual cells of rat uterus. *Reproduction* **125** 817–825. (<https://doi.org/10.1530/rep.0.1250817>)
- Donato LJ & Noy N 2005 Suppression of mammary carcinoma growth by retinoic acid: proapoptotic genes are targets for retinoic acid receptor and cellular retinoic acid-binding protein II signaling. *Cancer Research* **65** 8193–8199. (<https://doi.org/10.1158/0008-5472.CAN-05-1177>)
- Donato LJ, Suh JH & Noy N 2007 Suppression of mammary carcinoma cell growth by retinoic acid: the cell cycle control gene Btg2 is a direct target for retinoic acid receptor signaling. *Cancer Research* **67** 609–615. (<https://doi.org/10.1158/0008-5472.CAN-06-0989>)
- Dong D, Ruuska S, Levinthal D & Noy N 1999. Distinct roles for cellular retinoic acid-binding proteins I and II in regulating signaling by retinoic acid. *Journal of Biological Chemistry* **274** 23695–23698. (<https://doi.org/10.1074/jbc.274.34.23695>)
- Duester G 2000 Families of retinoid dehydrogenases regulating vitamin A function: production of visual pigment and retinoic acid. *European Journal of Biochemistry* **267** 4315–4324. (<https://doi.org/10.1046/j.1432-1327.2000.01497.x>)
- Farjo KM, Moiseyev G, Nikolaeva O, Sandell LL, Trainor PA & Ma JX 2011 RDH10 is the primary enzyme responsible for the first step of embryonic Vitamin A metabolism and retinoic acid synthesis. *Developmental Biology* **357** 347–355. (<https://doi.org/10.1016/j.ydbio.2011.07.011>)
- Fazleabas AT, Donnelly KM, Mavrogianis PA & Verhage HG 1994 Retinol-binding protein in the baboon (*Papio anubis*) uterus: immunohistochemical characterization and gene expression. *Biology of Reproduction* **50** 1207–1215. (<https://doi.org/10.1095/biolreprod50.6.1207>)
- Ferrara N & Davis-Smyth T 1997 The biology of vascular endothelial growth factor. *Endocrine Reviews* **18** 4–25. (<https://doi.org/10.1210/edrv.18.1.0287>)
- Filant J & Spencer TE 2013 Endometrial glands are essential for blastocyst implantation and decidualization in the mouse uterus. *Biology of Reproduction* **88** 93. (<https://doi.org/10.1095/biolreprod.113.107631>)
- Frenz DA & Liu W 2000 Treatment with all-trans-retinoic acid decreases levels of endogenous TGF- β 1 in the mesenchyme of the developing mouse inner ear. *Teratology* **61** 297–304. ([https://doi.org/10.1002/\(SICI\)1096-9926\(200004\)61:4<297::AID-TERA9>3.0.CO;2-H](https://doi.org/10.1002/(SICI)1096-9926(200004)61:4<297::AID-TERA9>3.0.CO;2-H))
- Fritzsche B, Vermot J, Neumann U, Schmidt A, Schweigert FJ, Dollé P & Rühl R 2007 Regulation of expression of the retinoic acid metabolizing enzyme CYP26A1 in uteri of ovariectomized mice after treatment with ovarian steroid hormones. *Molecular Reproduction and Development* **74** 258–264. (<https://doi.org/10.1002/mrd.20526>)
- Fukunaka K, Saito T, Wataba K, Ashihara K, Ito E & Kudo R 2001 Changes in expression and subcellular localization of nuclear retinoic acid receptors in human endometrial epithelium during the menstrual cycle. *Molecular Human Reproduction* **7** 437–446. (<https://doi.org/10.1093/molehr/7.5.437>)
- Garry R, Hart R, Karthigasu KA & Burke C 2010 Structural changes in endometrial basal glands during menstruation. *BJOG* **117** 1175–1185. (<https://doi.org/10.1111/j.1471-0528.2010.02630.x>)
- Gellersen B & Brosens JJ 2014 Cyclic decidualization of the human endometrium in reproductive health and failure. *Endocrine Reviews* **35** 851–905. (<https://doi.org/10.1210/er.2014-1045>)
- Gellersen B, Kempf R, Telgmann R & DiMattia GE 1994 Nonpituitary human prolactin gene transcription is independent of Pit-1 and differentially controlled in lymphocytes and in endometrial stroma. *Molecular Endocrinology* **8** 356–373. (<https://doi.org/10.1210/me.8.3.356>)
- Giudice LC & Kao LC 2004 Endometriosis. *Lancet* **364** 1789–1799. ([https://doi.org/10.1016/S0140-6736\(04\)17403-5](https://doi.org/10.1016/S0140-6736(04)17403-5))
- Giudice LC, Telles TL, Lobo S & Kao L 2002 The molecular basis for implantation failure in endometriosis: on the road to discovery. *Annals of the New York Academy of Sciences* **955** 252–264. (<https://doi.org/10.1111/j.1749-6632.2002.tb02786.x>)
- Gonzalez IM, Martin PM, Burdsal C, Sloan JL, Mager S, Harris T & Sutherland AE 2012 Leucine and arginine regulate trophoblast motility through mTOR-dependent and independent pathways in the preimplantation mouse embryo. *Developmental Biology* **361** 286–300. (<https://doi.org/10.1016/j.ydbio.2011.10.021>)
- Granot I, Dekel N, Bechor E, Segal I, Fieldust S & Barash A 2000 Temporal analysis of connexin 43 protein and gene expression throughout the menstrual cycle in human endometrium. *Fertility and Sterility* **73** 381–386. ([https://doi.org/10.1016/S0015-0282\(99\)00531-2](https://doi.org/10.1016/S0015-0282(99)00531-2))
- Han BC, Xia HE, Sun J, Yang Y & Peng JP 2010 Retinoic acid-metabolizing enzyme cytochrome P450 26a1 (CYP26A1) is essential for implantation: functional study of its role in early pregnancy. *Journal of Cellular Physiology* **223** 471–479. (<https://doi.org/10.1002/jcp.22056>)
- Hao M, Shi Y & Dong M 2000 Measurements of interleukin-6, interleukin-8 and transforming growth factor-beta1 levels in peritoneal fluid of patients with endometriosis. *Zhonghua Fu Chan Ke Za Zhi* **35** 329–331.
- Harrison EH 2005 Mechanisms of digestion and absorption of dietary vitamin A. *Annual Review of Nutrition* **25** 87–103. (<https://doi.org/10.1146/annurev.nutr.25.050304.092614>)
- Hu J, Gray CA & Spencer TE 2004 Gene expression profiling of neonatal mouse uterine development. *Biology of Reproduction* **70** 1870–1876. (<https://doi.org/10.1095/biolreprod.103.026336>)
- Hu HY, Li HJ, He YL & Wang X 2016 ATRA alleviated endometrial fibrosis in a rabbit intrauterine adhesions model through downregulation of the TGF- β 1/smad4 signaling pathway. *International Journal of Clinical and Experimental Pathology* **9** 6171–6178.
- Huang F-J, Wu T-C & Tsai M-Y 2001 Effect of retinoic acid on implantation and postimplantation development of mouse embryos in vitro. *Human Reproduction* **16** 2171–2176. (<https://doi.org/10.1093/humrep/16.10.2171>)
- Huang CC, Orvis GD, Wang Y & Behringer RR 2012 Stromal-to-epithelial transition during postpartum endometrial regeneration. *PLoS ONE* **7** e44285. (<https://doi.org/10.1371/journal.pone.0044285>)
- Irwin JC, de las Fuentes L & Giudice LC 1994 Growth factors and decidualization in vitro. *Annals of the New York Academy of Sciences* **734** 7–18. (<https://doi.org/10.1111/j.1749-6632.1994.tb21730.x>)
- Jahn E, Classen-Linke I, Kusche M, Beier HM, Traub O, Grummer R & Winterhager E 1995 Expression of gap junction connexins in the human endometrium throughout the menstrual cycle. *Human Reproduction* **10** 2666–2670. (<https://doi.org/10.1093/oxfordjournals.humrep.a135764>)
- Jetten AM, De Luca LM, Nelson K, Schroeder W, Burlingame S & Fujimoto W 1996 Regulation of cornifin alpha expression in

- the vaginal and uterine epithelium by estrogen and retinoic acid. *Molecular and Cellular Endocrinology* **123** 7–15. ([https://doi.org/10.1016/0303-7207\(96\)03871-3](https://doi.org/10.1016/0303-7207(96)03871-3))
- Kambhampati S, Li Y, Verma A, Sassano A, Majchrzak B, Deb DK, Parmar S, Gafis N, Kalvakolanu DV, Rahman A et al. 2003 Activation of protein kinase C by all-transretinoic acid. *Journal of Biological Chemistry* **278** 32544–32551. (<https://doi.org/10.1074/jbc.M301523200>)
- Kastner P, Mark M, Ghyselinck N, Krezel W, Dupé V, Grondona JM & Chambon P 1997 Genetic evidence that the retinoid signal is transduced by heterodimeric RXR/RAR functional units during mouse development. *Development* **124** 313–326.
- Katzenellenbogen BS & Korach KS 1997 A new actor in the estrogen receptor drama – enter ER-beta. *Endocrinology* **138** 861–862. (<https://doi.org/10.1210/endo.138.3.5080>)
- Kawaguchi R, Yu J, Honda J, Hu J, Whitelegge H, Ping P, Wita P, Bok D & Sun H 2007 A membrane receptor for retinol binding protein mediates cellular uptake of vitamin A. *Science* **315** 820–825. (<https://doi.org/10.1126/science.1136244>)
- Kawaguchi R, Zhong M, Kassai M, Ter-Stepanian M & Sun H 2015. Vitamin A transport mechanism of the multitransmembrane cell-surface receptor STRA6. *Membranes* **5** 425–453. (<https://doi.org/10.3390/membranes5030425>)
- Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, Hummelshøj L, Prentice A, Saridogan E, ESIGf Endometriosis & EGD Group 2005 ESHRE guideline for the diagnosis and treatment of endometriosis. *Human Reproduction* **20** 2698–2704. (<https://doi.org/10.1093/humrep/dei135>)
- Kimber SJ 2005 Leukaemia inhibitory factor in implantation and uterine biology. *Reproduction* **130** 131–145. (<https://doi.org/10.1530/rep.1.00304>)
- Kitareewan S, Pitha-Rowe I, Sekula D, Lowrey CH, Nemeth MJ, Golub TR, Freemantle SJ & Dmitrovsky E 2002 UBE1L is a retinoid target that triggers PML/RARalpha degradation and apoptosis in acute promyelocytic leukemia. *PNAS* **99** 3806–3811. (<https://doi.org/10.1073/pnas.052011299>)
- Klemmt PA, Carver JG, Kennedy SH, Koninckx PR & Mardon HJ 2006 Stromal cells from endometriotic lesions and endometrium from women with endometriosis have reduced decidualization capacity. *Fertility and Sterility* **85** 564–572. (<https://doi.org/10.1016/j.fertnstert.2005.08.046>)
- Kommagani R, Szwarc MM, Kovanci E, Gibbons WE, Putluri N, Maity S, Creighton CJ, Sreekumar A, DeMayo FJ, Lydon JP, et al. 2013 Acceleration of the glycolytic flux by steroid receptor coactivator-2 is essential for endometrial decidualization. *PLoS Genetics* **9** e1003900. (<https://doi.org/10.1371/journal.pgen.1003900>)
- Kumarendran M, Loughney A, Prentice A, Thomas E & Redfern C 1996 Nuclear retinoid receptor expression in normal human endometrium throughout the menstrual cycle. *Molecular Human Reproduction* **2** 123–129. (<https://doi.org/10.1093/molehr/2.2.123>)
- Kupker W, Schultze-Mosgau A & Diedrich K 1998 Paracrine changes in the peritoneal environment of women with endometriosis. *Human Reproduction Update* **4** 719–723. (<https://doi.org/10.1093/humupd/4.5.719>)
- Kurita T, Lee KJ, Cooke PS, Taylor JA, Lubahn DB & Cunha GR 2000 Paracrine regulation of epithelial progesterone receptor by estradiol in the mouse female reproductive tract. *Biology of Reproduction* **62** 821–830. (<https://doi.org/10.1093/biolreprod/62.4.821>)
- Kuroda K, Venkatakrishnan R, Salker MS, Lucas ES, Shaheen F, Kuroda M, Blanks A, Christian M, Quenby S & Brosens JJ 2013 Induction of 11 beta-HSD 1 and activation of distinct mineralocorticoid receptor- and glucocorticoid receptor-dependent gene networks in decidualizing human endometrial stromal cells. *Molecular Endocrinology* **27** 192–202. (<https://doi.org/10.1210/me.2012-1247>)
- Lawn AM, Wilson EW & Finn CA 1971 The ultrastructure of human decidual and predecidual cells. *Journal of Reproduction and Infertility* **26** 85–90. (<https://doi.org/10.1530/jrf.0.0260085>)
- Laws MJ, Taylor RN, Sidell N, DeMayo FJ, Lydon JP, Gutstein DE, Bagchi MK & Bagchi IC 2008 Gap junction communication between uterine stromal cells plays a critical role in pregnancy-associated neovascularization and embryo survival. *Development* **135** 2659–2668. (<https://doi.org/10.1242/dev.019810>)
- Lessey BA, Lebovic DI & Taylor RN 2013 Eutopic endometrium in women with endometriosis: ‘ground zero’ for the study of implantation defects. *Seminars in Reproductive Medicine* **31** 109–124. (<https://doi.org/10.1055/s-0032-1333476>)
- Li XH & Ong DE 2003 Cellular retinoic acid-binding protein II gene expression is directly induced by estrogen, but not retinoic acid, in rat uterus. *Journal of Biological Chemistry* **278** 35819–35825. (<https://doi.org/10.1074/jbc.M302551200>)
- Li XH, Kakkad B & Ong DE 2004 Estrogen directly induces expression of retinoic acid biosynthetic enzymes, compartmentalized between the epithelium and underlying stromal cells in rat uterus. *Endocrinology* **145** 4756–4762. (<https://doi.org/10.1210/en.2004-0514>)
- Lin M, Zhang M, Abraham M, Smith SM & Napoli JL 2003 Mouse retinal dehydrogenase 4 (RALDH4), molecular cloning, cellular expression, and activity in 9-cis-retinoic acid biosynthesis in intact cells. *Journal of Biological Chemistry* **278** 9856–9861. (<https://doi.org/10.1074/jbc.M211417200>)
- Loudig O, Babichuk C, White J, Abu-Abed S, Mueller C & Petkovich M 2000 Cytochrome P450RAI(CYP26) promoter: a distinct composite retinoic acid response element underlies the complex regulation of retinoic acid metabolism. *Molecular Endocrinology* **14** 1483–1497. (<https://doi.org/10.1210/mend.14.9.0518>)
- Loudig O, Maclean GA, Dore NL, Luu L & Petkovich M 2005 Transcriptional co-operativity between distant retinoic acid response elements in regulation of Cyp26A1 inducibility. *Biochemical Journal* **392** 241–248. (<https://doi.org/10.1042/BJ20050874>)
- Loughney AD & Redfern CP 1995 Menstrual cycle related differences in the proliferative responses of cultured human endometrial stromal cells to retinoic acid. *Journal of Reproduction and Infertility* **105** 153–159. (<https://doi.org/10.1530/jrf.0.1050153>)
- Loughney A, Kumarendran M, Thomas E & Redfern C 1995 Variation in the expression of cellular retinoid binding proteins in human endometrium throughout the menstrual cycle. *Human Reproduction* **10** 1297–1304. (<https://doi.org/10.1093/oxfordjournals.humrep.a136137>)
- Ma L, Benson GV, Lim H, Dey SK & Maas RL 1998 Abdominal B (AbdB) Hoxa genes: regulation in adult uterus by estrogen and progesterone and repression in müllerian duct by the synthetic estrogen diethylstilbestrol (DES). *Developmental Biology* **197** 141–154. (<https://doi.org/10.1006/dbio.1998.8907>)
- Ma JJ, Han BC, Yang Y & Peng JP 2012 Retinoic acid synthesis and metabolism are concurrent in the mouse uterus during peri-implantation. *Cell and Tissue Research* **350** 525–537. (<https://doi.org/10.1007/s00441-012-1507-4>)
- Mantena SR, Kannan A, Cheon YP, Li Q, Johnson PF, Bagchi IC & Bagchi MK 2006 C/EBPbeta is a critical mediator of steroid hormone regulated cell proliferation and differentiation in the uterine epithelium and stroma. *PNAS* **103** 1870–1875. (<https://doi.org/10.1073/pnas.0507261103>)
- Matsumoto H & Sato E 2006 Uterine angiogenesis during implantation and decidualization in mice. *Reproductive Medicine and Biology* **5** 81–86. (<https://doi.org/10.1007/BF03016143>)
- Matsumoto H, Ma WG, Daikoku T, Zhao X, Paria BC, Das SK, Trzaskos JM & Dey SK 2002 Cyclooxygenase-2 differentially directs uterine angiogenesis during implantation in mice. *Journal of Biological Chemistry* **277** 29260–29267. (<https://doi.org/10.1074/jbc.M203996200>)

- Matsumoto H, Fukui E & Yoshizawa M 2007 Uterine angiogenesis during implantation in mice. *Journal of Mammalian Ova Research* **24** 45–49. (<https://doi.org/10.1274/jmor.24.45>)
- Mendelsohn C, Lohnes D, Décimo D, Lufkin T, LeMeur M, Chambon P & Mark M 1994 Function of the retinoic acid receptors (RARs) during development (II). Multiple abnormalities at various stages of organogenesis in RAR double mutants. *Development* **120** 2749–2771.
- Mowa CN & Iwanaga T 2000a Developmental changes of the oestrogen receptor- α and - β mRNAs in the female reproductive organ of the rat: an analysis by in situ hybridization. *Journal of Endocrinology* **167** 363–369. (<https://doi.org/10.1677/joe.0.1670363>)
- Mowa CN & Iwanaga T 2000b Differential distribution of oestrogen receptor- α and - β mRNAs in the female reproductive organ of rats as revealed by in situ hybridization. *Journal of Endocrinology* **165** 59–66. (<https://doi.org/10.1677/joe.0.1650059>)
- Nakagawa S, Fujii T, Yokoyama G, Kazanietz MG, Yamana H & Shirouzu K 2003 Cell growth inhibition by all-trans retinoic acid in SKBR-3 breast cancer cells: involvement of protein kinase C and extracellular signal-regulated kinase mitogen-activated protein kinase. *Molecular Carcinogenesis* **38** 106–116 (<https://doi.org/10.1002/mc.10150>)
- Nakajima T, Hayashi S, Iguchi T & Sato T 2011a The role of fibroblast growth factors on the differentiation of vaginal epithelium of neonatal mice. *Differentiation* **82** 28–37. (<https://doi.org/10.1016/j.diff.2011.03.005>)
- Nakajima T, Iguchi T & Sato T 2011b Involvement of activin signaling in abnormalities of mouse vagina exposed neonatally to diethylstilbestrol. *Cell and Tissue Research* **344** 527–538. (<https://doi.org/10.1007/s00441-011-1161-2>)
- Nakajima T, Iguchi T & Sato T 2016 Retinoic acid signaling determines the fate of uterine stroma in the mouse Müllerian duct. *PNAS* **113** 14354–14359. (<https://doi.org/10.1073/pnas.1608808113>)
- Napoli JL 1999 Interactions of retinoid binding proteins and enzymes in retinoid metabolism. *Biochimica et Biophysica Acta* **1440** 139–162. ([https://doi.org/10.1016/S1388-1981\(99\)00117-1](https://doi.org/10.1016/S1388-1981(99)00117-1))
- Napoli JL 2012 Physiological insights into all-trans-retinoic acid biosynthesis. *Biochimica et Biophysica Acta* **1821** 152–167 (<https://doi.org/10.1016/j.bbali.2011.05.004>)
- Napoli JL 2016 Functions of intracellular retinoid binding-proteins. *Sub-Cellular Biochemistry* **81** 21–76. (https://doi.org/10.1007/978-94-024-0945-1_2)
- Nasu K, Yuge A, Tsuno A, Nishida M & Narahara H 2009 Involvement of resistance to apoptosis in the pathogenesis of endometriosis. *Histology and Histopathology* **24** 1181–1192. (<https://doi.org/10.14670/HH-24.1181>)
- Niwa H, Burdon T, Chambers I & Smith A 1998 Self-renewal of pluripotent embryonic stem cells is mediated via activation of STAT3. *Genes and Development* **12** 2048–2060. (<https://doi.org/10.1101/gad.12.13.2048>)
- Nnoaham KE, Hummelshoj L, Webster P, d'Hooghe T, de Cicco Nardone F, de Cicco Nardone C, Jenkinson C, Kennedy SH, Zondervan KT & World Endometriosis Research Foundation Global Study of Women's Health Consortium 2011 Impact of endometriosis on quality of life and work productivity: a multicenter study across ten countries. *Fertility and Sterility* **96** 366.e8–373.e8. (<https://doi.org/10.1016/j.fertnstert.2011.05.090>)
- Noy N 2000 Retinoid-binding proteins: mediators of retinoid action. *Biochemical Journal* **348** 481–495. (<https://doi.org/10.1042/bj3480481>)
- Nozaki Y, Yamagata T, Sugiyama M, Ikoma S, Kinoshita K & Funauchi M 2006 Anti-inflammatory effect of all-trans-retinoic acid in inflammatory arthritis. *Journal of Clinical Immunology* **119** 272–279. (<https://doi.org/10.1016/j.clim.2005.11.012>)
- Oh SY, Grupen CG & Murray AW 1991 Phorbol ester induces phosphorylation and down-regulation of connexin 43 in WB cells. *Biochimica et Biophysica Acta* **1094** 243–245. ([https://doi.org/10.1016/0167-4889\(91\)90016-Q](https://doi.org/10.1016/0167-4889(91)90016-Q))
- Omwandho CO, Konrad L, Halis G, Oehmke F & Tinneberg HR. 2010 Role of TGF- β s in normal human endometrium and endometriosis. *Human Reproduction* **25** 101–109. (<https://doi.org/10.1093/humrep/dep382>)
- Oosterlynck DJ, Meuleman C, Waer M & Koninckx PR 1994 Transforming growth factor- β activity is increased in peritoneal fluid from women with endometriosis. *Obstetrics and Gynecology* **83** 287–292.
- Osteen KG, Igarashi TM & Bruner-Tran KL 2003 Progesterone action in the human endometrium: induction of a unique tissue environment which limits matrix metalloproteinase (MMP) expression. *Frontiers in Bioscience* **8** 78–86. (<https://doi.org/10.2741/938>)
- Osteen KG, Bruner-Tran KL & Eisenberg E 2005 Endometrial biology and the etiology of endometriosis. *Fertility and Sterility* **84** 33–34. (<https://doi.org/10.1016/j.fertnstert.2005.01.124>)
- Ozaki R, Kuroda K, Ikemoto Y, Ochiai A, Matsumoto A, Kumakiri J, Kitade M, Itakura A, Muter J, Brosens JJ, et al. 2017 Reprogramming of the retinoic acid pathway in decidualizing human endometrial stromal cells. *PLoS ONE* **12** e0173035. (<https://doi.org/10.1371/journal.pone.0173035>)
- Pavone ME, Reierstad S, Sun H, Milad M, Bulun SE & Cheng YH 2010 Altered retinoid uptake and action contributes to cell survival in endometriosis. *Journal of Clinical Endocrinology and Metabolism* **95** E300–E309. (<https://doi.org/10.1210/jc.2010-0459>)
- Pavone ME, Dyson M, Reierstad S, Pearson E, Ishikawa H, Cheng YH & Bulun SE 2011 Endometriosis expresses a molecular pattern consistent with decreased retinoid uptake, metabolism and action. *Human Reproduction* **26** 2157–2164. (<https://doi.org/10.1093/humrep/der172>)
- Pavone ME, Malpani S, Dyson M & Bulun SE 2017 Altered retinoid signaling compromises decidualization in human endometriotic stromal cells. *Reproduction* **154** 107–116. (<https://doi.org/10.1530/REP-16-0592>)
- Pierchalski K, Taylor RN, Nezhat C, Jones JW, Napoli JL, Yang G, Kane MA & Sidell N 2014 Retinoic acid biosynthesis is impaired in human and murine endometriosis. *Biology of Reproduction* **91** 84. (<https://doi.org/10.1095/biolreprod.114.119677>)
- Ponnamperuma RM, Kirchhof SM, Trifiletti L & DeLuca LM 1999 Ovariectomy increases squamous metaplasia of the uterine horns and survival of SENCAR mice fed a vitamin A-deficient diet. *American Journal of Clinical Nutrition* **70** 502–508.
- Prentice A, Matthews CJ, Thomas EJ & Redfern CP 1992 The expression of retinoic acid receptors in cultured human endometrial stromal cells and effects of retinoic acid. *Human Reproduction* **7** 692–700. (<https://doi.org/10.1093/oxfordjournals.humrep.a137721>)
- Punyadeera C, Verbost P & Groothuis P 2003 Oestrogen and progesterone responses in human endometrium. *Journal of Steroid Biochemistry and Molecular* **84** 393–410. ([https://doi.org/10.1016/S0960-0760\(03\)00061-X](https://doi.org/10.1016/S0960-0760(03)00061-X))
- Quadro L, Blaner WS, Hamberger L, Novikoff PM, Vogel S, Piantadosi R, Gottesman ME & Colantuoni V 2004 The role of extrahepatic retinoid binding protein in the mobilization of retinoid stores. *Journal of Lipid Research* **45** 1975–1982. (<https://doi.org/10.1194/jlr.M400137-JLR200>)
- Regidor P-A, Regidor M, Schindler AE & Winterhager E 1997 Aberrant expression pattern of gap junction connexins in endometriotic tissues. *Molecular Human Reproduction* **3** 375–381. (<https://doi.org/10.1093/molehr/3.5.375>)
- Rochette-Egly C 2014 Retinoic acid signaling and mouse embryonic stem cell differentiation: cross talk between genomic and non-genomic effects of RA. *Biochimica et Biophysica Acta* **1851** 66–75. (<https://doi.org/10.1016/j.bbali.2014.04.003>)
- Ross SA, McCaffery PJ, Drager UC & De Luca LM 2000 Retinoids in embryonal development. *Physiological Reviews* **80** 1021–1054. (<https://doi.org/10.1152/physrev.2000.80.3.1021>)
- Ruberte E, Friederich V, Morriss-Kay G & Chambon P 1992. Differential distribution patterns of CRABPI and CRABPII transcripts during mouse embryogenesis. *Development* **115** 973–987.

- Rühl R, Fritzsche B, Vermot J, Niederreither K, Neumann U, Schmidt A, Schweigert FJ & Dollé P 2006 Regulation of expression of the retinoic acid-synthesising enzymes retinaldehyde dehydrogenases in the uteri of ovariectomised mice after treatment with oestrogen, gestagen and their combination. *Reproduction, Fertility and Development* **18** 339–345. (<https://doi.org/10.1071/rd05056>)
- Saito A, Sugawara A, Urano A, Kudo M, Kagechika H, Sato Y, Owada Y, Kondo H, Sato M, Kurabayashi M, et al. 2007 All-trans retinoic acid induces in vitro angiogenesis via retinoic acid receptor: possible involvement of paracrine effects of endogenous vascular endothelial growth factor signaling. *Endocrinology* **148** 1412–1423. (<https://doi.org/10.1210/en.2006-0900>)
- Sawatsri S, Desai N, Rock JA & Sidell N 2000 Retinoic acid suppresses interleukin-6 production in human endometrial cells. *Fertility and Sterility* **73** 1012–1019. ([https://doi.org/10.1016/S0015-0282\(00\)00483-0](https://doi.org/10.1016/S0015-0282(00)00483-0))
- Schug TT, Berry DC, Shaw NS, Travis SN & Noy N 2007 Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors. *Cell* **129** 723–733. (<https://doi.org/10.1016/j.cell.2007.02.050>)
- Schug TT, Berry DC, Toshkov IA, Cheng L, Nikitin AY & Noy N 2008 Overcoming retinoic acid-resistance of mammary carcinomas by diverting retinoic acid from PPAR α to RAR. *PNAS* **105** 7546–7551. (<https://doi.org/10.1073/pnas.0709981105>)
- Sharpe-Timms KL 2001 Endometrial anomalies in women with endometriosis. *Annals of the New York Academy of Sciences* **943** 131–147. (<https://doi.org/10.1111/j.1749-6632.2001.tb03797.x>)
- Sidell N, Han SW & Parthasarathy S 2002 Regulation and modulation of abnormal immune responses in endometriosis. *Annals of the New York Academy of Sciences* **955** 159–173. (<https://doi.org/10.1111/j.1749-6632.2002.tb02777.x>)
- Sidell N, Feng Y, Hao L, Wu J, Yu J, Kane MA, Napoli JL & Taylor RN 2010 Retinoic acid is a cofactor for translational regulation of vascular endothelial growth factor in human endometrial stromal cells. *Molecular Endocrinology* **24** 148–160. (<https://doi.org/10.1210/me.2009-0155>)
- Sokalska A, Anderson M, Villanueva J, Ortega I, Bruner-Tran KL, Osteen KG & Duleba AJ 2013 Effects of simvastatin on retinoic acid system in primary human endometrial stromal cells and in a chimeric model of human endometriosis. *Journal of Clinical Endocrinology and Metabolism* **98** E463–E471. (<https://doi.org/10.1210/jc.2012-3402>)
- Solan JL & Lampe PD 2009 Connexin43 phosphorylation: structural changes and biological effects. *Biochemical Journal* **419** 261–272. (<https://doi.org/10.1042/BJ20082319>)
- Spencer TE, Dunlap KA & Filant J 2012 Comparative developmental biology of the uterus: insights into mechanisms and developmental disruption. *Molecular and Cellular Endocrinology* **354** 34–53. (<https://doi.org/10.1016/j.mce.2011.09.035>)
- Stahl W & Sies H 1998 The role of carotenoids and retinoids in gap junctional communication. *International Journal for Vitamin and Nutrition Research* **68** 354–359.
- Stewart CL 1994 Leukaemia inhibitory factor and the regulation of pre-implantation development of the mammalian embryo. *Molecular Reproduction and Development* **39** 233–238. (<https://doi.org/10.1002/mrd.1080390217>)
- Stilley JA, Birt JA & Sharpe-Timms KL 2012 Cellular and molecular basis for endometriosis-445 associated infertility. *Cell and Tissue Research* **349** 849–862. (<https://doi.org/10.1007/s00441-011-1309-0>)
- Suzuki A, Watanabe H, Mizutani T, Sato T, Ohta Y & Iguchi T 2006 Global gene expression in mouse vaginae exposed to diethylstilbestrol at different ages. *Experimental Biology and Medicine* **231** 632–640. (<https://doi.org/10.1177/153537020623100518>)
- Suzuki A, Urushitani H, Sato T, Kobayashi T, Watanabe H, Ohta Y & Iguchi T 2007 Gene expression change in the Müllerian duct of the mouse fetus exposed to diethylstilbestrol in utero. *Experimental Biology and Medicine* **232** 503–514.
- Tang B, Culler S & Gurpide E 1993 Cyclic adenosine 3', 5'-monophosphate induces prolactin expression in stromal cells isolated from human proliferative endometrium. *Endocrinology* **133** 2197–2203. (<https://doi.org/10.1210/endo.133.5.8404671>)
- Tanmahasamut P & Sidell N 2005 Up-regulation of gap junctional intercellular communication and connexin43 expression by retinoic acid in human endometrial stromal cells. *Journal of Clinical Endocrinology and Metabolism* **90** 4151–4156. (<https://doi.org/10.1210/jc.2004-0663>)
- Taylor RN 2010 Classical and neo-classical concepts in the etiology of endometriosis. In *Proceedings of the IFFS 20th World Congress on Fertility and Sterility*, pp 70–73. Eds B Hedon, L Mettler & H-R Tinneberg. Munich, Germany: Lukon.
- Taylor RN, Kane MA & Sidell N 2015 Pathogenesis of endometriosis: roles of retinoids and inflammatory pathways. *Seminars in Reproductive Medicine* **33** 246–256. (<https://doi.org/10.1055/s-0035-1554920>)
- Toma S, Isnardi L, Raffo P, Dastoli G, De Francisci E, Riccardi L, Palumbo R & Bollag W 1997 Effects of ALL-trans-retinoic acid and 13-cis-retinoic acid on breast-cancer cell lines: growth inhibition and apoptosis induction. *International Journal of Cancer* **70** 619–627. ([https://doi.org/10.1002/\(SICI\)1097-0215\(19970304\)70:5<619::AID-IJC21>3.0.CO;2-6](https://doi.org/10.1002/(SICI)1097-0215(19970304)70:5<619::AID-IJC21>3.0.CO;2-6))
- Ulven SM, Gundersen TE, Weedon MS, Landaas VO, Sakhi AK, Fromm SH, Geronimo BA, Moskaug JO & Blomhoff R 2000 Identification of endogenous retinoids, enzymes, binding proteins, and receptors during early postimplantation development in mouse: important role of retinal dehydrogenase type 2 in synthesis of all-trans-retinoic acid. *Developmental Biology* **220** 379–391. (<https://doi.org/10.1006/dbio.2000.9634>)
- Vallet JL, Christenson RK, Bartol FF & Wiley AA 1995 Effect of treatment with retinyl palmitate, progesterone, oestradiol and tamoxifen on secretion of a protein similar to retinol-binding protein during uterine gland development in neonatal pigs. *Journal of Reproduction and Infertility* **103** 189–197. (<https://doi.org/10.1530/jrf.0.1030189>)
- Verma AK 1988 Inhibition of tumor promoter 12-O-tetradecanoylphorbol-13-acetate-induced synthesis of epidermal ornithine decarboxylase messenger RNA and diacylglycerol-promoted mouse skin tumor formation by retinoic acid. *Cancer Research* **48** 2168–2173.
- Vermot J, Fraulob V, Dollé P & Niederreither K 2000 Expression of enzymes synthesizing (aldehyde dehydrogenase 1 and retinaldehyde dehydrogenase 2) and metabolizing (Cyp26) retinoic acid in the mouse female reproductive system. *Endocrinology* **141** 3638–3645. (<https://doi.org/10.1210/endo.141.10.7696>)
- Wang H & Dey SK 2006 Roadmap to embryo implantation: clues from mouse models. *Nature Reviews Genetics* **7** 185–199. (<https://doi.org/10.1038/nrg1808>)
- Wang X, Sperkova Z & Napoli JL 2001 Analysis of mouse retinal dehydrogenase type 2 promoter and expression. *Genomics* **74** 245–250. (<https://doi.org/10.1006/geno.2001.6546>)
- Wardlaw SA, Bucco RA, Zheng WL & Ong DE 1997 Variable expression of cellular retinol- and cellular retinoic acid-binding proteins in the rat uterus and ovary during the estrous cycle. *Biology of Reproduction* **56** 125–132. (<https://doi.org/10.1095/biolreprod56.1.125>)
- Warn-Cramer BJ, Lampe PD, Durata WE, Kanemitsu MY, Loo LWM, Eckhart W & Lau AF 1996 Characterization of the mitogen-activated protein kinase phosphorylation sites on the connexin-43 gap junction protein. *Journal of Biological Chemistry* **271** 3779–3786. (<https://doi.org/10.1074/jbc.271.7.3779>)
- Wieser F, Wu J, Shen Z, Taylor RN & Sidell N 2012 Retinoic acid suppresses growth of lesions, inhibits peritoneal cytokine secretion, and promotes macrophage differentiation in an immunocompetent mouse model of endometriosis. *Fertility and Sterility* **97** 1430–1437. (<https://doi.org/10.1016/j.fertnstert.2012.03.004>)
- Wilson JG & Warkany J 1948 Malformations in the genito-urinary tract induced by maternal vitamin A deficiency in the rat. *American*

- Journal of Clinical Nutrition* **83** 357–407. (<https://doi.org/10.1002/aja.1000830303>)
- Winterhager E & Kidder GM 2015 Gap junction connexins in female reproductive organs: implications for women's reproductive health. *Human Reproduction Update* **21** 340–352. (<https://doi.org/10.1093/humupd/dmv007>)
- Winterhager E, Grümmer R, Mavrogianis PA, Jones CJP, Hastings JM & Fazleabas AT 2009 Connexin expression pattern in the endometrium of baboons is influenced by hormonal changes and the presence of endometriotic lesions. *Molecular Human Reproduction* **15** 645–652. (<https://doi.org/10.1093/molehr/gap060>)
- Wu L & Ross AC 2010 Acidic retinoids synergize with vitamin A to enhance retinol uptake and STRA6, LRAT, and CYP26B1 expression in neonatal lung. *Journal of Lipid Research* **51** 378–387. (<https://doi.org/10.1194/jlr.M001222>)
- Wu J, Taylor RN & Sidell N 2013 Retinoic acid regulates gap junction intercellular communication in human endometrial stromal cells through modulation of the phosphorylation status of connexin 43. *Journal of Cellular Physiology* **228** 903–910. (<https://doi.org/10.1002/jcp.24241>)
- Xia HF, Ma JJ, Sun J, Yang Y & Peng JP 2010 Retinoic acid metabolizing enzyme CYP26A1 is implicated in rat embryo implantation. *Human Reproduction* **25** 2985–2998. (<https://doi.org/10.1093/humrep/deq268>)
- Yamagata Y, Nishino K, Takaki E, Sato S, Maekawa R, Nakai A & Sugino N 2014 Genome-wide DNA methylation profiling in cultured eutopic and ectopic endometrial stromal cells. *PLoS ONE* **9** e83612. (<https://doi.org/10.1371/journal.pone.0083612>)
- Yamagata Y, Takaki E, Shinagawa M, Okada M, Jozaki K, Lee L, Sato S, Maekawa R, Taketani T, Asada H *et al.* 2015 Retinoic acid has the potential to suppress endometriosis development. *Journal of Ovarian Research* **8** 49. (<https://doi.org/10.1186/s13048-015-0179-6>)
- Yamanouchi H, Umezue T & Tomooka Y 2010 Reconstruction of oviduct and demonstration of epithelial fate determination in mice. *Biology of Reproduction* **82** 528–533. (<https://doi.org/10.1095/biolreprod.109.078329>)
- Yamasaki H, Krutovskikh V, Mesnil M, Tanaka T, Zaidan-Dagli ML & Omori Y 1999 Role of connexin (gap junction) genes in cell growth control and carcinogenesis. *Comptes Rendus de l'Académie des Sciences – Series III – Sciences de la Vie* **322** 151–159. ([https://doi.org/10.1016/s0764-4469\(99\)80038-9](https://doi.org/10.1016/s0764-4469(99)80038-9))
- Yu J, Wu J, Bagchi IC, Bagchi MK, Sidell N & Taylor RN 2011 Disruption of gap junctions reduces biomarkers of decidualization and angiogenesis and increases inflammatory mediators in human endometrial stromal cell cultures. *Molecular and Cellular Endocrinology* **344** 25–34. (<https://doi.org/10.1016/j.mce.2011.04.011>)
- Yu J, Berga SL, Zou W, Sun HY, Johnston-MacAnanny E, Yalcinkaya T, Sidell N, Bagchi IC, Bagchi MK & Taylor RN 2014a Gap junction blockade induces apoptosis in human endometrial stromal cells. *Molecular Reproduction and Development* **81** 666–675. (<https://doi.org/10.1002/mrd.22334>)
- Yu J, Boicea A, Barrett KL, James CO, Bagchi IC, Bagchi MK, Nezhat C, Sidell N & Taylor RN 2014b Reduced connexin 43 in eutopic endometrium and cultured endometrial stromal cells from subjects with endometriosis. *Molecular Human Reproduction* **20** 260–270. (<https://doi.org/10.1093/molehr/gat087>)
- Yu J, Berga SL, Johnston-MacAnanny EB, Sidell N, Bagchi IC, Bagchi MK & Taylor RN 2016 Endometrial stromal decidualization responds reversibly to hormone stimulation and withdrawal. *Endocrinology* **157** 2432–2446. (<https://doi.org/10.1210/en.2015-1942>)
- Zhang S, Lin H, Kong S, Wang S, Wang H, Wang H & Armant DR 2013 Physiological and molecular determinants of embryo implantation. *Molecular Aspects of Medicine* **34** 939–980. (<https://doi.org/10.1016/j.mam.2012.12.011>)
- Zheng WL & Ong DE 1998 Spatial and temporal patterns of expression of cellular retinol-binding protein and cellular retinoic acid-binding proteins in rat uterus during early pregnancy. *Biology of Reproduction* **58** 963–970. (<https://doi.org/10.1095/biolreprod58.4.963>)
- Zheng WL, Sierra-Rivera E, Luan J, Osteen KG & Ong DE 2000 Retinoic acid synthesis and expression of cellular retinol-binding protein and cellular retinoic acid-binding protein type II are concurrent with decidualization of rat uterine stromal cells. *Endocrinology* **141** 802–808. (<https://doi.org/10.1210/endo.141.2.7323>)

Received in final form 19 December 2017

Accepted 3 January 2018

Accepted Preprint published online 3 January 2018