REVIEW

Germline and mosaic mutations causing pituitary tumours: genetic and molecular aspects

Sara Pepe^{1,2}, Márta Korbonits¹ and Donato Iacovazzo¹¹Centre for Endocrinology, William Harvey Research Institute, Barts and the London School of Medicine, Queen Mary University of London, London, UK²Department of Medical Biotechnologies, University of Siena, Siena, ItalyCorrespondence should be addressed to M Korbonits: m.korbonits@qmul.ac.uk

Abstract

While 95% of pituitary adenomas arise sporadically without a known inheritable predisposing mutation, in about 5% of the cases they can arise in a familial setting, either isolated (familial isolated pituitary adenoma or FIPA) or as part of a syndrome. FIPA is caused, in 15–30% of all kindreds, by inactivating mutations in the *AIP* gene, encoding a co-chaperone with a vast array of interacting partners and causing most commonly growth hormone excess. While the mechanisms linking AIP with pituitary tumorigenesis have not been fully understood, they are likely to involve several pathways, including the cAMP-dependent protein kinase A pathway via defective G inhibitory protein signalling or altered interaction with phosphodiesterases. The cAMP pathway is also affected by other conditions predisposing to pituitary tumours, including X-linked acro gigantism caused by duplications of the *GPR101* gene, encoding an orphan G stimulatory protein-coupled receptor. Activating mosaic mutations in the *GNAS* gene, coding for the G α stimulatory protein, cause McCune–Albright syndrome, while inactivating mutations in the regulatory type 1 α subunit of protein kinase A represent the most frequent genetic cause of Carney complex, a syndromic condition with multi-organ manifestations also involving the pituitary gland. In this review, we discuss the genetic and molecular aspects of isolated and syndromic familial pituitary adenomas due to germline or mosaic mutations, including those secondary to *AIP* and *GPR101* mutations, multiple endocrine neoplasia type 1 and 4, Carney complex, McCune–Albright syndrome, DICER1 syndrome and mutations in the *SDHx* genes underlying the association of familial paragangliomas and pheochromocytomas with pituitary adenomas.

Key Words

- genetics
- pituitary
- pituitary adenoma
- mutation

Journal of Endocrinology
(2019) **240**, R21–R45

Introduction

The human pituitary gland consists of an anterior lobe, which derives from the oral ectoderm, and a posterior lobe, which originates from the neuroectoderm. The anterior pituitary contains five types of endocrine cells, including the somatotroph (producing growth

hormone (GH)), lactotroph (producing prolactin (PRL)), gonadotroph (producing the gonadotropins, LH and FSH), corticotroph (producing adrenocorticotrophin (ACTH)) and the thyrotroph (producing thyrotropin (TSH)) cells. The anterior pituitary also contains a non-endocrine cell

population represented by the folliculostellate cells, which have a sustentacular function to the hormone-producing cells (Devnath & Inoue 2008).

Pituitary adenomas (PAs) are usually benign tumours arising from the endocrine cells of the anterior pituitary. These tumours are quite common and they are found in approximately 15–20% of the general population in radiological or autopsy studies (Ezzat *et al.* 2004, Daly *et al.* 2009), and they represent the third most common intracranial neoplasm after meningiomas and gliomas (Aflori & Korbonits 2014). However, most of these tumours have no clinical relevance and often represent incidental findings (Freda *et al.* 2011). Clinically relevant pituitary tumours are rarer, occurring in about 0.1% of the general population (Daly *et al.* 2006a, Fontana & Gaillard 2009, Cannavo *et al.* 2010, Fernandez *et al.* 2010, Raappana *et al.* 2010, Gruppette *et al.* 2013, Agustsson *et al.* 2015). Although histologically benign, PAs can cause significant morbidity due to hormone excess, hypopituitarism and tumour mass effects on the surrounding structures, such as the optic pathway, the cavernous sinuses and the brain. The most common PAs are represented by prolactinomas (45–65%), followed by non-functioning PAs (NFPAs) (15–37%), somatotroph (9–15%), corticotroph (2–6%) and thyrotroph PAs (0–1%).

Pituitary tumours are believed to be monoclonal in origin (Herman *et al.* 1990). The exact molecular pathogenesis is still not clear; however, several mechanisms have been described, including, among others, dysregulation of cell cycle regulators (Jacks *et al.* 1992, Kiyokawa *et al.* 1996) or alterations of growth factors (Zhou *et al.* 2014). Somatic mutations can also occur, including activating *GNAS* mutations (found in 10–50% of somatotroph PAs (Peverelli *et al.* 2014)) or somatic mutations in the *USP8* gene causing activation of the EGF signalling pathway (found in 20–60% of corticotroph PAs) (Ma *et al.* 2015, Reincke *et al.* 2015, Ballmann *et al.* 2018).

While most PAs arise sporadically, about 5% occur in a familial setting (Daly *et al.* 2006b). Familial PAs are often distinct from their sporadic counterpart, as they can present an aggressive behaviour, are frequently resistant to treatment and they often arise at an earlier age. Familial PAs can develop as part of a syndromic condition, such as multiple endocrine neoplasia type 1 (MEN1), multiple endocrine neoplasia type 4 (MEN4), Carney complex (CNC), McCune–Albright syndrome (MAS), pheochromocytoma/paraganglioma with PA (3PAs) and DICER1 syndrome. However, as seen in the case of familial isolated PA (FIPA), PAs can develop in the absence of other clinical manifestations, as is the case for

patients harbouring mutations in the aryl hydrocarbon receptor-interacting protein (*AIP*) gene and in patients with X-linked acrogigantism (XLAG), a rare condition of early-onset pituitary gigantism due to duplications involving the *GPR101* gene. Nevertheless, in the majority of cases of FIPA, the causative genetic mutations still remain to be identified.

Owing to either incomplete penetrance or to *de novo* mutations, variants in genes associated with FIPA or syndromic conditions can also be identified in patients with sporadic PAs, especially in those with early-onset disease. The recognition of these patients is particularly important, as it can allow to identify unaffected carriers who will benefit from regular clinical screening which could result in early diagnosis and possibly improved treatment outcomes (Hernandez-Ramirez *et al.* 2015).

In this review, we aim to discuss the genetic causes of familial and sporadic pituitary tumours, focusing on germline and somatic mosaic mutations causing FIPA and syndromic conditions predisposing to pituitary tumours, including MEN1, MEN4, CNC, MAS, 3PAs and DICER1 syndrome (summarised in Table 1). As the clinical features of these conditions have been extensively reviewed elsewhere (Vasilev *et al.* 2011, Beckers *et al.* 2013, Caimari & Korbonits 2016, Marques & Korbonits 2017), here we will aim to focus on the genetic aspects and the mechanisms linking monogenic mutations with PA pathogenesis.

Familial isolated pituitary adenoma

FIPA is an inherited condition characterised by the occurrence of PAs in two or more members of the same family with no other associated manifestations (Beckers *et al.* 2013). It is estimated to account for about 2% of all PAs (Daly *et al.* 2006b). In a recent study looking systematically at the prevalence of familial PAs among patients with functioning pituitary tumours, FIPA was identified in 10/262 patients (3.8%) (Marques *et al.* 2017). FIPA is a highly clinically heterogeneous condition and can include families where affected family members have the same PA subtype (homogeneous FIPA) or families with different PAs (heterogeneous FIPA). Most homogeneous FIPA kindreds present with prolactinomas or somatotroph PAs, followed by NFPAs and, rarely, corticotroph PAs, while in heterogeneous FIPA families, all possible combinations of different PA subtypes can be observed, with the association of somatotroph PAs and prolactinomas being the most common.

Table 1 Germline and mosaic mutations causing pituitary tumours.

Syndrome	Gene (inheritance pattern)	Germline or mosaic	Location	Penetrance for pituitary disease	Main clinical characteristics
FIPA	<i>AIP</i> (AD)	Germline	11q13.2	15–30%	Young-onset (typically in the second decade) somatotroph or mixed somatotroph–lactotroph PAs and prolactinomas. Responsible for 15–30% of FIPA kindreds and up to 20% of young-onset PAs (typically causing gigantism or early-onset acromegaly) Early-onset (<4 years) gigantism
MEN1	<i>GPR101</i> (X-linked)	Germline or somatic mosaic in males with sporadic disease	Xq26.3	100%	
	<i>MEN1</i> (AD)	Germline	11q13.1	30–40%	Hyperparathyroidism, PAs (mostly prolactinomas and NFPAs), GEP NETs, other neoplasms
MEN4	<i>CDKN1B</i> (AD)	Germline	12p13.1	Unknown	MEN1-like phenotype
Carney complex	<i>PRKAR1A</i> (AD)	Germline	17q24.2	10% (symptomatic acromegaly)	Skin pigmented lesions, cardiac and cutaneous myxomas, multiple non-endocrine and endocrine neoplasms including pituitary hyperplasia and PAs (mostly somatotroph and lactotroph or mixed, very rarely corticotroph PAs)
McCune–Albright syndrome	<i>Unknown gene</i> <i>PRKACB</i>	Germline Germline	2p16 1p31.1	Unknown Unknown	Same as for <i>PRKAR1A</i> Described in one case with CNC phenotype (Forlino <i>et al.</i> 2014)
	<i>GNAS</i> (not inheritable)	Somatic mosaic	20q13.32	20%	Café-au-lait spots, polyostotic fibrous dysplasia, precocious puberty, GH excess in about 20% of patients
Phaeochromocytoma/paraganglioma with pituitary adenoma	<i>SDHA</i> (AD)	Germline	5p15.33	<1%	Familial PPGL
	<i>SDHB</i> (AD)	Germline	1p36.13	<1%	Familial PPGL
	<i>SDHC</i> (AD)	Germline	1q23.3	<1%	Familial PPGL
	<i>SDHD</i> (AD)	Germline	11q23.1	<1%	Familial PPGL
	<i>MAX</i> (AD)	Germline	14q23.3	Unknown	Familial PPGL
DICER1 syndrome	<i>DICER1</i> (AD)	Germline or somatic mosaic	14q32.13	<1%	Early-onset pituitary blastomas (ACTH-secreting)

AD, autosomal dominant; CNC, Carney complex; FIPA, familial isolated pituitary adenoma; GEP NET, gastroenteropancreatic neuroendocrine tumour; GPCR, G protein-coupled receptor; MEN1, multiple endocrine neoplasia type 1; MEN4, multiple endocrine neoplasia type 4; NFPA, non-functioning pituitary adenoma; PPGL, phaeochromocytoma and paraganglioma.

Most FIPA cases have no known genetic cause, while *AIP* mutations can be identified in 15–30% of FIPA families (Vierimaa *et al.* 2006, Daly *et al.* 2007, Leontiou *et al.* 2008, Hernandez-Ramirez *et al.* 2015). Owing to the low penetrance of the disease, *AIP* mutations can also be identified in subjects with early-onset PAs, and typically among those with gigantism and early-onset acromegaly (Tichomirowa *et al.* 2011, Cuny *et al.* 2013, Hernandez-Ramirez *et al.* 2015). Very rarely, duplications of Xq26.3 involving the *GPR101* gene have been identified in families with XLAG (Trivellin *et al.* 2014, Gordon *et al.* 2016). To this date, most of the reported XLAG patients presented as isolated cases due to *de novo* germline or somatic mosaic mutations, and only three cases of familial XLAG have been so far described (Trivellin *et al.* 2014, Gordon *et al.* 2016).

Aryl hydrocarbon receptor-interacting protein

The *AIP* gene maps to chromosome 11q13.2, consists of six exons and encodes a 330 amino acid protein. The AIP protein is characterised by an N-terminal immunophilin-like domain and a C-terminal tetratricopeptide (TPR) domain containing three TPR motifs a C-terminal alpha helix. The TPR domain of AIP is considered important to mediate the binding between AIP and its numerous interacting partners, in line with the role of AIP as a co-chaperone (Morgan *et al.* 2012). The best characterised function of AIP is to form, together with the heat shock protein 90 (HSP90), a protein complex which regulates the nuclear translocation of the aryl hydrocarbon receptor (Ma & Whitlock 1997). However, AIP has several other interacting partners, including other members of the heat shock protein family, growth factor receptors, nuclear receptors and viral proteins (Trivellin & Korbonits 2011).

Despite *AIP* being ubiquitously expressed, no other manifestations other than PAs have been consistently associated with mutations in the *AIP* gene. Interestingly, in the normal human pituitary gland, AIP is exclusively found in somatotroph and lactotroph cells, while its expression has been described in all PA subtypes and is particularly abundant in NFPAs (Leontiou *et al.* 2008). The mechanisms underlying the pituitary-specific pro-tumorigenic effects of *AIP* mutations remain to be elucidated, but are likely to involve different pathways. The pro-tumorigenic effects of *AIP* mutations depend on the loss of its tumour suppressor function. *AIP* mutations in fact result, in most cases, in the premature truncation of the coding sequence (nonsense and frameshift mutation) or in highly unstable proteins with reduced

half-life (missense mutations or segmental duplications) (Hernandez-Ramirez *et al.* 2016, Salvatori *et al.* 2017). Moreover, loss of heterozygosity (LOH) at the *AIP* locus is often found in *AIP*-related PAs (Gadelha *et al.* 1999, Vierimaa *et al.* 2006), confirming that the tumorigenic process depends on the loss of AIP. Several lines of evidence suggest a link between AIP and the cyclic AMP (cAMP)-dependent protein kinase A (PKA) pathway, which plays a central role in regulating GH expression and proliferation of somatotroph cells (Formosa & Vassallo 2014). The binding of GHRH to its receptor on the somatotroph cell determines the activation of a G stimulatory protein with consequent increase in cAMP levels and activation of PKA. The phosphorylation of the cAMP response element-binding protein (CREB) and the CREB-binding protein is responsible for the activation of the GH promoter mediated by PIT1, a transcription factor involved with pituitary development and pituitary hormone expression (Cohen *et al.* 1999). The cAMP-dependent PKA pathway and the effects of AIP on this pathway are summarised in Fig. 1. In GH3 mammosomatotroph cells, overexpression of *AIP* inhibits the cAMP response to forskolin, an adenylate cyclase activator, while *AIP* knockdown leads to enhanced cAMP production (Formosa *et al.* 2013). The exact mechanisms linking AIP with the cAMP-dependent PKA pathway remain to be fully elucidated. In human *AIP* mutation-positive PAs, the expression of the G inhibitory protein $G\alpha_{i-2}$ was found to be reduced compared to *AIP* mutation-negative tumours (Tuominen *et al.* 2015), and the expression of *AIP* was found to be positively associated with that of $G\alpha_{i-2}$ also in sporadic *AIP* mutation-negative PAs (Ritvonen *et al.* 2017). Moreover, while knockdown of $G\alpha_{i-2}$ and $G\alpha_{i-3}$ led to a significant increase in cAMP in *AIP* WT cells, this effect was not observed in *AIP*-knockout cell lines (Tuominen *et al.* 2015), suggesting that the loss of AIP can affect G inhibitory protein function. However, other mechanisms could also link AIP and the cAMP signalling pathway. For instance, AIP has been found to interact with both the catalytic (PRKACA) and the regulatory (PRKAR1A) subunits of PKA (Scherthaner-Reiter *et al.* 2018). An interaction between AIP and PRKACA was demonstrated in the presence of HSP90, and cytoplasmic co-localisation of AIP and PRKACA was observed (Hernandez-Ramirez *et al.* 2018, Scherthaner-Reiter *et al.* 2018), suggesting that the AIP/HSP90 complex could regulate PKA localisation and potentially affect the interaction between the catalytic and regulatory subunits of PKA. Moreover, AIP has been shown to interact with members of the type 4 phosphodiesterases family,

Figure 1

The cAMP-dependent protein kinase A pathway in the somatotroph cell and genes involved in its regulation. Under normal conditions, GHRH released by the GHRH neurons in the arcuate nucleus (ARC) of the hypothalamus determines activation of the adenylate cyclase (AC) through its G stimulatory protein-coupled receptor (GHRH receptor, GHRH-R) in the somatotroph cell. The increased cAMP production causes the regulatory subunits (R) of protein kinase A (PKA) to dissociate from the catalytic subunits (C), which can translocate to the nucleus and phosphorylate its targets, including CREB. Phosphorylated CREB can bind to the promoter of *PIT1*. These events are required to promote the expression of GH and somatotroph cell proliferation. Loss of AIP has been shown to increase cAMP production via various possible mechanisms, including reduced expression of the G inhibitory protein G_{α_i2} (which exerts an inhibitory effect on AC and is also involved in mediating the inhibitory effects of somatostatin (SS) on GH secretion via somatostatin receptors (SSTR)), AIP interaction with phosphodiesterases type 4 (PDE4), as well as its interaction with members of the PKA complex. This pathway is also affected by other genetic conditions, including Carney complex (CNC) due, in most cases, to inactivating mutations in the regulatory type 1 α subunit of PKA, and McCune–Albright syndrome (MAS), caused by post-zygotic activating mutations of the G stimulatory protein G_{α_s} . While *GPR101* is not expressed in adult human somatotroph cells, the duplication of *GPR101* causing X-linked acrogigantism (XLAG) could affect the cAMP-dependent PKA pathway, as *GPR101* is a G_{α_s} -coupled constitutively active receptor and is significantly overexpressed in the tumours of affected patients. Potentially, *GPR101* could also play a role in regulating GHRH secretion in the arcuate nucleus, where *GPR101* is physiologically expressed at high levels. 3V, third ventricle.

such as PDE4A5 (Bolger *et al.* 2003) – enzymes involved in the degradation of cAMP. Interestingly, the expression of PDE4A4 (human homologue of PDE4A5) and PDE4A8 was found to be significantly reduced in *AIP* mutation-positive somatotroph adenomas (Bizzi *et al.* 2018), suggesting that reduced expression of PDE4 enzymes might contribute to the enhanced cAMP signalling observed as a consequence of the loss of AIP.

AIP mutation-positive PAs are often resistant to treatment with somatostatin analogues (SSAs), despite expressing somatostatin receptors at levels comparable to sporadic *AIP* mutation-negative PAs (Chahal *et al.* 2012). Moreover, SSA resistance has also been observed in sporadic tumours with reduced AIP protein expression independently of the expression of the somatostatin receptor subtype 2 (SSTR2) (Kasuki *et al.*

2012, Iacovazzo *et al.* 2016a, Ozkaya *et al.* 2018). Thus, mechanisms other than altered somatostatin receptor expression are likely to be involved in determining resistance to SSAs in patients harbouring an *AIP* mutation. Reduced G_{α_i} protein expression observed in *AIP*-related PAs could potentially underlie their resistance to SSAs, as G_{α_i} signalling is involved in mediating the anti-secretory effect of SSAs (Theodoropoulou & Stalla 2013). Moreover, *AIP* knockdown was found to reduce the mRNA expression of *ZAC1*, a putative tumour suppressor gene involved in the anti-proliferative and anti-secretory effects of SSAs (Chahal *et al.* 2012). Notably, a positive correlation was described between *ZAC1* protein expression and IGF-1 normalisation and tumour shrinkage in a group of 45 patients with acromegaly (Theodoropoulou *et al.* 2009). While the mechanisms linking AIP and *ZAC1* remain

to be elucidated, the downregulation of *ZAC1* observed following knockdown of *AIP* suggests that this could be one of the mechanisms underlying the SSA resistance often observed in *AIP*-related PAs.

While *AIP*-related PAs are often invasive and clinically aggressive, this is rarely observed in other monogenic conditions predisposing to PAs through dysregulation of the cAMP-PKA pathway, such as CNC or MAS. Considering the vast repertoire of *AIP*-interacting proteins, cAMP-independent mechanisms could contribute to the clinical phenotype of *AIP*-related PAs. *AIP* has been recently shown to interact with proteins involved in the organisation of the cytoskeleton (Hernandez-Ramirez *et al.* 2018), such as members of the tubulin family, and specifically TUBB and TUBB2A (Hernandez-Ramirez *et al.* 2018). Moreover, two isoforms of beta tubulin, *TUBB1* and *TUBB2B*, were found to be significantly downregulated at the mRNA level in *AIP*-related PAs compared with the normal human pituitary gland (Hernandez-Ramirez *et al.* 2018). A direct interaction was also demonstrated between *AIP* and NME1 (Hernandez-Ramirez *et al.* 2018), a protein with anti-metastatic properties involved in the regulation of cell migration and motility (Murakami *et al.* 2008). Interestingly, *NME1* knockdown was found to disrupt E-cadherin-mediated cell adhesion in human hepatoma and colon cancer cell lines, suggesting a critical role for NME1 in the control of intercellular adhesions and cell migration (Boissan *et al.* 2010). In one study, an inverse relationship between NME1 expression and PA invasiveness was demonstrated (Pan *et al.* 2005). *AIP*-related somatotroph PAs are typically sparsely granulated (Hernandez-Ramirez *et al.* 2015) – a tumour subtype which is characterised by decreased E-cadherin expression and increased invasiveness (Nishioka *et al.* 2003, Sano *et al.* 2004) – suggesting that the loss of *AIP* could contribute, through the alteration of the cytoskeleton organisation, to the invasive and aggressive phenotype often observed in *AIP*-related PAs.

Aip-deficient mouse models generally recapitulate the human phenotype (Raitila *et al.* 2010, Gillam *et al.* 2017). While constitutional *Aip*-knockout animals die *in utero* and display severe cardiovascular defects (Lin *et al.* 2007), *Aip*^{+/-} mice are viable and develop pituitary tumours with full penetrance by the age of 15 months (Raitila *et al.* 2010), although in the same mouse model, only pituitary hyperplasia without occurrence of PAs was observed in 3- and 12-month-old animals (Lecoq *et al.* 2016b). *Aip*^{+/-} mice develop PAs at a higher rate compared to WT mice, where incidental pituitary tumours, mostly prolactinomas, are also frequently observed. The majority of the tumours

found in *Aip*^{+/-} mice produce GH, although prolactinomas and mixed somatotroph–lactotroph adenomas were also seen (Raitila *et al.* 2010). LOH of the WT *Aip* allele was observed in two available tumour samples, confirming the findings in human tumours. Moreover, *Aip*^{+/-} mice had higher circulating IGF-1 levels, and their pituitary tumours showed increased cell proliferation, evaluated via immunohistochemistry for Ki-67, compared to PAs observed in WT mice (Raitila *et al.* 2010). More recently, another mouse model where *Aip* was deleted specifically in the somatotroph cells has been characterised (Gillam *et al.* 2017). *Aip*-knockout animals were found to be bigger than WT controls both in terms of body length and weight beginning at 12 weeks of age (Gillam *et al.* 2017). Visceral organs, including heart, liver and kidney, were found to be larger compared to those in WT mice, and both GH and IGF-1 levels were significantly increased by 18 weeks of age. Macroscopic tumours, evaluated by MRI, were visible in 80% of mutant mice by the age of 20 weeks. Histological examination showed somatotroph cell adenomas which were preceded by pituitary hyperplasia observed starting from the age of 18 weeks (Gillam *et al.* 2017). Interestingly, markedly reduced expression of the cyclin-dependent kinase inhibitor p27 was observed in the adenomatous tissue, suggesting that dysregulation of cell cycle regulators, similarly to what has been observed for sporadic human PAs (Bamberger *et al.* 1999), could contribute to the neoplastic transformation.

The disease penetrance in *AIP* mutation carriers is typically low. Studies on large families show a penetrance of 15–30% (Vierimaa *et al.* 2006, Naves *et al.* 2007, Chahal *et al.* 2011, Williams *et al.* 2014), suggesting that environmental or additional genetic factors could participate in determining the risk of developing *AIP*-related PAs. PAs in *AIP* mutation carriers arise at a younger age compared to their sporadic counterpart, presenting clinically in most cases between the second and third decades of life, are often macroadenomas and are frequently larger and more invasive compared to *AIP* mutation-negative PAs (Daly *et al.* 2007, 2010, Igreja *et al.* 2010, Hernandez-Ramirez *et al.* 2015). In some (Leontiou *et al.* 2008, Daly *et al.* 2010), albeit not all, studies (Hernandez-Ramirez *et al.* 2015), a male preponderance has been described. This finding could be potentially explained by an ascertainment bias due to the prevalent inclusion of patients with gigantism, a condition that is more common in males (Rostomyan *et al.* 2015). Owing to the young onset of the disease, about 30% of *AIP*-related PAs manifest clinically with gigantism, a manifestation of GH excess starting at an early age,

before the closure of the growth plates (Leontiou *et al.* 2008, Daly *et al.* 2010). Apoplexy is relatively frequent in AIP-related PAs (about 8–10% of all cases) and can represent the presenting feature of the disease (Xekouki *et al.* 2013). Poor responsiveness to SSAs is common in AIP-related somatotroph adenomas (Leontiou *et al.* 2008, Daly *et al.* 2010) and an increased prevalence of AIP mutations has been described among sporadic patients with acromegaly who are resistant to SSAs (Oriola *et al.* 2013).

The vast majority (80%) of AIP-related PAs are represented by somatotroph adenomas, followed by mixed somatotroph–lactotroph and more rarely prolactinomas (Stiles & Korbonits 2011), while non-functioning pituitary adenomas (NFPAs) are rare, accounting for less than 10% of cases (Daly *et al.* 2010, Igreja *et al.* 2010), although many of these tumours are silent somatotroph/lactotroph adenomas, as they can be found to express GH or prolactin (Daly *et al.* 2010, Villa *et al.* 2011). Corticotroph and thyrotroph PAs have been very rarely described in AIP mutation carriers (Daly *et al.* 2010, Cazabat *et al.* 2012).

Over 100 different mutations have been identified in the AIP gene (Daly *et al.* 2010, Hernandez-Ramirez *et al.* 2015), including nonsense, missense, frameshift, splicing and promoter mutations, deletions, insertions and segmental duplications. About 70% of these mutations lead to the loss of the C-terminal end of AIP, due to either nonsense or frameshift mutations resulting in premature stop codons (Hernandez-Ramirez *et al.* 2015). A minority of mutations is represented by large deletions (<10%) (Georgitsi *et al.* 2008), highlighting the need to employ dedicated techniques, such as multiplex ligation-dependent probe amplification, in order to correctly identify these mutations.

X-linked acrogigantism

XLAG is a condition of early-onset pituitary gigantism due to the germline or somatic mosaic duplication of the *GPR101* gene (Trivellin *et al.* 2014, Iacovazzo *et al.* 2016b, Iacovazzo & Korbonits 2018). XLAG patients present with marked GH excess, in most cases with associated hyperprolactinaemia, caused by mixed somatotroph–lactotroph adenomas associated, in some patients, with pituitary hyperplasia. In a minority of patients, the disease is due to pituitary hyperplasia in the absence of a PA. XLAG is very rare, with only 33 confirmed cases described so far in the medical literature (Trivellin *et al.* 2014, Beckers *et al.* 2015, 2017, Gordon *et al.* 2016, Iacovazzo *et al.* 2016b, Rodd *et al.* 2016). XLAG accounted for approximately 10 and 8% of cases in two large independent series of patients

with pituitary gigantism, respectively (Rostomyan *et al.* 2015, Iacovazzo *et al.* 2016b). Different from other forms of gigantism, including those linked with AIP mutations and those without a known genetic predisposing factor, where most affected patients are males, XLAG is characterised by a female preponderance, and 24/33 reported XLAG patients are females carrying germline duplications, while somatic mosaic mutations have been identified in the only four reported cases of male patients with sporadic disease (Daly *et al.* 2016b, Iacovazzo *et al.* 2016b, Rodd *et al.* 2016). In three independent families, mother-to-son transmission has been described, in all cases with full penetrance (Trivellin *et al.* 2014, Gordon *et al.* 2016). As no other clinical manifestations have been described in these kindreds, XLAG is considered as a rare cause of FIPA.

The *GPR101* gene (Xq26.3) encodes a G-protein-coupled receptor whose ligand is unknown. In mice, *Gpr101* mRNA was identified primarily in the central nervous system, and particularly in the hypothalamus and amygdala (Bates *et al.* 2006). In humans, *GPR101* was found to be highly expressed at the mRNA level in the nucleus accumbens, as well as in the medulla and the occipital lobe (Trivellin *et al.* 2016a). Interestingly, *Gpr101* was found to be expressed in about half of the neuronal cells expressing the anorexigenic neuropeptide pro-opiomelanocortin in mice (Nilaweera *et al.* 2007). In the same study, starvation was found to increase *GPR101* expression in the posterior hypothalamus, while decreased expression was seen in obese mice carrying the *ob* gene mutation, suggesting a possible role for GPR101 in regulating appetite and energy metabolism. While *GPR101* was found to be significantly overexpressed in the pituitary tumours of XLAG patients, it was not expressed in sporadic somatotroph PAs or in the adult human pituitary gland (Trivellin *et al.* 2014). On the contrary, GPR101 protein expression was described using immunohistochemistry in the foetal human pituitary and in pituitary samples obtained from adolescents, suggesting that its expression, at least in the pituitary gland, could be age dependent and induced during development and adolescence (Trivellin *et al.* 2016a). The expression pattern of *GPR101* in the pituitary gland also seems to be species specific. In the pituitary of the rhesus monkey, for example, *GPR101* was uniquely expressed, at the protein level, in gonadotroph cells, while in the rat pituitary gland, *GPR101* was found to be expressed only in a subpopulation of somatotroph cells (Trivellin *et al.* 2016a).

The mechanisms underlying the pathogenesis of XLAG remain to be determined. GPR101 is coupled with the G stimulatory protein, and is constitutively active,

as shown by the increased production of cAMP following its overexpression in HEK293 and GH3 cells (Bates *et al.* 2006, Trivellin *et al.* 2014). Thus, the activation of the cAMP-PKA pathway induced by the overexpression of *GPR101* could potentially underlie the development of pituitary hyperplasia and PAs in XLAG patients. Interestingly, elevated circulating GHRH levels have been described in some, although not all, XLAG patients, suggesting that *GPR101* could also have a role in the regulation of GHRH secretion (Glasker *et al.* 2011, Beckers *et al.* 2015, Daly *et al.* 2016a, Iacovazzo *et al.* 2016b). Notably, the GHRH receptor was found to be abundantly expressed in XLAG patients' hyperplastic and tumour pituitary samples (Trivellin *et al.* 2014), and this could possibly relate to increased hypothalamic secretion of GHRH, as GHRH was shown, at least *in vitro*, to induce the expression of its own receptor (Horikawa *et al.* 1996). GHRH administration was found to concomitantly stimulate the release of both GH and prolactin in XLAG patients, both *in vivo* (Moran *et al.* 1990) and *in vitro* (Daly *et al.* 2016a), and this effect was abolished by concomitant treatment with a GHRH receptor antagonist in cells cultured from the PA of an XLAG patient (Daly *et al.* 2016a). The implication of *GPR101* in the hypothalamic regulation of GHRH secretion is further supported by the finding that *GPR101* was found to be expressed at higher levels in the arcuate nucleus where, among others, GHRH neurons are localised (Bates *et al.* 2006). Potentially, the duplication of *GPR101* could affect pituitary somatotroph cells both directly, as a result of its constitutive activity and activation of the cAMP-PKA pathway, and indirectly through increased GHRH secretion by the hypothalamus (Fig. 1). In this latter scenario, *GPR101* could be involved in the hypothalamic regulation of the GHRH–GH axis, similarly to the action of *GPR54*, a G protein-coupled receptor expressed in the GnRH neurons which mediates the stimulatory effects of kisspeptin on GnRH release (Franssen & Tena-Sempere 2018).

The clinical features of XLAG patients are strikingly uniform. The disease is characterised by early onset of accelerated growth, in most cases observed during the first 2 years of life, and in all patients before the age of 4 (Beckers *et al.* 2015, Iacovazzo *et al.* 2016b). XLAG patients present markedly elevated GH levels resulting in significantly increased IGF-1 and height SDS, which are higher compared to patients with gigantism due to *AIP* mutations or to genetically undetermined cases (Rostomyan *et al.* 2015, Iacovazzo *et al.* 2016b). Other frequently observed features at diagnosis include acral enlargement, coarse facial features, increased appetite and,

less frequently, acanthosis nigricans, sleep apnoea/snoring and hyperhidrosis (Beckers *et al.* 2015, Iacovazzo *et al.* 2016b). The histopathological features of XLAG-related PAs are also peculiar: these tumours present a typical sinusoidal and lobular architecture with frequent calcifications and follicle-like structures (Iacovazzo *et al.* 2016b).

Most XLAG patients harbour microduplications of Xq26.3 (in average spanning a region of about 500kb) involving the *GPR101* gene as well as three other neighbouring genes (Trivellin *et al.* 2014). However, one patient with a typical clinical phenotype was found to carry a complex genomic rearrangement with two duplicated regions separated by a normal copy number segment (Iacovazzo *et al.* 2016b). The distal duplication in this patient has allowed to narrow down the genomic region shared by all patients to an area encompassing solely (in its entirety) the *GPR101* gene, confirming its pathogenic role (Iacovazzo *et al.* 2016b).

A missense variant in the *GPR101* gene (c.924C>G p.E308D; minor allele frequency in the GnomAD database 0.0036) was initially described in about 4% of a series of patients with acromegaly and was found to modestly increase cell proliferation and GH release when expressed in GH3 cells (Trivellin *et al.* 2014). However, further studies have failed to show an increased prevalence of this variant in patients with acromegaly (Ferrau *et al.* 2016, Iacovazzo *et al.* 2016b), suggesting it might not play a role in the pathogenesis of somatotroph PAs. A further variant (c.1098C>A p.D366E) was described in one patient with sporadic acromegaly (Kamenicky *et al.* 2015). Although no *in vitro* studies are available, this variant was not identified in a series of almost 400 patients with acromegaly (Iacovazzo *et al.* 2016b). Other *GPR101* missense variants have been detected in patients with other PA subtypes, although their impact on the function of *GPR101* remains to be determined (Lecoq *et al.* 2016a, Trivellin *et al.* 2016b). No pathogenic *GPR101* mutations or copy number variations were identified in patients with congenital isolated GH deficiency (Castinetti *et al.* 2016).

Syndromic pituitary tumours

Multiple endocrine neoplasia type 1

MEN1 is a tumour predisposition syndrome inherited with an autosomal dominant pattern occurring with a prevalence between 1:10,000 and 1:100,000 (Pardi *et al.* 2015). Affected individuals develop mainly parathyroid hyperplasia or parathyroid adenomas – causing primary

hyperparathyroidism (in over 90% of patients by the age of 50 years) – gastroenteropancreatic neuroendocrine tumours (NETs, in approximately 60% of patients) and PAs (30–40% of cases). Other endocrine and non-endocrine tumours may also occur in this syndrome, such as bronchial and thymic NETs, facial angiofibromas, lipomas, collagenomas, adrenal cortical adenomas, meningiomas, ependymomas, breast cancer and, rarely, pheochromocytomas (Marini *et al.* 2006, Dreijerink *et al.* 2014, Thakker 2014, Maxwell *et al.* 2016). A diagnosis of MEN1 can be established in (i) an individual carrying a pathogenic MEN1 mutation, (ii) a patient with two or more main MEN1 manifestations or (iii) a patient with one MEN1-associated manifestation and a first-degree relative affected with MEN1 (Thakker *et al.* 2012). In 90% of cases, MEN1 is due to germline heterozygous mutations in the MEN1 gene. Most MEN1 patients have a positive family history for MEN1-associated manifestations, while *de novo* mutations occur in approximately 10% of the patients (Chandrasekharappa *et al.* 1997, Bassett *et al.* 1998). The MEN1 gene is located on the long arm of chromosome 11 (11q13) and acts as a tumour suppressor gene: heterozygous inactivating mutations in this gene predispose to the occurrence of tumours and in about 90% of MEN1-related tumours LOH at 11q13 can be identified (Larsson *et al.* 1988, Dong *et al.* 1997).

MEN1 encodes a protein named menin, a scaffold protein located mostly in the nucleus, involved in several cellular processes, including transcriptional regulation, genome stability, cell division and proliferation (Thakker 2014). The first identified direct partner of menin was JunD, a component of the AP1 transcription factor complex (Agarwal *et al.* 1999) which acts as a negative regulator of RAS-dependent cell proliferation and protects cells from p53-dependent senescence and apoptosis (Pfarr *et al.* 1994, Weitzman *et al.* 2000). It has been reported that menin represses JunD-activated transcription via recruitment of histone deacetylases through association with the corepressor mDin3A, suggesting a role of menin as a repressor at the transcriptional level (Kim *et al.* 2003). Interestingly, patients carrying mutations in the JunD-interacting domain of menin present a higher mortality risk (Thevenon *et al.* 2013). It has also been shown that menin serves as a molecular adaptor to allow the interaction between the mixed lineage leukaemia (MLL) protein and the transcriptional coactivator lens epithelium-derived growth factor, which is needed for the association of the MLL complex with chromatin and the expression of MLL target genes (Yokoyama & Cleary 2008). Menin recruits MLL to the promoters of the

cyclin-dependent kinase inhibitor 1B (CDKN1B) and 1C (CDKN1C) (Milne *et al.* 2005, Wu & Hua 2011), promoting the transcription of these genes coding, respectively, for p27 and p57. The predominant expression of these genes, which control cell cycle progression at the G1 phase, in endocrine tissues might explain the selectivity of MEN1 tumorigenesis for endocrine organs. Moreover, cyclin-dependent kinase 4 (CDK4) has also been described as a target of MEN1 (Gillam *et al.* 2015). CDK4 regulates the cell cycle during G1/S transition, and its activation may be related to tumorigenesis in pituitary and pancreatic tissues (Gillam *et al.* 2015), as shown by the evidence that mice with heterozygous deletion of the *Men1* gene and concomitant knockout of *Cdk4* do not develop pituitary or pancreatic tumours (Gillam *et al.* 2015).

Murine *Men1* heterozygous knockout models develop a phenotype similar to that of MEN1 patients, with hyperplasia and tumours mainly of the parathyroids, pancreatic islets and anterior pituitary (Crabtree *et al.* 2001). Constitutional homozygous *Men1*-knockout mice die at an early embryonic stage (Crabtree *et al.* 2001), while conditional tissue-specific disruption of menin leads to pancreatic and pituitary tumorigenesis (Biondi *et al.* 2004).

To date, more than 1500 MEN1 mutations have been described (Lemos & Thakker 2008, Concolino *et al.* 2016). Most of these are represented by frameshift, missense and nonsense mutations (Lemos & Thakker 2008, Concolino *et al.* 2016), and they are distributed throughout the whole gene. A clear genotype–phenotype correlation has not been demonstrated (Kouvaraki *et al.* 2002, Verges *et al.* 2002, Horiuchi *et al.* 2013, de Laat *et al.* 2015). Approximately 30–40% of MEN1 patients develop PAs (Verges *et al.* 2002, Trouillas *et al.* 2008, de Laat *et al.* 2015), which can represent the first manifestation of the disease in about 15–30% of all patients. PAs are more commonly diagnosed in female patients. Lactotroph PAs are the most common PA subtype in MEN1 (40–60%), followed by NFPAs (15–40%), somatotroph PAs (5–10%) and, rarely, corticotroph or thyrotroph adenomas (Verges *et al.* 2002, Trouillas *et al.* 2008, de Laat *et al.* 2015). PAs in MEN1 are frequently macroadenomas, and they arise at a younger age compared to sporadic PAs and can be multiple.

Considering that PAs can represent the first disease manifestation and the rate of *de novo* mutations, screening for MEN1 should be considered in patients with childhood-onset pituitary macroadenomas, especially prolactinomas, as a relatively high frequency (6%) of MEN1 mutations has been shown in paediatric PA patients (Cuny *et al.* 2013).

Multiple endocrine neoplasia type 4

A small percentage of patients showing MEN1 clinical features do not harbour mutations in the *MEN1* gene. The characterisation of the phenotype (named MENX) of a rat strain harbouring a spontaneous *Cdkn1b* mutation prompted studies in patients with an MEN1-like phenotype. *CDKN1B* mutations have been identified in rare cases of such patients without identifiable *MEN1* mutations, and this condition has been named MEN4 (Pellegata *et al.* 2006). To date, 19 cases have been reported (reviewed in Alrezk *et al.* 2017): most of these patients developed primary hyperparathyroidism, either isolated or associated with NETs, mostly gastroenteropancreatic. Seven of the reported patients developed PAs, including four with a somatotroph PA, one with an NFPA, one with a corticotroph PA and one with a prolactinoma. One of the patients with a somatotroph tumour presented with gigantism due to a somatotroph macroadenoma diagnosed at the age of 5 years (Sambugaro *et al.* 2015). It should be noted that, in the case of two *AIP* mutation-negative FIPA kindreds harbouring two distinct *CDKN1B* variants (Tichomirowa *et al.* 2012), for one of the identified variants segregation with the PA phenotype could not be assessed, while for the second, only one of the two affected family members carried the variant, therefore making it an unlikely cause for their familial PA.

CDKN1B is located on chromosome 12q13 and encodes for p27, a tumour suppressor gene involved in cell cycle regulation (Chu *et al.* 2008). p27 is a member of the cyclin-dependent kinase inhibitors family and negatively regulates the cyclin E/cyclin-dependent kinase 2 complex preventing transition from the G1 to the S phase of the cell cycle (Sheaff *et al.* 1997). Interestingly, *Cdkn1b*-knockout mice develop hyperplasia of the intermediate lobe of the pituitary gland, and about 50% these animals develop pituitary tumours originating from the intermediate lobe (Nakayama *et al.* 1996). Development of pituitary tumours was also observed in *Cdkn1b*^{+/-} animals challenged with either irradiation or carcinogens, although no deletions or mutations of the WT allele were detected in these tumours, suggesting that p27 does not conform to the two-hit inactivation hypothesis and that tumorigenesis depends on haploinsufficiency rather than complete loss of the gene product (Fero *et al.* 1998). Reduced p27 protein expression was detected in all human PA subtypes (Bamberger *et al.* 1999), and especially in corticotroph PAs and pituitary carcinomas (Lidhar *et al.* 1999). Interestingly, p27 expression was significantly reduced in PAs compared to the normal pituitary cells of

the same subtype (Lidhar *et al.* 1999). The mechanisms underlying downregulation of p27 in human PAs remain to be determined; in a study including 48 PA patients, no differences were observed among the various PA subtypes and the normal pituitary in terms of expression of *CDKN1B* transcriptional regulators and specific miRNAs (Martins *et al.* 2016).

The *CDKN1B* mutations described so far include frameshift, nonsense, missense as well as 5' UTR mutations leading to reduced p27 expression (Pellegata *et al.* 2006, Georgitsi *et al.* 2007, Agarwal *et al.* 2009, Molatore *et al.* 2010, Costa-Guda *et al.* 2011, Belar *et al.* 2012, Tichomirowa *et al.* 2012, Occhi *et al.* 2013, Tonelli *et al.* 2014, Elston *et al.* 2015, Sambugaro *et al.* 2015, Borsari *et al.* 2017). Considering the rarity of MEN4, penetrance or potential genotype–phenotype correlations remain to be established.

Carney complex

CNC is a rare multiple neoplasia syndrome inherited with an autosomal dominant manner. CNC is characterised by the presence of pigmented lesions of the skin, cardiac and cutaneous myxomas and multiple non-endocrine and endocrine neoplasms, including pituitary hyperplasia and PAs (Carney *et al.* 1985). The most common endocrine manifestation observed in CNC patients is ACTH-independent Cushing's syndrome due to primary pigmented nodular adrenocortical disease (Stratakis *et al.* 1993, Bertherat *et al.* 2009, Rothenbuhler & Stratakis 2010, Courcoutsakis *et al.* 2013). This condition is observed in about 25% of CNC patients and occurs more often in females (Stratakis *et al.* 1993). Other endocrine manifestations observed in CNC include testicular tumours, especially large-cell calcifying Sertoli cell tumours, observed in about one-third of affected males at presentation, thyroid nodules (mostly follicular adenomas) which occur in up to 75% of CNC patients and, occasionally, also differentiated thyroid cancer (both papillary and follicular). About two-thirds of CNC patients show elevation of GH or IGF-1, often with associated hyperprolactinaemia, although symptomatic acromegaly occurs only in about 10% of CNC patients, usually by the third decade of life (Bertherat *et al.* 2009, Correa *et al.* 2015). Most CNC patients with acromegaly present with pituitary hyperplasia, typically affecting the mammosomatotroph cells, expressing both GH and prolactin (Stergiopoulos *et al.* 2004) which can be accompanied, in some patients, by one or multiple areas of adenomatous transformation. Most PAs observed in CNC

are represented by somatotroph or mixed somatotroph–lactotroph microadenomas (Stergiopoulos *et al.* 2004, Stratakis *et al.* 2004), although large macroadenomas have also been described. While most cases of Cushing's syndrome in CNC patients are ACTH independent, two cases of ACTH-dependent Cushing's disease have been described in patients with CNC harbouring a *PRKARIA* mutation (Hernandez-Ramirez *et al.* 2017b, Kiefer *et al.* 2017). In both cases, LOH at the *PRKARIA* locus has been described in the pituitary tumour, supporting a pathogenic role for the *PRKARIA* mutation in causing Cushing's disease in these patients.

The genetic background of CNC is heterogeneous. About 70% of cases are due to heterozygous inactivating mutations in the *PRKARIA* gene (17q24.2), coding for the regulatory subunit type 1 alpha of PKA. PKA is a cAMP-dependent protein kinase implicated in several cellular processes including hormone release, transcriptional regulation, cell cycle progression, cell proliferation and apoptosis (Bossis & Stratakis 2004). The PKA enzyme complex is a tetramer formed of two catalytic and two regulatory components. In the presence of cAMP, the enzymatic complex dissociates releasing the two catalytically active subunits (McKnight *et al.* 1988) (Fig. 1). To date, four regulatory subunits (R1 α , R1 β , R2 α and R2 β) and three catalytic subunits (C α , C β and C γ) have been identified and, depending on the tissue availability of each subunits, several combinational PKA configurations exist (Skalhegg & Tasken 2000). Two major enzymatic complexes have been identified, named PKA type I and II. Type I PKA contains either R1 α or R1 β regulatory subunits and is considered the main subtype that mediates response to cAMP in mammalian cells (Gamm *et al.* 1996). Loss-of-function *PRKARIA* mutations lead to increased cAMP-dependent PKA activity which drives tumour formation in tissues affected by CNC (Casey *et al.* 2000, Salpea *et al.* 2014). Interestingly, *PRKARIA* does not seem to behave like a 'classical' tumour suppressor gene (Bossis & Stratakis 2004). First, while LOH at the 17q24 locus has been shown in many CNC-related tumours (Kirschner *et al.* 2000a), in some cases LOH was not detected (Groussin *et al.* 2002), suggesting that haploinsufficiency might be sufficient for tumour development. Moreover, *PRKARIA* seems to behave like an oncogene in selected tissues. For instance, increased expression was described in several human malignancies, including renal and breast cancer (Fossberg *et al.* 1978, Handschin & Eppenberger 1979). Overexpression of *PRKARIA* was also shown to promote growth advantages in different cell lines, including Chinese hamster ovary cells and in breast epithelial

cell lines (Tortora *et al.* 1994a,b). Thus, *PRKARIA* could function as both an oncogene and tumour suppressor gene in a cell context-dependent way.

Homozygous deletion of *Prkar1a* is lethal in mice during embryogenesis (Amieux *et al.* 2002), while *Prkar1a*^{+/-} mice were found to develop CNC-related tumours, including Schwannomas and thyroid neoplasms, although no PAs were observed in these animals (Kirschner *et al.* 2005). In contrast, mice with pituitary-specific homozygous deletion of *Prkar1a* under the GHRH receptor promoter developed pituitary tumours of the Pit1 lineage expressing GH, prolactin and TSH, and had higher circulating levels of GH compared to WT mice (Yin *et al.* 2008).

To date, more than 125 *PRKARIA* mutations have been described (Correa *et al.* 2015), most of which are represented by nonsense and frameshift mutations. In the majority of cases, the predicted mutant protein products are not identified as a result of nonsense mRNA-mediated decay (Kirschner *et al.* 2000a,b). Large deletions can be found in about 20% of the cases where *PRKARIA* mutations cannot be detected by Sanger sequencing, and are often associated with a more severe phenotype (Horvath *et al.* 2008, Salpea *et al.* 2014). About 70% of CNCs arise in a familial setting, while 30% present sporadically, due to *de novo* mutations (Correa *et al.* 2015).

A second genetic locus at 2p16 has been associated with CNC; however, the responsible gene is not yet known (Stratakis 2016). There are no phenotypic differences between CNC patients with mutations at either locus. A single case of a CNC patient harbouring a triplication of the catalytic beta subunit of PKA (*PRKACB*) has been described (Forlino *et al.* 2014). This patient presented at the age of 19 years with acromegaly, spotty skin hyperpigmentation and multiple myxomas. Interestingly, PKA activity measured in the patient's lymphocytes was found to be increased to levels comparable to those seen in two *PRKARIA* mutated patients (Forlino *et al.* 2014), suggesting that overexpression of the C β catalytic subunit can affect PKA activity in a way similar to that observed in case of *PRKARIA* mutations.

McCune-Albright syndrome

Somatic activating mutations in the *GNAS* gene (20q13.32), encoding the cAMP pathway associated G protein Gs α , represent the only recurrent mutation found in somatotroph adenomas (Valimaki *et al.* 2015, Ronchi *et al.* 2016) and can be identified at a rate of 10–50% (Peverelli *et al.* 2014). These missense mutations are known to occur at only one of two residues, Arg201 (more commonly) or,

rarely, Gln227, which represent critical sites for GTPase activity. Mutations at these sites cause loss of the GTPase activity with consequent permanent activation of the adenylate cyclase and constitutive activation of the cAMP-dependent PKA pathway (Fig. 1). This results in increased cell proliferation in cAMP-responsive tissue, including the pituitary gland. As such, *GNAS* is considered a proto-oncogene, activated by these point mutations into the *gsp* oncogene. When these mutations occur at an early post-zygotic stage, the resulting somatic mosaicism underlies a syndromic condition known as MAS. MAS is a rare disorder with an estimated prevalence between 1:100,000 and a 1:1,000,000 (Boyce & Collins 1993) and is defined by the occurrence of polyostotic fibrous dysplasia, café-au-lait skin macules and endocrinopathies, including precocious puberty (especially in females), hyperthyroidism, testicular lesions (Leydig and/or Sertoli cell hyperplasia), growth hormone excess or, more rarely, neonatal hypercortisolism. The clinical manifestations of MAS are extremely variable and depend on the degree of mosaicism. The probability of detecting a *GNAS* mutation by standard PCR is high in affected tissues, while it can be very low in leukocyte-derived DNA, especially in subjects with only one manifestation of MAS (Lumbroso *et al.* 2004), although the use of next-generation sequencing can increase the mutation detection rates (Narumi *et al.* 2013). The *GNAS* gene is paternally imprinted in several tissues, including the pituitary gland, and most somatotroph adenomas have been found to occur in patients harbouring the mutation on the maternal allele (Hayward *et al.* 2001, Mantovani *et al.* 2004).

Pituitary involvement in MAS manifests with GH excess which can be present in about 20% of patients (Salenave *et al.* 2014). This is in most cases associated with hyperprolactinaemia. PAs can be found in 30–50% of affected patients, with the other patients most likely having pituitary hyperplasia without an adenoma (Galland *et al.* 2006). The age at onset is variable with a mean age of 24 years (Salenave *et al.* 2014). While cases of young onset disease have been described, the final stature in MAS patients is often normal, and this might be explained by the high prevalence of associated precocious puberty and increased levels of sex steroids.

Phaeochromocytoma/paraganglioma with pituitary adenoma

Germline heterozygous mutations in genes encoding succinate dehydrogenase subunits (*SDHx*) and the SDH complex assembly factor 2 protein (*SDHAF2*) have been described in patients with hereditary phaeochromocytoma

and paraganglioma (PPGL) (Baysal *et al.* 2000, Niemann & Muller 2000, Astuti *et al.* 2001, Hao *et al.* 2009, Bayley *et al.* 2010, Burnichon *et al.* 2010).

The first description of PPGL coexisting with a PA dates back to 1952 (Iversen 1952), but only recently a causative link between genes predisposing to PPGL and PAs has been established, following the description of a patient carrying an *SDHD* mutation having bilateral phaeochromocytomas and concomitant acromegaly due to a somatotroph PA. This patient's pituitary tumour showed loss of heterozygosity at the *SDHD* locus and reduced protein expression of both *SDHD* and *SDHB* (Xekouki *et al.* 2012). Other reports (Benn *et al.* 2006, Dwight *et al.* 2013, Varsavsky *et al.* 2013, Gill *et al.* 2014, Papathomas *et al.* 2014, Denes *et al.* 2015, Xekouki *et al.* 2015, Tufton *et al.* 2017, Maher *et al.* 2018), including a study showing that *Sdhb*^{+/-} mice develop pituitary lactotroph hyperplasia (Xekouki *et al.* 2015), have further strengthened the link between germline *SDHx* mutations and PAs, and this has allowed the definition of a novel clinical entity called 3PAs (phaeochromocytoma/paraganglioma with PAs). Notably, only one case of a double somatic mutation (detected by loss of *SDHB* and *SDHA* immunostaining and confirmed by sequencing) was described in 1/309 sporadic PAs, implying that this is an extremely rare event (Gill *et al.* 2014).

The SDH enzymatic complex is composed of two subunits which form the catalytic core (*SDHA* and *SDHB*) and two subunits which are responsible for anchoring the complex to the mitochondrial membrane (*SDHC* and *SDHD*). The SDH complex is responsible for the reversible enzymatic conversion of succinate into fumarate within the citric acid cycle (Bardella *et al.* 2011). The mechanisms linking loss of SDH function with tumorigenesis remain to be fully determined but are likely to be multifactorial (Fig. 2). Disruption of the SDH complex as a result of loss-of-function *SDHx* mutations leads to the accumulation of succinate, which in turn causes an inhibition of prolyl-hydroxylases, leading to stabilisation of the hypoxia-inducible factor 1 α (HIF1 α) and transcription of HIF-responsive genes, some of which are involved in tumorigenesis, including, among others, VEGF and TGF (Selak *et al.* 2005, Cervera *et al.* 2008, Guzy *et al.* 2008). These findings are corroborated by the evidence that gene expression profiling in *SDHx*-related paragangliomas overlaps with that observed in tumours due to *VHL* (encoding a component of the ubiquitin ligase complex that mediates the degradation of HIFs) and *EPAS1* (*HIF2A*) mutations (Comino-Mendez *et al.* 2013). Interestingly, HIF1 α was found to exert an anti-apoptotic role in a

Figure 2

Mechanisms involved with SDH-related tumorigenesis. The SDH enzymatic complex mediates the reversible enzymatic conversion of succinate into fumarate within the citric acid cycle (here represented schematically). Inactivating mutations in the *SDHx* genes are responsible for familial paragangliomas and pheochromocytomas, and a small subset of patients carrying an *SDHx* mutation develop PAs. The accumulation of succinate as a result of a loss-of-function *SDHx* mutation determines inhibition of prolyl-hydroxylases, which results in the stabilisation of the hypoxia-inducible factor 1 α (HIF1 α), and increased expression of HIF1 α -responsive genes. Indirect effects (dashed lines) of loss of SDH function include increased production of reactive oxygen species (ROS), which also have an inhibitory effect on prolyl hydroxylase activity and may cause genomic instability as a result of oxidative stress. Moreover, *SDHx* mutations are associated with a hypermethylator phenotype which results in the silencing of genes involved with epithelial-to-mesenchymal (EMT) transition and cell invasiveness. α -KG, alpha-ketoglutarate; OAA, oxaloacetic acid.

human PA cell line in hypoxic conditions (Yoshida *et al.* 2006) and hypoxia was found to induce invasiveness of the same cell line *in vitro* (Yoshida & Teramoto 2007). *HIF1A* knockdown and HIF1 α inhibition were shown to increase the sensitivity of human PA cells to temozolomide, an alkylating agent employed for the treatment of aggressive PAs and pituitary carcinomas, both *in vitro* and in PA xenografts (Chen *et al.* 2013). Altogether, these data support a role for the HIF1 α pathway in pituitary tumorigenesis and might provide a mechanistic link between *SDHx* mutations and PAs.

In addition, loss of SDH activity leads to increased intracellular production of reactive oxygen species, which can also contribute to the inhibition of prolyl-hydroxylases and to the stabilisation of HIF1 α (Niecknig *et al.* 2012). Moreover, reactive oxygen species promote a condition of chronic metabolic oxidative stress and genomic instability (Ishii *et al.* 2005, Slane *et al.* 2006). Whether this could as well contribute to the SDH-related tumorigenesis is yet to be determined, also considering that *SDH*-mutated paragangliomas were

found to harbour a low rate of somatic mutations or copy number alterations (Castro-Vega *et al.* 2015).

Interestingly, SDH-deficient tumours present with a significantly greater genomic methylation level compared to SDH-proficient neoplasms, as it was shown in gastrointestinal stromal tumours (Killian *et al.* 2013). Similar findings were shown in paragangliomas, where *SDHx*- and particularly *SDHB*-related tumours showed a hypermethylator phenotype (Letouze *et al.* 2013). Hypermethylated tumours were characterised by younger age at diagnosis and a worse prognosis. *Sdhb* knockout chromaffin cells displayed increased 5-methylcytosine and increased H3K9 and H3K27 methylation (Letouze *et al.* 2013). These methylome changes were associated with downregulation of several genes, including genes associated with neuroendocrine differentiation, the tumour suppressor gene *RBP1*, known to be downregulated in several human malignancies (Esteller *et al.* 2002, Mendoza-Rodriguez *et al.* 2013) and *KRT19* (encoding cytokeratin-19), a marker of epithelial-to-mesenchymal transition. Moreover, *Sdhb*-knockout mouse chromaffin

cells presented mesenchymal changes reminiscent of epithelial-to-mesenchymal transition (Loriot *et al.* 2015) with increased invasiveness and enhanced cell migration, and expression of *KRT19* by lentiviral transduction partially rescued the invasive phenotype of these cells and enhanced cell adherence (Loriot *et al.* 2015). While the hypermethylation secondary to *SDHx* mutations seem to play a pivotal role in mediating tumorigenesis in paragangliomas, no data are available whether this mechanism could also be involved in *SDHx*-associated PAs.

The penetrance of pituitary tumours in patients carrying *SDHx* mutations is estimated to be low (<1%). However, this might be an underestimation, considering that subjects carrying *SDHx* mutations are not routinely screened for pituitary tumours. Among 18 cases of *SDHx*-related PAs which were confirmed by genetic testing (Benn *et al.* 2006, Xekouki *et al.* 2012, 2015, Dwight *et al.* 2013, Varsavsky *et al.* 2013, Papathomas *et al.* 2014, Denes *et al.* 2015, Tufton *et al.* 2017, Maher *et al.* 2018), data regarding family history are available from 16 patients. Among these, 14 patients had a positive family history of PPGL (and PAs in two kindreds), while only two patients presented with sporadic disease. Most patients with *SDHx*-related PAs were diagnosed with PPGL (in most instances, PPGL were diagnosed first or simultaneously to the pituitary tumour), while five PAs occurred in patients without a personal history of PPGL. Among the 16 patients with available clinical data, most (10) were affected by prolactinomas, while somatotroph or NFPAs occurred in three cases each. Most of the reported cases were macroadenomas, in some cases displaying an aggressive behaviour, including a non-functioning pituitary carcinoma in a patient harbouring an *SDHB* mutation (Tufton *et al.* 2017). Notably, *SDHx*-related PAs showed peculiar histopathology features, with typical intracytoplasmic vacuoles (Denes *et al.* 2015, Tufton *et al.* 2017, Maher *et al.* 2018). As *SDHx* mutations are extremely rare in patients with sporadic PAs (Gill *et al.* 2014, Xekouki *et al.* 2015), sequencing of *SDHx* genes should be reserved for patients with a personal or family history of paraganglioma or pheochromocytoma. Considering the aggressive phenotype of *SDHx*-related PAs and the potential risk of malignancy, patients carrying *SDHx* mutations should be screened for pituitary tumours, although, owing to the small number of reported cases, frequency and modalities of screening remain to be established.

PAs have been recently reported in patients with pheochromocytomas harbouring mutations in the *MAX* gene (14q23.3) (Roszko *et al.* 2017, Daly *et al.*

2018, Kobza *et al.* 2018), including three patients with prolactinomas and two affected with acromegaly. Interestingly, three of these reported cases carried large deletions that were missed by Sanger sequencing. *MAX* is one of several genes causing predisposition to familial PPGL (Comino-Mendez *et al.* 2011) and encodes a protein which acts as an interacting partner for MYC and MXD1, transcription factors involved in the regulation of cell proliferation and apoptosis (Atchley & Fitch 1995). While the role of *MAX* in PA pathogenesis has not been investigated, these reports expand the knowledge on the genetic background of the 3PAs association and suggest that genes other than *SDHx* could be involved in its pathogenesis.

DICER1 syndrome

The DICER1 syndrome, or pleuropulmonary blastoma (PPB)-familial tumour and dysplasia syndrome, is a rare autosomal dominant disorder due to germline heterozygous mutations in the *DICER1* gene.

DICER1 syndrome is characterised by a variety of cancerous and benign tumours, including pleuropulmonary blastoma, ovarian sex cord-stromal tumours (mostly Sertoli-Leydig cell tumour), cystic nephroma, nodular hyperplasia of the thyroid, differentiated thyroid cancer, pituitary blastoma, nasal chondromesenchymal hamartoma, ciliary body medulloepithelioma, renal sarcoma, genitourinary embryonal rhabdomyosarcoma and pinealoblastoma (Doros *et al.* 1993, Schultz *et al.* 2018).

The first case of pituitary blastoma was characterised in 2008 in a 13-month-old female with ACTH-dependent Cushing's disease (Scheithauer *et al.* 2008), although a link with *DICER1* mutations was only established more recently (de Kock *et al.* 2014). The term blastoma was employed as these neoplasms presented the appearances of pituitary embryonic tissue with an aggressive clinical behaviour (Scheithauer *et al.* 2008). The histopathological features of pituitary blastomas are typical and include Rathke-like epithelial cells forming rosettes or gland-like structures admixed with secretory cells disposed in lobules rather than acini and positive for ACTH and, less frequently, also for GH. Pituitary blastomas are very rare in the setting of DICER1 syndrome (<1%). These aggressive tumours usually arise in young children (median age at presentation is 8 months with a range from 7 to 24 months), and they present clinically with severe ACTH-dependent Cushing's disease and, in some cases, ophthalmoplegia. The condition can be fatal in about 40% of the cases (Scheithauer *et al.* 2008, de Kock *et al.*

2014). Out of 12 pituitary blastoma patients available for genetic testing, *DICER1* mutations were identified in 11 patients (de Kock *et al.* 2014), suggesting that these rare tumours represent a pathognomonic feature of the *DICER1* syndrome.

The *DICER1* gene is located on the long arm of chromosome 14 (14q32.13). This gene encodes a cytoplasmic endoribonuclease responsible for processing precursor into mature miRNAs, which modulate mRNA expression at the post-transcriptional level (Krol *et al.* 2010). The pathogenesis in *DICER1* syndrome normally relies on a germline loss-of-function mutation (more often represented by a nonsense or frameshift mutation) followed by a second somatic 'hit', often involving the RNase IIIb catalytic domain of *DICER1* (Heravi-Moussavi *et al.* 2012, de Kock *et al.* 2014). Somatic mosaic mutations, often affecting the RNase IIIb catalytic domain, have also been identified using high-sensitivity detection systems (Brenneman *et al.* 2015, de Kock *et al.* 2016). Interestingly, these mutations appeared to be accompanied by second somatic mutations represented by truncating *DICER1* mutations outside the RNase IIIb domain or by LOH. Mutations affecting the RNase IIIb domain of *DICER1* lead to loss of its enzymatic activity and loss of miRNAs generated from the 5p strand of miRNA precursors (Gurtan *et al.* 2012, Heravi-Moussavi *et al.* 2012, Anglesio *et al.* 2013). *In vitro*, *DICER1* mutations were shown to lead to a reduction of 5p-derived miRNAs in ovarian Sertoli-Leydig cell tumours and to promote cell proliferation in a granulosa cell line via deregulation of the let-7 miRNA family (Wang *et al.* 2015), miRNAs with important roles in cell differentiation and proliferation (Bussing *et al.* 2008). Interestingly, in mice lacking epithelial *Dicer1*, increased *Fgf9* expression in the lung epithelium, possibly mediated by downregulation of miR-140, resulted in hyperplastic changes resembling those observed in pleuropulmonary blastoma, the primary manifestation of *DICER1* syndrome (Yin *et al.* 2015). The occurrence of pituitary blastomas in *DICER1* syndrome is likely related to miRNA deregulation. For instance, let-7 miRNAs have been shown to be downregulated in PAs (Bottoni *et al.* 2007, Amaral *et al.* 2009, Qian *et al.* 2009). Among its targets, these miRNAs regulate the expression of *HMGA2*, which is often overexpressed in prolactinomas (Finelli *et al.* 2002). Moreover, mice transgenic for *Hmga2* develop PAs, especially lactotroph and somatotroph tumours, supporting a role for this oncogene in PA pathogenesis (Fedele *et al.* 2002). However, the molecular mechanisms linking *DICER1* with pituitary blastomas still remain to be determined.

Other germline mutations linked with pituitary tumours

Recently, novel genes have been implicated with the occurrence of both sporadic and familial PAs. In one FIPA kindred with two cases of acromegaly and two NFPA, exome sequencing revealed a heterozygous missense mutation in the *CDH23* gene (10q22.1) (Zhang *et al.* 2017), encoding a cadherin member previously implicated in the pathogenesis of a subtype of Usher syndrome (Usher syndrome type 1D), an autosomal recessive condition of hearing impairment, vestibular dysfunction and retinitis pigmentosa (Bolz *et al.* 2001). The *CDH23* c.4136G>T p.R1379L missense variant was found to segregate with the PA phenotype and was predicted to alter the formation of hydrogen bonds and impair the calcium-binding ability and stability of one of the extracellular cadherin domains. In 3 of 11 other FIPA families, three *CDH23* missense variants were detected and found to co-segregate with the phenotype. All these variants were rare (minor allele frequency <0.05%) and predicted to be pathogenic by at least one *in silico* prediction tool employed. Out of 125 patients with sporadic PAs of different subtypes, 15 harboured rare *CDH23* variants predicted to be potentially pathogenic. All potentially pathogenic *CDH23* variants identified in this study were found to affect extracellular cadherin domains, and the frequency of these variants in the PA cohort was significantly higher compared to 260 local healthy control individuals (Zhang *et al.* 2017). No *in vitro* functional studies have been performed, and the mechanisms how *CDH23* mutations could lead to PA remain unclear. Of note, Usher syndrome patients or unaffected heterozygous mutation carriers are not known to be at increased risk of developing PAs, and the highly polymorphic nature of *CDH23* (and other Usher syndrome-related genes) can make the interpretation of genetic variants in this gene challenging (Le Quesne Stabej *et al.* 2012). Thus, further studies will be needed to confirm the role of *CDH23* in PA pathogenesis.

The *CABLES1* gene (18q11.2) is a cell cycle regulator involved in the negative regulation of cell cycle progression in corticotroph cells in response to glucocorticoids (Roussel-Gervais *et al.* 2016). *Cables1* knockdown was found to stimulate the growth of a corticotroph cell line (AtT-20 cells) and counteracted the inhibitory effects of glucocorticoids on cell growth. Interestingly, *CABLES1* expression was lost in about 50% of a series of 31 corticotroph adenomas, and this was strongly associated with loss of p27 expression (Roussel-Gervais *et al.* 2016). *CABLES1* was previously shown to maintain p21 protein

stability by antagonising its proteasomal degradation (Shi *et al.* 2015), while *Cables1*-knockout mouse embryonic fibroblasts displayed reduced p27 and p16 expression (Kirley *et al.* 2005), supporting a potential broader role for CABLES1 as a cell cycle regulator. Recently, four patients harbouring potentially pathogenic missense *CABLES1* variants were found in a cohort of 181 sporadic patients (2.2%) with Cushing's disease (Hernandez-Ramirez *et al.* 2017a). These variants affected residues located within or in proximity with the predicted cyclin-dependent kinase 3-binding domains of CABLES1. All four patients carrying these variants had corticotroph macroadenomas (one patient harboured a silent corticotroph PA) with high proliferation index and an aggressive behaviour, with two patients requiring more than one operation. None of these patients had a family history of Cushing's disease or other PAs. Tamoxifen-inducible chimeric CABLES1 proteins were produced and, while WT *CABLES1* inhibited cell growth when expressed in AtT-20 corticotroph cells in the presence of tamoxifen, this effect was lost in cells expressing the four mutant forms of *CABLES1*, supporting their pathogenic role. Further confirmatory studies will be necessary to assess the occurrence of *CABLES1* mutations in patients with Cushing's disease or other PAs.

Concluding remarks

While most PAs occur sporadically, about 5% of all PAs occur in a familial setting as a result of a genetic predisposing mutation. More commonly, familial PAs occur without other associated manifestations as FIPA – two genes, *AIP* and very rarely *GPR101*, are known to be responsible for this condition, while the causative gene(s) in the majority of FIPA kindreds are yet to be identified. PAs can also occur as part of syndromic conditions and can sometimes represent the first manifestation of the disease. Remarkably, somatotroph PAs and prolactinomas represent the most common PA subtypes associated with a predisposing genetic mutation. While the molecular mechanisms linking these mutations with pituitary tumorigenesis have not always been uncovered, several lines of evidence confirm the involvement of the cAMP-dependent PKA pathway, which plays a central role in regulating hormone secretion and proliferation in cells of the PIT1 lineage. In some cases, more than one pathway might be affected, as seems to be the case for *AIP*-related PAs.

Significant advances have been achieved in the field of pituitary genetics in recent years, although further

studies will be needed to better elucidate the molecular mechanisms linking genetic mutations and pituitary tumours. With the use of pangenomic techniques, novel genes involved in PA pathogenesis are expected to be discovered, and this will broaden our understanding of the mechanisms underlying PA formation.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

D I is supported by a George Alberti Research Training Fellowship funded by Diabetes UK (16/0005395). We acknowledge the support of the Medical Research Council for our studies on familial pituitary adenomas (MR/M018539/1).

References

- Aflorei ED & Korbonits M 2014 Epidemiology and etiopathogenesis of pituitary adenomas. *Journal of Neurooncology* **117** 379–394. (<https://doi.org/10.1007/s11060-013-1354-5>)
- Agarwal SK, Guru SC, Heppner C, Erdos MR, Collins RM, Park SY, Saggari S, Chandrasekharappa SC, Collins FS, Spiegel AM, *et al.* 1999 Menin interacts with the AP1 transcription factor JunD and represses JunD-activated transcription. *Cell* **96** 143–152. ([https://doi.org/10.1016/S0092-8674\(00\)80967-8](https://doi.org/10.1016/S0092-8674(00)80967-8))
- Agarwal SK, Mateo CM & Marx SJ 2009 Rare germline mutations in cyclin-dependent kinase inhibitor genes in multiple endocrine neoplasia type 1 and related states. *Journal of Clinical Endocrinology and Metabolism* **94** 1826–1834. (<https://doi.org/10.1210/jc.2008-2083>)
- Agustsson TT, Baldvinsdottir T, Jonasson JG, Olafsdottir E, Steinthorsdottir V, Sigurdsson G, Thorsson AV, Carroll PV, Korbonits M & Benediktsson R 2015 The epidemiology of pituitary adenomas in Iceland, 1955–2012: a nationwide population-based study. *European Journal of Endocrinology* **173** 655–664. (<https://doi.org/10.1530/EJE-15-0189>)
- Alrezk R, Hannah-Shmouni F & Stratakis CA 2017 MEN4 and CDKN1B mutations: the latest of the MEN syndromes. *Endocrine-Related Cancer* **24** T195–T208. (<https://doi.org/10.1530/ERC-17-0243>)
- Amaral FC, Torres N, Saggari F, Neder L, Machado HR, Silva WA Jr, Moreira AC & Castro M 2009 MicroRNAs differentially expressed in ACTH-secreting pituitary tumors. *Journal of Clinical Endocrinology and Metabolism* **94** 320–323. (<https://doi.org/10.1210/jc.2008-1451>)
- Amieux PS, Howe DG, Knickerbocker H, Lee DC, Su T, Laszlo GS, Idzerda RL & McKnight GS 2002 Increased basal cAMP-dependent protein kinase activity inhibits the formation of mesoderm-derived structures in the developing mouse embryo. *Journal of Biological Chemistry* **277** 27294–27304. (<https://doi.org/10.1074/jbc.M200302200>)
- Anglesio MS, Wang Y, Yang W, Senz J, Wan A, Heravi-Moussavi A, Salamanca C, Maines-Bandiera S, Huntsman DG & Morin GB 2013 Cancer-associated somatic DICER1 hotspot mutations cause defective miRNA processing and reverse-strand expression bias to predominantly mature 3p strands through loss of 5p strand cleavage. *Journal of Pathology* **229** 400–409. (<https://doi.org/10.1002/path.4135>)
- Astuti D, Latif F, Dallol A, Dahia PL, Douglas F, George E, Skoldberg F, Husebye ES, Eng C & Maher ER 2001 Gene mutations in the succinate

- dehydrogenase subunit SDHB cause susceptibility to familial pheochromocytoma and to familial paraganglioma. *American Journal of Human Genetics* **69** 49–54. (<https://doi.org/10.1086/321282>)
- Atchley WR & Fitch WM 1995 Myc and Max: molecular evolution of a family of proto-oncogene products and their dimerization partner. *PNAS* **92** 10217–10221. (<https://doi.org/10.1073/pnas.92.22.10217>)
- Ballmann C, Thiel A, Korah HE, Reis AC, Saeger W, Stepanow S, Kohrer K, Reifenberger G, Knobbe-Thomsen CB, Knappe UJ, *et al.* 2018 USP8 mutations in pituitary cushing adenomas-targeted analysis by next-generation sequencing. *Journal of the Endocrine Society* **2** 266–278. (<https://doi.org/10.1210/js.2017-00364>)
- Bamberger CM, Fehn M, Bamberger AM, Ludecke DK, Beil FU, Saeger W & Schulte HM 1999 Reduced expression levels of the cell-cycle inhibitor p27Kip1 in human pituitary adenomas. *European Journal of Endocrinology* **140** 250–255. (<https://doi.org/10.1530/eje.0.1400250>)
- Bardella C, Pollard PJ & Tomlinson I 2011 SDH mutations in cancer. *Biochimica et Biophysica Acta* **1807** 1432–1443. (<https://doi.org/10.1016/j.bbmbio.2011.07.003>)
- Bassett JH, Forbes SA, Pannett AA, Lloyd SE, Christie PT, Wooding C, Harding B, Besser GM, Edwards CR, Monson JP, *et al.* 1998 Characterization of mutations in patients with multiple endocrine neoplasia type 1. *American Journal of Human Genetics* **62** 232–244. (<https://doi.org/10.1086/301729>)
- Bates B, Zhang L, Nawoschik S, Kodangattil S, Tseng E, Kopsco D, Kramer A, Shan Q, Taylor N, Johnson J, *et al.* 2006 Characterization of Gpr101 expression and G-protein coupling selectivity. *Brain Research* **1087** 1–14. (<https://doi.org/10.1016/j.brainres.2006.02.123>)
- Bayley JP, Kunst HP, Cascon A, Sampietro ML, Gaal J, Korpershoek E, Hinojar-Gutierrez A, Timmers HJ, Hoefsloot LH, Hermesen MA, *et al.* 2010 SDHAF2 mutations in familial and sporadic paraganglioma and pheochromocytoma. *Lancet Oncology* **11** 366–372. ([https://doi.org/10.1016/S1470-2045\(10\)70007-3](https://doi.org/10.1016/S1470-2045(10)70007-3))
- Baysal BE, Ferrell RE, Willett-Brozick JE, Lawrence EC, Myssiorek D, Bosch A, van der Mey A, Taschner PE, Rubinstein WS, Myers EN, *et al.* 2000 Mutations in SDHD, a mitochondrial complex II gene, in hereditary paraganglioma. *Science* **287** 848–851. (<https://doi.org/10.1126/science.287.5454.848>)
- Beckers A, Aaltonen LA, Daly AF & Karhu A 2013 Familial isolated pituitary adenomas (FIPA) and the pituitary adenoma predisposition due to mutations in the aryl hydrocarbon receptor interacting protein (AIP) gene. *Endocrine Reviews* **34** 239–277. (<https://doi.org/10.1210/er.2012-1013>)
- Beckers A, Lodish MB, Trivellin G, Rostomyan L, Lee M, Faucz FR, Yuan B, Choong CS, Caberg JH, Verrua E, *et al.* 2015 X-linked acrogigantism syndrome: clinical profile and therapeutic responses. *Endocrine-Related Cancer* **22** 353–367. (<https://doi.org/10.1530/ERC-15-0038>)
- Beckers A, Fernandes D, Fina F, Novak M, Abati A, Rostomyan L, Thiry A, Ouafik L, Pasture B, Pinhasi R, *et al.* 2017 Paleogenetic study of ancient DNA suggestive of X-linked acrogigantism. *Endocrine-Related Cancer* **24** L17–L20. (<https://doi.org/10.1530/ERC-16-0558>)
- Belar O, De La Hoz C, Perez-Nanclares G, Castano L, Gaztambide S & Spanish MENG 2012 Novel mutations in MEN1, CDKN1B and AIP genes in patients with multiple endocrine neoplasia type 1 syndrome in Spain. *Clinical Endocrinology* **76** 719–724. (<https://doi.org/10.1111/j.1365-2265.2011.04269.x>)
- Benn DE, Gimenez-Roqueplo AP, Reilly JR, Bertherat J, Burgess J, Byth K, Croxson M, Dahia PL, Elston M, Gimm O, *et al.* 2006 Clinical presentation and penetrance of pheochromocytoma/paraganglioma syndromes. *Journal of Clinical Endocrinology and Metabolism* **91** 827–836. (<https://doi.org/10.1210/jc.2005-1862>)
- Bertherat J, Horvath A, Groussin L, Grabar S, Boikos S, Cazabat L, Libe R, Rene-Corail F, Stergiopoulos S, Bourdeau I, *et al.* 2009 Mutations in regulatory subunit type 1A of cyclic adenosine 5'-monophosphate-dependent protein kinase (PRKAR1A): phenotype analysis in 353 patients and 80 different genotypes. *Journal of Clinical Endocrinology and Metabolism* **94** 2085–2091. (<https://doi.org/10.1210/jc.2008-2333>)
- Biondi CA, Gartside MG, Waring P, Loffler KA, Stark MS, Magnuson MA, Kay GF & Hayward NK 2004 Conditional inactivation of the MEN1 gene leads to pancreatic and pituitary tumorigenesis but does not affect normal development of these tissues. *Molecular and Cellular Biology* **24** 3125–3131. (<https://doi.org/10.1128/MCB.24.8.3125-3131.2004>)
- Bizzi MF, Pinheiro SVB, Bolger GB, Schweizer J, Giannetti AV, Dang MN, Ribeiro-Oliveira A Jr & Korbonits M 2018 Reduced protein expression of the phosphodiesterases PDE4A4 and PDE4A8 in AIP mutation positive somatotroph adenomas. *Molecular and Cellular Endocrinology* **476** 103–109. (<https://doi.org/10.1016/j.mce.2018.04.014>)
- Boissan M, De Wever O, Lizarraga F, Wendum D, Poincloux R, Chignard N, Desbois-Mouthon C, Dufour S, Nawrocki-Raby B, Birembaut P, *et al.* 2010 Implication of metastasis suppressor NM23-H1 in maintaining adherens junctions and limiting the invasive potential of human cancer cells. *Cancer Research* **70** 7710–7722. (<https://doi.org/10.1158/0008-5472.CAN-10-1887>)
- Bolger GB, Peden AH, Steele MR, MacKenzie C, McEwan DG, Wallace DA, Huston E, Baillie GS & Houslay MD 2003 Attenuation of the activity of the cAMP-specific phosphodiesterase PDE4A5 by interaction with the immunophilin XAP2. *Journal of Biological Chemistry* **278** 33351–33363. (<https://doi.org/10.1074/jbc.M303269200>)
- Bolz H, von Brederlow B, Ramirez A, Bryda EC, Kutsche K, Nothwang HG, Seeliger M, del CSCM, Vila MC, Molina OP, *et al.* 2001 Mutation of CDH23, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. *Nature Genetics* **27** 108–112. (<https://doi.org/10.1038/83667>)
- Borsari S, Pardi E, Pellegata NS, Lee M, Saponaro F, Torregrossa L, Basolo F, Paltrinieri E, Zatelli MC, Materazzi G, *et al.* 2017 Loss of p27 expression is associated with MEN1 gene mutations in sporadic parathyroid adenomas. *Endocrine* **55** 386–397. (<https://doi.org/10.1007/s12020-016-0941-6>)
- Bossis I & Stratakis CA 2004 Minireview: PRKAR1A: normal and abnormal functions. *Endocrinology* **145** 5452–5458. (<https://doi.org/10.1210/en.2004-0900>)
- Bottoni A, Zatelli MC, Ferracin M, Tagliati F, Piccin D, Vignali C, Calin GA, Negrini M, Croce CM & Degli Uberti EC 2007 Identification of differentially expressed microRNAs by microarray: a possible role for microRNA genes in pituitary adenomas. *Journal of Cellular Physiology* **210** 370–377. (<https://doi.org/10.1002/jcp.20832>)
- Boyce AM & Collins MT 1993 Fibrous dysplasia/McCune–Albright syndrome. In *GeneReviews*. Eds MP Adam, HH Ardinger, RA Pagon, SE Wallace, LJH Bean, K Stephens & A Amemiya. Seattle, WA, USA: University of Washington. (available at: <https://www.ncbi.nlm.nih.gov/books/NBK274564/>)
- Brenneman M, Field A, Yang J, Williams G, Doros L, Rossi C, Schultz KA, Rosenberg A, Ivanovich J, Turner J, *et al.* 2015 Temporal order of RNase IIIb and loss-of-function mutations during development determines phenotype in pleuropulmonary blastoma/DICER1 syndrome: a unique variant of the two-hit tumor suppression model. *F1000Research* **4** 214. (<https://doi.org/10.12688/f1000research.6746.1>)
- Burnichon N, Briere JJ, Libe R, Vescovo L, Riviere J, Tissier F, Jouanno E, Jeunemaitre X, Benit P, Tzagoloff A, *et al.* 2010 SDHA is a tumor suppressor gene causing paraganglioma. *Human Molecular Genetics* **19** 3011–3020. (<https://doi.org/10.1093/hmg/ddq206>)
- Bussing I, Slack FJ & Grosshans H 2008 let-7 microRNAs in development, stem cells and cancer. *Trends in Molecular Medicine* **14** 400–409. (<https://doi.org/10.1016/j.molmed.2008.07.001>)
- Caimari F & Korbonits M 2016 Novel genetic causes of pituitary adenomas. *Clinical Cancer Research* **22** 5030–5042. (<https://doi.org/10.1158/1078-0432.CCR-16-0452>)
- Cannavo S, Ferrau F, Ragonese M, Curto L, Torre ML, Magistri M, Marchese A, Alibrandi A & Trimarchi F 2010 Increased prevalence of acromegaly in a highly polluted area. *European Journal of Endocrinology* **163** 509–513. (<https://doi.org/10.1530/EJE-10-0465>)
- Carney JA, Gordon H, Carpenter PC, Shenoy BV & Go VL 1985 The complex of myxomas, spotty pigmentation, and endocrine

- overactivity. *Medicine* **64** 270–283. (<https://doi.org/10.1097/00005792-198507000-00007>)
- Casey M, Vaughan CJ, He J, Hatcher CJ, Winter JM, Weremowicz S, Montgomery K, Kucherlapati R, Morton CC & Basson CT 2000 Mutations in the protein kinase A R1alpha regulatory subunit cause familial cardiac myxomas and Carney complex. *Journal of Clinical Investigation* **106** R31–R38. (<https://doi.org/10.1172/JCI10841>)
- Castinetti F, Daly AF, Stratakis CA, Caberg JH, Castermans E, Trivellin G, Rostomyan L, Saveanu A, Jullien N, Reynaud R, *et al.* 2016 GPR101 mutations are not a frequent cause of congenital isolated growth hormone deficiency. *Hormone and Metabolic Research* **48** 389–393. (<https://doi.org/10.1055/s-0042-100733>)
- Castro-Vega LJ, Letouze E, Burnichon N, Buffet A, Disderot PH, Khalifa E, Loriot C, Elarouci N, Morin A, Menara M, *et al.* 2015 Multi-omics analysis defines core genomic alterations in pheochromocytomas and paragangliomas. *Nature Communication* **6** 6044. (<https://doi.org/10.1038/ncomms7044>)
- Cazabat L, Bouligand J, Salenave S, Bernier M, Gaillard S, Parker F, Young J, Guiochon-Mantel A & Chanson P 2012 Germline AIP mutations in apparently sporadic pituitary adenomas: prevalence in a prospective single-center cohort of 443 patients. *Journal of Clinical Endocrinology and Metabolism* **97** E663–E670. (<https://doi.org/10.1210/jc.2011-2291>)
- Cervera AM, Apostolova N, Crespo FL, Mata M & McCreath KJ 2008 Cells silenced for SDHB expression display characteristic features of the tumor phenotype. *Cancer Research* **68** 4058–4067. (<https://doi.org/10.1158/0008-5472.CAN-07-5580>)
- Chahal HS, Stals K, Unterlander M, Balding DJ, Thomas MG, Kumar AV, Besser GM, Atkinson AB, Morrison PJ, Howlett TA, *et al.* 2011 AIP mutation in pituitary adenomas in the 18th century and today. *New England Journal of Medicine* **364** 43–50. (<https://doi.org/10.1056/NEJMoa1008020>)
- Chahal HS, Trivellin G, Leontiou CA, Albani N, Fowkes RC, Tahir A, Igreja SC, Chapple JP, Jordan S, Lupp A, *et al.* 2012 Somatostatin analogs modulate AIP in somatotroph adenomas: the role of the ZAC1 pathway. *Journal of Clinical Endocrinology and Metabolism* **97** E1411–E1420. (<https://doi.org/10.1210/jc.2012-1111>)
- Chandrasekharappa SC, Guru SC, Manickam P, Olufemi SE, Collins FS, Emmert-Buck MR, Debelenko LV, Zhuang Z, Lubensky LA, Liotta LA, *et al.* 1997 Positional cloning of the gene for multiple endocrine neoplasia-type 1. *Science* **276** 404–407. (<https://doi.org/10.1126/science.276.5311.404>)
- Chen W, Xiao Z, Zhao Y, Huang L & Du G 2013 HIF-1alpha inhibition sensitizes pituitary adenoma cells to temozolomide by regulating MGMT expression. *Oncology Reports* **30** 2495–2501. (<https://doi.org/10.3892/or.2013.2689>)
- Chu IM, Hengst L & Slingerland JM 2008 The Cdk inhibitor p27 in human cancer: prognostic potential and relevance to anticancer therapy. *Nature Reviews Cancer* **8** 253–267. (<https://doi.org/10.1038/nrc2347>)
- Cohen LE, Hashimoto Y, Zanger K, Wondisford F & Radovick S 1999 CREB-independent regulation by CBP is a novel mechanism of human growth hormone gene expression. *Journal of Clinical Investigation* **104** 1123–1130. (<https://doi.org/10.1172/JCI7308>)
- Comino-Mendez I, Gracia-Aznarez FJ, Schiavi F, Landa I, Leandro-Garcia LJ, Leton R, Honrado E, Ramos-Medina R, Caronia D, Pita G, *et al.* 2011 Exome sequencing identifies MAX mutations as a cause of hereditary pheochromocytoma. *Nature Genetics* **43** 663–667. (<https://doi.org/10.1038/ng.861>)
- Comino-Mendez I, de Cubas AA, Bernal C, Alvarez-Escola C, Sanchez-Malo C, Ramirez-Tortosa CL, Pedrinaci S, Rapizzi E, Ercolino T, Bernini G, *et al.* 2013 Tumoral EPAS1 (HIF2A) mutations explain sporadic pheochromocytoma and paraganglioma in the absence of erythrocytosis. *Human Molecular Genetics* **22** 2169–2176. (<https://doi.org/10.1093/hmg/ddt069>)
- Concolino P, Costella A & Capoluongo E 2016 Multiple endocrine neoplasia type 1 (MEN1): an update of 208 new germline variants reported in the last nine years. *Cancer Genetics* **209** 36–41. (<https://doi.org/10.1016/j.cancergen.2015.12.002>)
- Correa R, Salpea P & Stratakis CA 2015 Carney complex: an update. *European Journal of Endocrinology* **173** M85–M97. (<https://doi.org/10.1530/EJE-15-0209>)
- Costa-Guda J, Marinoni I, Molatore S, Pellegata NS & Arnold A 2011 Somatic mutation and germline sequence abnormalities in CDKN1B, encoding p27Kip1, in sporadic parathyroid adenomas. *Journal of Clinical Endocrinology and Metabolism* **96** E701–E706. (<https://doi.org/10.1210/jc.2010-1338>)
- Courcousakis NA, Tatsi C, Patronas NJ, Lee CC, Prassopoulos PK & Stratakis CA 2013 The complex of myxomas, spotty skin pigmentation and endocrine overactivity (Carney complex): imaging findings with clinical and pathological correlation. *Insights into Imaging* **4** 119–133. (<https://doi.org/10.1007/s13244-012-0208-6>)
- Crabtree JS, Scacheri PC, Ward JM, Garrett-Beal L, Emmert-Buck MR, Edgemon KA, Lorang D, Libutti SK, Chandrasekharappa SC, Marx SJ, *et al.* 2001 A mouse model of multiple endocrine neoplasia, type 1, develops multiple endocrine tumors. *PNAS* **98** 1118–1123. (<https://doi.org/10.1073/pnas.98.3.1118>)
- Cuny T, Pertuit M, Sahnoun-Fathallah M, Daly A, Occhi G, Odou MF, Tabarin A, Nunes ML, Delemer B, Rohmer V, *et al.* 2013 Genetic analysis in young patients with sporadic pituitary macroadenomas: besides AIP don't forget MEN1 genetic analysis. *European Journal of Endocrinology* **168** 533–541. (<https://doi.org/10.1530/EJE-12-0763>)
- Daly AF, Rixhon M, Adam C, Dempegioti A, Tichomirowa MA & Beckers A 2006a High prevalence of pituitary adenomas: a cross-sectional study in the province of Liege, Belgium. *Journal of Clinical Endocrinology and Metabolism* **91** 4769–4775. (<https://doi.org/10.1210/jc.2006-1668>)
- Daly AF, Jaffrain-Rea ML, Ciccarella A, Valdes-Socin H, Rohmer V, Tamburrano G, Borson-Chazot C, Estour B, Ciccarella E, Brue T, *et al.* 2006b Clinical characterization of familial isolated pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* **91** 3316–3323. (<https://doi.org/10.1210/jc.2005-2671>)
- Daly AF, Vanbellinghen JF, Khoo SK, Jaffrain-Rea ML, Naves LA, Guitelman MA, Murat A, Emy P, Gimenez-Roqueplo AP, Tamburrano G, *et al.* 2007 Aryl hydrocarbon receptor-interacting protein gene mutations in familial isolated pituitary adenomas: analysis in 73 families. *Journal of Clinical Endocrinology and Metabolism* **92** 1891–1896. (<https://doi.org/10.1210/jc.2006-2513>)
- Daly AF, Tichomirowa MA & Beckers A 2009 The epidemiology and genetics of pituitary adenomas. *Best Practice and Research Clinical Endocrinology and Metabolism* **23** 543–554. (<https://doi.org/10.1016/j.beem.2009.05.008>)
- Daly AF, Tichomirowa MA, Petrossians P, Heliövaara E, Jaffrain-Rea ML, Barlier A, Naves LA, Ebeling T, Karhu A, Raappana A, *et al.* 2010 Clinical characteristics and therapeutic responses in patients with germ-line AIP mutations and pituitary adenomas: an international collaborative study. *Journal of Clinical Endocrinology and Metabolism* **95** E373–E383. (<https://doi.org/10.1210/jc.2009-2556>)
- Daly AF, Lysy PA, Desfilles C, Rostomyan L, Mohamed A, Caberg JH, Raverot V, Castermans E, Marbaix E, Maiter D, *et al.* 2016a GHRH excess and blockade in X-LAG syndrome. *Endocrine-Related Cancer* **23** 161–170. (<https://doi.org/10.1530/ERC-15-0478>)
- Daly AF, Yuan B, Fina F, Caberg JH, Trivellin G, Rostomyan L, de Herder WW, Naves LA, Metzger D, Cuny T, *et al.* 2016b Somatic mosaicism underlies X-linked acrogerism syndrome in sporadic male subjects. *Endocrine-Related Cancer* **23** 221–233. (<https://doi.org/10.1530/ERC-16-0082>)
- Daly AF, Castermans E, Oudijk L, Guitelman MA, Beckers P, Poterac I, Neggers S, Sacre N, van der Lely AJ, Bours V, *et al.* 2018 Pheochromocytomas and pituitary adenomas in three patients with MAX exon deletions. *Endocrine-Related Cancer* **25** L37–L42. (<https://doi.org/10.1530/ERC-18-0065>)
- de Kock L, Sabbaghian N, Plourde F, Srivastava A, Weber E, Bouron-Dal Soglio D, Hamel N, Choi JH, Park SH, Deal CL, *et al.* 2014 Pituitary

- blastoma: a pathognomonic feature of germ-line DICER1 mutations. *Acta Neuropathologica* **128** 111–122. (<https://doi.org/10.1007/s00401-014-1285-z>)
- de Kock L, Wang YC, Revil T, Badescu D, Rivera B, Sabbaghian N, Wu M, Weber E, Sandoval C, Hopman SM, *et al.* 2016 High-sensitivity sequencing reveals multi-organ somatic mosaicism causing DICER1 syndrome. *Journal of Medical Genetics* **53** 43–52. (<https://doi.org/10.1136/jmedgenet-2015-103428>)
- de Laat JM, Dekkers OM, Pieterman CR, Kluijfhout WP, Hermus AR, Pereira AM, van der Horst-Schrivers AN, Drent ML, Bisschop PH, Havekes B, *et al.* 2015 Long-term natural course of pituitary tumors in patients with MEN1: results from the DutchMEN1 Study Group (DMSG). *Journal of Clinical Endocrinology and Metabolism* **100** 3288–3296. (<https://doi.org/10.1210/JC.2015-2015>)
- Denes J, Swords F, Rattenberry E, Stals K, Owens M, Cranston T, Xekouki P, Moran L, Kumar A, Wassif C, *et al.* 2015 Heterogeneous genetic background of the association of pheochromocytoma/paraganglioma and pituitary adenoma: results from a large patient cohort. *Journal of Clinical Endocrinology and Metabolism* **100** E531–E541. (<https://doi.org/10.1210/jc.2014-3399>)
- Devnath S & Inoue K 2008 An insight to pituitary folliculo-stellate cells. *Journal of Neuroendocrinology* **20** 687–691. (<https://doi.org/10.1111/j.1365-2826.2008.01716.x>)
- Dong Q, Debelenko LV, Chandrasekharappa SC, Emmert-Buck MR, Zhuang Z, Guru SC, Manickam P, Skarulis M, Lubensky IA, Liotta LA, *et al.* 1997 Loss of heterozygosity at 11q13: analysis of pituitary tumors, lung carcinoids, lipomas, and other uncommon tumors in subjects with familial multiple endocrine neoplasia type 1. *Journal of Clinical Endocrinology and Metabolism* **82** 1416–1420. (<https://doi.org/10.1210/jcem.82.5.3944>)
- Doros L, Schultz KA, Stewart DR, Bauer AJ, Williams G, Rossi CT, Carr A, Yang J, Dehner LP, Messinger Y, *et al.* 1993 DICER1-related disorders. In *GeneReviews*. Eds MP Adam, HH Ardinger, RA Pagon, SE Wallace, LJH Bean, K Stephens & A Amemiya. Seattle, WA, USA: University of Washington. (available at: <https://www.ncbi.nlm.nih.gov/books/NBK196157/>)
- Dreijerink KM, Goudet P, Burgess JR, Valk GD & International Breast Cancer in MENSG 2014 Breast-cancer predisposition in multiple endocrine neoplasia type 1. *New England Journal of Medicine* **371** 583–584. (<https://doi.org/10.1056/NEJMc1406028>)
- Dwight T, Mann K, Benn DE, Robinson BG, McKelvie P, Gill AJ, Winship I & Clifton-Bligh RJ 2013 Familial SDHA mutation associated with pituitary adenoma and pheochromocytoma/paraganglioma. *Journal of Clinical Endocrinology and Metabolism* **98** E1103–E1108. (<https://doi.org/10.1210/jc.2013-1400>)
- Elston MS, Meyer-Rochow GY, Dray M, Swarbrick M & Conaglen JV 2015 Early onset primary hyperparathyroidism associated with a novel germline mutation in CDKN1B. *Case Reports in Endocrinology* **2015** 510985. (<https://doi.org/10.1155/2015/510985>)
- Esteller M, Guo M, Moreno V, Peinado MA, Capella G, Galm O, Baylin SB & Herman JG 2002 Hypermethylation-associated inactivation of the cellular retinol-binding-protein 1 gene in human cancer. *Cancer Research* **62** 5902–5905.
- Ezzat S, Asa SL, Couldwell WT, Barr CE, Dodge WE, Vance ML & McCutcheon IE 2004 The prevalence of pituitary adenomas: a systematic review. *Cancer* **101** 613–619. (<https://doi.org/10.1002/cncr.20412>)
- Fedeles M, Battista S, Kenyon L, Baldassarre G, Fidanza V, Klein-Szanto AJ, Parlow AF, Visone R, Pierantoni GM, Outwater E, *et al.* 2002 Overexpression of the HMGA2 gene in transgenic mice leads to the onset of pituitary adenomas. *Oncogene* **21** 3190–3198. (<https://doi.org/10.1038/sj.onc.1205428>)
- Fernandez A, Karavitaki N & Wass JA 2010 Prevalence of pituitary adenomas: a community-based, cross-sectional study in Banbury (Oxfordshire, UK). *Clinical Endocrinology* **72** 377–382. (<https://doi.org/10.1111/j.1365-2265.2009.03667.x>)
- Fero ML, Randel E, Gurley KE, Roberts JM & Kemp CJ 1998 The murine gene p27Kip1 is haplo-insufficient for tumour suppression. *Nature* **396** 177–180. (<https://doi.org/10.1038/24179>)
- Ferrau F, Romeo PD, Puglisi S, Ragonese M, Torre ML, Scaroni C, Occhi G, De Menis E, Arnaldi G, Trimarchi F, *et al.* 2016 Analysis of GPR101 and AIP genes mutations in acromegaly: a multicentric study. *Endocrine* **54** 762–767. (<https://doi.org/10.1007/s12020-016-0862-4>)
- Finelli P, Pierantoni GM, Giardino D, Losa M, Rodeschini O, Fedeles M, Valtorta E, Mortini P, Croce CM, Larizza L, *et al.* 2002 The high mobility group A2 gene is amplified and overexpressed in human prolactinomas. *Cancer Research* **62** 2398–2405.
- Fontana E & Gaillard R 2009 Epidemiology of pituitary adenoma: results of the first Swiss study. *Revue Médicale Suisse* **5** 2172–2174.
- Forlino A, Vetro A, Garavelli L, Ciccone R, London E, Stratakis CA & Zuffardi O 2014 PRKACB and Carney complex. *New England Journal of Medicine* **370** 1065–1067. (<https://doi.org/10.1056/NEJMc1309730>)
- Formosa R & Vassallo J 2014 cAMP signalling in the normal and tumorigenic pituitary gland. *Molecular and Cellular Endocrinology* **392** 37–50. (<https://doi.org/10.1016/j.mce.2014.05.004>)
- Formosa R, Xuereb-Anastasi A & Vassallo J 2013 Aip regulates cAMP signalling and GH secretion in GH3 cells. *Endocrine-Related Cancer* **20** 495–505. (<https://doi.org/10.1530/ERC-13-0043>)
- Fossberg TM, Doskeland SO & Ueland PM 1978 Protein kinases in human renal cell carcinoma and renal cortex. A comparison of isozyme distribution and of responsiveness to adenosine 3':5'-cyclic monophosphate. *Archives of Biochemistry and Biophysics* **189** 272–281.
- Franssen D & Tena-Sempere M 2018 The kisspeptin receptor: a key G-protein-coupled receptor in the control of the reproductive axis. *Best Practice and Research Clinical Endocrinology and Metabolism* **32** 107–123. (<https://doi.org/10.1016/j.beem.2018.01.005>)
- Freda PU, Beckers AM, Katznelson L, Molitch ME, Montori VM, Post KD, Vance ML & Endocrine Society 2011 Pituitary incidentaloma: an endocrine society clinical practice guideline. *Journal of Clinical Endocrinology and Metabolism* **96** 894–904. (<https://doi.org/10.1210/jc.2010-1048>)
- Gadelha MR, Prezant TR, Une KN, Glick RP, Moskal SF, 2nd, Vaisman M, Melmed S, Kineman RD & Frohman LA 1999 Loss of heterozygosity on chromosome 11q13 in two families with acromegaly/gigantism is independent of mutations of the multiple endocrine neoplasia type I gene. *Journal of Clinical Endocrinology and Metabolism* **84** 249–256. (<https://doi.org/10.1210/jcem.84.1.5370>)
- Galland F, Kamenicky P, Affres H, Reznik Y, Pontvert D, Le Bouc Y, Young J & Chanson P 2006 McCune-Albright syndrome and acromegaly: effects of hypothalamic pituitary radiotherapy and/or pegvisomant in somatostatin analog-resistant patients. *Journal of Clinical Endocrinology and Metabolism* **91** 4957–4961. (<https://doi.org/10.1210/jc.2006-0561>)
- Gamm DM, Baude EJ & Uhler MD 1996 The major catalytic subunit isoforms of cAMP-dependent protein kinase have distinct biochemical properties in vitro and in vivo. *Journal of Biological Chemistry* **271** 15736–15742. (<https://doi.org/10.1074/jbc.271.26.15736>)
- Georgitsi M, Raitila A, Karhu A, van der Luijt RB, Aalfs CM, Sane T, Vierimaa O, Mäkinen MJ, Tuppurainen K, Paschke R, *et al.* 2007 Germline CDKN1B/p27Kip1 mutation in multiple endocrine neoplasia. *Journal of Clinical Endocrinology and Metabolism* **92** 3321–3325. (<https://doi.org/10.1210/jc.2006-2843>)
- Georgitsi M, Heliovaara E, Paschke R, Kumar AV, Tischkowitz M, Vierimaa O, Salmela P, Sane T, De Menis E, Cannavo S, *et al.* 2008 Large genomic deletions in AIP in pituitary adenoma predisposition. *Journal of Clinical Endocrinology and Metabolism* **93** 4146–4151. (<https://doi.org/10.1210/jc.2008-1003>)
- Gill AJ, Toon CW, Clarkson A, Sioson L, Chou A, Winship I, Robinson BG, Benn DE, Clifton-Bligh RJ & Dwight T 2014 Succinate dehydrogenase deficiency is rare in pituitary adenomas. *American Journal of Surgical Pathology* **38** 560–566. (<https://doi.org/10.1097/PAS.0000000000000149>)

- Gillam MP, Nimbalkar D, Sun L, Christov K, Ray D, Kaldis P, Liu X & Kiyokawa H 2015 MEN1 tumorigenesis in the pituitary and pancreatic islet requires Cdk4 but not Cdk2. *Oncogene* **34** 932–938. (<https://doi.org/10.1038/onc.2014.3>)
- Gillam MP, Ku CR, Lee YJ, Kim J, Kim SH, Lee SJ, Hwang B, Koo J, Kineman RD, Kiyokawa H, *et al.* 2017 Somatotroph-specific aip-deficient mice display pretumorigenic alterations in cell-cycle signaling. *Journal of the Endocrine Society* **1** 78–95. (<https://doi.org/10.1210/js.2016-1004>)
- Glasker S, Vortmeyer AO, Lafferty AR, Hofman PL, Li J, Weil RJ, Zhuang Z & Oldfield EH 2011 Hereditary pituitary hyperplasia with infantile gigantism. *Journal of Clinical Endocrinology and Metabolism* **96** E2078–E2087. (<https://doi.org/10.1210/jc.2011-1401>)
- Gordon RJ, Bell J, Chung WK, David R, Oberfield SE & Wardlaw SL 2016 Childhood acromegaly due to X-linked acrogigantism: long term follow-up. *Pituitary* **19** 560–564. (<https://doi.org/10.1007/s11102-016-0743-0>)
- Groussin L, Kirschner LS, Vincent-Dejean C, Perlemoine K, Jullian E, Delemer B, Zacharieva S, Pignatelli D, Carney JA, Luton JP, *et al.* 2002 Molecular analysis of the cyclic AMP-dependent protein kinase A (PKA) regulatory subunit 1A (PRKAR1A) gene in patients with Carney complex and primary pigmented nodular adrenocortical disease (PPNAD) reveals novel mutations and clues for pathophysiology: augmented PKA signaling is associated with adrenal tumorigenesis in PPNAD. *American Journal of Human Genetics* **71** 1433–1442. (<https://doi.org/10.1086/344579>)
- Gruppetta M, Mercieca C & Vassallo J 2013 Prevalence and incidence of pituitary adenomas: a population based study in Malta. *Pituitary* **16** 545–553. (<https://doi.org/10.1007/s11102-012-0454-0>)
- Gurtan AM, Lu V, Bhutkar A & Sharp PA 2012 In vivo structure-function analysis of human Dicer reveals directional processing of precursor miRNAs. *RNA* **18** 1116–1122. (<https://doi.org/10.1261/rna.032680.112>)
- Guzy RD, Sharma B, Bell E, Chandel NS & Schumacker PT 2008 Loss of the SdhB, but not the SdhA, subunit of complex II triggers reactive oxygen species-dependent hypoxia-inducible factor activation and tumorigenesis. *Molecular and Cellular Biology* **28** 718–731. (<https://doi.org/10.1128/MCB.01338-07>)
- Handschin JC & Eppenberger U 1979 Altered cellular ratio of type I and type II cyclic AMP-dependent protein kinase in human mammary tumors. *FEBS Letters* **106** 301–304. ([https://doi.org/10.1016/0014-5793\(79\)80519-0](https://doi.org/10.1016/0014-5793(79)80519-0))
- Hao HX, Khalimonchuk O, Schraders M, Dephoure N, Bayley JP, Kunst H, Devilee P, Cremers CW, Schiffman JD, Bentz BG, *et al.* 2009 SDH5, a gene required for flavination of succinate dehydrogenase, is mutated in paraganglioma. *Science* **325** 1139–1142. (<https://doi.org/10.1126/science.1175689>)
- Hayward BE, Barlier A, Korbonits M, Grossman AB, Jacquet P, Enjalbert A & Bonthron DT 2001 Imprinting of the G(s)alpha gene GNAS1 in the pathogenesis of acromegaly. *Journal of Clinical Investigation* **107** R31–R36. (<https://doi.org/10.1172/JCI11887>)
- Heravi-Moussavi A, Anglesio MS, Cheng SW, Senz J, Yang W, Prentice L, Fejes AP, Chow C, Tone A, Kalloger SE, *et al.* 2012 Recurrent somatic DICER1 mutations in nonepithelial ovarian cancers. *New England Journal of Medicine* **366** 234–242. (<https://doi.org/10.1056/NEJMoa1102903>)
- Herman V, Fagin J, Gonsky R, Kovacs K & Melmed S 1990 Clonal origin of pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* **71** 1427–1433. (<https://doi.org/10.1210/jcem-71-6-1427>)
- Hernandez-Ramirez LC, Gabrovskaja P, Denes J, Stals K, Trivellin G, Tilley D, Ferrau F, Evanson J, Ellard S, Grossman AB, *et al.* 2015 Landscape of familial isolated and young-onset pituitary adenomas: prospective diagnosis in AIP mutation carriers. *Journal of Clinical Endocrinology and Metabolism* **100** E1242–E1254. (<https://doi.org/10.1210/jc.2015-1869>)
- Hernandez-Ramirez LC, Martucci F, Morgan RM, Trivellin G, Tilley D, Ramos-Guajardo N, Iacovazzo D, D'Acquisto F, Prodromou C & Korbonits M 2016 Rapid proteasomal degradation of mutant proteins is the primary mechanism leading to tumorigenesis in patients with missense AIP mutations. *Journal of Clinical Endocrinology and Metabolism* **101** 3144–3154. (<https://doi.org/10.1210/jc.2016-1307>)
- Hernandez-Ramirez LC, Gam R, Valdes N, Lodish MB, Pankratz N, Balsalobre A, Gauthier Y, Fauch FR, Trivellin G, Chittiboina P, *et al.* 2017a Loss-of-function mutations in the CABLES1 gene are a novel cause of Cushing's disease. *Endocrine-Related Cancer* **24** 379–392. (<https://doi.org/10.1530/ERC-17-0131>)
- Hernandez-Ramirez LC, Tatsi C, Lodish MB, Fauch FR, Pankratz N, Chittiboina P, Lane J, Kay DM, Valdes N, Dimopoulos A, *et al.* 2017b Corticotropinoma as a component of carney complex. *Journal of the Endocrine Society* **1** 918–925. (<https://doi.org/10.1210/js.2017-00231>)
- Hernandez-Ramirez LC, Morgan RML, Barry S, D'Acquisto F, Prodromou C & Korbonits M 2018 Multi-chaperone function modulation and association with cytoskeletal proteins are key features of the function of AIP in the pituitary gland. *Oncotarget* **9** 9177–9198. (<https://doi.org/10.18632/oncotarget.24183>)
- Horikawa R, Hellmann P, Cella SG, Torsello A, Day RN, Muller EE & Thorner MO 1996 Growth hormone-releasing factor (GRF) regulates expression of its own receptor. *Endocrinology* **137** 2642–2645. (<https://doi.org/10.1210/endo.137.6.8641220>)
- Horiuchi K, Okamoto T, Iihara M & Tsukada T 2013 Analysis of genotype-phenotype correlations and survival outcomes in patients with primary hyperparathyroidism caused by multiple endocrine neoplasia type 1: the experience at a single institution. *Surgery Today* **43** 894–899. (<https://doi.org/10.1007/s00595-012-0354-y>)
- Horvath A, Bossis I, Giatzakis C, Levine E, Weinberg F, Meoli E, Robinson-White A, Siegel J, Soni P, Groussin L, *et al.* 2008 Large deletions of the PRKAR1A gene in Carney complex. *Clinical Cancer Research* **14** 388–395. (<https://doi.org/10.1158/1078-0432.CCR-07-1155>)
- Iacovazzo D & Korbonits M 2018 X-linked acrogigantism. In *GeneReviews*. Eds MP Adam, HH Ardinger, RA Pagon, SE Wallace, LJH Bean, K Stephens & A Amemiya. Seattle, WA, USA: University of Washington. (available at: <https://www.ncbi.nlm.nih.gov/books/NBK476671/>)
- Iacovazzo D, Carlsen E, Lugli F, Chiloiro S, Piacentini S, Bianchi A, Giampietro A, Mormando M, Clear AJ, Doglietto F, *et al.* 2016a Factors predicting pasireotide responsiveness in somatotroph pituitary adenomas resistant to first-generation somatostatin analogues: an immunohistochemical study. *European Journal of Endocrinology* **174** 241–250. (<https://doi.org/10.1530/EJE-15-0832>)
- Iacovazzo D, Caswell R, Bunce B, Jose S, Yuan B, Hernandez-Ramirez LC, Kapur S, Caimari F, Evanson J, Ferrau F, *et al.* 2016b Germline or somatic GPR101 duplication leads to X-linked acrogigantism: a clinico-pathological and genetic study. *Acta Neuropathologica Communications* **4** 56. (<https://doi.org/10.1186/s40478-016-0328-1>)
- Igreja S, Chahal HS, King P, Bolger GB, Srirangalingam U, Guasti L, Chapple JP, Trivellin G, Gueorguiev M, Guegan K, *et al.* 2010 Characterization of aryl hydrocarbon receptor interacting protein (AIP) mutations in familial isolated pituitary adenoma families. *Human Mutation* **31** 950–960. (<https://doi.org/10.1002/humu.21292>)
- Ishii T, Yasuda K, Akatsuka A, Hino O, Hartman PS & Ishii N 2005 A mutation in the SDHC gene of complex II increases oxidative stress, resulting in apoptosis and tumorigenesis. *Cancer Research* **65** 203–209.
- Iversen K 1952 Acromegaly associated with pheochromocytoma. *Acta Medica Scandinavica* **142** 1–5. (<https://doi.org/10.1111/j.0954-6820.1952.tb13837.x>)
- Jacks T, Fazeli A, Schmitt EM, Bronson RT, Goodell MA & Weinberg RA 1992 Effects of an Rb mutation in the mouse. *Nature* **359** 295–300. (<https://doi.org/10.1038/359295a0>)
- Kamenicky P, Bouligand J & Chanson P 2015 Gigantism, acromegaly, and GPR101 mutations. *New England Journal of Medicine* **372** 1264. (<https://doi.org/10.1056/NEJMc1500340>)
- Kasuki L, Vieira Neto L, Wildemberg LE, Colli LM, de Castro M, Takiya CM & Gadelha MR 2012 AIP expression in sporadic somatotropinomas is a predictor of the response to octreotide LAR

- therapy independent of SSTR2 expression. *Endocrine-Related Cancer* **19** L25–L29. (<https://doi.org/10.1530/ERC-12-0020>)
- Kiefer FW, Winhofer Y, Iacovazzo D, Korbonits M, Wolfsberger S, Knosp E, Trautinger F, Hoftberger R, Krebs M, Luger A, *et al.* 2017 PRKAR1A mutation causing pituitary-dependent Cushing disease in a patient with Carney complex. *European Journal of Endocrinology* **177** K7–K12. (<https://doi.org/10.1530/EJE-17-0227>)
- Killian JK, Kim SY, Miettinen M, Smith C, Merino M, Tsokos M, Quezado M, Smith WI Jr, Jahromi MS, Xekouki P, *et al.* 2013 Succinate dehydrogenase mutation underlies global epigenomic divergence in gastrointestinal stromal tumor. *Cancer Discovery* **3** 648–657. (<https://doi.org/10.1158/2159-8290.CD-13-0092>)
- Kim H, Lee JE, Cho EJ, Liu JO & Youn HD 2003 Menin, a tumor suppressor, represses JunD-mediated transcriptional activity by association with an mSin3A-histone deacetylase complex. *Cancer Research* **63** 6135–6139.
- Kirley SD, Rueda BR, Chung DC & Zukerberg LR 2005 Increased growth rate, delayed senescence and decreased serum dependence characterize cables-deficient cells. *Cancer Biology and Therapy* **4** 654–658. (<https://doi.org/10.4161/cbt.4.6.1732>)
- Kirschner LS, Carney JA, Pack SD, Taymans SE, Giatzakis C, Cho YS, Cho-Chung YS & Stratakis CA 2000a Mutations of the gene encoding the protein kinase A type I- α regulatory subunit in patients with the Carney complex. *Nature Genetics* **26** 89–92. (<https://doi.org/10.1038/79238>)
- Kirschner LS, Sandrini F, Monbo J, Lin JP, Carney JA & Stratakis CA 2000b Genetic heterogeneity and spectrum of mutations of the PRKAR1A gene in patients with the carney complex. *Human Molecular Genetics* **9** 3037–3046. (<https://doi.org/10.1093/hmg/9.20.3037>)
- Kirschner LS, Kusewitt DF, Matyakhina L, Towns WH 2nd, Carney JA, Westphal H & Stratakis CA 2005 A mouse model for the Carney complex tumor syndrome develops neoplasia in cyclic AMP-responsive tissues. *Cancer Research* **65** 4506–4514. (<https://doi.org/10.1158/0008-5472.CAN-05-0580>)
- Kiyokawa H, Kineman RD, Manova-Todorova KO, Soares VC, Hoffman ES, Ono M, Khanam D, Hayday AC, Frohman LA & Koff A 1996 Enhanced growth of mice lacking the cyclin-dependent kinase inhibitor function of p27(Kip1). *Cell* **85** 721–732. ([https://doi.org/10.1016/S0092-8674\(00\)81238-6](https://doi.org/10.1016/S0092-8674(00)81238-6))
- Kobza AO, Dizon S & Arnaout A 2018 Case report of bilateral pheochromocytomas due to a novel MAX mutation in a patient known to have a pituitary prolactinoma. *AACE Clinical Case Reports*.
- Kouvaraki MA, Lee JE, Shapiro SE, Gagel RF, Sherman SI, Sellin RV, Cote GJ & Evans DB 2002 Genotype-phenotype analysis in multiple endocrine neoplasia type 1. *Archives of Surgery* **137** 641–647. (<https://doi.org/10.1001/archsurg.137.6.641>)
- Krol J, Loedige I & Filipowicz W 2010 The widespread regulation of microRNA biogenesis, function and decay. *Nature Reviews Genetics* **11** 597–610. (<https://doi.org/10.1038/nrg2843>)
- Larsson C, Skogseid B, Oberg K, Nakamura Y & Nordenskjold M 1988 Multiple endocrine neoplasia type 1 gene maps to chromosome 11 and is lost in insulinoma. *Nature* **332** 85–87. (<https://doi.org/10.1038/332085a0>)
- Le Quesne Stabej P, Saihan Z, Rangesh N, Steele-Stallard HB, Ambrose J, Coffey A, Emmerson J, Haralambous E, Hughes Y, Steel KP, *et al.* 2012 Comprehensive sequence analysis of nine Usher syndrome genes in the UK National Collaborative Usher Study. *Journal of Medical Genetics* **49** 27–36. (<https://doi.org/10.1136/jmedgenet-2011-100468>)
- Lecoq AL, Bouligand J, Hage M, Cazabat L, Salenave S, Linglart A, Young J, Guiochon-Mantel A, Chanson P & Kamenicky P 2016a Very low frequency of germline GPR101 genetic variation and no biallelic defects with AIP in a large cohort of patients with sporadic pituitary adenomas. *European Journal of Endocrinology* **174** 523–530. (<https://doi.org/10.1530/EJE-15-1044>)
- Lecoq AL, Zizzari P, Hage M, Decourtay L, Adam C, Viengchareun S, Veldhuis JD, Geoffroy V, Lombes M, Tolle V, *et al.* 2016b Mild pituitary phenotype in 3- and 12-month-old Aip-deficient male mice. *Journal of Endocrinology* **231** 59–69. (<https://doi.org/10.1530/JOE-16-0190>)
- Lemos MC & Thakker RV 2008 Multiple endocrine neoplasia type 1 (MEN1): analysis of 1336 mutations reported in the first decade following identification of the gene. *Human Mutation* **29** 22–32. (<https://doi.org/10.1002/humu.20605>)
- Leontiou CA, Gueorguiev M, van der Spuy J, Quinton R, Lolli F, Hassan S, Chahal HS, Igreja SC, Jordan S, Rowe J, *et al.* 2008 The role of the aryl hydrocarbon receptor-interacting protein gene in familial and sporadic pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* **93** 2390–2401. (<https://doi.org/10.1210/jc.2007-2611>)
- Letouze E, Martinelli C, Lorient C, Burnichon N, Abernill N, Ottolenghi C, Janin M, Menara M, Nguyen AT, Benit P, *et al.* 2013 SDH mutations establish a hypermethylator phenotype in paraganglioma. *Cancer Cell* **23** 739–752. (<https://doi.org/10.1016/j.ccr.2013.04.018>)
- Lidhar K, Korbonits M, Jordan S, Khalimova Z, Kaltsas G, Lu X, Clayton RN, Jenkins PJ, Monson JP, Besser GM, *et al.* 1999 Low expression of the cell cycle inhibitor p27Kip1 in normal corticotroph cells, corticotroph tumors, and malignant pituitary tumors. *Journal of Clinical Endocrinology and Metabolism* **84** 3823–3830. (<https://doi.org/10.1210/jcem.84.10.6066>)
- Lin BC, Sullivan R, Lee Y, Moran S, Glover E & Bradfield CA 2007 Deletion of the aryl hydrocarbon receptor-associated protein 9 leads to cardiac malformation and embryonic lethality. *Journal of Biological Chemistry* **282** 35924–35932. (<https://doi.org/10.1074/jbc.M705471200>)
- Lorient C, Domingues M, Berger A, Menara M, Ruel M, Morin A, Castro-Vega LJ, Letouze E, Martinelli C, Bemelmans AP, *et al.* 2015 Deciphering the molecular basis of invasiveness in Sdhb-deficient cells. *Oncotarget* **6** 32955–32965. (<https://doi.org/10.18632/oncotarget.5106>)
- Lumbroso S, Paris F, Sultan C & European Collaborative Study 2004 Activating Gsalpha mutations: analysis of 113 patients with signs of McCune-Albright syndrome – a European Collaborative Study. *Journal of Clinical Endocrinology and Metabolism* **89** 2107–2113. (<https://doi.org/10.1210/jc.2003-031225>)
- Ma Q & Whitlock JP Jr 1997 A novel cytoplasmic protein that interacts with the Ah receptor, contains tetrapeptide repeat motifs, and augments the transcriptional response to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Journal of Biological Chemistry* **272** 8878–8884. (<https://doi.org/10.1074/jbc.272.14.8878>)
- Ma ZY, Song ZJ, Chen JH, Wang YF, Li SQ, Zhou LF, Mao Y, Li YM, Hu RG, Zhang ZY, *et al.* 2015 Recurrent gain-of-function USP8 mutations in Cushing's disease. *Cell Research* **25** 306–317. (<https://doi.org/10.1038/cr.2015.20>)
- Maier M, Roncaroli F, Mendoza N, Meeran K, Canham N, Kosicka-Slawinska M, Bernhard B, Collier D, Drummond J, Skordilis K, *et al.* 2018 A patient with a germline SDHB mutation presenting with an isolated pituitary macroprolactinoma. *Endocrinology, Diabetes and Metabolism Case Reports* **2018** 18-0078. (<https://doi.org/10.1530/EDM-18-0078>)
- Mantovani G, Bondioni S, Lania AG, Corbetta S, de Sanctis L, Cappa M, Di Battista E, Chanson P, Beck-Peccoz P & Spada A 2004 Parental origin of Gsalpha mutations in the McCune-Albright syndrome and in isolated endocrine tumors. *Journal of Clinical Endocrinology and Metabolism* **89** 3007–3009. (<https://doi.org/10.1210/jc.2004-0194>)
- Marini F, Falchetti A, Del Monte F, Carbonell Sala S, Gozzini A, Luzi E & Brandi ML 2006 Multiple endocrine neoplasia type 1. *Orphanet Journal of Rare Diseases* **1** 38. (<https://doi.org/10.1186/1750-1172-1-38>)
- Marques P & Korbonits M 2017 Genetic aspects of pituitary adenomas. *Endocrinology and Metabolism Clinics of North America* **46** 335–374. (<https://doi.org/10.1016/j.eccl.2017.01.004>)
- Marques NV, Kasuki L, Coelho MC, Lima CHA, Wildemberg LE & Gadelha MR 2017 Frequency of familial pituitary adenoma syndromes among patients with functioning pituitary adenomas in a reference outpatient clinic. *Journal of Endocrinological Investigation* **40** 1381–1387. (<https://doi.org/10.1007/s40618-017-0725-8>)

- Martins CS, Camargo RC, Saggioro FP, Neder L, Machado HR, Moreira AC & de Castro M 2016 P27/CDKN1B translational regulators in pituitary tumorigenesis. *Hormone and Metabolic Research* **48** 840–846. (<https://doi.org/10.1055/s-0042-118613>)
- Maxwell JE, Sherman SK & Howe JR 2016 Translational diagnostics and therapeutics in pancreatic neuroendocrine tumors. *Clinical Cancer Research* **22** 5022–5029. (<https://doi.org/10.1158/1078-0432.CCR-16-0435>)
- McKnight GS, Clegg CH, Uhler MD, Chrvia JC, Cadd GG, Correll LA & Otten AD 1988 Analysis of the cAMP-dependent protein kinase system using molecular genetic approaches. *Recent Progress in Hormone Research* **44** 307–335.
- Mendoza-Rodriguez M, Arreola H, Valdivia A, Peralta R, Serna H, Villegas V, Romero P, Alvarado-Hernandez B, Paniagua L, Marrero-Rodriguez D, *et al.* 2013 Cellular retinol binding protein 1 could be a tumor suppressor gene in cervical cancer. *International Journal of Clinical and Experimental Pathology* **6** 1817–1825.
- Milne TA, Hughes CM, Lloyd R, Yang Z, Rozenblatt-Rosen O, Dou Y, Schnepf RW, Krankel C, Livolsi VA, Gibbs D, *et al.* 2005 Menin and MLL cooperatively regulate expression of cyclin-dependent kinase inhibitors. *PNAS* **102** 749–754. (<https://doi.org/10.1073/pnas.0408836102>)
- Molatore S, Marinoni I, Lee M, Pulz E, Ambrosio MR, degli Uberti EC, Zatelli MC & Pellegata NS 2010 A novel germline CDKN1B mutation causing multiple endocrine tumors: clinical, genetic and functional characterization. *Human Mutation* **31** E1825–E1835. (<https://doi.org/10.1002/humu.21354>)
- Moran A, Asa SL, Kovacs K, Horvath E, Singer W, Sagman U, Reubi JC, Wilson CB, Larson R & Pescovitz OH 1990 Gigantism due to pituitary mammosomatotroph hyperplasia. *New England Journal of Medicine* **323** 322–327. (<https://doi.org/10.1056/NEJM199008023230507>)
- Morgan RM, Hernandez-Ramirez LC, Trivellin G, Zhou L, Roe SM, Korbonits M & Prodromou C 2012 Structure of the TPR domain of AIP: lack of client protein interaction with the C-terminal alpha-7 helix of the TPR domain of AIP is sufficient for pituitary adenoma predisposition. *PLoS ONE* **7** e53339. (<https://doi.org/10.1371/journal.pone.0053339>)
- Murakami M, Meneses PI, Lan K & Robertson ES 2008 The suppressor of metastasis Nm23-H1 interacts with the Cdc42 Rho family member and the pleckstrin homology domain of oncoprotein Dbl-1 to suppress cell migration. *Cancer Biology and Therapy* **7** 677–688. (<https://doi.org/10.4161/cbt.7.5.5665>)
- Nakayama K, Ishida N, Shirane M, Inomata A, Inoue T, Shishido N, Horii I, Loh DY & Nakayama K 1996 Mice lacking p27(Kip1) display increased body size, multiple organ hyperplasia, retinal dysplasia, and pituitary tumors. *Cell* **85** 707–720. ([https://doi.org/10.1016/S0092-8674\(00\)81237-4](https://doi.org/10.1016/S0092-8674(00)81237-4))
- Narumi S, Matsuo K, Ishii T, Tanahashi Y & Hasegawa T 2013 Quantitative and sensitive detection of GNAS mutations causing McCune-Albright syndrome with next generation sequencing. *PLoS ONE* **8** e60525. (<https://doi.org/10.1371/journal.pone.0060525>)
- Naves LA, Daly AF, Vanbellinghen JF, Casulari LA, Spilioti C, Magalhaes AV, Azevedo ME, Giacomini LA, Nascimento PP, Nunes RO, *et al.* 2007 Variable pathological and clinical features of a large Brazilian family harboring a mutation in the aryl hydrocarbon receptor-interacting protein gene. *European Journal of Endocrinology* **157** 383–391. (<https://doi.org/10.1530/EJE-07-0533>)
- Niecknig H, Tug S, Reyes BD, Kirsch M, Fandrey J & Berchner-Pfannschmidt U 2012 Role of reactive oxygen species in the regulation of HIF-1 by prolyl hydroxylase 2 under mild hypoxia. *Free Radical Research* **46** 705–717. (<https://doi.org/10.3109/10715762.2012.669041>)
- Niemann S & Muller U 2000 Mutations in SDHC cause autosomal dominant paraganglioma, type 3. *Nature Genetics* **26** 268–270. (<https://doi.org/10.1038/81551>)
- Nilaweera KN, Ozanne D, Wilson D, Mercer JG, Morgan PJ & Barrett P 2007 G protein-coupled receptor 101 mRNA expression in the mouse brain: altered expression in the posterior hypothalamus and amygdala by energetic challenges. *Journal of Neuroendocrinology* **19** 34–45. (<https://doi.org/10.1111/j.1365-2826.2006.01502.x>)
- Nishioka H, Haraoka J & Akada K 2003 Fibrous bodies are associated with lower GH production and decreased expression of E-cadherin in GH-producing pituitary adenomas. *Clinical Endocrinology* **59** 768–772. (<https://doi.org/10.1046/j.1365-2265.2003.01921.x>)
- Occhi G, Regazzo D, Trivellin G, Boaretto F, Ciato D, Bobisse S, Ferasin S, Cetani F, Pardi E, Korbonits M, *et al.* 2013 A novel mutation in the upstream open reading frame of the CDKN1B gene causes a MEN4 phenotype. *PLoS Genetics* **9** e1003350. (<https://doi.org/10.1371/journal.pgen.1003350>)
- Oriola J, Lucas T, Halperin I, Mora M, Perales MJ, Alvarez-Escola C, Paz de MN, Diaz Soto G, Salinas I, Julian MT, *et al.* 2013 Germline mutations of AIP gene in somatotropinomas resistant to somatostatin analogues. *European Journal of Endocrinology* **168** 9–13. (<https://doi.org/10.1530/EJE-12-0457>)
- Ozkaya HM, Comunoglu N, Sayitoglu M, Keskin FE, Firtina S, Khodzhaev K, Apaydin T, Gazioglu N, Tanriover N, Oz B, *et al.* 2018 Germline mutations of aryl hydrocarbon receptor-interacting protein (AIP) gene and somatostatin receptor 1–5 and AIP immunostaining in patients with sporadic acromegaly with poor versus good response to somatostatin analogues. *Pituitary* **21** 335–346. (<https://doi.org/10.1007/s11102-018-0876-4>)
- Pan LX, Chen ZP, Liu YS & Zhao JH 2005 Magnetic resonance imaging and biological markers in pituitary adenomas with invasion of the cavernous sinus space. *Journal of Neurooncology* **74** 71–76. (<https://doi.org/10.1007/s11060-004-6150-9>)
- Papathomas TG, Gaal J, Corssmit EP, Oudijk L, Korpershoek E, Heimdal K, Bayley JP, Morreau H, van Dooren M, Papaspyrou K, *et al.* 2014 Non-pheochromocytoma (PCC)/paraganglioma (PGL) tumors in patients with succinate dehydrogenase-related PCC-PGL syndromes: a clinicopathological and molecular analysis. *European Journal of Endocrinology* **170** 1–12. (<https://doi.org/10.1530/EJE-13-0623>)
- Pardi E, Mariotti S, Pellegata NS, Benfani K, Borsari S, Saponaro F, Torregrossa L, Cappai A, Satta C, Mastinu M, *et al.* 2015 Functional characterization of a CDKN1B mutation in a Sardinian kindred with multiple endocrine neoplasia type 4 (MEN4). *Endocrine Connections* **4** 1–8. (<https://doi.org/10.1530/EC-14-0116>)
- Pellegata NS, Quintanilla-Martinez L, Siggelkow H, Samson E, Bink K, Hofler H, Fend F, Graw J & Atkinson MJ 2006 Germ-line mutations in p27Kip1 cause a multiple endocrine neoplasia syndrome in rats and humans. *PNAS* **103** 15558–15563. (<https://doi.org/10.1073/pnas.0603877103>)
- Peverelli E, Mantovani G, Lania AG & Spada A 2014 cAMP in the pituitary: an old messenger for multiple signals. *Journal of Molecular Endocrinology* **52** R67–R77. (<https://doi.org/10.1530/JME-13-0172>)
- Pfarr CM, Mechta F, Spyrou G, Lallemand D, Carillo S & Yaniv M 1994 Mouse JunD negatively regulates fibroblast growth and antagonizes transformation by ras. *Cell* **76** 747–760. ([https://doi.org/10.1016/0092-8674\(94\)90513-4](https://doi.org/10.1016/0092-8674(94)90513-4))
- Qian ZR, Asa SL, Siomi H, Siomi MC, Yoshimoto K, Yamada S, Wang EL, Rahman MM, Inoue H, Itakura M, *et al.* 2009 Overexpression of HMGA2 relates to reduction of the let-7 and its relationship to clinicopathological features in pituitary adenomas. *Modern Pathology* **22** 431–441. (<https://doi.org/10.1038/modpathol.2008.202>)
- Raappana A, Koivukangas J, Ebeling T & Pirila T 2010 Incidence of pituitary adenomas in Northern Finland in 1992–2007. *Journal of Clinical Endocrinology and Metabolism* **95** 4268–4275. (<https://doi.org/10.1210/jc.2010-0537>)
- Raitila A, Lehtonen HJ, Arola J, Heliovaara E, Ahlsten M, Georgitsi M, Jalanko A, Paetau A, Aaltonen LA & Karhu A 2010 Mice with inactivation of aryl hydrocarbon receptor-interacting protein (Aip) display complete penetrance of pituitary adenomas with aberrant ARNT expression. *American Journal of Pathology* **177** 1969–1976. (<https://doi.org/10.2353/ajpath.2010.100138>)

- Reincke M, Sbiera S, Hayakawa A, Theodoropoulou M, Osswald A, Beuschlein F, Meitinger T, Mizuno-Yamasaki E, Kawaguchi K, Saeki Y, *et al.* 2015 Mutations in the deubiquitinase gene USP8 cause Cushing's disease. *Nature Genetics* **47** 31–38. (<https://doi.org/10.1038/ng.3166>)
- Ritvonen E, Pitkanen E, Karppinen A, Vehkavaara S, Demir H, Paetau A, Schalin-Jantti C & Karhu A 2017 Impact of AIP and inhibitory G protein alpha 2 proteins on clinical features of sporadic GH-secreting pituitary adenomas. *European Journal of Endocrinology* **176** 243–252. (<https://doi.org/10.1530/EJE-16-0620>)
- Rodd C, Millette M, Iacovazzo D, Stiles CE, Barry S, Evanson J, Albrecht S, Caswell R, Bunce B, Jose S, *et al.* 2016 Somatic GPR101 duplication causing X-linked acrogigantism (XLAG)-diagnosis and management. *Journal of Clinical Endocrinology and Metabolism* **101** 1927–1930. (<https://doi.org/10.1210/jc.2015-4366>)
- Ronchi CL, Peverelli E, Herterich S, Weigand I, Mantovani G, Schwarzmayer T, Sbiera S, Allolio B, Honegger J, Appenzeller S, *et al.* 2016 Landscape of somatic mutations in sporadic GH-secreting pituitary adenomas. *European Journal of Endocrinology* **174** 363–372. (<https://doi.org/10.1530/EJE-15-1064>)
- Rostomyan L, Daly AF, Petrossians P, Nachev E, Lila AR, Lecoq AL, Lecomberri B, Trivellin G, Salvatori R, Moraitis AG, *et al.* 2015 Clinical and genetic characterization of pituitary gigantism: an international collaborative study in 208 patients. *Endocrine-Related Cancer* **22** 745–757. (<https://doi.org/10.1530/ERC-15-0320>)
- Roszkó KL, Blouch E, Blake M, Powers JF, Tischler AS, Hodin R, Sadow P & Lawson EA 2017 Case report of a prolactinoma in a patient with a novel MAX mutation and bilateral pheochromocytomas. *Journal of the Endocrine Society* **1** 1401–1407. (<https://doi.org/10.1210/js.2017-00135>)
- Rothenbuhler A & Stratakis CA 2010 Clinical and molecular genetics of Carney complex. *Best Practice and Research Clinical Endocrinology and Metabolism* **24** 389–399. (<https://doi.org/10.1016/j.beem.2010.03.003>)
- Roussel-Gervais A, Couture C, Langlais D, Takayasu S, Balsalobre A, Rueda BR, Zukerberg LR, Figarella-Branger D, Brue T & Drouin J 2016 The Cables1 gene in glucocorticoid regulation of pituitary corticotrope growth and cushing disease. *Journal of Clinical Endocrinology and Metabolism* **101** 513–522. (<https://doi.org/10.1210/jc.2015-3324>)
- Salenave S, Boyce AM, Collins MT & Chanson P 2014 Acromegaly and McCune-Albright syndrome. *Journal of Clinical Endocrinology and Metabolism* **99** 1955–1969. (<https://doi.org/10.1210/jc.2013-3826>)
- Salpea P, Horvath A, London E, Faucz FR, Vetro A, Levy I, Gourgari E, Dauber A, Holm IA, Morrison PJ, *et al.* 2014 Deletions of the PRKAR1A locus at 17q24.2–q24.3 in Carney complex: genotype-phenotype correlations and implications for genetic testing. *Journal of Clinical Endocrinology and Metabolism* **99** E183–E188. (<https://doi.org/10.1210/jc.2013-3159>)
- Salvatori R, Radian S, Diekmann Y, Iacovazzo D, David A, Gabrovská P, Grassi G, Bussell AM, Stals K, Weber A, *et al.* 2017 In-frame seven amino-acid duplication in AIP arose over the last 3000 years, disrupts protein interaction and stability and is associated with gigantism. *European Journal of Endocrinology* **177** 257–266. (<https://doi.org/10.1530/EJE-17-0293>)
- Sambugaro S, Di Ruvo M, Ambrosio MR, Pellegata NS, Bellio M, Guerra A, Buratto M, Foschini MP, Tagliati F, degli Uberti E, *et al.* 2015 Early onset acromegaly associated with a novel deletion in CDKN1B 5'UTR region. *Endocrine* **49** 58–64. (<https://doi.org/10.1007/s12020-015-0540-y>)
- Sano T, Rong QZ, Kagawa N & Yamada S 2004 Down-regulation of E-cadherin and catenins in human pituitary growth hormone-producing adenomas. *Frontiers of Hormone Research* **32** 127–132.
- Scheithauer BW, Kovacs K, Horvath E, Kim DS, Osamura RY, Ketterling RP, Lloyd RV & Kim OL 2008 Pituitary blastoma. *Acta Neuropathologica* **116** 657–666. (<https://doi.org/10.1007/s00401-008-0388-9>)
- Scherthaner-Reiter MH, Trivellin G & Stratakis CA 2018 Interaction of AIP with protein kinase A (cAMP-dependent protein kinase). *Human Molecular Genetics* **27** 2604–2613. (<https://doi.org/10.1093/hmg/ddy166>)
- Schultz KAP, Williams GM, Kamihara J, Stewart DR, Harris AK, Bauer AJ, Turner J, Shah R, Schneider K, Schneider KW, *et al.* 2018 DICER1 and associated conditions: identification of at-risk individuals and recommended surveillance strategies. *Clinical Cancer Research* **24** 2251–2261. (<https://doi.org/10.1158/1078-0432.CCR-17-3089>)
- Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, Pan Y, Simon MC, Thompson CB & Gottlieb E 2005 Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF- α prolyl hydroxylase. *Cancer Cell* **7** 77–85. (<https://doi.org/10.1016/j.ccr.2004.11.022>)
- Sheaff RJ, Groudine M, Gordon M, Roberts JM & Clurman BE 1997 Cyclin E-CDK2 is a regulator of p27Kip1. *Genes and Development* **11** 1464–1478. (<https://doi.org/10.1101/gad.11.11.1464>)
- Shi Z, Li Z, Li ZJ, Cheng K, Du Y, Fu H & Khuri FR 2015 Cables1 controls p21/Cip1 protein stability by antagonizing proteasome subunit α type 3. *Oncogene* **34** 2538–2545. (<https://doi.org/10.1038/onc.2014.171>)
- Skalhegg BS & Tasken K 2000 Specificity in the cAMP/PKA signaling pathway. Differential expression, regulation, and subcellular localization of subunits of PKA. *Frontiers in Bioscience* **5** D678–D693.
- Slane BG, Aykin-Burns N, Smith BJ, Kalen AL, Goswami PC, Domann FE & Spitz DR 2006 Mutation of succinate dehydrogenase subunit C results in increased O₂•, oxidative stress, and genomic instability. *Cancer Research* **66** 7615–7620. (<https://doi.org/10.1158/0008-5472.CAN-06-0833>)
- Stergiopoulos SG, Abu-Asab MS, Tsokos M & Stratakis CA 2004 Pituitary pathology in Carney complex patients. *Pituitary* **7** 73–82. (<https://doi.org/10.1007/s11102-005-5348-y>)
- Stiles CE & Korbonits M 2011 Familial isolated pituitary adenoma. In *Endotext*. Eds LJ De Groot, G Chrousos, K Dungan, KR Feingold, A Grossman, JM Hershman, C Koch, M Korbonits, R McLachlan, M New, *et al.* South Dartmouth (MA): MDText.com, Inc.
- Stratakis CA 2016 Carney complex: a familial lentiginosis predisposing to a variety of tumors. *Reviews in Endocrine and Metabolic Disorders* **17** 367–371. (<https://doi.org/10.1007/s11154-016-9400-1>)
- Stratakis CA, Salpea P & Raygada M 1993 Carney complex. In *GeneReviews*. Eds MP Adam, HH Ardinger, RA Pagon, SE Wallace, LJH Bean, K Stephens & A Amemiya. Seattle, WA, USA: University of Washington. (available at: <https://www.ncbi.nlm.nih.gov/books/NBK1286/>)
- Stratakis CA, Matyakhina L, Courkoutsakis N, Patronas N, Voutetakis A, Stergiopoulos S, Bossis I & Carney JA 2004 Pathology and molecular genetics of the pituitary gland in patients with the 'complex of spotty skin pigmentation, myxomas, endocrine overactivity and schwannomas' (Carney complex). *Frontiers of Hormone Research* **32** 253–264.
- Thakker RV 2014 Multiple endocrine neoplasia type 1 (MEN1) and type 4 (MEN4). *Molecular and Cellular Endocrinology* **386** 2–15. (<https://doi.org/10.1016/j.mce.2013.08.002>)
- Thakker RV, Newey PJ, Walls GV, Bilezikian J, Dralle H, Ebeling PR, Melmed S, Sakurai A, Tonelli F, Brandi ML, *et al.* 2012 Clinical practice guidelines for multiple endocrine neoplasia type 1 (MEN1). *Journal of Clinical Endocrinology and Metabolism* **97** 2990–3011. (<https://doi.org/10.1210/jc.2012-1230>)
- Theodoropoulou M & Stalla GK 2013 Somatostatin receptors: from signaling to clinical practice. *Frontiers in Neuroendocrinology* **34** 228–252. (<https://doi.org/10.1016/j.ynrne.2013.07.005>)
- Theodoropoulou M, Tichomirowa MA, Sievers C, Yassouridis A, Arzberger T, Hougrand O, Deprez M, Daly AF, Petrossians P, Pagotto U, *et al.* 2009 Tumor ZAC1 expression is associated with the response

- to somatostatin analog therapy in patients with acromegaly. *International Journal of Cancer* **125** 2122–2126. (<https://doi.org/10.1002/ijc.24602>)
- Thevenon J, Bourredjem A, Faivre L, Cardot-Bauters C, Calender A, Murat A, Giraud S, Niccoli P, Odou MF, Borson-Chazot F, *et al.* 2013 Higher risk of death among MEN1 patients with mutations in the JunD interacting domain: a Groupe d'étude des Tumeurs Endocrines (GTE) cohort study. *Human Molecular Genetics* **22** 1940–1948. (<https://doi.org/10.1093/hmg/ddt039>)
- Tichomirowa MA, Barlier A, Daly AF, Jaffrain-Rea ML, Ronchi C, Yaneva M, Urban JD, Petrossians P, Elenkova A, Tabarin A, *et al.* 2011 High prevalence of AIP gene mutations following focused screening in young patients with sporadic pituitary macroadenomas. *European Journal of Endocrinology* **165** 509–515. (<https://doi.org/10.1530/EJE-11-0304>)
- Tichomirowa MA, Lee M, Barlier A, Daly AF, Marinoni I, Jaffrain-Rea ML, Naves LA, Rodien P, Rohmer V, Faucz FR, *et al.* 2012 Cyclin-dependent kinase inhibitor 1B (CDKN1B) gene variants in AIP mutation-negative familial isolated pituitary adenoma kindreds. *Endocrine-Related Cancer* **19** 233–241. (<https://doi.org/10.1530/ERC-11-0362>)
- Tonelli F, Giudici F, Giusti F, Marini F, Cianferotti L, Nesi G & Brandi ML 2014 A heterozygous frameshift mutation in exon 1 of CDKN1B gene in a patient affected by MEN4 syndrome. *European Journal of Endocrinology* **171** K7–K17. (<https://doi.org/10.1530/EJE-14-0080>)
- Tortora G, Pepe S, Bianco C, Baldassarre G, Budillon A, Clair T, Cho-Chung YS, Bianco AR & Ciardiello F 1994a The RI alpha subunit of protein kinase A controls serum dependency and entry into cell cycle of human mammary epithelial cells. *Oncogene* **9** 3233–3240.
- Tortora G, Pepe S, Bianco C, Damiano V, Ruggiero A, Baldassarre G, Corbo C, Cho-Chung YS, Bianco AR & Ciardiello F 1994b Differential effects of protein kinase A sub-units on Chinese-hamster-ovary cell cycle and proliferation. *International Journal of Cancer* **59** 712–716. (<https://doi.org/10.1002/ijc.2910590521>)
- Trivellin G & Korbonits M 2011 AIP and its interacting partners. *Journal of Endocrinology* **210** 137–155. (<https://doi.org/10.1530/JOE-11-0054>)
- Trivellin G, Daly AF, Faucz FR, Yuan B, Rostomyan L, Larco DO, Scherthaner-Reiter MH, Szarek E, Leal LF, Caberg JH, *et al.* 2014 Gigantism and acromegaly due to Xq26 microduplications and GPR101 mutation. *New England Journal of Medicine* **371** 2363–2374. (<https://doi.org/10.1056/NEJMoa1408028>)
- Trivellin G, Bjelobaba I, Daly AF, Larco DO, Palmeira L, Faucz FR, Thiry A, Leal LF, Rostomyan L, Quezado M, *et al.* 2016a Characterization of GPR101 transcript structure and expression patterns. *Journal of Molecular Endocrinology* **57** 97–111. (<https://doi.org/10.1530/JME-16-0045>)
- Trivellin G, Correa RR, Batsis M, Faucz FR, Chittiboina P, Bjelobaba I, Larco DO, Quezado M, Daly AF, Stojilkovic SS, *et al.* 2016b Screening for GPR101 defects in pediatric pituitary corticotropinomas. *Endocrine-Related Cancer* **23** 357–365. (<https://doi.org/10.1530/ERC-16-0091>)
- Trouillas J, Labat-Moleur F, Sturm N, Kujas M, Heymann MF, Figarella-Branger D, Patey M, Mazucca M, Decullier E, Verges B, *et al.* 2008 Pituitary tumors and hyperplasia in multiple endocrine neoplasia type 1 syndrome (MEN1): a case-control study in a series of 77 patients versus 2509 non-MEN1 patients. *American Journal of Surgical Pathology* **32** 534–543. (<https://doi.org/10.1097/PAS.0b013e31815ade45>)
- Tufton N, Roncaroli F, Hadjijdemetriou I, Dang MN, Denes J, Guasti L, Thom M, Powell M, Baldeweg SE, Fersht N, *et al.* 2017 Pituitary carcinoma in a patient with an SDHB mutation. *Endocrine Pathology* **28** 320–325. (<https://doi.org/10.1007/s12022-017-9474-7>)
- Tuominen I, Heliovaara E, Raitila A, Rautiainen MR, Mehine M, Katainen R, Donner I, Aittomäki V, Lehtonen HJ, Ahlsten M, *et al.* 2015 AIP inactivation leads to pituitary tumorigenesis through defective Galphai-cAMP signaling. *Oncogene* **34** 1174–1184. (<https://doi.org/10.1038/ncr.2014.50>)
- Valimäki N, Demir H, Pitkanen E, Kaasinen E, Karppinen A, Kivipeltö L, Schalin-Jantti C, Aaltonen LA & Karhu A 2015 Whole-genome sequencing of growth hormone (GH)-secreting pituitary adenomas. *Journal of Clinical Endocrinology and Metabolism* **100** 3918–3927. (<https://doi.org/10.1210/jc.2015-3129>)
- Varsavsky M, Sebastian-Ochoa A & Torres Vela E 2013 Coexistence of a pituitary macroadenoma and multicentric paraganglioma: a strange coincidence. *Endocrinología y Nutrición* **60** 154–156. (<https://doi.org/10.1016/j.endonu.2012.02.009>)
- Vasilev V, Daly AF, Petrossians P, Zacharieva S & Beckers A 2011 Familial pituitary tumor syndromes. *Endocrine Practice* **17** (Supplement 3) 41–46. (<https://doi.org/10.4158/EP11064.RA>)
- Verges B, Boureille F, Goudet P, Murat A, Beckers A, Sassolas G, Cougand P, Chambe B, Montvernay C & Calender A 2002 Pituitary disease in MEN type 1 (MEN1): data from the France-Belgium MEN1 multicenter study. *Journal of Clinical Endocrinology and Metabolism* **87** 457–465. (<https://doi.org/10.1210/jcem.87.2.8145>)
- Vierimäa O, Georgitsi M, Lehtonen R, Vahteristo P, Kokko A, Raitila A, Tuppurainen K, Ebeling TM, Salmela PI, Paschke R, *et al.* 2006 Pituitary adenoma predisposition caused by germline mutations in the AIP gene. *Science* **312** 1228–1230. (<https://doi.org/10.1126/science.1126100>)
- Villa C, Lagonigro MS, Magri F, Koziak M, Jaffrain-Rea ML, Brauner R, Bouligand J, Junier MP, Di Rocco F, Sainte-Rose C, *et al.* 2011 Hyperplasia-adenoma sequence in pituitary tumorigenesis related to aryl hydrocarbon receptor interacting protein gene mutation. *Endocrine-Related Cancer* **18** 347–356. (<https://doi.org/10.1530/ERC-11-0059>)
- Wang Y, Chen J, Yang W, Mo F, Senz J, Yap D, Anglesio MS, Gilks B, Morin GB & Huntsman DG 2015 The oncogenic roles of DICER1 RNase IIIb domain mutations in ovarian Sertoli-Leydig cell tumors. *Neoplasia* **17** 650–660. (<https://doi.org/10.1016/j.neo.2015.08.003>)
- Weitzman JB, Fiette L, Matsuo K & Yaniv M 2000 JunD protects cells from p53-dependent senescence and apoptosis. *Molecular Cell* **6** 1109–1119. ([https://doi.org/10.1016/S1097-2765\(00\)00109-X](https://doi.org/10.1016/S1097-2765(00)00109-X))
- Williams F, Hunter S, Bradley L, Chahal HS, Storr HL, Akker SA, Kumar AV, Orme SM, Evanson J, Abid N, *et al.* 2014 Clinical experience in the screening and management of a large kindred with familial isolated pituitary adenoma due to an aryl hydrocarbon receptor interacting protein (AIP) mutation. *Journal of Clinical Endocrinology and Metabolism* **99** 1122–1131. (<https://doi.org/10.1210/jc.2013-2868>)
- Wu T & Hua X 2011 Menin represses tumorigenesis via repressing cell proliferation. *American Journal of Cancer Research* **1** 726–739.
- Xekouki P, Pacak K, Almeida M, Wassif CA, Rustin P, Nesterova M, de la Luz Sierra M, Matro J, Ball E, Azevedo M, *et al.* 2012 Succinate dehydrogenase (SDH) D subunit (SDHD) inactivation in a growth-hormone-producing pituitary tumor: a new association for SDH? *Journal of Clinical Endocrinology and Metabolism* **97** E357–E366. (<https://doi.org/10.1210/jc.2011-1179>)
- Xekouki P, Mastroyiannis SA, Avgeropoulos D, de la Luz Sierra M, Trivellin G, Gourgari EA, Lyssikatos C, Quezado M, Patronas N, Kanaka-Gantenbein C, *et al.* 2013 Familial pituitary apoplexy as the only presentation of a novel AIP mutation. *Endocrine-Related Cancer* **20** L11–L14. (<https://doi.org/10.1530/ERC-13-0218>)
- Xekouki P, Szarek E, Bullova P, Giubellino A, Quezado M, Mastroyiannis SA, Mastorakos P, Wassif CA, Raygada M, Rentia N, *et al.* 2015 Pituitary adenoma with paraganglioma/pheochromocytoma (3PAs) and succinate dehydrogenase defects in humans and mice. *Journal of Clinical Endocrinology and Metabolism* **100** E710–E719. (<https://doi.org/10.1210/jc.2014-4297>)
- Yin Z, Williams-Simons L, Parlow AF, Asa S & Kirschner LS 2008 Pituitary-specific knockout of the Carney complex gene *Pkrar1a* leads to

- pituitary tumorigenesis. *Molecular Endocrinology* **22** 380–387. (<https://doi.org/10.1210/me.2006-0428>)
- Yin Y, Castro AM, Hoekstra M, Yan TJ, Kanakamedala AC, Dehner LP, Hill DA & Ornitz DM 2015 Fibroblast growth factor 9 regulation by microRNAs controls lung development and links DICER1 loss to the pathogenesis of pleuropulmonary blastoma. *PLoS Genetics* **11** e1005242. (<https://doi.org/10.1371/journal.pgen.1005242>)
- Yokoyama A & Cleary ML 2008 Menin critically links MLL proteins with LEDGF on cancer-associated target genes. *Cancer Cell* **14** 36–46. (<https://doi.org/10.1016/j.ccr.2008.05.003>)
- Yoshida D & Teramoto A 2007 Elevated cell invasion is induced by hypoxia in a human pituitary adenoma cell line. *Cell Adhesion and Migration* **1** 43–51. (<https://doi.org/10.4161/cam.4080>)
- Yoshida D, Kim K, Noha M & Teramoto A 2006 Anti-apoptotic action by hypoxia inducible factor 1-alpha in human pituitary adenoma cell line, HP-75 in hypoxic condition. *Journal of Neurooncology* **78** 217–225. (<https://doi.org/10.1007/s11060-005-9017-9>)
- Zhang Q, Peng C, Song J, Zhang Y, Chen J, Song Z, Shou X, Ma Z, Peng H, Jian X, *et al.* 2017 Germline mutations in CDH23, encoding cadherin-related 23, are associated with both familial and sporadic pituitary adenomas. *American Journal of Human Genetics* **100** 817–823. (<https://doi.org/10.1016/j.ajhg.2017.03.011>)
- Zhou Y, Zhang X & Klibanski A 2014 Genetic and epigenetic mutations of tumor suppressive genes in sporadic pituitary adenoma. *Molecular and Cellular Endocrinology* **386** 16–33. (<https://doi.org/10.1016/j.mce.2013.09.006>)

Received in final form 31 October 2018

Accepted 7 November 2018